

## PURIFICATION AND CHARACTERIZATION OF CYTOCHROME $b_5$ FROM THE HOUSEFLY (*MUSCA DOMESTICA*)

GEOFFREY D. WHELOCK and JEFFREY G. SCOTT\*

Department of Entomology and Institute for Comparative and Environmental Toxicology, Comstock Hall, Cornell University, Ithaca, NY 14853-0999, USA (Tel: 607 255-7340)

(Received 8 June 1991)

**Abstract**—1. Isopropanol treatment of *Musca domestica* (house fly) microsomes extracted cytochrome  $b_5$  and resulted in a minimum 5.2-fold purification.

2. Cytochrome  $b_5$  was highly purified from the isopropanol extract using ion exchange and reverse phase high performance liquid chromatography.

3. A polyclonal antiserum was produced against the purified cytochrome  $b_5$ .

4. Purified house fly cytochrome  $b_5$  had a molecular mass of 19.7 kDa, slightly higher than those reported for rabbit and rat microsomal cytochrome  $b_5$  (16.6–16.7 kDa), but lower than cytochrome  $b_5$  from Mediterranean fruit fly (21 kDa).

5. Reduced house fly cytochrome  $b_5$  absorbed maximally at 424 nm, similar to other cytochromes  $b_5$ .

6. Amino acid analysis of house fly cytochrome  $b_5$  produces a higher percentage of hydrophobic amino acid residues and less acidic residues than other cytochromes  $b_5$ .

7. In immuno-blot analysis, house fly cytochrome  $b_5$  shared antigenic epitope(s) with the dipteran species *Musca autumnalis* (face fly), *Stomoxys calcitrans* (stable fly) and to a lesser extent, *Drosophila melanogaster* (fruit fly).

8. Anti-house fly cytochrome  $b_5$  did not recognize cytochrome  $b_5$  from *Apis mellifera* (honey bee), *Tricoplusia ni* (cabbage looper), *Tetranychus urticae* (two-spotted spider mite) or *Rattus rattus* (rat).

### INTRODUCTION

In the endoplasmic reticulum, cytochrome  $b_5$  transfers reducing equivalents from cytochrome  $b_5$  reductase or cytochrome P450 reductase to stearyl-CoA desaturase or to cytochrome P450, and thus is thought to participate in many important cellular oxidation reactions (Estabrook, 1978). However, its role as a donor of the second electron to cytochrome P450 remains unresolved (Hildebrandt and Estabrook, 1971).

The ability to isolate and study intact cytochrome  $b_5$  from insects is desirable since it is abundant in insects and may be involved with monooxygenase-dependent insecticide resistance (Scott *et al.*, 1990). House flies are an exceptionally good model to study the role of insect cytochrome  $b_5$  in monooxygenase reactions since other components of the system are well studied in house flies (Ronis and Hodgson, 1989).

Intact cytochrome  $b_5$  was first isolated from mammalian microsomes by Spatz and Strittmatter (1971), and several similar procedures for isolation were subsequently developed for mammalian tissues (Ozols, 1974; Carlsen *et al.*, 1988; Burchell, 1985). However, alternative procedures are required for house fly cytochrome  $b_5$  (Wheelock and Scott, 1989). Proteolytically solubilized cytochrome  $b_5$  was purified from armyworm (McFadden *et al.*, 1979). Intact (i.e. solubilized with detergent) cytochrome  $b_5$  has been purified to apparent homogeneity from only two

insects, the Mediterranean fruit fly *Ceratitis capitata* (Megias *et al.*, 1986) and the house fly. In the case of the house fly, cytochrome  $b_5$  was isolated in small yield as a by-product of cytochrome P450 isolation (Wheelock and Scott, 1989). The poor yield of cytochrome  $b_5$  was primarily due to the initial extraction procedure that was designed to optimize isolation of cytochrome P450. The present study resulted from experiments to optimize purification of cytochrome  $b_5$  from house fly microsomes in yields sufficient for antiserum production and characterization.

### MATERIALS AND METHODS

#### *Rearing and microsome preparation*

House flies were reared as previously described (Wheelock and Scott, 1989). Face flies, stable flies, fruit flies, honey bees, cabbage loopers and mites were obtained and reared as described by Wheelock *et al.* (1991). Male Wistar rats were obtained from Cornell research animal resources, Cornell University, Ithaca, NY.

Microsomes were prepared from house fly abdomens as previously described (Wheelock and Scott, 1990). Microsomes from other insects and mites were prepared as described by Wheelock *et al.* (1991). Rats were anesthetized and killed with carbon dioxide and microsomes prepared from the rat livers according to Wheelock and Scott (1990). All microsomes were resuspended in 5 ml phosphate buffer plus 20% glycerol (Wheelock and Scott, 1989) and stored at  $-80^{\circ}\text{C}$ .

#### *Assay of cytochrome $b_5$*

Cytochrome  $b_5$  was assayed according to Omura and Sato (1964) using sodium dithionite as reducing agent according to Capdevila and Agosin (1975). Cytochrome  $b_5$  was

\*Author to whom correspondence should be addressed.




Fig. 1. Denaturing gel electrophoresis. Lane 1, 6: molecular mass markers (top to bottom 99.7, 66, 45, 31, 21.5, 14.4 kDa). Lane 2: 50 pmol crude microsomal cytochrome  $b_5$ . Lane 3: 50 pmol cytochrome  $b_5$ , isopropanol extract. Lane 4: 100 pmol cytochrome  $b_5$ , ion exchange HPLC fraction. Lane 5: 100 pmol cytochrome  $b_5$ , reverse phase HPLC fraction.

always assayed in the presence of detergent. Crude microsomes were solubilized in 0.5% CHAPS (3-[(3-cholamidopropyl)-dimethylammonio]-1-propanosulfonate, Sigma, St Louis, MO) or 0.4% emulgen (emulgen 911, Kao Atlas, Tokyo) prior to assay. Chromatography fractions contained 0.4% emulgen.

#### Extraction and isolation of cytochrome $b_5$

Five milliliters of LPR strain house fly microsomes were diluted with an equal volume of isopropanol (HPLC grade, Fisher Scientific, Rochester, NY) and immediately mixed. The cytochrome  $b_5$  was extracted by gentle rocking at 4°C for 10 min, centrifuged at 8000  $g$  for 10 min and the supernatant saved. The pellet was reextracted with 2 ml 50% isopropanol, 10% glycerol and 40% phosphate buffer and centrifuged again. The supernatants were pooled and diluted with an equal volume of  $1.8 \times 10^7$  ohm deionized water, frozen to  $-80^\circ\text{C}$  and lyophilized overnight.

The lyophilized extract was diluted to 4.2 ml with 0.4% emulgen in water, filtered through a  $0.45 \mu\text{m}$  HPLC filter, divided in half, and chromatographed on a DEAE 5PW ion exchange high performance liquid chromatography (HPLC) column (Altex, San Ramon, CA) in two runs using conditions described previously (Wheelock and Scott, 1989). Cytochrome  $b_5$  from the DEAE 5PW HPLC step was injected onto a VYDAC C8 reverse phase HPLC column (Alltech, Deerfield, IL) in two runs and developed using conditions described previously (Wheelock and Scott, 1989).

#### Antisera production

Polyclonal anti-house fly cytochrome  $b_5$  was produced in a Flemish Giant/Chinchilla cross rabbit (Cornell research animal resources). For each immunization, purified cytochrome  $b_5$  from the reverse phase HPLC step (see above) was diluted 25- or 50-fold with phosphate buffer containing 20% glycerol and 0.4% emulgen and concentrated on a YM2 filter (Amicon, Danvers, MA) to approximately 0.5 ml. The purpose of this step was to dilute out the

isopropanol in the HPLC fraction prior to immunization. Concentrated antigen was diluted with an equal volume of Freund's adjuvant and injected subcutaneously at multiple sites along the rabbit's back. Freund's complete adjuvant was used for initial immunization and Freund's incomplete adjuvant was used for subsequent injections. Immunization schedule was as follows: 3.30, 3.83, 3.83 and 3.60 nmol purified cytochrome  $b_5$  on days 1, 14, 29 and 64 respectively followed by exsanguination on day 71. The IgG fraction was partially purified from the serum by two ammonium sulfate precipitations (Kunkel, 1988).

The anti-cytochrome  $b_5$  antiserum was made monospecific (adsorbed) using the following procedure. Molecular mass markers (Biorad, Danvers, MA) and protein from 3 mg of house fly microsomes were resolved by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) using a 10% gel according to Laemmli (1970). Protein was transferred to a nitrocellulose sheet according to Towbin *et al.* (1979) with the specific conditions of Wheelock and Scott (1990). Extreme left and right ends of the sheet, containing the molecular mass markers, were removed and stained with Amido Black 10B to estimate the position of cytochrome  $b_5$ . Under these conditions house fly cytochrome  $b_5$  has an apparent molecular mass of 19.7 kDa and runs near the bottom of the gel (Wheelock and Scott, 1989). The lower portion of the sheet containing the cytochrome  $b_5$  was discarded and the remaining sheet was rinsed three times in phosphate buffered saline (Wheelock and Scott, 1990) and incubated with 0.5 ml antiserum at room temperature in a sealed plastic bag. After 4 hr the unbound, absorbed antiserum was recovered, combined with a 0.5 ml phosphate buffer rinse and stored at 4°C with 0.01% thimerosal.

#### Amino acid analysis

Purified cytochrome  $b_5$  fractions (0.37–1.9 nmol) from reverse phase HPLC described above were diluted 1:1 with deionized water and lyophilized overnight in a Speed Vac Concentrator (Savant, Farmingdale, NY). Glycerol and

Table 1. Recovery of cytochrome  $b_5$ 

Fraction	Yield nmol $b_5$	% Recovery at each step
Crude microsomes	$7.04 \pm 0.89$	100%
Isopropanol extract	$3.82 \pm 0.56$	54.4 (47–60)
DEAE 5PW HPLC	$2.60 \pm 0.30$	75.2 (65–80)
C8 HPLC	$2.33 \pm 0.14$	93.6 (88–100)

Yields are averages  $\pm$  SD from five separate purifications using 5 ml microsomes starting material each, uncorrected for samples removed at each step for assays. Recoveries are average % recovery after that step compared with the activity found at the end of the previous step corrected for removal of material for assay. Values in parentheses are ranges.

emulgen were removed from the residue by three washes of 4 ml ethyl ether. The remaining cytochrome  $b_5$  was dissolved in 100  $\mu$ l 1:2 85% formic acid: absolute ethanol and analyzed for amino acids using the Waters Pico-Tag System<sup>®</sup>. Separate analyses in the presence of thiodiglycol were done to measure tryptophan. In two out of four analyses, a large peak, tentatively considered to be unremoved emulgen, obscured CYS and/or ILE peaks. CYS and ILE values reported by the integrator were discarded from these analyses. All analyses were done by the Biotechnology Analytical and Synthetic Facility at Cornell University, Ithaca, NY.

#### Other methods

Protein in chromatography fractions was assayed using the detergent-tolerant method of Dulley and Grieve (1975). For analysis of purity and immuno-blotting, SDS-PAGE was performed according to Laemmli (1970) in 15% gels. Chromatography fractions, too ionic or too dilute for SDS-PAGE, were concentrated by precipitation in 10% ice-cold trichloroacetic acid, resuspension in lysis buffer (Laemmli, 1970) and neutralization with 1M Trizma-hydroxide (Sigma). Under these conditions emulgen in the chromatography fractions co-precipitated with cytochrome  $b_5$  and trapped excess trichloroacetic acid, which therefore required excess neutralization and the reintroduction of ions. This produced abnormally broad bands in SDS-PAGE with the chromatography fractions. Electrophoresed samples for immuno-blot analysis were transferred to nitrocellulose according to Towbin *et al.* (1979). Nitrocellulose sheets were probed with absorbed anti-cytochrome  $b_5$  antiserum at 1:500 dilution using the conditions of Wheelock and Scott (1990). Rocket immuno-electrophoresis was performed according to Wheelock and Scott (1990) with the

Table 2. Amino acid composition of house fly cytochrome  $b_5$ 

Residue	Number
ASX	14
THR	11
SER	13
GLX	17
PRO	7
GLY	11
ALA	12
VAL	16
CYS	0
MET	8
ILE	10
LEU	12
TYR	5
PHE	11
HIS	3
LYS	18
ARG	4
TRP	0

ASX + GLX: 18%  
HIS + LYS + ARG: 14.5%  
APOLAR: 42.2%

Average of four determinations. TRP: average of two determinations done separately. CYS: two determinations. ILE: three determinations. Apolar residues: MET, VAL, LEU, ILE, PRO, TRP, PHE, ALA.

following changes: agarose contained 0.5% CHAPS (Sigma) and 1.5% (unabsorbed) or 5.0% (absorbed, diluted) anti-serum.

## RESULTS

### Extraction of cytochrome $b_5$

We have previously shown that house fly cytochrome  $b_5$  is eluted in 38% isopropanol by C8 reverse phase HPLC and can be directly assayed when diluted 1:1 with phosphate buffer (Wheelock and Scott, 1989). This suggested to us that cytochrome  $b_5$  would be soluble in, and recoverable from, high concentrations of isopropanol. In the present study, we found that extraction of the microsomes in 50% isopropanol resulted in an apparent 51.3% recovery of cytochrome  $b_5$  ( $n = 10$ , range = 43–62). Additionally, the spec. act. of the isopropanol extract was  $2.29 \pm 0.068$  nmol cytochrome  $b_5$ /mg protein ( $n = 5$ ), compared with a specific content of  $0.438 \pm 0.053$  ( $n = 5$ ) for crude microsomes. This implies a 5.2-fold purification by isopropanol extraction compared with the detergent solubilized dithionite assayed crude microsomes. However, these values are most likely conservative, since assay of crude house fly microsomes with dithionite results in overestimation of cytochrome  $b_5$  due to cytochrome P450 reduction (Estabrook and Werringloer, 1978; Ozols, 1974). In crude LPR microsomes, assay of cytochrome  $b_5$  results in up to a 2-fold overestimation while partially purified cytochrome  $b_5$  fractions (which do not contain cytochrome P450) are not subject to the same artifact (unpubl. observns). Thus, isopropanol extraction of cytochrome  $b_5$  yields may approach 100% with a purification of about 10-fold. Furthermore, SDS-PAGE analysis of the extract showed significant enrichment for cytochrome  $b_5$ , suggesting that this was a suitable method for optimization of cytochrome  $b_5$  purification from house flies (Fig. 1).




Fig. 2. Amicon membrane concentrated, purified cytochrome  $b_5$  (4.15 nmol/ml). Scan represents the difference spectra between oxidized and reduced cytochrome  $b_5$ .




Fig. 3. Immunological detection of cytochrome  $b_5$ . Lane 1: 50 pmol purified cytochrome  $b_5$ . Lane 2: pellet after isopropanol extraction of microsomes, equivalent to 50 pmol cytochrome  $b_5$  in original sample. Lane 3: 50 pmol cytochrome  $b_5$ , crude microsomes.

#### Purification of cytochrome $b_5$ from isopropanol extract

Isolation of house fly cytochrome  $b_5$  as a by-product of cytochrome P450 purification consists of phenyl 5PW hydrophobic interaction HPLC, DEAE 5PW ion exchange HPLC, followed by C8 reverse phase HPLC (Wheelock and Scott, 1989). When the isopropanol extract of microsomes was used as the starting material, almost complete purification was achieved with only ion exchange HPLC and apparent homogeneity was achieved after reverse phase HPLC (Fig. 1). Thus, the original hydrophobic interaction HPLC step could be eliminated. Retention times of isopropanol-extracted cytochrome  $b_5$  on DEAE-5PW and C8 HPLC were identical with those previously reported (Wheelock and Scott, 1989) and are not reported here. Recoveries of cytochrome  $b_5$  were high throughout the isolation procedure, allowing a yield of 2.33 nmol of cytochrome  $b_5$  per run (Table 1), 2.7-fold higher than previous results (Wheelock and Scott, 1989). Purified cytochrome  $b_5$  showed an absorbance peak typical of cytochrome  $b_5$  at 424 nm (Fig. 2) as found in crude house fly microsomes (Lee and Scott, 1989). This shows the isolated material was, in fact, cytochrome  $b_5$ , and that the cytochrome  $b_5$  heme is not removed using this procedure (Burchell, 1985). Amino acid analysis of house fly cytochrome  $b_5$  showed the following percentages of amino acids: 42.2% apolar (MET, VAL, LEU, ILE, PRO, TRP, PHE, ALA), 18% acidic (taken as ASX and GLX) and 14.5% basic (HIS, LYS, ARG). There were no cysteine or tryptophan

residues detected (Table 2). *N*-Terminal amino acid analysis was unsuccessful due to *N*-terminal blockage (data not shown). The molecular mass of house fly cytochrome  $b_5$  was estimated at 19.7 kDa (Fig. 1) in confirmation of our previous determination of  $19.7 \pm 0.76$  using a different purification technique and SDS-PAGE system (Wheelock and Scott, 1989).

#### Anti-cytochrome $b_5$

The purified cytochrome  $b_5$  from the reverse phase HPLC step was used as antigen to immunize a rabbit. The resulting antiserum reacted with cytochrome  $b_5$  both as purified protein and microsomal cytochrome  $b_5$  in immuno-blotting (Fig. 3). Immuno-blot analysis of microsomes from different animals revealed that the antiserum only recognized dipteran cytochrome  $b_5$  among the types tested (Fig. 4). In order of decreasing band intensity, the antiserum recognized *Musca domestica* (LPR and S+ house fly strains), *Musca autumnalis* (face fly) and *Stomoxys calcitrans* (stable fly). In addition, the antiserum showed an extremely faint reaction with *Drosophila melanogaster* microsomes (Canton-S and Hikone-R strains). There were no reactions with microsomes from *Apis mellifera* (honey bee), *Tricoplusia ni* (cabbage looper), *Tetranychus urticae* (two-spotted spider mite) or *Rattus rattus* (rat). It should be noted that cytochrome  $b_5$  was unassayable in mite microsomes. Although similar amounts of protein were loaded onto the gel compared with other samples, it is not known whether the lack of immuno-staining in this sample was due to lack of recognition by the antiserum or lack of an adequate amount of cytochrome  $b_5$ .

The absorbed antiserum did not form an immune precipitate with house fly cytochrome  $b_5$  using rocket immuno-electrophoresis, unlike the unabsorbed antiserum (Fig. 5). Typically, rocket immuno-electrophoresis requires at least three to four epitope/antibody reactions per antigen to form the required insoluble immune-precipitate lattice. Conversely, immuno-blot analysis need only have one epitope/antibody reaction per antigen for a positive response (Kunkel, 1988). Therefore, the antigenic site(s) unique to house fly cytochrome  $b_5$  and recognized by the absorbed antiserum are probably limited in number, since the absorbed antiserum did not form an immune precipitate while the unabsorbed antiserum did.

#### DISCUSSION

We report a procedure to extract house fly cytochrome  $b_5$  in high yield that allows reduction of chromatographic steps needed to purify it from three to two, and increases the yield 2.7-fold over our previously reported procedure. We were able to isolate quantities sufficient to produce a polyclonal antibody to undamaged cytochrome  $b_5$  holoprotein and to afford some preliminary characterization. Difficulties in producing an immune response in rabbits with cytochrome  $b_5$  have been reported, especially with immunizations utilizing little protein (Carlsen *et al.*, 1988). Therefore, the significant




Fig. 4. Immuno-blot analysis of microsomal cytochrome  $b_5$  from several animals. Microsomal loadings expressed as protein content: (1) *S. calcitrans*, 100  $\mu$ g; (2) Canton-S strain *D. melanogaster*, 72  $\mu$ g; (3) Hikone-R strain *D. melanogaster*, 74  $\mu$ g; (4) *M. autumnalis*, 105  $\mu$ g; (5) *A. mellifera*, 73  $\mu$ g; (6) LPR strain *M. domestica*, 121  $\mu$ g; (7) *T. ni*, 75  $\mu$ g; (8) *T. urticae*, 75  $\mu$ g; (9) *R. rattus*, 365  $\mu$ g; (10) S+ strain *M. domestica*, 143  $\mu$ g; (11) LPR strain *M. domestica*, 121  $\mu$ g. Cytochrome  $b_5$  in *T. urticae* was unassayable. Other loadings contained 25 pmol cytochrome  $b_5$  except (7) *T. ni* (8 pmol) and (9) *R. rattus*, which contained 50 pmol.




Fig. 5. Rocket immuno-electrophoresis of house fly cytochrome  $b_5$ . Slide A: unabsorbed antiserum. Slide B: absorbed antiserum. Picomoles purified cytochrome  $b_5$  in parentheses: well 1 (1.5), well 2 (3.0), well 3 (6.0), well 4 (12). Picomoles crude microsomal cytochrome  $b_5$ : well 5 (5.2), well 6 (5.2), well 7 (2.6), well 8 (2.6).

improvement in yield and the demonstration of adequate immune response represent fundamental improvements over our previous technique for isolating house fly cytochrome  $b_5$  (Wheelock and Scott, 1989).

The molecular mass for house fly cytochrome  $b_5$  was previously reported as  $19.7 \pm 0.76$  kDa (Wheelock and Scott, 1989) and was confirmed here as 19.7 (Fig. 1). The dipteran Mediterranean fruit fly cytochrome  $b_5$  has a value of 21.0 kDa (Megias *et al.*, 1985), similar to house fly. In addition, immuno-blot analysis of stable fly and face fly also detected putative cytochromes  $b_5$  at the same  $M_r$  as house fly cytochrome  $b_5$ . In contrast, mammalian cytochromes  $b_5$  have masses in the range 16–19.0 kDa (Spatz and Strittmatter, 1971), with rabbit liver (16.7 kDa) and rat liver cytochrome  $b_5$  (16.6 kDa) being typical (Spatz and Strittmatter, 1971; Burchell, 1985). Whether the higher masses seen with the dipteran cytochrome  $b_5$  are typical of dipterans in general or of insects in general requires further study. The only other purified insect cytochrome  $b_5$ , that from armyworm (McFadden *et al.*, 1979), was proteolytically truncated so that the reported mass of 11.4 kDa does not represent the entire molecule.

House fly cytochrome  $b_5$  appeared to be more hydrophobic, as judged by amino acid analysis (Table 2), than other insect or mammal cytochromes  $b_5$ . We have previously remarked on the unusually hydrophobic behavior of house fly cytochrome  $b_5$  during size-exclusion chromatography (Wheelock and Scott, 1989). However, the difference in per cent apolar amino acids between house fly and other insect cytochromes  $b_5$  was greater (42.2% vs 30.8–31.1%, respectively) than the same comparison between house fly and mammalian cytochromes  $b_5$  (42.2% vs 34.3–39.7%, respectively) (Table 2, Megias *et al.*, 1985; McFadden *et al.*, 1979; Ozols, 1972, 1974; Spatz and Strittmatter, 1971).

The consistent richness of ASX and GLX residues compared with basic residues HIS, LYS and ARG in cytochromes  $b_5$  has been noted by others (Megias *et al.*, 1985; Ozols, 1974). Compared with Mediterranean fruit fly (Megias *et al.*, 1985), armyworm (McFadden *et al.*, 1979), human (Ozols, 1972), rabbit (Spatz and Strittmatter, 1971), porcine, bovine and equine (Ozols, 1974) cytochromes  $b_5$ , house fly cytochrome  $b_5$  had the lowest [ASX + GLX/HIS + LYS + ARG] ratio (i.e. 1.24, compared with a range of 2.13–2.48 for the insects and 1.41–1.76 for the mammals).

In summary, the molecular mass and absorbance maximum of house fly cytochrome  $b_5$  are typical of other cytochromes  $b_5$ . However, the amino acid analysis reported here indicates that house fly cytochrome  $b_5$  has somewhat unique characteristics compared with other cytochromes  $b_5$ . In addition, the lack of cross-reactivity between anti-house fly cytochrome  $b_5$  and non-dipteran cytochrome  $b_5$  (Fig. 4) also suggests divergence between fly cytochrome  $b_5$  and other cytochromes  $b_5$ . Whether these characteristics translate into unusual functional properties for house fly cytochrome  $b_5$  remains to be seen.

**Acknowledgements**—We thank Dr Y. Konno for preparing the non-house fly, insect microsome samples and R. Sherwood for performing the amino acid analyses. This study was supported by a grant from the US Department of Agriculture and Hatch Project 139414.

## REFERENCES

- Burchell A. (1985) A simple method for purification of rat hepatic microsomal cytochrome  $b_5$ . *Biochem. J.* **226**, 339–341.
- Capdevila J. and Agosin M. (1975) Soluble cytochrome P-450 from housefly microsomes. Partial purification and characterization of two hemoprotein forms. *J. biol. Chem.* **250**, 1049–1060.
- Carlsen J., Christiansen K. and Jensen H. M. (1988) Rat hepatic microsomal cytochrome  $b_5$ . *Biochem. J.* **256**, 1051–1054.
- Dulley J. R. and Grieve P. A. (1975) A simple technique for eliminating interference by detergents in the Lowry method of protein determination. *Analyt. Biochem.* **64**, 136–141.
- Estabrook R. W. (1978) Microsomal electron-transport reactions: an overview. In *Methods in Enzymology* (Edited by Fleischer S. and Packer L.), Vol. 52, pp. 43–47. Academic Press, New York.
- Estabrook R. W. and Werringloer J. (1978) The measurement of difference spectra: application to the cytochromes of microsomes. In *Methods in Enzymology* (Edited by Fleischer S. and Packer L.), Vol. 52, pp. 212–220. Academic Press, New York.
- Hildebrandt A. and Estabrook R. W. (1971) Evidence for the participation of cytochrome  $b_5$  in hepatic microsomal mixed-function oxidation reactions. *Archs Biochem. Biophys.* **143**, 66–79.
- Kunkel J. G. (1988) Analytic immunologic techniques. *Immunological Techniques in Insect Biology* (Edited by Gilbert L. I. and Miller T. A.). Springer, New York.
- Laemmli U. K. (1970) Cleavage of structural proteins during assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- Lee S. S. T. and Scott J. G. (1989) Microsomal cytochrome P450 monooxygenases in the house fly (*Musca domestica* L.): biochemical changes associated with pyrethroid resistance and phenobarbital induction. *Pestic. Biochem. Physiol.* **35**, 1–10.
- McFadden K. A., Crankshaw D. L., Hetnarski H. K. and Wilkinson C. F. (1979) Purification and properties of cytochrome  $b_5$  from the larval midgut of the southern armyworm (*Spodoptera eridania*). *Insect Biochem.* **9**, 301–308.
- Megias A., Sabrido A. and Municio A. M. (1986) Cytochrome  $b_5$  from the insect *Ceratitis capitata*. *Biochem. biophys. Acta* **872**, 116–124.
- Omura T. and Sato R. (1964) The carbon monoxide binding pigment of liver microsomes. I. Evidence for its hemoprotein nature. *J. biol. Chem.* **239**, 2370–2378.
- Ozols J. (1972) Cytochrome  $b_5$  from a normal human liver. *J. biol. Chem.* **247**, 2241–2254.
- Ozols J. (1974) Cytochrome  $b_5$  from microsomal membranes of equine, bovine and porcine livers. Isolation and properties of preparations containing the membranous segment. *Biochemistry* **13**, 426–439.
- Ronis M. J. J. and Hodgson E. (1989) Cytochrome P-450 monooxygenases in insects. *Xenobiotica* **19**, 1077–1092.
- Scott J. G., Lee S. S. T. and Shono T. (1990) Biochemical changes in the cytochrome P450 monooxygenases of seven insecticide-resistant house fly (*Musca domestica* L.) strains. *Pestic. Biochem. Physiol.* **36**, 124–127.
- Spatz L. and Strittmatter P. (1971) A form of cytochrome  $b_5$  that contains an additional hydrophobic sequence of

- 40 amino acid residues. *Proc. natn. Acad. Sci. USA* **68**, 1042–1046.
- Towbin H., Staehelin T. and Gordon J. (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc. natn. Acad. Sci. USA* **76**, 4350–4354.
- Wheelock G. D., Konno Y. and Scott J. G. (1991) Expression of P450<sub>pr</sub> in house flies. *J. Biochem. Toxic.* **6**, 239–246.
- Wheelock G. D. and Scott J. G. (1989) Simultaneous purification of a cytochrome *P*-450 and cytochrome *b*<sub>5</sub> from the house fly, *Musca domestica* L. *Insect Biochem.* **19**, 481–488.
- Wheelock G. D. and Scott J. G. (1990) Immunological detection of cytochrome *P*450 from insecticide susceptible and resistant house flies. *Insecticide Biochem. Physiol.* **38**, 130–139.