

STRUCTURE NOTE

Rhesus macaque: A tight homodimeric CD8 $\alpha\alpha$

Lili Zong,^{1,2} Yong Chen,^{1,3,4} Hao Peng,¹ Feng Gao,¹ Aikichi Iwamoto,^{3,5,6} and George F. Gao^{1,3*}

¹ CAS Key Laboratory of Pathogenic Microbiology and Immunology (CASPMI), Institute of Microbiology, Chinese Academy of Sciences (CAS), Beijing 100101, China

² Department of Obstetrics and Gynaecology, Zhujiang Hospital, Nanfang Medical University, Guangzhou 510280, China

³ China-Japan Joint Laboratory of Molecular Immunology and Molecular Microbiology, Institute of Microbiology, Chinese Academy of Sciences (CAS), Beijing 100101, China

⁴ College of Life Sciences, Graduate University, Chinese Academy of Sciences (GUCAS), Beijing 100049, China

⁵ Research Center for Asian Infectious Diseases, The Institute of Medical Science, The University of Tokyo, Tokyo, Japan

⁶ Division of Infectious Diseases, Advanced Clinical Research Center, The Institute of Medical Science, The University of Tokyo, Tokyo, Japan

Key words: rhesus macaque; crystal structure; MHC binding; CD8; dimer; HIV; vaccine.

INTRODUCTION

Simian immunodeficiency virus (SIV) infection of rhesus macaque (*macaca mulatta*) is widely used as an animal model for human immunodeficiency virus (HIV) infection^{1–3} as well as other human diseases. It is known that the host cytotoxic T lymphocyte (CTL) responses provide powerful protection against HIV infection, and CTL-based immunization is currently believed to be the most promising approach toward vaccine development.⁴

As a coreceptor of T cell receptor (TCR) on the surface of CTLs, CD8 molecules stabilize the interaction of the TCR with major histocompatibility complex (MHC) by binding to the MHC class I (MHCI) molecule on the surface of antigen-presenting cells. In the absence of CD8 interaction, MHCI-restricted immune responses are hampered.⁵ In addition, recent data indicate that CD8 has the ability to bind to a nonclassical MHC class I-like molecule, TL antigen, independently of TCR and CD3, expanding the function of CD8 further to an immunomodulator.^{6–8} Moreover, soluble forms of CD8 can disrupt activation of some T cell clones with higher efficacy than anti-CD8 antibodies.^{9,10}

In both human and mouse, the functions of CD8 involved in immune responses have been extensively studied.^{11,12} The crystal structures of the human HLA-A*0201-CD8 $\alpha\alpha$ complex,^{13,14} murine MHC H-2K^b-CD8 $\alpha\alpha$ complex,¹⁵ TL antigen-CD8 $\alpha\alpha$ complex,¹⁶ and

the murine CD8 $\alpha\beta$ heterodimer¹⁷ have been solved. For macaque monkeys, however, little is known on the structures of the CTL-related molecules (e.g., TCR, MHC, and CD8). Only the structure of MHC allele Mamu-A*01 has been recently solved in our laboratory.¹⁸

In this article, we present the crystal structure of rhesus macaque CD8 $\alpha\alpha$ (rCD8 $\alpha\alpha$) homodimer and discuss the relatedness and uniqueness of rCD8 $\alpha\alpha$ structure with that of human/mouse CD8 $\alpha\alpha$ homodimer. Strikingly, with two Thr43 residues in C–C' loop, rCD8 $\alpha\alpha$ shows a unique extra hydrogen bond in the homodimeric interface indicating a tighter homodimeric interaction.

Grant sponsor: Ministry of Science and Technology (MOST), China (Basic Research Program 973); Grant number: 2006CB504204; Grant sponsor: National Natural Science Foundation (NSFC), China; Grant number: 30671903; Grant sponsor: Chinese Academy of Sciences (CAS, Knowledge Innovation Project); Grant number: KSCX2-SW-227; Grant sponsor: NSFC; Grant number: 30525010; Grant sponsor: Postdoctoral Fund of China; Grant number: 20070410649; Grant sponsor: Japan Ministry of Education, Culture, Sports, Science and Technology (MEXT, The China-Japan Joint Laboratory of Molecular Immunology and Molecular Microbiology).

Lili Zong and Yong Chen contributed equally to this work.

*Correspondence to: George F. Gao, Center for Molecular Immunology, Institute of Microbiology, Chinese Academy of Sciences (CAS), Datun Road, ChaoYang District, Beijing 100101, China. E-mail: gaof@im.ac.cn

Received 28 September 2008; Revised 10 November 2008; Accepted 12 November 2008

Published online 19 November 2008 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/prot.22331

MATERIALS AND METHODS

Expression and purification of rCD8 $\alpha\alpha$ homodimer

Rhesus macaque CD8 (rCD8) alpha chain nucleotides covering amino acids 1–120 of the ectodomain were synthesized based on the sequence of Indian origin rhesus (GeneBank ID: 698329). Inclusion bodies of rCD8 α were prepared, and rCD8 $\alpha\alpha$ homodimer was renatured and purified by using the protocols described earlier.^{13,14}

Crystallization, data collection, and processing

All crystallization attempts were performed at 18°C by the hanging drop vapor diffusion method. Ideal rCD8 $\alpha\alpha$ crystals grew from a 1:1 mixture of the protein solution (10 mg/mL) with crystallization reagent of 0.05M potassium phosphate monobasic, 20% w/v polyethylene glycol 8000. Data were collected using a Rigaku MicroMax007 rotating-anode X-ray generator (Cu K α ; λ = 1.5418 Å) equipped with an R-Axis VII++ image-plate detector. Data were processed and scaled using HKL2000.¹⁹

Structure solution, refinement, and analysis

Data were analyzed by molecular replacement²⁰ using Molrep in the CCP4 package,²¹ taking human CD8 $\alpha\alpha$ as the search probe (PDB code: 1AKJ).¹⁴ Final rounds of refinement resulted in a final Rcryst of 21.3% (R_{free} = 25.7%) for all data between 35.0 and 2.20 Å.

Buried surface areas were calculated using SURFACE²¹ with a 1.4 Å probe radius. The PyMOL Molecular Graphics System (DeLano Scientific, <http://www.pymol.org>) was used to prepare figures. Geometry of the refined structure was validated according to Ramachandran plot criteria.²² The data collection and refinement statistics of the structure are shown in Table I.

Accession number

Atomic coordinates of rhesus macaque CD8 $\alpha\alpha$ homodimer have been deposited in the Protein Data Bank (PDB, <http://www.rcsb.org/pdb>) under accession code: 2Q3A.

RESULTS AND DISCUSSION

Overall structure of rhesus macaque CD8 $\alpha\alpha$ homodimer

The crystals contained two CD8 $\alpha\alpha$ molecules as a dimer in a hand-shaking mode per crystallographic asymmetric unit. Belonging to the V set of Ig folds,¹⁴ the overall structure comparison of rCD8 $\alpha\alpha$ homodimer with the human counterpart is shown in Figure 1(A).

Table I

X-Ray Diffraction Data Processing and Refinement Statistics

| | |
|---|---|
| Data collecting | |
| Space group | P2 ₁ 2 ₁ 2 ₁ |
| Unit cell dimensions (<i>a</i> , <i>b</i> , <i>c</i>) | 46.54, 56.26, 82.31 |
| Unit cell dimensions (α , β , γ) | 90.00, 90.00, 90.00 |
| Resolution range (Å) | 40.00–2.20 (2.32–2.20) ^a |
| Total number of reflections | 138,752 |
| Number of unique reflections | 17,807 |
| Number of molecule in the asymmetric unit | 2 |
| Average redundancy | 6.60 (5.90) |
| Completeness (%) | 99.6 (100.0) |
| R_{merge} (%) | 9.6 (29.5) |
| I/σ | 12.8 (5.7) |
| Refinement | |
| Resolution (Å) | 35.85–2.20 |
| R -factor (%) | 21.3 |
| R_{free} ^b (%) | 25.7 |
| RMS deviations from restraint target values: | |
| Bond lengths (Å) | 0.006 |
| Bond angles (°) | 1.33 |
| Ramachandran plot Quality: | |
| Residues in most favored regions | 165 [84.2%] |
| Residues in additional allowed | 27 [13.8%] |
| Residues in generously allowed | 4 [2.0%] |
| Residues in disallowed regions | 0 [0%] |

^aValues in parentheses are given for the highest resolution shell.

^b R_{free} is calculated over reflections in a test set (5%) not included in atomic refinement.

Each CD8 α molecule is primarily composed of β structure arranged into two antiparallel β sheets. Short regions of 3_{10} helix are found between the E and F strands, which are not commonly found in CD8 β molecules.

All residues corresponding to the HLA-A2-CD8 $\alpha\alpha$ interface remain the same in MHC Mamu-A*01 and rCD8 $\alpha\alpha$.¹⁸ The interaction of HLA-CD8 is mainly based on charge complementarity and exhibits relatively low affinity (KD = 100–223 μ M) and rapid kinetics.^{8,23} The molecular surfaces of Mamu-A*01 and rCD8 $\alpha\alpha$ show similar complementarities (data not shown), indicating their interaction.

Structural comparison of rhesus macaque with human or murine CD8 $\alpha\alpha$ homodimer

Superposition of the final structure of rCD8 $\alpha\alpha$ dimer shows closer resemblance to human CD8 $\alpha\alpha$ homodimer (1AKJ, in complex with HLA-A2)¹⁴ than to murine CD8 $\alpha\alpha$ (mCD8 $\alpha\alpha$) (1BQH, in complex with Kd)¹⁵ [Fig. 1(A,B)]. Root mean square deviation (RMSD) of rCD8 $\alpha\alpha$ and hCD8 $\alpha\alpha$ dimer results in 0.831 Å for all C α atoms, which is much smaller than that of rCD8 $\alpha\alpha$ with mCD8 dimer of 1.634 Å.

The comparison of r/h/m CD8 $\alpha\alpha$ reveals some differences in loop regions, especially in complementarity




Figure 1

Crystal structure of rCD8 $\alpha\alpha$ homodimer and its superposition with human and mouse CD8 $\alpha\alpha$ structures. The ribbon diagram of each CD8 $\alpha\alpha$ was drawn and color-coded as: rCD8 $\alpha\alpha$, green and cyan; hCD8 $\alpha\alpha$, blue and magenta; mCD8 $\alpha\alpha$, yellow and orange. (A) Superposition of rCD8 $\alpha\alpha$ and hCD8 $\alpha\alpha$. The extra hydrogen bond region of rCD8 $\alpha\alpha$ was boxed. (B) Superposition of rCD8 $\alpha\alpha$ and mCD8 $\alpha\alpha$. The extra hydrogen bond region of rCD8 $\alpha\alpha$ was boxed. (C) The transition of A'- β strand in mCD8 $\alpha\alpha$ and A' loop in both rCD8 $\alpha\alpha$ /hCD8 $\alpha\alpha$ as the residue changes (Ala16 to Trp13). (D) Thr43 residues in C-C' loop of rCD8 $\alpha\alpha$ homodimer (cyan) form extra hydrogen bonds. Hydrogen bond between two main-chains is shown as red dashed line.

determining regions (CDRs), which are involved in MHCs binding. Interestingly, though CDR1 and CDR2 are structurally variable, CDR3 has almost the same conformation among r/h/m CD8 $\alpha\alpha$ molecules [Fig. 1(A,B)].

As one of the elements commonly found in many Ig domains, the first β strand of each domain is split into two shorter strands (A and A').²⁰ rCD8 $\alpha\alpha$ homodimer can also be characterized by a *cis*-proline (Pro7) at the transition point at the A strand. However, at the place of the A' strand, a loop is located in rhesus macaque as well as human CD8 α molecules as the result of the big side

chain of the residue Trp13, which is equivalent to Ala16 in mCD8 α molecules [Fig. 1(C)].

Corresponding to Ala42 in hCD8 α and Ser48 in mCD8 α , Thr43 residues in rCD8 α form an additional hydrogen bond to each other. This extra H-bond enhances the dimeric interaction and brings the C-C' loop of the two molecules into a much closer position [Fig. 1(D)]. As a result, the interface size of the two rCD8 α subunits ($\sim 2274 \text{ \AA}^2$, total buried solvent-accessible surface area) is significantly larger than that of human CD8 α interface ($\sim 2038 \text{ \AA}^2$) indicating a tighter homodimeric interaction of CD8 $\alpha\alpha$ in rhesus macaque.

ACKNOWLEDGMENTS

The authors thank Dr. Fuliang Chu for his valuable suggestions for this project and Mr. Christopher Pannell for his critical reading of the manuscript.

REFERENCES

- Allen TM, Sidney J, del Guercio ME, Glickman RL, Lensmeyer GL, Wiebe DA, DeMars R, Pauza CD, Johnson RP, Sette A, Watkins DI. Characterization of the peptide binding motif of a rhesus MHC class I molecule (Mamu-A*01) that binds an immunodominant CTL epitope from simian immunodeficiency virus. *J Immunol* 1998;160:6062–6071.
- Matano T, Shibata R, Siemon C, Connors M, Lane HC, Martin MA. Administration of an anti-CD8 monoclonal antibody interferes with the clearance of chimeric simian/human immunodeficiency virus during primary infections of rhesus macaques. *J Virol* 1998;72:164–169.
- Schmitz JE, Kuroda MJ, Santra S, Sasseville VG, Simon MA, Lifton MA, Racz P, Tenner-Racz K, Dalesandro M, Scallan BJ, Ghayeb J, Forman MA, Montefiori DC, Rieber EP, Letvin NL, Reimann KA. Control of viremia in simian immunodeficiency virus infection by CD8⁺ lymphocytes. *Science* 1999;283:857–860.
- Kaufmann SH, McMichael AJ. Annulling a dangerous liaison: vaccination strategies against AIDS and tuberculosis. *Nat Med* 2005;11: S33–S44.
- Fung-Leung WP, Schilham MW, Rahemtulla A, Kundig TM, Vollenweider M, Potter J, van Ewijk W, Mak TW. CD8 is needed for development of cytotoxic T cells but not helper T cells. *Cell* 1991;65:443–449.
- Cole DK, Gao GF. CD8: adhesion molecule, co-receptor and immuno-modulator. *Cell Mol Immunol* 2004;1:81–88.
- Leishman AJ, Naidenko OV, Attinger A, Koning F, Lena CJ, Xiong Y, Chang HC, Reinherz E, Kronenberg M, Cheroutre H. T cell responses modulated through interaction between CD8 $\alpha\alpha$ and the nonclassical MHC class I molecule, TL. *Science* 2001;294:1936–1939.
- Wyer JR, Willcox BE, Gao GF, Gerth UC, Davis SJ, Bell JI, van der Merwe PA, Jakobsen BK. T cell receptor and coreceptor CD8 $\alpha\alpha$ bind peptide-MHC independently and with distinct kinetics. *Immunity* 1999;10:219–225.
- Cole DK, Rizkallah PJ, Boulter JM, Sami M, Vuidepot AL, Glick M, Gao F, Bell JI, Jakobsen BK, Gao GF. Computational design and crystal structure of an enhanced affinity mutant human CD8 $\alpha\alpha$ coreceptor. *Proteins* 2007;67:65–74.
- Sewell AK, Gerth UC, Price DA, Purbhoo MA, Boulter JM, Gao GF, Bell JI, Phillips RE, Jakobsen BK. Antagonism of cytotoxic T-lymphocyte activation by soluble CD8. *Nat Med* 1999;5:399–404.
- Gao GF, Jakobsen BK. Molecular interactions of coreceptor CD8 and MHC class I: the molecular basis for functional coordination with the T-cell receptor. *Immunol Today* 2000;21:630–636.
- Gao GF, Rao Z, Bell JI. Molecular coordination of $\alpha\beta$ T-cell receptors and coreceptors CD8 and CD4 in their recognition of peptide-MHC ligands. *Trends Immunol* 2002;23:408–413.
- Gao GF, Gerth UC, Wyer JR, Willcox BE, O'Callaghan CA, Zhang Z, Jones EY, Bell JI, Jakobsen BK. Assembly and crystallization of the complex between the human T cell coreceptor CD8 α homodimer and HLA-A2. *Protein Sci* 1998;7:1245–1249.
- Gao GF, Tormo J, Gerth UC, Wyer JR, McMichael AJ, Stuart DI, Bell JI, Jones EY, Jakobsen BK. Crystal structure of the complex between human CD8 $\alpha(\alpha)$ and HLA-A2. *Nature* 1997;387:630–634.
- Kern PS, Teng MK, Smolyar A, Liu JH, Liu J, Hussey RE, Spoerl R, Chang HC, Reinherz EL, Wang JH. Structural basis of CD8 coreceptor function revealed by crystallographic analysis of a murine CD8 $\alpha\alpha$ ectodomain fragment in complex with H-2Kb. *Immunity* 1998;9:519–530.
- Liu Y, Xiong Y, Naidenko OV, Liu JH, Zhang R, Joachimiak A, Kronenberg M, Cheroutre H, Reinherz EL, Wang JH. The crystal structure of a TL/CD8 $\alpha\alpha$ complex at 2.1 Å resolution: implications for modulation of T cell activation and memory. *Immunity* 2003;18:205–215.
- Chang HC, Tan K, Ouyang J, Parisini E, Liu JH, Le Y, Wang X, Reinherz EL, Wang JH. Structural and mutational analyses of a CD8 $\alpha\beta$ heterodimer and comparison with the CD8 $\alpha\alpha$ homodimer. *Immunity* 2005;23:661–671.
- Chu F, Lou Z, Chen YW, Liu Y, Gao B, Zong L, Khan AH, Bell JI, Rao Z, Gao GF. First glimpse of the peptide presentation by rhesus macaque MHC class I: crystal structures of Mamu-A*01 complexed with two immunogenic SIV epitopes and insights into CTL escape. *J Immunol* 2007;178:944–952.
- Otwinowski Z, Minor W. Processing of X-ray diffraction data collected in oscillation mode. *Methods Enzymol* 1997;276:307–326.
- Chen Y, Chu F, Gao F, Zhou B, Gao GF. Stability engineering, biophysical, and biological characterization of the myeloid activating receptor immunoglobulin-like transcript 1 (ILT1/LIR-7/LILRA2). *Protein Expr Purif* 2007;56:253–260.
- CCP4. The CCP4 suite: programs for protein crystallography. *Acta Crystallogr D Biol Crystallogr* 1994;50:760–763.
- Lovell SC, Davis IW, Arendall WB, III, de Bakker PI, Word JM, Prisant MG, Richardson JS, Richardson DC. Structure validation by C α geometry: π , ψ and C β deviation. *Proteins* 2003;50:437–450.
- Gao GF, Willcox BE, Wyer JR, Boulter JM, O'Callaghan CA, Maenaka K, Stuart DI, Jones EY, Van Der Merwe PA, Bell JI, Jakobsen BK. Classical and nonclassical class I major histocompatibility complex molecules exhibit subtle conformational differences that affect binding to CD8 $\alpha\alpha$. *J Biol Chem* 2000;275:15232–15238.