

Published in final edited form as:

Bioorg Med Chem Lett. 2010 February 1; 20(3): 1302. doi:10.1016/j.bmcl.2009.10.018.

Syntheses and antibacterial activity studies of new oxazolidinones from nitroso Diels-Alder chemistry

Shanshan Yan^a, Marvin J. Miller^a, Timothy A. Wencewicz^a, and Ute Möllmann^b
^aDepartment of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, IN 46556, USA
^bLeibniz Institute for Natural Product Research and Infection Biology – Hans Knöell Institute,
Beutenbergstrasse 11a, D-07745 Jena, Germany

Abstract

A series of novel oxazolidinone antibiotics having [2.2.1] and [2.2.2] bicyclic oxazine moieties at the C-5 side chain of the A ring was synthesized by nitroso Diels—Alder reactions, from three linezolid analogs containing morpholine, piperazine and thiomorpholine, respectively, as the C-ring components. Subsequent N—O bond cleavage generated oxazolidinones with 4-amino cyclo-2- en-1-ol substituents. The *in vitro* antibacterial activities of these oxazolidinone analogs were evaluated.

Oxazolidinones are a new class of synthetic antimicrobial agents which are now clinically useful. They inhibit bacterial protein synthesis by binding to the 50S ribosomal subunit at the translation step. The most promising feature of these compounds is their oral activity against multidrug-resistant Gram-positive bacteria, including methicillin-resistant Staphylococcus aureus (MRSA) as well as select anaerobic organisms. Linezolid (Figure 1) is the first oxazolidinone drug approved for the treatment of Gram-positive bacterial infections in humans. ^{2b,3} Unfortunately, resistance to linezolid has already been observed in Gram-positive bacteria such as S. aureus and Enterococcus faecium.⁴ More recently, resistant strains of MRSA, E. coli and other bacteria have also been identified.⁵ Thus, there is significant need for the rapid development of novel antibacterial agents. Four Types of chemical modifications of linezolid and oxazolidinone-type antibiotics have been reported, including modifications on each of the A-, B- and C-rings as well as the C-5 side chain of the A ring substructure. ^{2b,6} The structureactivity- relationship studies of oxazolidinone antibiotics show that the stereochemistry (S) at the C-5 position of the A ring is critical in terms of biological activity. While an acetamide substituent on the 5-methyl group is found to produce optimal antibacterial activity, modification of the C-5 side chain would still benefit the development of new oxazolidinone antibiotics. In this regard, a frequently employed strategy involves the addition of heterocycles, such as triazoles, ⁷ and pyridones. ⁸ However, to the best of our knowledge, the incorporation of bicyclic oxazines into this class of antibiotics has not been disclosed.

Syntheses of interesting biologically relevant molecules using nitroso Diels–Alder chemistry (NDA) is of growing interest. As a synthetic tool for the formation of 4-amino cyclopent-2-en-1-ol derivatives in one step, acylnitroso Diels–Alder reactions are useful for the creation of unique structural and functional diversity. We and others have demonstrated that the nitroso

Correspondence to: Marvin J. Miller.

^{© 2009} Elsevier Ltd. All rights reserved.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

bicyclic oxazine **3** derived from NDA reaction between *in situ* generated acylnitroso **2** and 1,3-cyclopentadiene can undergo various chemical transformations (Scheme 1). We envisioned that modification at the A-ring C-5 side chain of linezolid-type oxazolidinone antibiotics using nitroso Diels—Alder chemistry would generate novel analogs with incorporation of bicyclic oxazine functionalities. These new analogs were anticipated to not only enhance the available set of oxazolidinone antibiotics, but also provide new scaffolds for further transformations. Herein we report the syntheses and biological evaluation of a series of oxazolidinone antibiotics in which the C-5 side chain has been modified by nitroso Diels—Alder reactions and subsequent chemical transformations.

To begin our investigation, three types of oxazolidinone precursors 14–16 were synthesized from commercial piperazine (4), morpholine (6) and thiomorpholine (7), respectively, as shown in Scheme 2. The syntheses began by separate S_NAr reactions of 3,4-difluoronitrobenzene with mono Boc-protected piperazine, 5, morpholine, 6, and thiomorpholine, 7, under basic conditions to afford para-substituted nitrobenzene derivatives 8–10 in moderate to good yields. Palladium catalyzed hydrogenolyses of 8 and 9, followed by reactions with CbzCl gave protected anilines 11–12. The synthesis of thiomorpholine analog 13 required modification to avoid poisoning of the catalyst by the sulfur atom during the hydrogenation step. Thus, use of indium powder and ammonium chloride 10 successfully reduced the nitro group of 10 and gave compound 13 in 62% yield after Cbz protection. Then, as reported, 11 compounds 11–13 were treated with nBuLi and R-glycidyl butyrate sequentially at -78 °C to form oxazolidinone analogs 14–16 in 70–80% yields with defined stereochemistry at the A-ring C-5 position.

With the desired aza-, oxa-, and thia-analogs, 14–16, in hand, introduction of the N-hydroxy carbamate moiety at the C-5 side chain and subsequent nitroso Diels-Alder reactions were explored (Scheme 3). Reactions of 14 and 15 with O-benzylhydroxylamine hydrochloride (OBHA·HCl) and carbonyldiimidazole (CDI) generated protected N-hydroxy carbamates 17 and 18 in 96% and 94% yields, respectively. Hydrogenolyses afforded the desired free Nhydroxy carbamates, 20 and 21, which were directly used in nitroso Diels-Alder reactions without further purification. Toward this end, N-hydroxy carbamate 20 was chosen as the model substrate. The cycloaddition reaction was carried out by in situ oxidation of 20 in the presence of 1,3-cyclohexadiene. Among several oxidants attempted, including sodium periodate (NaIO₄), Dess-Martin and FeCl₃/H₂O₂, ¹² we found that use of sodium periodate gave cycloadduct 23b in the best yield (56%) as one regioisomer after chromatography purification. For the thiomorpholine series, in order to again avoid catalyst poisoning during the subsequent deprotection step, O-p-methoxybenzyl hydroxylamine hydrochloride (PMBONH₂·HCl)¹³ instead of OBHA·HCl was used in the coupling reaction with **16**. The PMB group was removed under acidic conditions to give N-hydroxy carbamate 22 in good yield. Using the optimized NDA reaction conditions, a series of new oxazolidinones 23a-25b with bicyclic oxazines at the C-5 side chain were synthesized. Among them, cycloadducts 25a and 25b were obtained with the sulfur atom in the thiomorpholinyl ring oxidized. N-O bond cleavage reactions of these cycloadducts were also carried out, with intent to generate the corresponding oxazolidinones with additional 1,4-amino alcohol substituents. Thus, Mo (CO)₆¹⁴ successfully reduced the N–O bond of compounds 23a–25b to generate the corresponding oxazolidinone analogs, 26a-28b, that contain aminocycloalkenol functional groups, in moderate yields. In addition, under Mo(CO)₆ conditions, the sulfoxide group of 25a and 25b was also reduced. For comparison during the biological testing, an oxazolidinone analog 29 with a C-5 acetate substituent was synthesized from compound 15 (Scheme 4).

With the biological profile of parent linezolid, 1, known for comparison, all the oxazolidinone analogs 23a–28b, 29, as well as precursors 14–16, and ciprofloxacin, as a positive control, were subjected to broad antibacterial studies against various strains of Gram-positive and

Gram-negative bacteria as well as *Mycobacterium vaccae*, using agar diffusion assays (Table 1). ¹⁵ Compound **29** was found to be comparable to linezolid (**1**) itself for all tested organisms, as expected. Interestingly, all the oxazolidinone precursors **14–16** were roughly equipotent *in vitro* with linezolid (**1**) against several Gram-positive organisms, including *Bacillus subtilis*, *S. aureus*, *E. faecalis* and *M. luteus*. They also exhibited antimycobacterial activity and could potentially be useful for treatment of *M. tuberculosis* as they induced large inhibition zones against *Mycobacterium vaccae*, a common model for *M. tuberculosis*. Most oxazolidinone analogs with bicyclic oxazine or aminocycloalkenol moieties at the C-5 side chain, generated from NDA chemistry, had antimicrobial profiles similar to linezolid (**1**), but at a generally diminished level of activity. In general, oxazolidinone analogs substituted with [2.2.1] bicyclic oxazines (**23a–25a**) were more active than those with [2.2.2] bicyclic oxazines (**23b–25b**). Among them, compound **25a** derived from thiomorpholine exhibited the best activity. Analogs **26a–28b**, derived from reduction of the N–O bonds were relatively inactive compared to their parent cycloadducts.

In summary, we have synthesized a series of novel oxazolidinone antibiotics with [2.2.1] and [2.2.2] bicyclic oxazine as well as aminocycloalkenol moieties at the C-5 side chain through nitroso Diels–Alder chemistry. These oxazolidinone analogs exhibited *in vitro* antibacterial profiles similar to that of linezolid.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge the NIH (GM 075885) for support of this research. We thank Uta Wohlfeld (HKI) for performing antibacterial assays. We also thank the Lizzadro Magnetic Resonance Research Center at Notre Dame for NMR facility and Nonka Sevova for mass spectroscopic analyses. TAW acknowledges the University of Notre Dame Chemistry-Biochemistry-Biology Interface (CBBI) Program and NIH Training Grant T32GM075762 for a fellowship.

References and notes

- (a) Shinabarger DL, Marotti KR, Murray RW, Lin AH, Melchoir EP, Swaney SM, Dunyak DS, Demyan WF, Buysse JM. Antimicrob. Agents Chemother 1997;41:2132. [PubMed: 9333037] (b) Kloss P, Xiong L, Mankin AS, Shinabarger DL, Mankin AS. J. Mol. Biol 1999;294:93. [PubMed: 10556031]
- (a) Hutchinson DK. Curr. Top. Med. Chem 2003;3:1021. [PubMed: 12678835] (b) Barbachyn MR, Ford CW. Angew. Chem. Int. Ed 2003;42:2010. (c) Wookey A, Turner PJ, Greenhalgh JM, Eastwood M, Clarke J, Sefton C. Clin. Microbiol. Infect 2004;10:247. [PubMed: 15008947] (d) Phillips OA, Rotimi VO, Jamal WJ, Shahin M, Verghese TL. J. Chemother 2003;15:113. [PubMed: 12797385] (e) Moellering RC Jr. Ann. Intern. Med 2003;138:135. [PubMed: 12529096]
- 3. Brickner SJ, Hutchinson DK, Barbachyn MR, Manninen PR, Ulanowicz DA, Garmon SA, Grega KC, Hendges SK, Toops DS, Ford CW, Zurenko GE. J. Med. Chem 1996;39:673. [PubMed: 8576909]
- 4. Gonzales RD, Schreckenberger PC, Graham MB, Kelkar S, DenBesten K, Quinn JP. Lancet 2001;357:1179. [PubMed: 11323048]
- (a) Tsiodras S, Gold HS, Sakoulas G, Eliopoulos GM, Wennersten C, Venkataraman L, Moellering RC, Ferraro MJ. Lancet 2001;358:207. [PubMed: 11476839] (b) Xiong L, Kloss P, Douthwaite S, Anderson NM, Swaney S, Shinabarger DC, Mankin AS. J. Bacterial 2000;182:5325. (c) Mutnick AH, Enne V, Jones RN. Ann. Phermacother 2003;37:769.
- (a) Brickner S. Curr. Pharm. Des 1996;2:175. (b) Ford CW, Barbachyn MR. Angew. Chem. Int. Ed 2003;42:2010. (c) Hutchinson DK. Expert. Opin. Ther. Patents 2004;14:1309. (d) Renslo AR, Luehr GW, Gordeev MF. Bioorg. Med. Chem 2006;14:4227. [PubMed: 16527486] (e) Prasad JV. Curr. Opin. Microbiol 2007;10:454. [PubMed: 17928263]

(a) Phillips OA, Udo EE, Ali AAM, Samuel SM. Bioorg. Med. Chem 2005;13:4113. [PubMed: 15911326]
 (b) Phillips OA, Udo EE, Ali AAM, Samuel SM. Eur. J. Med. Chem 2007;42:214. [PubMed: 17140706]

- 8. Paget SD, Boggs CM, Foleno BD, Goldschmidt RM, Hlasta DJ, Weidner-Wells MA, Werblood HM, Bush K, Macielag MJ. Bioorg. Med. Chem. Lett 2006;16:4537. [PubMed: 16806921]
- 9. For reviews, see: (a) Streith J, Defoin A. Synthesis 1994;11:1107. (b) Vogt PF, Miller MJ. Tetrahedron 1998;54:1317. (c) Yamamoto H, Momiyama N. Chem. Commun 2005:3514. (d) Yamamoto Y, Yamamoto H. Eur. J. Org. Chem 2006:2031. (e) Samarakoon T, Hanson RR. Chemtracts 2007;20:220.
- 10. Banik BK, Banik I, Becker FF. Org. Synth 2005;81:188.
- 11. Tucker JA, Allwine DA, Grega KC, Barbachyn MR, Klock JL, Adamski JL, Brickner SJ, Hutchinson DK, Ford CW, Zurenko GE, Conradi RA, Burton PS, Jensen RM. J. Med. Chem 1998;41:3727. [PubMed: 9733498]
- 12. Adamo MFA, Bruschi S. J. Org. Chem 2007;72:2666. [PubMed: 17326685]
- 13. For synthesis of PMBONH₂·HCl, see: Zhu G, Yang FL, Balachandran R, Hook P, Vallee RB, Curran DP, Day BW. J. Med. Chem 2006;49:2063. [PubMed: 16539395]
- 14. Cicchi S, Goti A, Brandi A, Guarna A, De Sarlos FD. Tetrahedron Lett 1990;31:3351.
- 15. The *in vitro* biological testing of *N*-hydroxy carbamates **20–22** (>90% pure based on LC/MS) showed these compounds also retained antimicrobial profiles similar to linezolid (1), but at a generally diminished level of activity. For detailed data, see Table 1 in Supplementary Content.

C-5 side chain

Figure 1. Linezolid antibiotic (1).

$$\begin{array}{c|c}
 & O \\
 & N \end{array}$$

Scheme 1. Acylnitroso Diels–Alder (NDA) reaction

Scheme 2.

Reagents and conditions: (a) Boc₂O, DCM, 50%; (b) 3,4-difluoronitrobenzene, Hunig's base, CH₃CN, 49% (for **8**), 84% (for **9**), 70% (for **10**); (c) (i) Pd/C, H₂, MeOH, (ii) CbzCl, NaHCO₃, THF/H₂O, 89% (for **11**), 92% (for **12**); (d) (i) In, NH₄Cl, EtOH, reflux, (ii) CbzCl, NaHCO₃, THF/H₂O, 62% (for **13**); (e) (i) *n*BuLi, THF, -78 °C, (ii) *R*-glycidyl butyrate, 71% (for **14**), 70% (for **15**), 80% (for **16**).

Scheme 3.

Reagents and conditions: (a) (i) CDI, CH₃CN (ii) OBHA·HCl, imidazole, 96% (for **17**), 94% (for **18**); (b) CDI, CH₃CN, (ii) PMBONH₂·HCl, imidazole, 92% (for **19**); (c) Pd/C, H₂, MeOH; (d) (i) 10% TFA, DCM, (ii) sat. Na₂CO₃; (e) NaIO₄, 1,3-cyclopentadiene, MeOH/H₂O, 0°C, 50% (for **23a**, from **17**), 49% (for **24a**, from **18**), 40% (for **25a**, from **19**); (e) NaIO₄, 1,3-cyclopentadiene, MeOH/H₂O, 0°C, 50% (for **23a**, from **17**), 49% (for **24a**, from **18**), 40% (**25a**, from **19**); (f) NaIO₄, 1,3-cyclohexadiene, MeOH/H₂O, 0°C, 56% (for **23b**, from **17**), 47% (for **24b**, from **18**), 42% (**25b**, from **19**); (g) Mo(CO)₆, CH₃CN/H₂O, 80°C, 50% (for **26a**), 43% (for **26b**), 42% (for **27a**), 62% (for **27b**), 54% (for **28a**), 55% (for **28b**).

Scheme 4. Reagents and conditions: (a) Ac₂O, Pyridine, DMAP, DCM, 75%.

Table 1

Yan et al.

Antimicrobial activity in the agar diffusion assay (Diameter of inhibition zone, measured in mm).

	Gram-positive bacteria	acteria					Gram-negative bacteria	eria		
Compds	Bacillus subtilis	Micrococcus luteus	Staphylococcus aureus	cus aureus	Enterococcus faecalis	us faecalis	Escherichia coli	Pseudomonas aeruginosa	s aeruginosa	Mycobacterium vaccae
	ATCC 6633	ATCC 10240	$SG 511^a$	134/93 (MRSA)	ATCC 49532	1528 (VRE)	ATCC 25922	$\kappa_{199/WT}$	$\mathrm{K799/61}^{b}$	IMET 10670
1	31/35P	38	27/36P	LN	25	L	0	0	10P	42
29	26.5	38/46p	26/35P	LN	24	Z	0	0	17h	43
14	23/26P	34.5	24/33P	LN	21	Z	0	0	10P	L
15	26/29P	38	23/32P	LN	21	Z	0	0	11P	43
16	26/29P	41	24/35P	LN	21.5	Z	0	0	11P	43
23a	12/18P	16	13/19p	19	19	16	0	14/20p	0	19p
23b	18P	LN	16P	14p	Z	12P	IN	LN	0	13
24a	17/30P	L	15/21p	25	N	19	L	LN	0	30
24b	19P	L	16P	18p	N	16	IN	LN	0	15
25a	21/28P	22/32P	18/22p	LN	19	N	0	0	11P	35
25b	17P	15p	15p	LN	17	Z	0	0	10P	18
26a	13h	12P	14p	LN	12.5	N	0	0	10P	12h
26b	0	0	0	LN	12p	N	0	0	12P	0
27a	14P	L	15P	14p	N	12P	L	LN	0	18
27b	13P	L	14P	14P	N	12P	IN	LN	0	0
28a	14P	15h	12.5p	LN	12.5	N	0	0	10P	18
28b	12h	14	0	LN	12	Z	0	0	10P	0
cipro	29.5	0	24	0	13	15	31/40P	35	39	19p

a wild type.

 $\frac{b}{\text{permeability mutant.}}$

p, partially clear inhibition zone/colonies in the inhibition zone.

P, unclear inhibition zone/many colonies in the inhibition zone.

h, faint indication of inhibition zone.

Page 10

Exactly 50 µl of a 2.0 mM solution of each compound dissolved in 1:9 DMSO:MeOH was filled in 9 mm wells in agar media (Standard I Nutrient Agar, Serva or Mueller Hinton II Agar, Becton, Dickinson and Company). Inhibition zones read after incubation at 37 °C for 24 h. Cipro (ciprofloxacin) was dissolved in H2O to give 5 µg/mL solution.