



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### An Ion-Pair Template for Rotaxane Formation and its Exploitation in an Orthogonal Interaction Anion-Switchable Molecular Shuttle

**Citation for published version:**

Barrell, MJ, Leigh, DA, Lusby, PJ & Slawin, AMZ 2008, 'An Ion-Pair Template for Rotaxane Formation and its Exploitation in an Orthogonal Interaction Anion-Switchable Molecular Shuttle' *Angewandte Chemie International Edition*, vol 47, no. 42, pp. 8036-8039., 10.1002/anie.200802745

**Digital Object Identifier (DOI):**

[10.1002/anie.200802745](https://doi.org/10.1002/anie.200802745)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

*Angewandte Chemie International Edition*

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



This is the peer-reviewed version of the following article:

Barrell, M. J., Leigh, D. A., Lusby, P. J., & Slawin, A. M. Z. (2008). An Ion-Pair Template for Rotaxane Formation and its Exploitation in an Orthogonal Interaction Anion-Switchable Molecular Shuttle. *Angewandte Chemie-International Edition*, 47(42), 8036-8039.

which has been published in final form at <http://dx.doi.org/10.1002/anie.200802745>

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for self-archiving (<http://olabout.wiley.com/WileyCDA/Section/id-817011.html>).

Manuscript received: 11/06/2008; Article published: 15/09/2008

## An Ion-Pair Template for Rotaxane Formation and its Exploitation in an Orthogonal Interaction Anion-Switchable Molecular Shuttle\*\*

Michael J. Barrell,<sup>1</sup> David A. Leigh,<sup>1,\*</sup> Paul J. Lusby<sup>1,\*</sup> and Alexandra M. Z. Slawin<sup>2</sup>

<sup>[1]</sup>EaStCHEM, School of Chemistry, Joseph Black Building, University of Edinburgh, West Mains Road, Edinburgh, EH9 3JJ, UK.

<sup>[2]</sup>School of Chemistry, University of St. Andrews, Purdie Building, St. Andrews, Fife KY16 9ST, UK.


<sup>[\*]</sup>Corresponding authors; P.J.L. e-mail: [paul.lusby@ed.ac.uk](mailto:paul.lusby@ed.ac.uk), fax: (+44) 131-650-6453, website: <http://www.catenane.net>; D.A.L. e-mail: [david.leigh@ed.ac.uk](mailto:david.leigh@ed.ac.uk)

<sup>[\*\*]</sup>We thank the EPSRC National Mass Spectrometry Service Centre (Swansea, U.K.) for accurate mass data. This work was supported by the EPSRC. P.J.L. is a Royal Society University Research Fellow. D.A.L. is an EPSRC Senior Research Fellow and holds a Royal Society-Wolfson Research Merit Award.

### Supporting information:

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

### Graphical abstract:



### Keywords:

Rotaxanes; molecular shuttles; template synthesis; palladium; anion recognition


## Abstract

Chloride Control: The serendipitous discovery of a molecular recognition motif triggered by the formation of an anion-palladium coordination bond is exploited as both an efficient rotaxane-forming template and as the basis of a chloride-switchable molecular shuttle.

## Main text

Despite significant advances<sup>[1]</sup> in anion-template methods for the construction of mechanically interlocked molecules,<sup>[2]</sup> the use of anions to induce changes in the relative positions of the components of catenanes and rotaxanes has proved particularly challenging,<sup>[3]</sup> especially in comparison to the widespread success achieved with other stimuli.<sup>[4-6]</sup> The few examples of anion-switchable molecular shuttles developed to date employ competition between the same types of weak interaction in both states of the molecule to achieve switching (solvation effects<sup>[3d,e]</sup> or the anion out-competing macrocycle hydrogen bonding acceptor groups for donor groups on the thread<sup>[3e,f]</sup>). Other features of anions, such as the propensity of halides to form strong coordination bonds to various transition metals, have yet to be exploited.<sup>[7]</sup> Here we report on the serendipitous discovery of a new efficient template for rotaxane formation and its use in the assembly of a chloride-switchable molecular shuttle which exhibits excellent positional integrity (>98%) of the ring in both states due to orthogonal modes of binding: direct intercomponent metal-ligand coordination in one state and a combination of tight ion-pairing, aromatic stacking interactions and CH $\cdots$ O and CH $\cdots$ Cl hydrogen bonding in the other.

The development of the new rotaxane-forming template was prompted by the chance observation that chloride displacement of the acetonitrile ligand of [(**L1**)Pd(CH<sub>3</sub>CN)] using benzyl pyridinium chloride (**1.Cl**) was accompanied by encapsulation of the organic cation by the anionic PdCl-coordinated macrocycle [(**L1**)PdCl]<sup>-</sup> (Figure 1a).<sup>[8]</sup> The threaded nature of the complex {[(**L1**)PdCl].**1**} in CDCl<sub>3</sub> was clearly apparent from <sup>1</sup>H NMR spectroscopy (a distinct upfield shift in the pyridinium resonances with respect to those in **1.Cl** caused by shielding by the benzylic groups of the macrocycle, see Supporting Information) and was also found to persist in the solid state (Figure 1b and 1c) from X-ray crystallography of single crystals grown from a saturated CH<sub>2</sub>Cl<sub>2</sub>/EtOAc solution.<sup>[9]</sup>




**Figure 1.** An allosteric anion-activated template for threading based on tight ion-pairing reinforced through multiple other non-covalent interactions. a) Synthesis of pseudorotaxane  $\{[(L1)PdCl]·1\}$ . Conditions:  $CH_2Cl_2$ , 1 h, quantitative yield (employing **1**.PF<sub>6</sub> instead of **1**.Cl, or using DMSO instead of  $CH_2Cl_2$ , does not lead to pseudorotaxane formation). b) Side-on and c) face-on views of the X-ray crystal structure of  $\{[(L1)PdCl]·1\}$ .<sup>[9]</sup> Nitrogen atoms are shown in blue, oxygen red, palladium silver, chlorine and the carbons of the macrocycle green, pyridinium purple, and other carbons gray. Selected bond lengths [Å] and angles [°]: N1-Pd 2.04, N2-Pd 1.93, N3-Pd 2.03, Cl-Pd 2.32, O1-H3 2.47, O2-H1 2.57, O2-H1 2.37, Pd-Cl 3.12, N1-Pd-N3 160.8, N2-Pd-Cl 176.4.

The solid state structure of  $\{[(\mathbf{L1})\text{PdCl}]\cdot\mathbf{1}\}$  indicates that a broad range of noncovalent interactions are responsible for the assembly of the threaded architecture. In addition to the tight ion pair<sup>[10]</sup> (the pyridinium nitrogen atom is within 4 Å of atoms in the first coordination sphere of the formally negatively charged metal complex), aromatic stacking interactions between the aromatic rings of host and guest, aryl- and alkyl-CH $\cdots$ O hydrogen bonding between the polyether oxygen atoms and protons on carbons adjacent to the pyridinium nitrogen (C–O distances 3.19–3.36 Å), and CH $\cdots$ Cl(Pd) second coordination sphere interactions<sup>[7]</sup> (C–Cl distance 3.72 Å) all apparently contribute to the stability of the interpenetrated structure. Many of these primarily electrostatic interactions should be much weaker in more polar environments and, indeed, the <sup>1</sup>H NMR spectrum of  $\{[(\mathbf{L1})\text{PdCl}]\cdot\mathbf{1}\}$  (see Supporting Information) shows the complex is largely unthreaded in [D<sub>6</sub>]DMSO. The chloride anion is a vital component of the assembly process:  $[(\mathbf{L1})\text{Pd}(\text{CH}_3\text{CN})]$  did not form a threaded complex when reacted with  $\mathbf{1}\cdot\text{PF}_6$  in CH<sub>2</sub>Cl<sub>2</sub>, a result which suggested that the recognition motif could also be used as the basis of an anion-selective trigger.


A pyridinium chloride salt suitable for rotaxane synthesis ( $\mathbf{2}\cdot\text{Cl}$ ) was prepared in four steps from 4-pyridinepropanol (see Supporting Information). Treatment of  $[(\mathbf{L1})\text{Pd}(\text{CH}_3\text{CN})]$  with  $\mathbf{2}\cdot\text{Cl}$  in dichloromethane for 1 h, followed by reaction with  $\mathbf{3}$  via a Cu(I)-catalyzed azide-alkyne 1,3-cycloaddition (CuAAC)<sup>[11]</sup> in the presence of *tris*-(benzyltriazolylmethyl)amine (TBTA)<sup>[12]</sup> and DIPEA (diisopropylethylamine) led to  $[(\mathbf{L2})\text{PdCl}]$  in 64% yield (Scheme 1). Mass spectrometry of this product was strongly suggestive of a rotaxane—the major peak, corresponding to  $[(\mathbf{L2})\text{Pd}]^+$ , does not fragment to the intact macrocycle and axle as would be expected for a non-interlocked salt—and the covalent capture of the threaded structure was confirmed when removal of palladium also did not cause separation of the components (see Supporting Information). Comparison of the <sup>1</sup>H NMR spectrum of  $[(\mathbf{L2})\text{PdCl}]$  (Figure 2b) with the chloride salt of the thread (Figure 2a), shows significant upfield shifts to the pyridinium resonances H<sub>d</sub>, H<sub>e</sub> and H<sub>f</sub>, and adjacent protons (H<sub>a</sub> and H<sub>c</sub>). Other signals of the thread exhibit little change, indicating that the anionic PdCl-macrocycle is located overwhelmingly over the pyridinium station, i.e. the structure is *pyrdm*<sup>[13]</sup>- $[(\mathbf{L2})\text{PdCl}]$  (Scheme 1).

The triazole group introduced by the CuAAC reaction can also act as a ligating station for the palladium-macrocycle.<sup>[14]</sup> Treatment of *pyrdm*- $[(\mathbf{L2})\text{PdCl}]$  with AgPF<sub>6</sub> (1.1 equiv., Scheme 1 step c) smoothly precipitated AgCl giving quantitative conversion to a new rotaxane,  $[(\mathbf{L2})\text{Pd}]\text{PF}_6$ , in which the chloride ligand had been replaced by the non-coordinative PF<sub>6</sub> counter-ion. Comparison of the <sup>1</sup>H NMR spectrum of  $[(\mathbf{L2})\text{Pd}]\text{PF}_6$  (Figure 2c) with that of the PF<sub>6</sub> salt of the thread (Figure 2d), shows that the signals of the pyridinium station (H<sub>d-f</sub>) now occur at similar chemical shifts in thread and rotaxane, while the triazole resonance (H<sub>j</sub>) and adjacent protons (e.g. H<sub>i</sub>) are shifted upfield in the rotaxane indicating that removal of the chloride from the palladium center is accompanied by translocation of the palladium-macrocycle component to give *triazole*- $[(\mathbf{L2})\text{Pd}]\text{PF}_6$ . Simple addition of tetrabutylammonium chloride to *triazole*- $[(\mathbf{L2})\text{Pd}]\text{PF}_6$  in

chloroform (Scheme 1, step d) reverses this process, producing a product with identical physical and spectroscopic properties to the original rotaxane, *pyrdm*-[(**L2**)PdCl].



**Scheme 1.** Synthesis and operation of chloride-switchable molecular shuttle [(**L2**)Pd] $^+$ : a)  $\text{CH}_2\text{Cl}_2$ , 1 h; b) **3** (1.1 equiv),  $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$  (0.2 equiv), TBTA (0.25 equiv), DIPEA (1 equiv),  $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{CN}$  (7:1), 18 h, 64% (from **2-Cl**); c)  $\text{AgPF}_6$  (1.1 equiv), acetone, 18 h, quantitative; d)  $\text{Bu}_4\text{NCl}$  (1.5 equiv),  $\text{CHCl}_3$ , quantitative.



**Figure 2.** Partial  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 298 K) spectra of a) thread chloride salt, b) *pyrdm*-[(**L2**)PdCl], c) *triazole*-[(**L2**)Pd]PF<sub>6</sub>, d) thread PF<sub>6</sub> salt. The assignments correspond to the lettering shown in Scheme 1. Signals shown in gray are due to impurities and residual solvents.

The chance discovery of a molecular recognition motif that is triggered by the formation of an anion-palladium coordination bond has been exploited as both an efficient rotaxane-forming template and as the basis for a chloride-switchable molecular shuttle. The use of wholly different and orthogonal binding modes in the two states of the shuttle leads to exceptional positional integrity of the ring in both forms. Anion-activated allosteric templates could lead to new developments in sensors, logic gates, transport agents and molecular machines.

### Experimental Section

Experimental procedure for the preparation of *pyrdm*-[(**L2**)PdCl]: **2**·Cl (59 mg, 0.066 mmol) and [(**L1**)Pd(CH<sub>3</sub>CN)] (50 mg, 0.073 mmol) were stirred for 1 h in CH<sub>2</sub>Cl<sub>2</sub> (7 mL) prior to the addition of **3** (43 mg, 0.073 mmol) and DIPEA (11  $\mu\text{L}$ , 0.066 mmol). A solution of TBTA (9 mg, 0.017 mmol) and Cu(CH<sub>3</sub>CN)<sub>4</sub>PF<sub>6</sub> (6 mg, 0.0147 mmol) in CH<sub>3</sub>CN (1 mL) was added and the reaction mixture allowed to stir

for a further 18 h. After this time the volatile compounds were removed under reduced pressure and the residue taken up in CH<sub>2</sub>Cl<sub>2</sub> (10 mL). The resulting solution was washed with NH<sub>4</sub>Cl (3 x 10 mL) and the aqueous phase re-extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 5 mL). The combined organic extracts were concentrated under reduced pressure and subjected to flash chromatography on silica gel (0-3% MeOH in CH<sub>2</sub>Cl<sub>2</sub> as eluent) to give *pyrdm*-[(**L2**)PdCl] as a bright yellow solid (89 mg, 64%). For compound characterization, full synthetic details for all precursors and the switching experiments, see the Supporting Information.



## Notes and references

- [1] For a recent review on the anion-template synthesis of interlocked molecules, see: M. S. Vickers, P. D. Beer, *Chem. Soc. Rev.* **2007**, *36*, 211–225.
- [2] a) G. M. Hübner, J. Gläser, C. Seel, F. Vögtle, *Angew. Chem.* **1999**, *111*, 395–398; *Angew. Chem. Int. Ed.* **1999**, *38*, 383–386; b) J. A. Wisner, P. D. Beer, M. G. B. Drew, *Angew. Chem.* **2001**, *113*, 3718–3721; *Angew. Chem. Int. Ed.* **2001**, *40*, 3606–3609; c) J. A. Wisner, P. D. Beer, M. G. B. Drew, M. R. Sambrook, *J. Am. Chem. Soc.* **2002**, *124*, 12469–12476; d) M. R. Sambrook, P. D. Beer, J. A. Wisner, R. L. Paul, A. R. Cowley, *J. Am. Chem. Soc.* **2004**, *126*, 15364–15365; e) K.-Y. Ng, A. R. Cowley, P. D. Beer, *Chem. Commun.* **2006**, 3676–3678.
- [3] For co-conformational switching induced by intramolecular anion-hydrogen bonding (involving deprotonation of a phenol), see: a) C. M. Keaveney, D. A. Leigh, *Angew. Chem.* **2004**, *116*, 1242–1244; *Angew. Chem. Int. Ed.* **2004**, *43*, 1222–1224; b) J. Berná, A. M. Brouwer, S. M. Fazio, N. Haraszkiewicz, D. A. Leigh, C. M. Lennon, *Chem. Commun.* **2007**, 1910–1912; c) K.-Y. Ng, V. Felix, S. M. Santos, N. H. Reesa, P. D. Beer, *Chem. Commun.* **2008**, 1281–1283. For co-conformational switching induced by anion solvation effects, see: d) B. W. Laursen, S. Nygaard, J. O. Jeppesen, J. F. Stoddart, *Org. Lett.* **2004**, *6*, 4167–4170; e) C.-F. Lin, C.-C. Lai, Y.-H. Liu, S.-M. Peng, S.-H. Chiu, *Chem. Eur. J.* **2007**, *13*, 4350–4355. For co-conformational switching induced by intermolecular anion-hydrogen bonding, see: ref [3e] and f) Y.-L. Huang, W.-C. Hung, C.-C. Lai, Y.-H. Liu, S.-M. Peng, S.-H. Chiu, *Angew. Chem.* **2007**, *119*, 6749–6753; *Angew. Chem. Int. Ed.* **2007**, *46*, 6629–6633.
- [4] For examples of light-switchable molecular shuttles, see: a) A. C. Benniston, A. Harriman, *Angew. Chem.* **1993**, *105*, 1553–1555; *Angew. Chem. Int. Ed. Engl.* **1993**, *32*, 1459–1461; b) H. Murakami, A. Kawabuchi, K. Kotoo, M. Kunitake, N. Nakashima, *J. Am. Chem. Soc.* **1997**, *119*, 7605–7606; c) A. M. Brouwer, C. Frochot, F. G. Gatti, D. A. Leigh, L. Mottier, F. Paolucci, S. Roffia, G. W. H. Worpel, *Science* **2001**, *291*, 2124–2128; d) C. A. Stanier, S. J. Alderman, T. D. W. Claridge, H. L. Anderson, *Angew. Chem.* **2002**, *114*, 1847–1850; *Angew. Chem. Int. Ed.* **2002**, *41*, 1769–1772; e) A. Altieri, G. Bottari, F. Dehez, D. A. Leigh, J. K. Y. Wong, F. Zerbetto, *Angew. Chem.* **2003**, *115*, 2398–2402; *Angew. Chem. Int. Ed.* **2003**, *42*, 2296–2300; f) G. Bottari, D. A. Leigh, E. M. Pérez, *J. Am. Chem. Soc.* **2003**, *125*, 13360–13361; g) E. M. Pérez, D. T. F. Dryden, D. A. Leigh, G. Teobaldi, F. Zerbetto, *J. Am. Chem. Soc.* **2004**, *126*, 12210–12211; h) H. Murakami, A. Kawabuchi, R. Matsumoto, T. Ido, N. Nakashima, *J. Am. Chem. Soc.* **2005**, *127*, 15891–15899; i) J.-P. Collin, D. Jouvenot, M. Koizumi, J.-P. Sauvage, *Eur. J. Inorg. Chem.* **2005**, 1850–1855; j) D.-H. Qu, Q.-C. Wang, H. Tian, *Angew. Chem.* **2005**, *117*, 5430–5433; *Angew. Chem. Int. Ed.* **2005**, *44*, 5296–5299; k) S. Schmidt-Schäffer, L. Grubert, U. W. Grummt, K. Buck, W. Abraham, *Eur. J. Org. Chem.* **2006**, 378–398; l) V. Balzani, M. Clemente-León, A. Credi, B. Ferrer, M. Venturi, A. H. Flood, J. F. Stoddart, *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 1178–1183; m) W. Abraham, K. Buck, M. Orda-Zgadaj, S. Schmidt-Schäffer, U.-W. Grummt, *Chem. Commun.* **2007**,

3094–3096; n) V. Serreli, C.-F. Lee, E. R. Kay, D. A. Leigh, *Nature* **2007**, *445*, 523–527; o) W. Zhou, D. Chen, J. Li, J. Xu, J. Lv, H. Liu, Y. Li, *Org. Lett.* **2007**, *9*, 3929–3932; For a recent review, see: p) S. Saha, J. F. Stoddart, *Chem. Soc. Rev.* **2007**, *36*, 77–92.

- [5] For examples of pH-switchable molecular shuttles, see: a) R. A. Bissell, E. Córdova, A. E. Kaifer, J. F. Stoddart, *Nature* **1994**, *369*, 133–137; b) M.-V. Martinez-Diaz, N. Spencer, J. F. Stoddart, *Angew. Chem.* **1997**, *109*, 1991–1994; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 1904–1907; c) P. R. Ashton, R. Ballardini, V. Balzani, I. Baxter, A. Credi, M. C. T. Fyfe, M. T. Gandolfi, M. Gómez-López, M.-V. Martinez-Díaz, A. Piersanti, N. Spencer, J. F. Stoddart, M. Venturi, A. J. P. White, D. J. Williams, *J. Am. Chem. Soc.* **1998**, *120*, 11932–11942; d) A. M. Elizarov, S.-H. Chiu, J. F. Stoddart, *J. Org. Chem.* **2002**, *67*, 9175–9181; e) K.-W. Cheng, C.-C. Lai, P.-T. Chianga, S.-H. Chiu, *Chem. Commun.* **2006**, 2854–2856; f) Y. Tokunaga, T. Nakamura, M. Yoshioka, Y. Shimomura, *Tetrahedron Lett.* **2006**, *47*, 5901–5904; g) D. A. Leigh, A. R. Thomson, *Org. Lett.* **2006**, *8*, 5377–5379; h) J. D. Crowley, D. A. Leigh, P. J. Lusby, R. T. McBurney, L.-E. Perret-Aebi, C. Petzold, A. M. Z. Slawin, M. D. Symes, *J. Am. Chem. Soc.* **2007**, *129*, 15085–15090; i) S. J. Vella, J. Tiburcio, S. J. Loeb, *Chem. Commun.* **2007**, 4752–4754; j) D. Tuncel, Ö. Özsar, H. B. Tiftika, B. Salih, *Chem. Commun.* **2007**, 1369–1371; k) W. Zhou, J. Li, X. He, C. Li, J. Lv, Y. Li, S. Wang, H. Liu, D. Zhu, *Chem. Eur. J.* **2008**, *14*, 754–763.
- [6] For examples of redox-switchable molecular shuttles, see: a) N. Armaroli, V. Balzani, J.-P. Collin, P. Gaviña, J.-P. Sauvage, B. Ventura, *J. Am. Chem. Soc.* **1999**, *121*, 4397–4408; b) A. Altieri, F. G. Gatti, E. R. Kay, D. A. Leigh, D. Martel, F. Paolucci, A. M. Z. Slawin, J. K. Y. Wong, *J. Am. Chem. Soc.* **2003**, *125*, 8644–8654; c) G. Cooke, J. F. Garety, B. Jordan, N. Kryvokhyzha, A. Parkin, G. Rabani, V. M. Rotello, *Org. Lett.* **2006**, *8*, 2297–2300; d) T. D. Nguyen, Y. Liu, S. Saha, K. C.-F. Leung, J. F. Stoddart, J. I. Zink, *J. Am. Chem. Soc.* **2007**, *129*, 626–634; e) S. Nygaard, K. C.-F. Leung, I. Aprahamian, T. Ikeda, S. Saha, B. W. Laursen, S.-Y. Kim, S. W. Hansen, P. C. Stein, A. H. Flood, J. F. Stoddart, J. O. Jeppesen, *J. Am. Chem. Soc.* **2007**, *129*, 960–970; f) S. Saha, A. H. Flood, J. F. Stoddart, S. Impellizzeri, S. Silvi, M. Venturi, A. Credi, *J. Am. Chem. Soc.* **2007**, *129*, 12159–12171; g) I. Aprahamian, W. R. Dichtel, T. Ikeda, J. R. Heath, J. F. Stoddart, *Org. Lett.* **2007**, *9*, 1287–1290; h) F. Durola, J.-P. Sauvage, *Angew. Chem.* **2007**, *119*, 3607–3610; *Angew. Chem. Int. Ed.* **2007**, *46*, 3537–3540; i) G. Fioravanti, N. Haraszkiewicz, E. R. Kay, S. M. Mendoza, C. Bruno, M. Marcaccio, P. G. Wiering, F. Paolucci, P. Rudolf, A. M. Brouwer, D. A. Leigh, *J. Am. Chem. Soc.* **2008**, *130*, 2593–2601; j) Y.-L. Zhao, I. Aprahamian, A. Trabolsi, N. Erina, J. F. Stoddart, *J. Am. Chem. Soc.* **2008**, *130*, 6348–6350.
- [7] For the exploitation of second sphere coordination of metal-halide and thiocyanate bonds in rotaxane and catenane assembly, see: a) B. A. Blight, K. A. Van Noortwyk, J. A. Wisner, M. C. Jennings, *Angew. Chem.* **2005**, *117*, 1523–1528; *Angew. Chem. Int. Ed.* **2005**, *44*, 1499–1504; b) B. A. Blight, J. A. Wisner, M. C. Jennings, *Chem. Commun.* **2006**, 4593–4595; c) B. A. Blight, J. A. Wisner, M. C. Jennings, *Angew.*

*Chem.* **2007**, *119*, 2893–2896; *Angew. Chem. Int. Ed.* **2007**, *46*, 2835–2838; d) B. A. Blight, X. Wei, J. A. Wisner, M. C. Jennings, *Inorg. Chem.* **2007**, *46*, 8445–8447.

- [8] In a two step process, carbon monoxide and chloride coordination to rhodium has been used to change the shape and charge of a ‘weak-link approach’ macrocycle to promote subsequent pseudorotaxane formation with viologens, see: J. Kuwabara, C. L. Stern, C. A. Mirkin, *J. Am. Chem. Soc.* **2007**, *129*, 10074–10075.
- [9] The X-ray crystal data and experimental details of the structural refinement for {[**(L1)**PdCl].**1**} are provided in the Supporting Information. CCDC-689628 contains the supplementary data for this paper. This data can be obtained free of charge from the Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).
- [10] For other examples of rotaxane formation aided by ion-pairing, see: a) D. J. Hoffart, J. Tiburcio, A. de la Torre, L. K. Knight, S. J. Loeb, *Angew. Chem.* **2008**, *120*, 103–107; *Angew. Chem. Int. Ed.* **2008**, *47*, 97–101; b) E. Lestini, K. Nikitin, H. Müller-Bunz, D. Fitzmaurice, *Chem. Eur. J.* **2008**, *14*, 1095–1106.
- [11] C. W. Tornøe, C. Christensen, M. Meldal, *J. Org. Chem.* **2002**, *67*, 3057–3064; b) V. V. Rostovtsev, L. G. Green, V. V. Fokin, K. B. Sharpless, *Angew. Chem.* **2002**, *114*, 2708–2711; *Angew. Chem. Int. Ed.* **2002**, *41*, 2596–2599.
- [12] T. R. Chan, R. Hilgraf, K. B. Sharpless, V. V. Fokin, *Org. Lett.* **2004**, *6*, 2853–2855.
- [13] The prefix indicates the position of the macrocycle on the thread: *pyrdm*- non-covalent bonding to the pyridinium group; *triazole*- coordinated through the palladium ion to the triazole ring.
- [14] V. Aucagne, J. Berná, J. D. Crowley, S. M. Goldup, K. D. Hänni, D. A. Leigh, P. J. Lusby, V. E. Ronaldson, A. M. Z. Slawin, A. Viterisi, D. B. Walker, *J. Am. Chem. Soc.* **2007**, *129*, 11950–11963.