

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/26769878>

Synthesis, structure, and antimycobacterial activity of 6-[1(3H)-isobenzofuranylidene-methyl]purines and analogs. Bioorg Med Chem

ARTICLE in BIOORGANIC & MEDICINAL CHEMISTRY · SEPTEMBER 2009

Impact Factor: 2.79 · DOI: 10.1016/j.bmc.2009.08.012 · Source: PubMed

CITATIONS

16

READS

24

3 AUTHORS, INCLUDING:



Vebjørn Bakken

University of Oslo

30 PUBLICATIONS 2,138 CITATIONS

SEE PROFILE



Synthesis, structure, and antimycobacterial activity of 6-[1(3H)-isobenzofuranylidene-methyl]purines and analogs

Morten Brændvang, Vebjørn Bakken, Lise-Lotte Gundersen *

Department of Chemistry, University of Oslo, PO Box 1033, Blindern, N-0315 Oslo, Norway

ARTICLE INFO

Article history:

Received 7 April 2009

Revised 4 August 2009

Accepted 6 August 2009

Available online 12 August 2009

Keywords:

Purines

Cross-couplings

Antimycobacterials

X-ray

Ab initio calculations

ABSTRACT

6-Benzofuryl-, styryl, benzyl, and furfuryl purines as well as 6-[1(3H)-isobenzofuranylidene-methyl]purines have been synthesized and their activities against *Mycobacterium tuberculosis* (*Mtb*) determined. Several compounds displayed profound antimycobacterial activity in combination with low toxicity towards mammalian cells. NMR and X-ray crystallography were employed to determine the detailed structures and the results were supported by quantum chemical calculations.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

Tuberculosis (TB) still claims ca. two-million deaths per year world-wide and resistance to existing drugs is a growing problem.¹ We have previously reported selective antibacterial activity against *Mycobacterium tuberculosis* (*Mtb*) for certain 9-benzylpurines.² Fig. 1 shows examples of purines with profound antimycobacterial activity as well as a summary of SAR knowledge. Previous results have led us to believe that an aryl group in the purine 6-position is a requirement for significant antimycobacterial effect,^{2a} and among the 6-aryl- and 6-heteroaryl purines examined, especially high activities are found for 6-(2-furyl)purines,^{2e} for instance compounds **1a–1d** (Fig. 1). The more bulky 6-(benzofur-2-yl)purine **2a** is less active than the corresponding furyl purine **1a**.^{2d} However, we recently found MIC values for antimycobacterial activity of 6-[1(3H)-isobenzofuranylidene-methyl]purines **3a** and **4a** only one titer step higher than 6-furyl purine **1a**.³ Since we have shown that compound **4a** easily isomerizes into the *Z*-isomer **3a**,³ it may be that compound **4a** also isomerizes into compound **3a** in the antimycobacterial assay.

The profound, and somewhat unexpected, effect on *Mtb* growth found for the *Z*-6-[1(3H)-isobenzofuranylidene-methyl]purine **3a**, prompted us to synthesize and determine antimycobacterial activity for the related compound **3b** (Fig. 1) where the substitution pattern elsewhere in the molecule is optimized according to our current SAR knowledge, as well as the optimized 6-(benzofur-2-

yl)purine **2b** (Fig. 1) and the styryl- **5**, benzyl- **6**, and furfuryl purine **7**. The structural relationship between compounds **5–7** and the (isobenzofuranylidene-methyl)purine **3b**, is illustrated in Figure 2. It was previously shown that compound **3a** exists only as a (isobenzofuranylidene-methyl)purine, and not as an aromatic, but quinoid, benzo[c]furylmethyl purine.³ On the other hand, the simple furyl derivative **7**, may exist as tautomer **7b** rather than **7a**. Hence, we included a structural study of the synthesized compounds employing NMR, X-ray, and quantum chemical calculations in order to reveal their detailed structures and broaden our SAR knowledge.

2. Synthesis

Z-6-[1(3H)-isobenzofuranylidene-methyl]purine **3b** was synthesized essentially as reported for the preparation of compound **3a** before.³ When the dichloropurine **8** was reacted with the alkyne **9** under Sonogashira coupling conditions, cyclization took place to give mainly the *E*-isomer **4b**. The mixture of isomers was treated with TFA to give the pure *Z*-6-[1(3H)-isobenzofuranylidene-methyl]purine **3b** (Scheme 1).

The other target molecules, **2b** and **5–7**, were available by regioselective Stille- or Negishi-coupling on the dichloropurine **8** (Scheme 2, Table 1) a synthetic strategy employed by us in the synthesis of several 6-substituted purines.^{1,4} Furfuryl halides are regarded as unstable and difficult to utilize for synthetic purposes.⁵ However, we found that the desired furfurylzinc chloride could be generated from furfuryl chloride and Rieke zinc following the protocol previously used for generation of benzylzinc chloride.⁶

* Corresponding author. Tel.: +47 22857019; fax: +47 22855507.
E-mail address: l.l.gundersen@kjemi.uio.no (L.-L. Gundersen).




Figure 1. Structures of antimycobacterial purines and previously reported MIC values,^{2a,d,e,3} summary of SAR for antimycobacterial 9-benzylpurines, and target molecules in the study.

This constitutes the first formation and synthetic application of a furfurylmethyl reagent carrying no stabilizing groups on the furyl ring.

3. Structure

In line with our previous observations on compound **3a**,³ no quinoid benzo[*c*]furylmethylpurine tautomer could be detected in the ¹H NMR spectrum of the Z-6-[1(3*H*)-isobenzofuranylidene-methyl]purine **3b** or the isomer **4b**. On the other hand, ¹H NMR revealed that compound **7** was present only as the furfuryl-purine **7b**. Electronic structure calculations supported these findings (see [Supplementary data](#)). The furfuryl-purine **7** also crystallized as the **7b** tautomer ([Fig. 3](#)). The benzyl group in crystalline compound **7b** is oriented quite different from what is previously found by X-ray crystallography for the benzylic substituent in the 6-furyl-purine **1d**⁷ and related structures,⁸ for further discussions, see [Supplementary data](#).

4. Antimycobacterial activity

The novel purine derivatives **2b**, **3b**, **5**, **6**, and **7b** were screened for antibacterial activity against *M. tuberculosis* H₃₇Rv in vitro and the IC₉₀ and IC₅₀ values are presented in [Table 2](#) together with comparable data for the compounds **1d** and **3a** synthesized earlier. Previously determined MIC values (*Mtb*) for compounds **1a–1d**, **2a**, **3a**, and **4a** are displayed in [Figure 1](#).

Both the 6-furyl-**1d** and 6-benzofuryl-purine **2b** were found to be highly potent inhibitors of *Mtb* (IC₉₀ <0.20 μg/mL, <0.6 μM). As expected, both (isobenzofuranylidene-methyl)purines **3a** and **3b** also displayed considerable growth inhibition (IC₉₀ 3–5 μg/mL, 8–12 μM). However, the furfuryl-purine **7b** was essentially inactive (IC₉₀ >100 μg/mL) whereas the benzyl-**6** and styryl-purine **5** were only slightly weaker inhibitors (IC₉₀ 6–11 μg/mL, 16–30 μM) compared to the (isobenzofuranylidene-methyl)purines **3**. None of the novel compounds described herein could compete with 6-furyl-**1d** or 6-benzofuryl-purine **2b** where the aryl substituent is connected






Figure 2. Target molecules **5–7** and their structural relationship with target molecule **3b**.



Scheme 1. Reagents and conditions: (a) $(\text{Ph}_3\text{P})_2\text{PdCl}_2$, CuI, $(i\text{-Pr})_2\text{NH}$, DMF, 60 °C; (b) TFA, CH_2Cl_2 .



Scheme 2. Reagents and conditions: (a) R-Met, Pd-cat, 50 °C, for more details, see Table 1.

Table 1
Reaction conditions applied in synthesis of compounds 5–7

–Met	Pd-catalyst	Solvent	Time (h)	Yield (%)
–SnBu ₃	$[(2\text{-Furyl})_3\text{P}]_4\text{Pd}$	DMF	8	78, 2b
–SnBu ₃	$[(2\text{-Furyl})_3\text{P}]_4\text{Pd}$	DMF	7	77, 5
–ZnCl	$(\text{Ph}_3\text{P})_4\text{Pd}$	THF	5	69, 6
–ZnCl	$(\text{Ph}_3\text{P})_4\text{Pd}$	THF	3	59, 7

directly to C-6 in the purine, with respect to inhibition of *Mtb*. However, and in contrast to our previous believe (Fig. 1),² also other substituents (i.e., styryl- or isobenzofuranylidene-methyl-) in the 6-position may result in purines with a considerable antimycobacterial activity. The compounds displaying IC_{90} values against *Mtb* lower than 10 $\mu\text{g}/\text{mL}$ were also screened for toxicity towards mammalian cells (VERO cells, Table 2). All compounds examined, except **3a**, showed IC_{50} against VERO cells >40 $\mu\text{g}/\text{mL}$. We have previously reported that compound **1c** (Fig. 1) showed virtually no cross resistance against a panel of drug-resistant *Mtb* strains.^{2b} We assume that the closely related structures reported herein act by the same, currently unknown, mechanism of action as purine **1c**, so we have reasons to believe that cross resistance will not be an issue for the novel compounds described herein.

5. Experimental

The ^1H NMR spectra were recorded at 300 MHz with a Bruker Avance DPX 300 instrument, and the ^1H decoupled ^{13}C NMR spec-

tra were recorded 75 MHz using the same instrument. Mass spectra under electron impact conditions (EI) were recorded at 70 eV ionizing voltage with a VG Prospec instrument, and are presented as m/z (% rel. int.). Elemental analyses were performed by Ilse Beetz Mikroanalytisches Laboratorium, Kronach, Germany. Melting points were determined with a C. Reichert melting point apparatus or a Büchi Melting Point B-545 apparatus and are uncorrected. DMF was distilled from BaO and stored over 4 Å mol. sieve, dichloromethane was distilled from CaH_2 , and THF from Na/benzophenone. Antimycobacterial activity was determined as previously reported.² The following compounds were prepared according to literature procedures: 2-chloro-6-(2-furyl)-9-(4-methoxyphenylmethyl)-9H-purine **1d**,^{2d} (Z)-9-benzyl-6-[isobenzofuran-1(3H)-ylidenemethyl]-9H-purine **3a**,² 2,6-dichloro-9-(4-methoxyphenylmethyl)-9H-purine **8**,^{2d} (E)-tributyl(phenylethenyl)stannane,⁹ tributyl(benzo[b]fur-2-yl)stannane,¹⁰ and furfuryl chloride.¹¹ Benzylzinc chloride was generated from benzyl chloride and Rieke zinc¹² according to a published procedure⁶ and the concentration of the solution was determined by hydrolysis and iodolysis.¹³ Furfurylzinc chloride was generated from furfuryl chloride following the same protocol.

5.1. X-ray crystallographic analysis for compound 7b

Crystals of **7b** suitable for X-ray crystallography were obtained from benzene solution at +4 °C. X-ray data were collected on a Siemens SMART CCD diffractometer¹⁴ using graphite monochromated Mo $\text{K}\alpha$ radiation ($\lambda = 0.71073 \text{ \AA}$). Data collection method: ω -scan, step 0.3°, crystal to detector distance 5 cm. Data reduction and cell determination were carried out with the SAINT and XPREP programs. Absorption corrections were applied by the use of the SADABS program.¹⁵ The structure was determined and refined using the SHELX program package.¹⁶ The non-hydrogen atoms were refined with isotropic thermal parameters; H atoms were positioned geometrically and allowed to ride and rotate (for the CH_3 group) on their carrier atoms, with C–H bond lengths of 0.95 (aromatic C–H), 0.99 (CH_2) or 0.98 Å (CH_3) and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for CH_2 and aromatic C–H or $1.5U_{\text{eq}}(\text{C})$ for CH_3 . Crystal structure data for **7b** are available from the Cambridge Crystallographic Data Center, CCDC no. 643827.




Figure 3. X-ray structure of compound **7b**, the gas-phase equilibrium structure of compound **7b**, and X-ray structure of compound **1d**.⁷

Table 2

Activity against *M. tuberculosis* for purines **1d**, **2b**, **3a**, **3b**, **5**, **6**, and **7b** as well as cytotoxicity against VERO cells for selected compounds^a

Compound	IC ₉₀ <i>M. tuberculosis</i> H ₃₇ Rv (μg/mL), μM values in brackets ^b	IC ₅₀ <i>M. tuberculosis</i> H ₃₇ Rv (μg/mL), μM values in brackets ^b	IC ₅₀ VERO cells (μg/mL) ^c
1d	<0.20 (<0.59)	<0.20 (<0.59)	>40
2b	<0.20 (<0.51)	<0.20 (<0.51)	>40
3a	2.7 (7.9)	1.5 (4.3)	17
3b	4.9 (12)	<0.20 (<0.48)	>40
5	6.1 (16)	3.0 (8.0)	>40
6	11 (30)	7.4 (20)	n.d.
7b	>100 (>280)	>100 (>280)	n.d.

^a Structures of compounds are shown in Schemes 1 and 2.

^b IC₉₀ ampicillin 0.13 μg/mL (0.22 μM) and IC₅₀ ampicillin 0.07 μg/mL (0.12 μM).

^c EC₅₀ hyamine 0.01 μg/mL.

5.2. Crystal data for C₁₈H₁₅ClN₄O₂ **7b**

$M = 354.79$, monoclinic, $P2(1)/n$ $a = 4.5528(6)$ Å, $b = 18.867(3)$ Å, $c = 18.955(3)$ Å, $\beta = 96.486(2)^\circ$, $V = 1617.8(4)$ Å³, $Z = 4$, $D_x = 1.457$ Mg m⁻³, $\mu = 0.26$ mm⁻¹, $T = 105(2)$ K, measured 16,774 reflections in 2θ range 6.2–60.6°, $R_{\text{int}} = 0.032$. To 128 parameters refined against 4518 F^2 , $R = 0.0333$ for 3293 $I_o > 2\sigma(I_o)$ and 0.0553 for all data.

5.3. Antimycobacterial data

The purines were screened for antimycobacterial activities essentially as described before.^{2,17} Compounds were tested in 10 twofold dilutions, from 100 to 0.19 μg/mL, against *M. tuberculosis* H₃₇Rv (ATCC 27294) in BACTEC 12B medium using the Microplate Alamar Blue Assay (MABA). The IC₉₀ and IC₅₀ values are determined from the dose–response curve as the IC₉₀ using the curve fitting program XLFIT, formula 205.

5.4. Activity against VERO cells

The compounds were screened for mammalian cell cytotoxicity to VERO cells essentially as described before;² after 72 h exposure, viability is assessed using the CellTiter 96® Non-Radioactive Cell Proliferation Assay (MTT) reagent from Promega. Cytotoxicity is determined from the dose–response curve as the EC₅₀ using the curve fitting program XLFIT, formula 205.

5.5. 6-(Benzofuran-2-yl)-2-chloro-9-(4-methoxyphenylmethyl)-9H-purine (**2b**)

A mixture of tris(dibenzylideneacetone)dipalladium, Pd₂dba₃ (28 mg, 0.030 mmol), and tri(2-furyl)phosphine (51 mg, 0.22 mmol) in dry DMF (4 mL) was stirred at ambient temperature for 5 min, before a solution of 2,6-dichloro-9-(4-methoxyphenylmethyl)-9H-purine **8** (309 mg, 1.0 mmol) in DMF (4 mL) was added. After an additional 5 min, tributyl(benzo[b]fur-2-yl)stannane (575 mg, ca. 1.20 mmol, ca. 85% pure) was introduced, the resulting mixture stirred at 50 °C for 8 h and evaporated in vacuo. A satd solution of potassium fluoride in methanol (40 mL) was added to the residue, the mixture was stirred over night and evaporated in vacuo together with a small amount of silica gel. The residue was added on top of a silica gel column, and the product purified by flash chromatography eluting with CH₂Cl₂–acetone (29:1), followed by CH₂Cl₂–acetone (19:1); yield 303 mg (78%), mp 189 °C, colorless crystals. ¹H NMR (300 MHz, CDCl₃) δ = 3.75 (s, 3H, OCH₃), 5.32 (s, 2H, NCH₂), 6.87 (d, J = 8.7 Hz, 2H, Ar), 7.24–7.30 (m, 3H, Ar and benzofuryl), 7.38–7.43 (m, 1H, benzofuryl), 7.67–7.70 (m, 2H, benzofuryl), 8.02 (s, 1H, 8-H) and 8.28 (s, 1H, 3-H in benzofuryl) ppm. ¹³C NMR (75 MHz, CDCl₃) δ = 47.1 (NCH₂), 55.3 (OCH₃), 112.4 (CH in benzofuryl), 114.6 (CH in Ar), 115.2 (CH in benzofuryl), 122.4

(CH in benzofuryl), 123.6 (CH in benzofuryl), 126.4 (C-1 in Ar), 127.2 (CH in benzofuryl), 128.1 (C in benzofuryl), 128.4 (C-5), 129.6 (CH in Ar), 145.1 (C-8), 147.6 (C-6/C-2), 149.7 (C-2 in benzofuryl), 153.7 (C-4), 154.5 (C-2/C-6), 156.0 (C in benzofuryl) and 159.9 (C-4 in Ar) ppm. MS (EI): m/z (%): 392/390 (15/42) [M]⁺, 234 (1), 195 (1), 181 (2), 122 (9), 121 (100). HRMS (EI): calcd for C₂₁H₁₅ClN₄O₂ 390.0884, found 390.0886. C₂₁H₁₅ClN₄O₂ (390.8): C, 64.54; H, 3.87; N, 14.34. Found: C, 64.50; H, 3.93; N, 14.20.

5.6. 2-Chloro-6-[(Z)-1(3H)-isobenzofuranylidene-methyl]-9-(methoxyphenylmethyl)-9H-purine (**3b**)

2,6-Dichloro-9-(4-methoxyphenylmethyl)-9H-purine **8** (155 mg, 0.50 mmol) was added to a stirring solution of CuI (9.5 mg, 0.050 mmol), bis(triphenylphosphine)palladium(II) chloride (18 mg, 0.025 mmol), and dry diisopropylamine (420 μL, 3.00 mmol) in dry DMF (2.5 mL) under N₂. The solution was heated to 60 °C before 2-ethynylbenzenemethanol **9** (79 mg, 0.60 mmol) dissolved in DMF (1 mL) was added dropwise over 1 h. After stirring for additional 4 h at 60 °C, the reaction mixture evaporated in vacuo with a small amount of silica. The residue was added on top of a silica gel column, and the product, as a ca. 7:3 *E/Z* mixture was isolated by flash chromatography eluting with CH₂Cl₂–acetone (19:1) followed by CH₂Cl₂–acetone (9:1). The isomeric mixture was dissolved in CH₂Cl₂ (30 mL) and trifluoroacetic acid (74 μL, 0.96 mmol) was added. After stirring for 1 h at ambient temperature the reaction mixture was washed with satd aq NaHCO₃ (2 × 20 mL), water (20 mL), brine (20 mL), dried (MgSO₄) and evaporated in vacuo. The product was purified by flash chromatography on silica gel eluting with CH₂Cl₂–acetone (9:1); yield 101 mg (50%), pale yellow crystals. ¹H NMR (300 MHz, CDCl₃) δ = 3.76 (s, 3H, OCH₃), 5.28 (s, 2H, NCH₂), 5.71 (s, 2H, OCH₂), 6.72 (s, 1H, =CH), 6.87 (d, J = 8.7 Hz, 2 H, Ar), 7.25 (d, J = 8.7 Hz, 2H, Ar), 7.40–7.48 (m, 3H, Ar'), 7.79–7.82 (m, 1H, Ar') and 7.84 (s, 1H, 8-H) ppm. ¹³C NMR (75 MHz, CDCl₃) δ = 46.8 (NCH₂), 55.3 (OCH₃), 77.2 (OCH₂), 88.6 (=CH), 114.5 (CH in Ar), 121.3 (CH in Ar'), 121.8 (CH in Ar'), 126.9 (C-1 in Ar), 128.5 (CH in Ar'), 129.0 (C-5), 129.6 (CH in Ar), 131.1 (CH in Ar'), 133.9 (C in Ar'), 141.5 (C in Ar'), 142.7 (C-8), 152.1 (C-4), 154.6 (C-2/C-6), 156.2 (C-6/C-2), 159.8 (C-4 in Ar) and 165.7 (=C) ppm. MS (EI): m/z (%): 406/404 (9/25) [M]⁺, 285 (3), 284 (2), 283 (8), 140 (2), 121 (100). HRMS (EI): calcd for C₂₂H₁₇ClN₄O₂ 404.1040, found 404.1030. C₂₂H₁₇ClN₄O₂ (404.9): C, 65.27; H, 4.23; N, 13.84. Found: C, 64.94; H, 4.16; N, 13.81.

5.7. (E)-2-Chloro-9-(4-methoxybenzyl)-6-(2-phenylethenyl)-9H-purine (**5**)

The product was formed by Stille coupling between 2,6-dichloro-9-(4-methoxyphenylmethyl)-9H-purine **8** (309 mg, 1.0 mmol) and (E)-tributyl(phenylethenyl)stannane (1.12 g, 1.20 mmol, ca. 42% purity) as described for compound **3b** above. The reaction time was 7 h, and the product purified by flash chromatography on silica gel eluting with CH₂Cl₂–acetone (29:1), followed by CH₂Cl₂–acetone (19:1) and finally CH₂Cl₂–acetone (9:1); yield 292 mg (77%), colorless foam. ¹H NMR (300 MHz, CDCl₃) δ = 3.75 (s, 3H, OCH₃), 5.30 (s, 2H, NCH₂), 6.88 (d, J = 8.8 Hz, 2H, Ar), 7.26 (d, J = 8.8 Hz, 2H, Ar), 7.32–7.42 (m, 3H, Ph), 7.59 (d, J = 16.1 Hz, 1H, CH=), 7.66–7.69 (m, 2H, Ph), 7.94 (s, 1H, 8-H) and 8.42 (d, J = 16.1 Hz, 1H, CH=) ppm. ¹³C NMR (75 MHz, CDCl₃) δ = 47.0 (NCH₂), 55.3 (OCH₃), 114.5 (CH in Ar), 121.4 (CH=), 126.6 (C-1 in Ar), 128.1 (CH in Ph), 128.8 (CH in Ph), 129.6 (CH in Ar), 129.8 (CH in Ph), 130.0 (C-5), 135.7 (C-1 in Ph), 141.7 (CH=), 144.3 (C-8), 153.4 (C-4), 154.3 (C-2), 155.8 (C-6) and 159.9 (C-4 in Ar) ppm. MS (EI): m/z (%): 378/376 (13/36) [M]⁺, 257 (1), 255 (4), 122 (9), 121 (100). HRMS (EI): calcd for C₂₁H₁₇ClN₄O 376.1091, found 376.1093. C₂₁H₁₇ClN₄O (376.8): C, 66.93; H, 4.55; N, 14.87. Found: C, 66.76; H, 4.46; N, 14.71.

5.8. 6-Benzyl-2-chloro-9-(4-methoxyphenylmethyl)-9H-purine (6)

2,6-Dichloro-9-(4-methoxyphenylmethyl)-9H-purine **8** (309 mg, 1.0 mmol) was added to a solution of tetrakis(triphenylphosphine)palladium(0) [generated in situ from tris(diphenylmethylideneacetone)dipalladium chloroform adduct (23 mg, 0.025 mmol) and triphenylphosphine (53 mg, 0.20 mmol)] in dry THF (4 mL) at ambient temperature under N₂. After 10 min, a solution of benzylzinc chloride (1.83 mL, 1.52 mmol, 0.83 M) in THF was added and the mixture stirred at 50 °C for 5 h. Satd aq NH₄Cl (10 mL) was added and the resulting mixture was extracted with EtOAc (4 × 25 mL). The combined organic extracts were washed with brine (2 × 20 mL), dried (MgSO₄), and evaporated in vacuo with a small amount on silica. The residue was added on top of a silica gel column, and the product purified by flash chromatography eluting with CH₂Cl₂–acetone (29:1), followed by CH₂Cl₂–acetone (19:1) and finally CH₂Cl₂–acetone (9:1); yield 250 mg (69%), colorless oil. ¹H NMR (300 MHz, CDCl₃) δ = 3.74 (s, 3H, OCH₃), 4.43 (CH₂), 5.25 (s, 2H, NCH₂), 6.85 (d, *J* = 8.7 Hz, 2H, Ar), 7.11–7.27 (m, 2H, Ar and 3H, Ph), 7.45–7.48 (m, 2H, Ph) and 7.92 (s, 1H, 8-H) ppm. ¹³C NMR (75 MHz, CDCl₃) δ = 39.5 (CH₂), 47.0 (NCH₂), 55.3 (OCH₃), 114.5 (CH in Ar), 126.4 (C-1 in Ar), 126.7 (CH in Ph), 128.5 (CH in Ph), 129.3 (CH in Ph), 129.6 (CH in Ar), 131.7 (C-5), 137.0 (C-1 in Ph), 144.4 (C-8), 152.8 (C-4), 154.1 (C-2), 159.9 (C-4 in Ar) and 162.8 (C-6) ppm. MS (EI): *m/z* (%): 366/364 (24/60) [M]⁺, 245 (9), 243 (27), 207 (2), 180 (3), 153 (2), 121 (100). HRMS (EI): calcd for C₂₀H₁₇ClN₄O 364.1091, found 364.1090. C₂₀H₁₇ClN₄O (363.8): C, 65.84; H, 4.70; N, 15.36. Found: C, 65.83; H, 4.84; N, 15.26.

5.9. 2-Chloro-6-(furan-2-ylmethyl)-9-(4-methoxyphenylmethyl)-9H-purine (7b)

The product was formed by Negishi coupling between 2,6-dichloro-9-(4-methoxyphenylmethyl)-9H-purine **8** (309 mg, 1.0 mmol) and furfurylzinc chloride (3.9 mL, 1.60 mmol, 0.41 M) as described for compound **6** above. The reaction time was 3 h, and the product purified by flash chromatography on silica gel eluting with CH₂Cl₂–acetone (29:1), followed by CH₂Cl₂–acetone (19:1) and finally CH₂Cl₂–acetone (9:1); yield 208 mg (59%), mp 135 °C, colorless crystals. ¹H NMR (300 MHz, CDCl₃) δ = 3.74 (s, 3H, OCH₃), 4.46 (s, 2H, CH₂), 5.27 (s, 2H, NCH₂), 6.22 (dd, *J* = 3.2 and 0.7 Hz, 1H, 3-H in furyl), 6.25 (dd, *J* = 3.2 and 1.9 Hz, 1H, 4-H in furyl), 6.85 (d, *J* = 8.7 Hz, 2H, Ar), 7.25 (d, *J* = 8.7 Hz, 2H, Ar), 7.28 (dd, *J* = 1.8 and 0.8 Hz, 1H, 5-H in furyl) and 7.94 (s, 1H, 8-H) ppm. ¹³C NMR (75 MHz, CDCl₃) δ = 32.2 (CH₂), 47.0 (NCH₂), 55.3 (OCH₃), 107.6 (C-3 in furyl), 110.5 (C-4 in furyl), 114.5 (CH in Ar), 126.4 (C-1 in Ar), 129.6 (CH in Ar), 131.7 (C-5), 141.9 (C-5 in furyl), 144.7 (C-8), 149.9 (C-2 in furyl), 152.9 (C-4), 154.1 (C-2), 159.8 (C-6) and 159.9 (C-4 in Ar) ppm. MS (EI): *m/z* (%): 356/354 (15/43) [M]⁺, 121 (100), 91 (3), 78 (5), 77 (5). HRMS (EI): calcd for

C₁₈H₁₅ClN₄O₂ 354.0884, found 354.0873. C₁₈H₁₅ClN₄O₂ (354.8): C, 60.94; H, 4.26; N, 15.79. Found: C, 60.92; H, 4.28; N, 15.71.

Acknowledgments

Antimycobacterial data were provided by the Tuberculosis Antimicrobial Acquisition and Coordinating Facility (TAACF) through a research and development contract with the US National Institute of Allergy and Infectious Diseases. We are grateful for all help provided by Dr. Cecil Kwong and co-workers. This work has received support through a grant of computer time from the Research Council of Norway (Grant No. NN1118K). NFR is also greatly acknowledged for partial financing of the Bruker Avance instrument used in this study. We thank Dr. Osamu Sekiguchi, Department of Chemistry, University of Oslo for the single-crystal data collection and initial refinement and Tom Chr. Berg for the synthesis of compound **3a**.

Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.bmc.2009.08.012.

References and notes

- See for instance: Bhowruth, V.; Dover, L. G.; Besra, G. S. *Prog. Med. Chem.* **2007**, 45, 169, and references therein.
- (a) Bakkestuen, A. K.; Gundersen, L.-L.; Langli, G.; Liu, F.; Nolsøe, J. M. N. *Bioorg. Med. Chem. Lett.* **2000**, 10, 1207; (b) Gundersen, L.-L.; Nissen-Meyer, J.; Spilsberg, B. J. *Med. Chem.* **2002**, 45, 1383; (c) Andresen, G.; Gundersen, L.-L.; Nissen-Meyer, J.; Rise, F.; Spilsberg, B. *Bioorg. Med. Chem. Lett.* **2002**, 12, 567; (d) Bakkestuen, A. K.; Gundersen, L.-L.; Utenova, B. T. J. *Med. Chem.* **2005**, 48, 2710; (e) Brændvang, M.; Gundersen, L.-L. *Bioorg. Med. Chem.* **2005**, 13, 6360; (f) Brændvang, M.; Gundersen, L.-L. *Bioorg. Med. Chem.* **2007**, 15, 7144.
- Berg, T. C.; Bakken, V.; Gundersen, L.-L.; Petersen, D. *Tetrahedron* **2006**, 62, 6121.
- Langli, G.; Gundersen, L.-L.; Rise, F. *Tetrahedron* **1996**, 52, 5625.
- Takanishi, K.; Hirokazu, U.; Kuwajima, I. *Tetrahedron Lett.* **1987**, 28, 2281.
- Zhu, L.; Wehmeyer, R. M.; Rieke, R. D. J. *Org. Chem.* **1991**, 56, 1445.
- (a) Brændvang, M.; Gundersen, L.-L. *Acta. Crystallogr., Sect. C* **2007**, 63, o274–o276; (b) Brændvang, M.; Gundersen, L.-L. *Acta. Crystallogr., Sect. E* **2007**, 63, o3036.
- Mazumdar, P. A.; Das, A. K.; Bakkestuen, A. K.; Gundersen, L.-L.; Bertolasi, V. *Acta. Crystallogr., Sect. E* **2001**, 57, o1052.
- Labadie, J. W.; Tueting, D.; Stille, J. K. J. *Org. Chem.* **1983**, 48, 4634.
- Liebeskind, L. S.; Wang, J. J. *Org. Chem.* **1993**, 58, 3550.
- Chaudhari, S. S.; Akamanchi, K. G. *Synlett* **1999**, 1763.
- (a) Chen, T.-A.; Wu, X.; Rieke, R. D. J. *Am. Chem. Soc.* **1995**, 117, 233–244; (b) Rieke, R. D.; Hanson, M. V.; Brown, J. D.; Niu, Q. J. *Org. Chem.* **1996**, 61, 2726.
- Knochel, P.; Jones, P.; Langer, F. In *Organozinc Reagents: A Practical Approach*; Knochel, P., Jones, P., Eds.; Oxford University Press: Oxford, 1999; p 14.
- SMART (Version 5.625) and SAINT+ (Version 6.02). Area-Detector Control and Integration Software, Bruker Analytical X-ray Instruments, Madison, WI, 1998.
- Sheldrick, G. M. SADABS. Program for Empirical Correction of Area Detector Data, University of Göttingen, Germany, 1996.
- Sheldrick, G. M. SHELXL (release 97-2)—Programs for Crystal Structure Analysis Structure, University of Göttingen, Germany, 1997.
- Collins, L. A.; Franzblau, S. G. *Antimicrob. Agents Chemother.* **1997**, 41, 1004.