

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/9085819>

# 6-Amino-3-Pyridinols: Towards Diffusion-Controlled Chain-Breaking Antioxidants

ARTICLE *in* ANGEWANDTE CHEMIE INTERNATIONAL EDITION · OCTOBER 2003

Impact Factor: 11.26 · DOI: 10.1002/anie.200351881 · Source: PubMed

CITATIONS

94

READS

39

6 AUTHORS, INCLUDING:



**Maikel Wijtmans**

VU University Amsterdam

40 PUBLICATIONS 1,069 CITATIONS

SEE PROFILE



**Luca Valgimigli**

University of Bologna

112 PUBLICATIONS 3,347 CITATIONS

SEE PROFILE



**Gino DiLabio**

University of British Columbia - Okanagan

124 PUBLICATIONS 4,835 CITATIONS

SEE PROFILE

## **6-Amino-3-Pyridinols: Towards Diffusion-Controlled Chain-Breaking Antioxidants\*\***

*Maikel Wijtmans, Derek A. Pratt,\* Luca Valgimigli,\* Gino A. DiLabio, Gian Franco Pedulli and Ned A. Porter\**

[\*] D. A. Pratt, Prof. Dr. N. A. Porter, M. Wijtmans  
Department of Chemistry  
Vanderbilt University  
Nashville, Tennessee, USA 37235  
Fax: 1-615-343-5478  
E-mail derek.a.pratt@vanderbilt.edu, n.porter@vanderbilt.edu

Dr. L. Valgimigli, Prof. Dr. G. F. Pedulli  
Department of Organic Chemistry "A. Mangini"  
University of Bologna  
Via San Donato 15, I-40127 Bologna, Italy  
Fax: +39-051-244064  
E-mail: valgimig@alma.unibo.it


Dr. G. A. DiLabio  
National Institute for Nanotechnology  
Nation Research Council of Canada  
W6-010 ECERF, 9107 116th Street, Edmonton, AB, T6G 2V4

[\*\*] D.A.P. thanks NSERC Canada and Vanderbilt University for their support. This project was in part supported by the National Science Foundation as well as the University of Bologna and the MIUR Research project "Free Radical Research in Chemistry and Biology: Fundamental Aspects and Applications in Environment and Material Science".


Peroxyl radicals ( $\text{LOO}^\bullet$ ) are the chain-propagating species in the rate-determining step of lipid peroxidation (Eq. 1). Antioxidants, most commonly substituted phenols ( $\text{ArOH}$ ) such as **1-3**, effectively intercept peroxy radicals by transferring the phenolic H-atom to the propagating radical (Eq. 2), at a rate ( $k_{inh}[\text{ArOH}]$ ) faster than that of chain propagation ( $k_p[\text{L-H}]$ ). The best-known example of a phenolic antioxidant is  $\alpha$ -tocopherol ( $\alpha$ -TOH, **2**), the most potent form of Vitamin E, Nature's primary defense against radical chain oxidation.<sup>[1]</sup> Recently, considerable effort has been devoted to the development of antioxidants more effective than  $\alpha$ -TOH,<sup>[2],[3],[4],[5]</sup> since the *in-vivo* peroxidation of lipids has been suggested to be a major factor in the development of several degenerative diseases.<sup>[6]</sup>



The bond dissociation enthalpy (BDE) of the phenolic O-H plays a central role in determining antioxidant efficacy with compounds having lower O-H BDEs generally being better antioxidants.<sup>[7],[8]</sup> Substitution with electron donating (ED) groups at the *o*- and *p*- positions to the phenolic O-H leads to compounds having lower BDEs. However good ED substituents also lead to a decrease in the ionization potential (IP) of the phenol, thereby rendering the resulting antioxidant directly reactive with oxygen. This results in limited air-stability of the antioxidant and in the potential generation of toxic products in biological environments. Calculations predicted that incorporation of two nitrogen atoms at the 3 and 5 positions of the phenolic ring significantly raises the IP and greatly improves the air stability of the antioxidant while only minimally lowering the O-H BDE.<sup>[4]</sup> Experiments confirmed this prediction: thus, for example, the 5-pyrimidinol **4** is perfectly stable in air and has a  $k_{inh}$  about twice that of  $\alpha$ -TOH.<sup>[4]</sup>






Calculations also predicted that 3-pyridinol would have a lower O-H BDE than the analogous 5-pyrimidinol and an IP intermediate to those of phenol and 5-pyrimidinol.<sup>[4]</sup> We therefore expected that 6-amino-3-pyridinols such as **5** would have lower O-H BDEs than the analogous 5-pyrimidinols (e.g. **4**) while maintaining improved air-stability compared to the corresponding phenols. Furthermore, the free 5-position in the pyridine-skeleton allows for fusion of aliphatic rings (i.e. to give **6** and **7**); structural changes that have been shown to increase  $k_{inh}$  significantly in the series **1**, **2** and **3** because of improved stereoelectronics.<sup>[2]</sup> In this communication we show by theory the effects of substitution on the O-H BDE (antioxidant activity) and IP (air stability) of a series of 3-pyridinols. We also report here the synthesis and experimental investigation of three new 3-pyridinol antioxidants that were selected on the basis of our calculations.



The calculated O-H BDEs and IPs of several substituted phenols, 3-pyridinols and 5-pyrimidinols were obtained using density functional theory models ((RO)B3LYP/6-311+G(2d,2p)//AM1/AM1 and B3LYP/6-31G(d)//AM1/AM1, respectively) and the values are given in Table 1.<sup>[9],[10]</sup>

**Table 1.** Calculated Substituent Effects on Gas Phase O-H BDEs at 298K and Adiabatic IPs at 0K of Substituted Phenols, 3-Pyridinols and 5-Pyrimidinols.<sup>[a]</sup>


						
	BDE	IP	BDE	IP	BDE	IP
Substitution						
R <sup>1</sup> =H, R <sup>2</sup> =H	87.1 (0.0)	195.4 (0.0)	88.2 (0.0)	206.4 (0.0)	89.6 (0.0)	219.7 (0.0)
R <sup>1</sup> =H, R <sup>2</sup> =CH <sub>3</sub>	84.6 (-2.5)	186.9 (-8.5)	85.4 (-2.8)	196.6 (-9.8)	86.8 (-2.8)	209.3 (-10.4)
R <sup>1</sup> =CH <sub>3</sub> , R <sup>2</sup> =CH <sub>3</sub>	80.4 (-6.7)	178.3 (-17.1)	81.1 (-7.1)	186.3 (-20.1)	83.2 (-6.4)	198.0 (-22.7)
R <sup>1</sup> =H, R <sup>2</sup> =OCH <sub>3</sub>	81.0 (-6.1)	176.5 (-18.9)	82.0 (-6.2)	186.1 (-20.3)	83.6 (-6.0)	198.1 (-21.6)
R <sup>1</sup> =CH <sub>3</sub> , R <sup>2</sup> =OCH <sub>3</sub>	77.0 (-10.1)	169.2 (-26.2)	78.1 (-10.1)	177.4 (-29.0)	79.8 (-9.8)	188.3 (-31.4)
R <sup>1</sup> =H, R <sup>2</sup> =N(CH <sub>3</sub> ) <sub>2</sub>	77.0 (-10.1)	157.7 (-37.7)	77.0 (-11.2)	164.6 (-41.8)	78.3 (-11.3)	174.6 (-45.1)
Compounds						
<b>8</b> , <b>5</b> , <b>4</b> (monocyclic)	72.3 (-14.8)	152.3 (-43.1)	73.5 (-14.7)	157.7 (-48.7)	74.1 (-15.5)	167.0 (-52.7)
<b>9</b> , <b>6</b> (fused 6-ring)	71.2 (-15.9)	148.3 (-47.1)	73.3 (-14.9)	154.6 (-51.8)		
<b>10</b> , <b>7</b> (fused 5-ring)	70.5 (-16.6)	145.2 (-50.3)	72.4 (-15.8)	152.3 (-54.1)		
<b>2</b> (α-TOH)	74.8 (-12.3)	159.3 (-36.1)				

[a] Data for phenols and 5-pyrimidinols are from Ref. [4] except for **8**, **9** and **10**. All values are in kcal mol<sup>-1</sup>. Substituent effects (relative to the unsubstituted parent) are in parentheses.

Clear trends can be seen from Table 1. For all three classes, the O-H BDE and IP values decrease by increasing the electron density into the aromatic ring. The order for the calculated values of these properties is always phenol < 3-pyridinol < 5-pyrimidinol. The calculations suggest that the highest antioxidant activities in conjunction with reasonable air-stability (IP values comparable to **2**)

are expected from pyridinols **5-7**. Therefore, we selected these three molecules as targets for synthesis and experimental investigations.

**Scheme 1.** Synthetic Approach to 3-Pyridinols **5-7**.




Key: (a) dibromodimethylhydantoin (DBDMH),  $\text{CH}_2\text{Cl}_2$ ,  $-40^\circ\text{C}$ , 40 min, 67 %; (b)  $\text{HCOOH}$ , aq.  $\text{H}_2\text{CO}$ , reflux, 18 h, 100 % for **11**, 77 % for **12**, 84 % for **13**; (c) acrylic acid, pyridine, reflux, 24 h, 30 % after cryst.; (d) polyphosphoric acid,  $125^\circ\text{C}$ , 40 min, 75 %; (e)  $\text{BH}_3\cdot\text{THF}$ , THF, reflux, 18 h, 83 %; (f) DBDMH,  $\text{CH}_2\text{Cl}_2$ ,  $-78^\circ\text{C}$ , 10-30 min, 93 % for **12**, 83 % for **13**; (g)  $\text{Tf}_2\text{O}$ ,  $\text{Et}_3\text{N}$ ,  $\text{CH}_2\text{Cl}_2$ ,  $0^\circ\text{C}$ , 1 h, 98 %; (h) 15 M aq.  $\text{KOH}$ , reflux, 4 h, 95 %; (i)  $(\text{PhO})_2\text{PON}_3$ , *t*-BuOH, reflux, 30 h, 79 %; (j) ethereal  $\text{HCl}$ , 24 h, 70 % (k) **14**,  $\text{Et}_3\text{N}$ , DMF, 1 d, 93 %; (l)  $\text{Ac}_2\text{O}$ , DMAP,  $100^\circ\text{C}$ , 20 h, 91 %; (m)  $\text{Ph}_2\text{O}$ , reflux, 10 h, 82 %; (n)  $\text{NaOH}$ , MeOH, reflux, 20 h, 100 %; (o) 1. *n*-BuLi, THF,  $-78^\circ\text{C}$ , 30 min; 2. dry 2-nitro-*m*-xylene, THF,  $-78^\circ\text{C}$ , 1 to 3 h, 63 % for **5**, 25 % for **6**, 27 % for **7**.

There are few synthetic approaches to 6-amino-3-pyridinol structures described in the literature and **5-7** have never been reported.<sup>[11]</sup> We devised a synthetic sequence in which the reactive -OH moiety is introduced at low temperatures in the last step, thus minimizing potential decomposition of product

and/or intermediates. Only a brief description of the syntheses will be presented here; a forthcoming paper will report our extensive synthetic efforts in detail. Construction of the appropriate pyridine substructure involved a three-step Friedel-Crafts approach for **6** and a seven-step intramolecular Diels-Alder sequence for **7**. Construction of the pyridine was followed by bromination, methylation and hydroxylation (Scheme 1). The pyridinols **5-7** were obtained as yellow or orange solids.

Table 2 collects the O-H BDEs that were measured experimentally by radical equilibration EPR studies.<sup>[7b]</sup> The absolute experimental values (in benzene) were slightly higher (2 – 3 kcal mol<sup>-1</sup>) than the calculated (gas-phase) values (Table 1), an issue already addressed in previous works.<sup>[4]</sup> However, the corresponding  $\Delta\text{BDE}$  ( $\text{BDE}_{\text{PyrOH}} - \text{BDE}_{\alpha\text{-TOH}}$ ) shows good agreement between experimental and calculated substituent effects for the 3-pyridinol series. The stability of the 3-pyridinols towards air was examined by monitoring their typical fluorescence and UV absorbance (see Supporting Information, SI) in aerated *tert*-butylbenzene solution (0.3 mM) at 37 °C. The simple 3-pyridinol **5** was stable over a 24 h period while **7** showed significant decomposition over the same time course, the compound **6** showing stability intermediate of **5** and **7**, in parallel with the calculated IPs for these compounds (Table 1).

Two different experimental approaches were used to determine absolute  $k_{inh}$  for the 3-pyridinols. First, we utilized a peroxy radical clock based on the antioxidant-dependent trapping of bisallylic (11-OOH) and conjugated (9- and 13-OOH) hydroperoxide products in initiator-induced methyl linoleate autoxidations.<sup>[12]</sup> At low concentrations of antioxidant (10 – 100 mM), there is a linear correlation between concentration of antioxidant and ratio of trapping of 11-OO• vs 9- and 13-OO• with the slope of the line directly related to  $k_{inh}$  of the antioxidant.<sup>[12]</sup> Thus, by measuring the ratio of linoleate hydroperoxide products at various concentrations of an antioxidant it is possible to determine its  $k_{inh}$ .



3-Pyridinol **5** proved more effective than **4** or **2** in trapping 11-OO<sup>•</sup> and a value of  $k_{inh} = 1.16 (\pm 0.04) \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$  was determined by this method (see SI). The total yield of hydroperoxide products formed from linoleate in the presence of **5** was ca. eight-fold lower than that observed for identical oxidations inhibited by **2** (data not shown). Interestingly, **6** and **7** proved to be such good inhibitors that we were unable to obtain reliable  $k_{inh}$  values for these compounds using the radical clock method.<sup>[13]</sup> Controlled styrene autoxidations inhibited by **5** – **7** were well behaved, however,<sup>[2]</sup> and allowed us to investigate the antioxidant activity of these compounds (see SI). The stoichiometric factor  $n$  (number of chains broken by one molecule of antioxidant) was determined to be 2 for all three compounds, similar to phenols and pyrimidinols.<sup>[4]</sup> The inhibition rate constants  $k_{inh}$  were determined from the slope of the inhibited oxygen consumption plots,<sup>[2],[4],[14]</sup> and the values obtained are presented in Table 2 together with previous data for **2** and **4**, reported for comparison.<sup>[4],[12]</sup>

**Table 2.** Solution Phase O-H BDEs and Inhibition Rate Constants for 3-Pyridinols **5-7**.

Antioxidant	O-H BDE (kcal mol <sup>-1</sup> ) <sup>[a]</sup>	$\Delta$ (O-H BDE) (kcal mol <sup>-1</sup> ) <sup>[b]</sup>	$k_{inh}$ by clock ( $\times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ ) <sup>[c]</sup>	$k_{inh}$ by O <sub>2</sub> uptake ( $\times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ ) <sup>[d]</sup>	$k_{inh}/k_{inh}(\mathbf{2})$ <sup>[e]</sup>
<b>5</b>	77.0 $\pm$ 0.5	-1.3 / -1.3	1.16 $\pm$ 0.04	1.6 $\pm$ 0.6	5.0
<b>6</b>	76.3 $\pm$ 0.6	-2.0 / -1.5	N.D. <sup>[f]</sup>	8.8 $\pm$ 3.2	28
<b>7</b>	75.4 $\pm$ 0.7	-2.9 / -2.4	N.D. <sup>[f]</sup>	28.0 $\pm$ 18 <sup>[g]</sup>	88
<b>4</b>	78.2 $\pm$ 0.3	-0.1 / -0.7	0.65 $\pm$ 0.08	0.86 $\pm$ 0.05	2.1
<b>2</b> ( $\alpha$ -TOH)	78.3 $\pm$ 0.3	0.0 / 0.0	0.38 <sup>[h]</sup>	0.32 <sup>[h]</sup>	1.0

[a] From EPR equilibration studies. [b] Experimental BDE (78.3 kcal mol<sup>-1</sup>) or calculated BDE (74.8 kcal mol<sup>-1</sup>) for  $\alpha$ -TOH, **2**, as reference. Values are experimental/calculated  $\Delta$ BDE. [c] In benzene at 37



°C by methyl linoleate radical clock. [d] In chlorobenzene by inhibited styrene autoxidation at 30 °C for **2**, **5**, **6**, **7**; in benzene at 50 °C for **4**. [e]  $k_{inh}$  from O<sub>2</sub> uptake were used. [f] See text. [g] The rather large error (=2SD) is the result of the large scatter in the measurements due to the dramatic inhibition by **7**; initiation rates as low as  $1 \times 10^{-10} \text{ s}^{-1}$  had to be used in order to obtain measurable rates of oxygen consumption (as low as  $1 \times 10^{-9} \text{ M}^{-1} \text{ s}^{-1}$ ) with concentrations of **7** as low as  $2 \times 10^{-7} \text{ M}$ . [h] Value used to calibrate the radical clock. [h] Value from ref. [5a], re-confirmed in ref. [6]. The value of  $0.41 \pm 0.04$  at 50 °C is used for comparison with **4**.

The 6-amino-3-pyridinol described here are a novel class of phenolic antioxidants that are more effective than any other phenolic class reported to date. The vacant 5-position on the pyridine ring of these compounds allows for fusion of an additional aliphatic ring, a substitution that further lowers the O-H BDE. Indeed, the pyridinol **6** and **7** are, to the best of our knowledge, the fastest chain-breaking antioxidants ever reported (previously 3,7-dimethoxy-phenothiazine with  $k_{inh} = 5.5 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ ).<sup>[15],[16]</sup> The air-stability of **7** is moderate but its  $k_{inh}$  exceeds  $10^8 \text{ M}^{-1} \text{ s}^{-1}$ , thus approaching the diffusion-controlled limit for a bimolecular reaction.

**Keywords:** lipids · antioxidants · autoxidation · pyridines · ab initio calculations

---

[1] a) G. W. Burton, K. U. Ingold, *J. Am. Chem. Soc.* **1981**, *103*, 6472; b) V. R. Bowry, K. U. Ingold, *Acc. Chem. Res.* **1999**, *32*, 27.

[2] G. W. Burton, T. Doba, E. J. Gabe, L. Hughes, F. Lee, L. Prasad, K. U. Ingold, *J. Am. Chem. Soc.* **1985**, *107*, 7053.

[3] A. Watanabe, N. Noguchi, A. Fujisawa, T. Kodama, K. Tamura, O. Cynshi, E. Niki, *J. Am. Chem. Soc.* **2000**, *122*, 5438.

[4] a) D. A. Pratt, G. A. DiLabio, G. Brigati, G. F. Pedulli, L. Valgimigli, *J. Am. Chem. Soc.* **2001**, *123*, 4625. b) L. Valgimigli, G. Brigati, G. F. Pedulli, G. A. DiLabio, M. Mastragostino, C. Arbizzani, D. A. Pratt, *Chem. – Eur. J.* **2003**, *9*, in press.


- 
- [5] M. C. Foti, E. R. Johnson, M. R. Vinqvist, J. S. Wright, L. R. C. Barclay, K. U. Ingold, *J. Org. Chem.* **2002**, *67*, 5190.
- [6] a) D. P. S. Steinberg, T. E. Carew, J. C. Khoo, J. L. Witztum, *New England J. Med.* **1989**, *320*, 915.  
b) P. Holvoet, D. Collen, *FASEB J.* **1994**, *8*, 1279. c) D. Steinberg, *Lancet* **1995**, *346*, 36.
- [7] a) D. D. M. Wayner, E. Lusztyk, K. U. Ingold, P. Mulder, *J. Org. Chem.* **1996**, *61*, 6430. b) M. Lucarini, P. Pedrielli, G. F. Pedulli, S. Cabiddu, C. Fattuoni, *J. Org. Chem.* **1996**, *61*, 9263.
- [8] G. Brigati, M. Lucarini, V. Mugnaini, G. F. Pedulli, *J. Org. Chem.* **2002**, *67*, 4828.
- [9] G. A. DiLabio, D. A. Pratt, A. D. LoFaro, J. S. Wright, *J. Phys. Chem. A* **1999**, *103*, 1653.
- [10] G. A. DiLabio, D. A. Pratt, J. S. Wright, *J. Org. Chem.* **2000**, *65*, 2195.
- [11] a) C. A. A. van Boeckel, L. P. C. Delbressine, F. M. Kaspersen, *Recl. Trav. Chim. Pays-bas* **1985**, *104*, 259. b) H. van de Poël, G. Guillaumet; M. C. Viaud-Massuard, *Heterocycles* **2002**, *57*, 55.
- [12] K. A. Tallman, D. A. Pratt, N. A. Porter, *J. Am. Chem. Soc.* **2001**, *123*, 11827.
- [13] The very low oxidation yields led to almost undetectable HPLC peaks.
- [14] R. Amorati, G. F. Pedulli, L. Valgimigli, O. A. Attanasi, P. Filippone, C. Fiorucci, R. Saladino, *J. Chem. Soc. Perkin Trans. 2*, **2001**, 2142.
- [15] M. Lucarini, P. Pedrielli, G. F. Pedulli, L. Valgimigli, D. Gigmes, P. J. Tordo, *J. Am. Chem. Soc.* **1999**, *121*, 11546.
- [16] This compound is quite reactive with air, much more so than **6** or **7**; quantitatively this is indicated by our calculated IP that is 7.8 and 5.5 kcal mol<sup>-1</sup> lower than **6** and **7**, respectively.

## SUPPORTING INFORMATION


for

### **6-Amino-3-Pyridinols: Towards Diffusion-Controlled Chain-Breaking Antioxidants\*\***


*Maikel Wijtmans, Derek A. Pratt,\* Luca Valgimigli,\* Gino A. DiLabio, Gian Franco Pedulli and  
Ned A. Porter\**



**Figure S1.** Determination of  $k_{inh}$  by the peroxy radical clock method. Plotted are the values for  $[11\text{-OOH}] / ([9\text{-OOH}] + [13\text{-OOH}])$  versus concentration of antioxidant in methyl linoleate autoxidations in benzene (200 mM) at 37 °C initiated by MeOAMVN (10 mM). Data for **2**, **4** and **5** are shown with the corresponding linear correlations from which the rate constants in Table 2 were derived.



**Figure S2.** A typical oxygen uptake plot for an inhibited styrene autoxidation. This plot shows the oxygen consumption during autoxidation of styrene (8.6M) in chlorobenzene at 30°C initiated by AMVN (1.25 mM) in the presence of **6** (1.1  $\mu$ M).



**Figure S3.** Decay of UV signal for 3-pyridinols **5**, **6** and **7** (@ 330 nm) and  $\alpha$ -tocopherol **2** (@ 304 nm) in aerated *t*-butylbenzene (300  $\mu$ M) at 37  $^{\circ}$ C.