

doi:10.1016/j.bmcl.2009.12.089 | How to Cite or Link Using DOI  
Copyright © 2009 Elsevier Ltd All rights reserved.

## Novel bis-2,2,6,6-tetramethylpiperidine (bis-TMP) and bis-mecamylamine antagonists at neuronal nicotinic receptors mediating nicotine-evoked dopamine release

Zhenfa Zhang<sup>a</sup>, Marharyta Pivavarchyk<sup>a</sup>, Karunai Leela Subramanian<sup>a</sup>, A. Gabriela Deaciuc<sup>a</sup>, Linda P. Dwoskin<sup>a</sup> and Peter A. Crooks<sup>a</sup>, , 

<sup>a</sup> College of Pharmacy, Department of Pharmaceutical Sciences, University of Kentucky, Lexington, KY 40536-0082, United States

Received 17 November 2009;

revised 21 December 2009;

accepted 24 December 2009.


Available online 4 January 2010.

### Abstract

By linking two or three mecamylamine or 2,2,6,6-tetramethylpiperidine (TMP) molecules together via a linear lipophilic bis-methylene linker or a specially designed conformationally restricted tris-linker, a series of bis- and tris-tertiary amine analogs has been synthesized and evaluated as potent antagonists at nAChRs mediating nicotine-evoked [<sup>3</sup>H]dopamine release from rat striatal slices. Compounds **7e**, **14b** and **16** demonstrated high potency in decreasing nicotine-evoked [<sup>3</sup>H]dopamine release ( $IC_{50}$  = 2.2, 46, and 107 nM, respectively). The preliminary structure–activity data obtained with these new analogs suggest the importance of the length of the methylene linker in the bis-analog series. Such bis-tertiary amino analogs may provide a new strategy for the design of drugable ligands that have high inhibitory potency against nAChRs mediating nicotine-evoked dopamine release in striatum, which have been suggested to be target receptors of interest in the development of potential smoking cessation therapies.

### Graphical abstract

Compounds **7e** and **14b** demonstrated high potency in decreasing nicotine-evoked [<sup>3</sup>H]dopamine release from rat striatal slices ( $IC_{50}$  = 2.2 and 46 nM, respectively). Such bis-tertiary amino analogs may provide a new strategy for the design of drugable ligands that have high inhibitory potency against nAChRs mediating nicotine-evoked dopamine release in striatum, which have been suggested to be target receptors of interest in the development of potential smoking cessation therapies.



**Keywords:** Nicotinic acetylcholine receptor; Quaternary ammonium; Dopamine release; Nicotine addiction

## Article Outline

[Acknowledgements](#)

[References](#)





Figure 1.

Structures of mecamylamine (1), bPiDDB (2), TMP (3), BTMPS (4), and tPy3PiB (5).


[View Within Article](#)



Scheme 1.

Synthesis of bis-TMP analogs **7a–7e**, and bis-S-(+)-mecamylamine analogs **14a** and **14b**.

[View Within Article](#)



Scheme 2.

Synthesis of tris-mecamylamine analog **16**.

[View Within Article](#)




Figure 2.

S(-)-Nicotine-evoked fractional [ $^3\text{H}$ ]DA release from rat striatal slices superfused with 100 nM **14a** (bMecD), **14b** (bMecDD) and **16** (tMecBPY). Data are expressed as mean  $\pm$  SEM fractional release as a percent of basal fractional release, that is, percent of samples prior to the addition of analog or nicotine. Control represents the amount of fractional release evoked by S(-)-nicotine in the absence of analog;  $n = 3$  rats/analog.

[View Within Article](#)




Figure 3.

Analog **16** (tMecBPY) inhibited S(-)-nicotine-evoked [ $^3\text{H}$ ]DA overflow from rat striatal slices in a concentration-dependent manner. Control represents [ $^3\text{H}$ ]DA overflow in response to 10  $\mu\text{M}$  nicotine in the absence of analog and is expressed as a percent of tissue [ $^3\text{H}$ ] content, mean  $\pm$  S.E.M,  $n = 4$  rats.

[View Within Article](#)

Table 1. Inhibition of nicotine-evoked [ $^3\text{H}$ ]DA release from superfused rat striatal slices<sup>a</sup>



<sup>a</sup> Percentage of inhibition at 100 nM are presented unless otherwise specified. Each value represents data from at least three independent experiments, each performed in duplicate.

<sup>b</sup>  $\text{IC}_{50}$  and  $I_{\text{max}}$  from full concentration response assays; data from 4 to 6 independent experiments.

<sup>c</sup> Not determined.

[View Within Article](#)



Corresponding author. Tel.: +1 859 257 1718; fax: +1 859 257 7585.