

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/6275764>

Synthesis and in vitro activity of 2-thiazolylylhydrazone derivatives compared with the activity of clotrimazole against clinical isolates of *Candida* spp.

ARTICLE in BIOORGANIC & MEDICINAL CHEMISTRY LETTERS · SEPTEMBER 2007

Impact Factor: 2.42 · DOI: 10.1016/j.bmcl.2007.05.078 · Source: PubMed

CITATIONS

29

READS

51

13 AUTHORS, INCLUDING:



Elias Maccioni

Università degli studi di Cagliari

68 PUBLICATIONS 736 CITATIONS

SEE PROFILE



Daniela Secci

Sapienza University of Rome

110 PUBLICATIONS 2,113 CITATIONS

SEE PROFILE



Simone Carradori

Università degli Studi G. d'Annunzio Chieti ...

92 PUBLICATIONS 931 CITATIONS

SEE PROFILE



Alessandra Zicari

Sapienza University of Rome

34 PUBLICATIONS 283 CITATIONS


SEE PROFILE



Vol. 17 · No. 16 · 15 August 2007 · ISSN 0960-894X

Bioorganic & Medicinal Chemistry Letters

The Tetrahedron Journal for Research at the Interface
of Chemistry and Biology



Editor-in-Chief
DALE L. BOGER

Available online at
 ScienceDirect
www.sciencedirect.com

This article was published in an Elsevier journal. The attached copy is furnished to the author for non-commercial research and education use, including for instruction at the author's institution, sharing with colleagues and providing to institution administration.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>

Synthesis and in vitro activity of 2-thiazolyldrazone derivatives compared with the activity of clotrimazole against clinical isolates of *Candida* spp.

Franco Chimenti,^a Bruna Bizzarri,^{a,*} Elias Maccioni,^b Daniela Secci,^a Adriana Bolasco,^a Rossella Fioravanti,^a Paola Chimenti,^a Arianna Granese,^a Simone Carradori,^a Daniela Rivanera,^c Daniela Lilli,^c Alessandra Zicari^d and Simona Distinto^b

^aDipartimento di Studi di Chimica e Tecnologia delle Sostanze Biologicamente Attive Università degli Studi di Roma “La Sapienza”, P.le A. Moro 5, 00185 Rome, Italy

^bDipartimento Farmaco Chimico Tecnologico, Università degli Studi di Cagliari, Via Ospedale 72, 09124 Cagliari, Italy

^cDipartimento di Scienze di Sanità Pubblica, Università “La Sapienza”, P.le A. Moro 5, 00185 Rome, Italy

^dDipartimento di Medicina Sperimentale, Università “La Sapienza”, V.le Regina Elena 324, 00161 Rome, Italy

Received 17 April 2007; revised 23 May 2007; accepted 24 May 2007

Available online 27 May 2007

Abstract—In this paper, we report on the synthesis of a novel series of 2-thiazolyldrazone derivatives and the influence of the substituents on the thiazole ring on antifungal activity. All synthesized compounds were screened for their in vitro activities against 22 clinical isolates of *Candida* spp., representing six different species, compared to clotrimazole as a reference compound. Some of the tested compounds were found to possess significant antifungal activity when compared to clotrimazole, in particular compound **14** which exhibited higher potency against most of the *Candida* spp. considered. The compounds that were most active as anti-*Candida* agents were also submitted to cytotoxic screening by the Trypan Blue dye exclusion assay and in general they were shown to induce low cytotoxic effects.

© 2007 Elsevier Ltd. All rights reserved.

Although *Candida* spp. are present as commensal flora in 30–60% of healthy individuals,¹ they can become infective depending on predisposing conditions related to the host, such as systemic disease leading to immunosuppression, and local conditions.² As a matter of fact, the pathogenicity of *Candida* spp. is affected by several virulence factors, such as the ability to adhere to epithelial and endothelial cells,³ germination, extracellular proteinases and phospholipases, and phenotypic switching.⁴

Over the past few decades, the frequency of systematic fungal infections has progressively increased, due to the larger population of immunologically compromised hosts (AIDS, cancer, and transplant), thanks to advances

in supportive therapy and an increasing elderly population.

In addition, the widespread use of antimicrobial agents for prophylaxis and treatment has led to less effective antibiotics, not only because many induced side effects but also due to the emergence of drug resistant microorganisms.^{5–7}

Large-scale surveillance for fungal bloodstream infections has been performed worldwide by a number of organizations, including the European Confederation of Medical Mycology (ECMM),⁸ the Centres for Disease Control and Prevention (CDC),⁹ and the National Epidemiology of Mycoses Survey (NEMIS).¹⁰ These studies have demonstrated an increasing incidence of drug-resistant fungal pathogens. As a matter of fact, a significant number of yeast species (*Candida glabrata*, *Candida krusei*, *Candida guilliermondii*, *Candida lusitanae*) have been shown to exhibit primary resistance to amphotericin B, while *C. glabrata* and *C. krusei* have

Keywords: 2-Thiazolyldrazone derivatives; Antifungal agents; *Candida* spp.

* Corresponding author. Tel.: +39 06 49913975; fax: +39 06 49913772; e-mail: bruna.bizzarri@uniroma1.it

infections today can reach 90% in immunocompromised patients especially.¹⁵


Apart from this, antifungal chemotherapy could be associated with a low toxicological profile, determining termination of the treatment.¹⁶

Another important issue is the clinical outcome due to several factors such as host defenses, the virulence of the fungal pathogen, drug efficacy/toxicity, and drug interactions.^{17,18} Besides, pharmacokinetic studies have shown a variability of at least 50% between individuals;^{19,20} in particular, plasma levels for voriconazole can vary from 74% to 100%.²¹ These variations could account for the lack of efficacy and the toxicity.

As a consequence of the toxicity of the currently used polyene antifungal drugs and the emergence of candidal species resistant to azole-based agents, there is an urgent need for investigating alternative drug therapies.


It has recently been reported in the literature that phenyl-thiazole analogues can be used as efficient antifungal agents.²² Moreover, a large number of substituted benzothiazole,²³ thiazole derivatives of triazoles,²⁴ thiazolyldiazones²⁵ have been shown to exhibit significant antimicrobial activity against a variety of fungal strains.

Moving from these literature indications and pursuing our research in the field,^{26,27} in this report we describe the synthesis and antimicrobial evaluation of a new series of 2-thiazolylhydrazones derivatives.



Scheme 1. Synthesis of 2-thiazolylhydrazone derivatives **1–20**.
Reagents: (i) isopropyl alcohol, AcOH; (ii) isopropyl alcohol.

Table 1. Chemical and physical data of derivatives **1–20**



Compound	R	R ¹	M.W.	Mp (°C)	% Yield
1	Cyclopentyl	4-CH ₃	352.27 ^a	218–220	71
2	Cyclopentyl	4-OCH ₃	368.27 ^a	214–217	88
3	Cyclopentyl	3-NO ₂	383.27 ^a	187–190	69
4	Cyclopentyl	4-NO ₂	383.27 ^a	219–220	80
5	Cyclopentyl	4-CN	363.28 ^a	207–208	67
6	Cyclopentyl	4-Cl	372.70 ^a	223–225	58
7	Cyclopentyl	4-Br	417.16 ^a	213–215	66
8	Cyclopentyl	4-C ₆ H ₅	414.35 ^a	235–237	58
9	Cyclohexyl-2-CH ₃	4-CH ₃	380.34 ^a	160–162	64
10	Cyclohexyl-2-CH ₃	4-OCH ₃	396.34 ^a	135–137	94
11	Cyclohexyl-2-CH ₃	3-NO ₂	411.31 ^a	165–166	60
12	Cyclohexyl-2-CH ₃	4-Br	445.21 ^a	130–133	71
13	Cyclohexyl-2-CH ₃	4-C ₆ H ₅	442.41 ^a	150–152	87
14	Cyclohexyl-3-CH ₃	4-CH ₃	380.34 ^a	171–174	95
15	Cyclohexyl-3-CH ₃	4-OCH ₃	396.34 ^a	162–165	84
16	Cyclohexyl-3-CH ₃	4-CN	391.32 ^a	189–190	98
17	Cyclohexyl-3-CH ₃	4-Cl	400.76 ^a	195–199	84
18	Cyclohexyl-3-CH ₃	4-F	339.85 ^b	192–195	77
19	Cyclohexyl-4-CH ₃	4-Cl	400.76 ^a	177–179	92
20 ²⁸	Cycloheptyl	H	285.41	184–185	97

^a Chloridrate.^b Bromidrate.

The cytotoxic activity of selected compounds that showed good activity against *Candida* species was evaluated: we incubated them in the presence of an immortalized hybrid cell line displaying an endothelial

phenotype, EAhy 926, derived from the fusion of human umbilical vein endothelial cells (HUVEC) with a lung carcinoma cells and determined their cell viability by the Trypan Blue dye exclusion assay.

Table 2. ^1H NMR data of derivatives 1–20

Compound	^1H NMR δ (ppm)
1 ^a	1.80 (q, 2H, cyclopentyl), 1.92 (q, 2H, cyclopentyl), 2.35 (s, 3H, 4'-CH ₃ -phenyl), 2.50 (t, 2H, cyclopentyl), 2.62 (t, 2H, cyclopentyl), 6.67 (s, 1H, C ₅ H-thiaz.), 7.22 (d, $J = 7.9$ Hz, 2H, Ar), 7.55 (d, $J = 7.9$ Hz, 2H, Ar), 12.13 (br s, 1H, NH, D ₂ O exch.)
2 ^a	1.85 (q, 2H, cyclopentyl), 1.90 (q, 2H, cyclopentyl), 2.50 (t, 2H, cyclopentyl), 2.63 (t, 2H, cyclopentyl), 3.83 (s, 3H, 4'-OCH ₃ -phenyl), 6.51 (s, 1H, C ₅ H-thiaz.), 6.96 (d, $J = 7.9$ Hz, 2H, Ar), 7.63 (d, $J = 7.9$ Hz, 2H, Ar), 12.13 (br s, 1H, NH, D ₂ O exch.)
3 ^a	1.90 (q, 2H, cyclopentyl), 1.97 (q, 2H, cyclopentyl), 2.58 (t, 2H, cyclopentyl), 2.68 (t, 2H, cyclopentyl), 6.96 (s, 1H, C ₅ H-thiaz.), 7.77 (t, 1H, Ar), 8.20 (t, 1H, Ar), 8.31 (d, $J = 7.9$ Hz, 1H, Ar), 8.53 (d, $J = 1.68$ Hz, 1H, Ar), 12.13 (br s, 1H, NH, D ₂ O exch.)
4 ^a	1.91 (q, 2H, cyclopentyl), 2.01 (q, 2H, cyclopentyl), 2.57 (t, 2H, cyclopentyl), 2.66 (t, 2H, cyclopentyl), 6.97 (s, 1H, C ₅ H-thiaz.), 7.94 (d, $J = 8.8$ Hz, 2H, Ar), 8.35 (d, $J = 8.8$ Hz, 2H), 12.40 (br s, 1H, NH, D ₂ O exch.)
5 ^b	1.83 (q, 2H, cyclopentyl), 1.87 (q, 2H, cyclopentyl), 2.37 (t, 4H, cyclopentyl), 7.66 (s, 1H, C ₅ H-thiaz.), 7.96 (d, $J = 8.4$ Hz, 2H, Ar), 8.12 (d, $J = 8.4$ Hz, 2H, Ar), 10.83 (br s, 1H, NH, D ₂ O exch.)
6 ^b	1.72 (q, 2H, cyclopentyl), 1.78 (q, 2H, cyclopentyl), 2.37 (t, 4H, cyclopentyl), 7.30 (s, 1H, C ₅ H-thiaz.), 7.44 (d, $J = 8.4$ Hz, 2H, Ar), 7.84 (d, $J = 8.4$ Hz, 2H, Ar), 10.70 (br s, 1H, NH, D ₂ O exch.)
7 ^b	1.72 (q, 2H, cyclopentyl), 1.80 (q, 2H, cyclopentyl), 2.37 (t, 4H, cyclopentyl), 7.31 (s, 1H, C ₅ H-thiaz.), 7.57 (d, $J = 8.4$ Hz, 2H, Ar), 7.77 (d, $J = 8.4$ Hz, 2H, Ar), 10.65 (br s, 1H, NH, D ₂ O exch.)
8 ^a	1.87 (q, 2H, cyclopentyl), 1.94 (q, 2H, cyclopentyl), 2.52 (t, 2H, cyclopentyl), 2.66 (t, 2H, cyclopentyl), 6.72 (s, 1H, C ₅ H-thiaz.), 7.38 (d, $J = 7.1$ Hz, 1H, Ar), 7.47 (t, 2H, Ar), 7.59 (d, $J = 7.1$ Hz, 2H, Ar), 7.69 (d, $J = 7.9$ Hz, 2H, Ar), 7.77 (d, $J = 7.9$ Hz, 2H, Ar), 12.24 (br s, 1H, NH, D ₂ O exch.)
9 ^a	1.16 (d, $J = 6.3$ Hz, 3H, CH ₃), 1.24–1.42 (m, 1H, cyclohexyl), 1.57–1.66 (m, 2H, cyclohexyl), 1.79–1.86 (m, 1H, cyclohexyl), 1.98–2.03 (m, 2H, cyclohexyl), 2.17–2.27 (m, 1H, cyclohexyl), 2.41 (s, 3H, 4'-CH ₃ -phenyl), 2.43–2.49 (m, 1H, cyclohexyl), 3.04–3.10 (m, 1H, cyclohexyl), 6.66 (s, 1H, C ₅ H-thiaz.), 7.28 (d, $J = 8.4$ Hz, 2H, Ar), 7.60 (d, $J = 4$ Hz, 2H, Ar), 12.50 (s, 1H, NH, D ₂ O exch.)
10 ^a	1.11 (d, $J = 6.3$ Hz, 3H, CH ₃), 1.17–1.37 (m, 1H, cyclohexyl), 1.48–1.61 (m, 2H, cyclohexyl), 1.75–1.81 (m, 1H, cyclohexyl), 1.95–1.99 (m, 2H, cyclohexyl), 2.11–2.21 (m, 1H, cyclohexyl), 2.35–2.44 (m, 1H, cyclohexyl), 2.97–3.03 (m, 1H, cyclohexyl), 3.81 (s, 3H, 4'-OCH ₃ -phenyl), 7.53 (s, 1H, C ₅ H-thiaz.), 6.94 (d, $J = 8.8$ Hz, 2H, Ar), 7.60 (d, $J = 8.8$ Hz, 2H, Ar), 12.37 (s, 1H, NH, D ₂ O exch.)
11 ^b	1.08 (d, $J = 6.5$ Hz, 3H, CH ₃), 1.16–1.26 (m, 1H, cyclohexyl), 1.37–1.59 (m, 2H, cyclohexyl), 1.69–2.02 (m, 4H, cyclohexyl), 2.31–2.42 (m, 1H, cyclohexyl), 2.91–2.96 (m, 1H, cyclohexyl), 7.56 (s, 1H, C ₅ H-thiaz.), 7.70 (t, 1H, Ar), 8.12 (d, $J = 8.1$ Hz, 1H, Ar), 8.27 (d, $J = 8.1$ Hz, 1H, Ar), 8.68 (s, 1H, Ar), 10.95 (s, 1H, NH, D ₂ O exch.)
12 ^a	1.15 (d, $J = 6.3$ Hz, 3H, CH ₃), 1.26–1.38 (m, 1H, cyclohexyl), 1.54–1.65 (m, 2H, cyclohexyl), 1.75–1.81 (m, 1H, cyclohexyl), 1.92–1.98 (m, 2H, cyclohexyl), 2.13–2.23 (m, 1H, cyclohexyl), 2.39–2.47 (m, 1H, cyclohexyl), 2.97–3.02 (m, 1H, cyclohexyl), 6.76 (s, 1H, C ₅ H-thiaz.), 7.59 (s, 4H, Ar), 12.41 (s, 1H, NH, D ₂ O exch.)
13 ^a	1.13 (d, $J = 6.3$ Hz, 3H, CH ₃), 1.26–1.39 (m, 1H, cyclohexyl), 1.54–1.61 (m, 2H, cyclohexyl), 1.79–1.83 (m, 1H, cyclohexyl), 1.89–1.97 (m, 2H, cyclohexyl), 2.17–2.22 (m, 1H, cyclohexyl), 2.38–2.41 (m, 1H, cyclohexyl), 3.01–3.05 (m, 1H, cyclohexyl), 6.75 (s, 1H, C ₅ H-thiaz.), 7.36 (d, $J = 7.1$ Hz, 1H, Ar), 7.46 (t, 2H, Ar), 7.59 (d, $J = 7.1$ Hz, 2H, Ar), 7.69 (d, $J = 8.4$ Hz, 2H, Ar), 7.78 (d, $J = 8.4$ Hz, 2H, Ar), 12.48 (s, 1H, NH, D ₂ O exch.)
14 ^a	1.16 (d, $J = 6.4$ Hz, 3H, CH ₃), 1.35 (s, 1H, cyclohexyl), 1.60–1.62 (m, 2H, cyclohexyl), 1.81 (s, 2H, cyclohexyl), 1.98–1.99 (m, 2H, cyclohexyl), 2.20–2.22 (m, 1H, cyclohexyl), 2.42 (d, $J = 6.1$ Hz, 3H, 4'-CH ₃ -phenyl), 3.03–3.04 (m, 1H, cyclohexyl), 6.63 (s, 1H, C ₅ H-thiaz.), 7.28 (s, 2H, Ar), 7.60 (s, 2H, Ar), 12.52 (s, 1H, NH, D ₂ O exch.), 13.56 (s, 1H, NH, D ₂ O exch.)
15 ^a	1.06 (d, $J = 6.2$ Hz, 3H, CH ₃), 1.20–1.21 (m, 1H, cyclohexyl), 1.81–1.95 (m, 6H, cyclohexyl), 2.52 (t, 1H, cyclohexyl), 3.05–3.06 (m, 1H, cyclohexyl), 3.85 (s, 3H, 4'-OCH ₃ -phenyl), 6.52 (s, 1H, C ₅ H-thiaz.), 6.99 (d, $J = 8.0$ Hz, 2H, Ar), 7.65 (d, $J = 8.6$ Hz, 2H, Ar), 12.44 (s, 1H, NH, D ₂ O exch.), 13.58 (s, 1H, NH, D ₂ O exch.)
16 ^a	1.19 (d, $J = 6.4$ Hz, 3H, CH ₃), 1.57–1.61 (m, 3H, cyclohexyl), 1.83–1.85 (m, 1H, cyclohexyl), 2.00–2.03 (m, 2H, cyclohexyl), 2.19–2.21 (m, 1H, cyclohexyl), 2.50–2.52 (m, 1H, cyclohexyl), 3.00–3.01 (m, 1H, cyclohexyl), 6.87 (s, 1H, C ₅ H-thiaz.), 7.79 (d, $J = 7.8$ Hz, 2H, Ar), 7.86 (s, 2H, Ar), 12.48 (s, 1H, NH, D ₂ O exch.), 14.01 (br s, 1H, NH, D ₂ O exch.)
17 ^a	1.64 (d, $J = 6.5$ Hz, 3H, CH ₃), 1.25–1.26 (m, 1H, cyclohexyl), 1.69–1.71 (m, 1H, cyclohexyl), 1.98–1.99 (m, 2H, cyclohexyl), 2.20–2.21 (m, 2H, cyclohexyl), 2.50–2.52 (m, 2H, cyclohexyl), 2.95–2.96 (m, 1H, cyclohexyl), 6.68 (s, 1H, C ₅ H-thiaz.), 7.47 (d, $J = 8.6$ Hz, 2H, Ar), 7.67 (d, $J = 8.5$ Hz, 2H, Ar), 12.50 (s, 1H, NH, D ₂ O exch.), 13.80 (s, 1H, NH, D ₂ O exch.)
18 ^a	1.16 (d, $J = 6.5$ Hz, 3H, CH ₃), 1.58–1.59 (m, 3H, cyclohexyl), 1.98–1.99 (m, 1H, cyclohexyl), 2.05–2.07 (m, 2H, cyclohexyl), 2.25–2.27 (m, 1H, cyclohexyl), 2.40–2.41 (m, 1H, cyclohexyl), 3.00–3.01 (m, 1H, cyclohexyl), 6.62 (s, 1H, C ₅ H-thiaz.), 7.18 (d, $J = 6.1$ Hz, 2H, Ar), 7.70 (d, $J = 5.3$ Hz, 2H, Ar), 12.71 (s, 1H, NH, D ₂ O exch.), 14.30 (s, 1H, NH, D ₂ O exch.)
19 ^a	0.99 (d, $J = 6.1$ Hz, 3H, CH ₃), 1.20–1.21 (m, 2H, cyclohexyl), 1.60–1.61 (m, 1H, cyclohexyl), 2.00–2.01 (t, 2H, cyclohexyl), 2.20–2.30 (m, 2H, cyclohexyl), 2.60–2.61 (m, 1H, cyclohexyl), 3.09–3.10 (m, 1H, cyclohexyl), 6.68 (s, 1H, C ₅ H-thiaz.), 7.47–7.51 (m, 2H, Ar), 7.71 (d, $J = 6.6$ Hz, 2H, Ar), 12.55 (s, 1H, NH, D ₂ O exch.), 13.75 (s, 1H, NH, D ₂ O exch.)
20 ^a	1.58–1.63 (m, 6H, cycloheptyl), 1.70–1.71 (m, 2H, cycloheptyl), 2.55–2.56 (m, 2H, cycloheptyl), 2.70–2.72 (m, 2H, cycloheptyl), 6.80 (s, 1H, C ₅ H-thiaz.), 7.48–7.50 (m, 3H, Ar), 7.73 (d, $J = 7.9$ Hz, 2H, Ar), 12.40 (s, 1H, NH, D ₂ O exch.), 13.80 (br s, 1H, NH, D ₂ O exch.)

^a CDCl₃.

^b DMSO-*d*₆.

2-Thiazolylhydrazone derivatives **1–20** were synthesized as reported in our previous communications.^{28,29}

Cyclic ketones or aryl aldehydes reacted directly with thiosemicarbazide and the obtained thiosemicarbazones subsequently reacted with α -halogenoketones to yield the 4-substituted thiazole ring derivatives as shown in Scheme 1. In the synthesis of all compounds isopropyl alcohol proved to be the best solvent for our purpose. As a matter of fact, the reaction products precipitate on cooling down and can be filtered and purified by crystallization from ethanol or ethanol/isopropanol. All synthesized compounds were fully characterized by analytical and spectral data as listed in Tables 1 and 2.

All synthesized compounds were evaluated for antifungal activity and compared with the reference compound clotrimazole (Table 3).^{30,31}

The newly prepared compounds were dissolved in dimethylsulfoxide (DMSO) and their in vitro activity was evaluated against a total of 22 strains of *Candida* species. The included isolates were *Candida albicans* (8 strains), *Candida glabrata* (4), *Candida krusei* (3), *Candida tropicalis* (3), *Candida sakei* (2), and *Candida parapsilosis* (2). The isolates were collected from specimens of patients at the 'Azienda Policlinico Umberto I' of Rome 'La Sapienza' University and were obtained from haematology/oncology and surgery departments, which also included an intensive care unit. In particular, the samples were isolated from the upper and lower respiratory tract, blood, and indwelling venous catheters; the intensive care

unit accounted for 65% of the cases (15/22 isolates). The isolates were identified by conventional methodologies. Prior to testing, each isolate was subcultured on a qualified medium to ensure purity and optimal growth.

The data reported in Table 3 show that cyclopentyl derivatives **1–6** had good anti-*C. albicans* and anti-*C. glabrata* activity. In particular compounds **1**, **2**, and **6** were very active against both strains, while compounds **3**, **4**, and **5** showed good activity only against *C. glabrata*. Furthermore, cyclohexyl substituted derivatives **9–19** showed good anti-*C. albicans* activity, in particular with compounds **9–11** and **14–19**. Some derivatives, especially **9** and **10**, also showed good anti-*C. glabrata* activity. Cycloheptyl derivative **20** showed good anti-*C. albicans* activity. Among all synthesized compounds, we found the antifungal activity showed by compound **14**, 2-(3-methylcyclohexyl)-1-[4-(4-methylphenyl)-2-thiazolyl]hydrazone, particularly interesting. Compound **14** was more active than the clotrimazole reference compound against *C. albicans*, *C. tropicalis*, *C. krusei*, *C. parapsilosis*, and *C. sakei*.

The cytotoxic profile of the compounds showed that the derivatives were less toxic at concentrations below 0.5 $\mu\text{g/mL}$ with a percentage of viable cells of 73.0 and 94.7 (Table 4).^{32–34}

In conclusion, a series of novel 2-thiazolylhydrazone derivatives was assessed for antifungal activity on 22 *Candida* strains. Compound **14** was found to have good antifungal activity and may be used as a good reference for identifying features of the structure that could be important for antifungal activity.

Table 3. Minimal inhibitory concentration (MIC)^a of compounds **1–20** and clotrimazole against 22 strains of *Candida* species

Compound	Tested fungi (MIC ^a $\mu\text{g/mL}$)					
	<i>C. albicans</i> (8 strains)	<i>C. glabrata</i> (4 strains)	<i>C. tropicalis</i> (3 strains)	<i>C. krusei</i> (3 strains)	<i>C. parapsilosis</i> (2 strains)	<i>C. sakei</i> (2 strains)
1	0.25–2	0.50–2	4–16	64–128	128	16–32
2	0.50–2	2–4	64–128	32–64	8–16	32
3	1–8	0.50–2	64–128	>128	>128	16–32
4	8–32	0.50–2	64–128	32–64	>128	16
5	2–8	0.50–2	32–64	32–64	128	32
6	0.25–2	0.50–2	32–64	64–128	8–16	16
7	2–8	2–8	128–>128	>128	128	32
8	16–64	>128	64–128	>128	>128	32
9	0.25–2	0.50–2	2–4	64	8	32
10	0.50–2	0.25–2	8	16	8	32
11	0.50–2	16–64	64	>128	8–16	16–32
12	4–8	2–4	32–64	64–128	8–16	16
13	32–64	64–128	64	16	16	16–32
14	0.50–2	32–64	0.50–2	0.50–2	0.50–1	2–4
15	0.50–2	>128	2–4	2–4	1–2	8–16
16	0.50–2	>128	2–4	4–8	0.50–1	4–8
17	0.50–2	>128	2–4	4–8	1	2
18	0.50–4	>128	4–16	4–16	1	8
19	0.50–2	32–64	2–4	05–1	8	2
20	0.50–2	>128	8–16	8–16	8–16	16–32
Clotrimazole^b	0.50–4	4–8	4–8	4–8	4–8	8–16

^a Range value.

^b Clotrimazole.

Table 4. Cytotoxic effect of selected compounds tested on EAhy 926 cells after 24 h of incubation at 37 °C, using Trypan Blue exclusion test, expressed as cell survival fraction^a (%)

Compound	Concentration ^b (μg/mL)			
	50	5	0.5	0.05
1	75.0 ± 3.2	80.0 ± 4.1	82.0 ± 4.6	85.4 ± 5.0
2	66.0 ± 3.2	76.6 ± 4.0	82.4 ± 2.8	85.0 ± 2.2
5	76.0 ± 4.6	81.0 ± 3.2	84.7 ± 2.8	88.9 ± 6.0
6	64.9 ± 4.2	81.0 ± 3.0	89.2 ± 2.8	86.0 ± 2.6
9	64.0 ± 3.2	63.6 ± 3.6	78.5 ± 4.2	85.7 ± 3.6
10	75.0 ± 4.0	83.3 ± 3.6	84.0 ± 3.2	86.0 ± 4.8
11	83.3 ± 3.6	85.7 ± 4.2	86.0 ± 3.8	88.8 ± 2.2
14	38.5 ± 4.3	72.7 ± 3.1	83.3 ± 3.6	84.0 ± 2.0
15	54.5 ± 3.2	72.7 ± 3.8	80.0 ± 1.9	83.3 ± 2.2
16	25.0 ± 1.9	50.0 ± 2.5	68.7 ± 2.9	68.8 ± 1.4
17	28.6 ± 2.8	50.0 ± 1.9	70.0 ± 4.2	73.0 ± 3.1
18	52.9 ± 2.7	60.0 ± 1.5	76.9 ± 1.7	80.0 ± 2.0
20	54.5 ± 5.0	80.0 ± 3.9	87.5 ± 1.8	94.7 ± 2.5

^a Cells incubated with culture medium alone represented the controls and the cell viability was always greater than 97%.

^b Data represent the arithmetic means ± SD of at least three independent experiments.

Moreover other tested compounds may be good candidates for further investigation.

Acknowledgments

This work was supported by grants from MURST. We acknowledge Anton Gerada, a professional translator Fellow of the Institute of Translation and Interpreting of London and Member of AIIC (Association Internationale des Interprètes de Conférences—Geneva) for revision of the manuscript.

References and notes

- Haberland-Carrodegua, C.; Allen, C. M.; Beck, F. M.; Bueschiling, W. J.; Koletar, S. L.; Sundstron, P. *J. Oral Pathol. Med.* **2002**, *31*, 99.
- Maza, J. L.; Elguezabal, N.; Prado, C.; Ellacuria, J.; Soler, I.; Pontón, J. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol. Endod.* **2002**, *94*, 589.
- Sugita, T.; Kurosaka, S.; Yajitate, M.; Sato, H.; Nishikawa, A. *Microbiol. Immunol.* **2002**, *46*, 881.
- Gannoum, M. *Clin. Microbiol. Rev.* **2000**, *13*, 122.
- Moudgal, V.; Little, T.; Bolkov, D.; Vazquez, J. A. *Antimicrob. Agents Chemother.* **2005**, *49*, 767.
- Hernandez, S.; Lopez-Ribot, J. L.; Najvar, L. K.; McCarthy, D. I.; Bocanegra, R.; Graybill, J. R. *Antimicrob. Agents Chemother.* **2004**, *48*, 1382.
- Krogh-Madsen, M.; Arendrup, M. C.; Heslet, L.; Knudsen, J. D. *Clin. Infect. Dis.* **2006**, *42*, 938.
- Tortorano, A. M.; Kibbler, C.; Peman, J.; Bernhardt, H.; Klingspor, L.; Grillot, R. *Int. J. Antimicrob. Agents* **2006**, *27*, 359.
- Pfaller, M. A.; Jones, R. N.; Doern, G. V.; Fluit, A. C.; Verhoef, J.; Sader, H. S.; Messer, S. A.; Houston, A.; Coffman, S.; Hollis, R. J. *Diagn. Microbiol. Infect. Dis.* **1999**, *35*, 19.
- Blumberg, H. M.; Jarvis, W. R.; Soucie, J. M.; Edwards, J. E.; Patterson, J. E.; Pfaller, M. A.; Rangel-Frausto, M. S.; Rinaldi, M. G.; Saiman, L.; Wiblin, R. T.; Wenzel, R. P. *Clin. Infect. Dis.* **2001**, *33*, 177.
- Moran, G. P.; Sanglard, D.; Donnelly, S. M.; Shanley, D. B.; Sullivan, D. J.; Coleman, D. C. *Antimicrob. Agents Chemother.* **1998**, *42*, 1819.
- Sanglard, D.; Odds, F. C. *Lancet Infect. Dis.* **2002**, *2*, 73.
- White, T. C.; Marr, K. A.; Bowden, R. A. *Clin. Microbiol. Rev.* **1998**, *35*, 382.
- Hossain, M. A.; Ghannoum, M. A. *Exp. Opin. Investig. Drugs* **2000**, *9*, 1797.
- Grossi, P.; Farina, C.; Fiocchi, R.; Dalla Gasperina, D. *Transplantation* **2000**, *70*, 112.
- Groll, A. H.; Piscitelli, S. C.; Walsh, T. J. *Adv. Pharmacol.* **1998**, *44*, 343.
- Kontoyiannis, D. P.; Lewis, R. E. *Lancet* **2002**, *359*, 1135.
- White, T. C.; Marr, K. A.; Bowden, R. A. *Clin. Microbiol. Rev.* **1998**, *11*, 382.
- Courtney, R.; Radwanski, E.; Lim, J.; Laughlin, M. *Antimicrob. Agents Chemother.* **2004**, *48*, 804.
- Csajka, C.; Decosterd, L. A.; Buclin, T.; Pagani, J. L.; Fattinger, K.; Bille, J.; Biollaz, J. *Eur. J. Clin. Pharmacol.* **2001**, *57*, 723.
- Hoffman, H. L.; Rathbun, R. C. *Exp. Opin. Investig. Drugs* **2002**, *11*, 409.
- De Logu, A.; Saddi, M.; Cardia, M. C.; Borgna, R.; Sanna, C.; Saddi, B.; Maccioni, E. *J. Antimicrob. Chemother.* **2005**, *55*, 692.
- Turan-Zitouni, G.; Demirayak, S.; Ozdemir, A.; Kaplancikli, Z. A.; Yildiz, M. T. *Eur. J. Med. Chem.* **2003**, *39*, 267.
- Turan-Zitouni, G.; Kaplancikli, Z. A.; Yildiz, M. T.; Chevallet, P.; Kaya, D. *Eur. J. Med. Chem.* **2005**, *40*, 607.
- Cukurovali, A.; Yilmaz, I.; Gur, S.; Kazaz, C. *Eur. J. Med. Chem.* **2006**, *41*, 201.
- Chimenti, F.; Bizzarri, B.; Manna, F.; Bolasco, A.; Secci, D.; Chimenti, P.; Granese, A.; Rivanera, D.; Lilli, D.; Scaltrito, M. M.; Brenciaglia, M. I. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 603.
- Chimenti, F.; Bizzarri, B.; Bolasco, A.; Secci, D.; Chimenti, P.; Carradori, S.; Granese, A.; Rivanera, D.; Lilli, D.; Scaltrito, M. M.; Brenciaglia, M. I. *Eur. J. Med. Chem.* **2006**, *41*, 208.
- Maccioni, E.; Cardia, M. C.; Bonsignore, L.; Plumitallo, A.; Pellerano, M. L.; De Logu, A. *Il Farmaco* **2002**, *57*(10), 809.
- Maccioni, E.; Cardia, M. C.; Distinto, S.; Bonsignore, L.; De Logu, A. *Il Farmaco* **2003**, *58*(9), 951.
- Antifungal activity*: All synthesized derivatives 1–20 were evaluated for their antifungal activity dissolved in dimethylsulfoxide (DMSO). The in vitro antifungal activities of the compounds were determined with the broth microdilution method with Sabouraud dextrose broth (BBL Microbiology Systems, Cockeysville, MD) as recommended by the NCCLS.³¹ Microtiter plates containing serial dilutions of each compound were inoculated with each organism to yield the appropriate density (10³/mL) in a 100 μL final volume; each plate included positive controls (fungi without a compound) and a negative control (medium only). The plates were incubated for 24 h at 37 °C. The MIC for all isolates was defined as the lowest concentration of antifungal agent that completely inhibited growth of the organism, as detected by the unaided eye.
- National Committee for Clinical Laboratory Standards 2003. Performance standards for antimicrobial susceptibility testing. Supplement M100-S13. National Committee for Clinical Laboratory Standards, Wayne, PA.

32. *In vitro* cytotoxicity: The cytotoxicity of the newly synthesized compounds under investigation was tested against EAhy, human cell line obtained from a hybridoma between HUVEC and epithelial cells from a lung carcinoma. The viability of cells exposed to test compounds was estimated by the Trypan Blue dye exclusion assay. Cell lines were maintained as adherent type cultures under humidified atmosphere in 5% CO₂ at 37 °C, Dulbecco's modified Eagle's culture medium (high glucose) supplemented with 2 mM L-Glutamine, HAT supplement and containing antibiotic mixture. Experiments were performed in cells grown to 60–70% confluency.³³ The stock solutions of the investigated compounds were prepared in sterile dimethylsulfoxide (DMSO) and the successive dilutions were made in culture medium; the DMSO percent present in culture medium never exceeded 0.5%. EAhy cells in the exponential phase of growth (1×10^5 /mL) were seeded into 24-well microplate. and incubated

for 24 h with four different concentrations of the compounds (50–0.05 µg/mL). Some plates containing cells alone or cells and DMSO represented the controls. After the incubation period, cells were mechanically scraped off from the plates and an aliquot was diluted (1:1) with a solution 0.4% Trypan Blue Stain. After few minutes at rt cells were counted under an optical microscope in a Thoma hemocytometer chamber by two different operators. On the basis that Trypan Blue is a vital dye³⁴ and can enter and interact with the cells unless the plasmatic membrane is damaged, blue stained cells were considered as dead. Values are expressed as % of viable cells. Cell viability in control samples was always 97–98%.

33. Edgell, C. J. S.; McDonald, C. C.; Graham, J. B. *Proc. Natl. Acad. Sci. U.S.A.* **1983**, *80*, 3734.
34. Freshney, R. In *Culture of Animal Cells: A Manual of Basic Technique*; Alan R. Liss, Inc.: New York, 1987; p 117.