

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/267571997>

Quinoline: A promising antitubercular target

ARTICLE *in* BIOMEDECINE [?] PHARMACOTHERAPY · OCTOBER 2014

Impact Factor: 2.02 · DOI: 10.1016/j.biopha.2014.10.007

CITATIONS

7

READS

267

2 AUTHORS, INCLUDING:



Siddappa A Patil

Jain University

74 PUBLICATIONS 641 CITATIONS

SEE PROFILE



Available online at
ScienceDirect
 www.sciencedirect.com

Elsevier Masson France
EM|consulte
 www.em-consulte.com/en



Review

Quinoline: A promising antitubercular target



Rangappa S. Keri*, Siddappa A. Patil

Centre for Nano and Material Sciences, Jain University, Jain Global Campus, Jakkasandra post, Kanakapura Road, Ramanagara District, Karnataka 562112, India

ARTICLE INFO

Article history:

Received 13 September 2014

Accepted 16 October 2014

Keywords:

Quinoline

Synthesis

Anti-tubercular

Tubercle Bacillus

Mycobacterium tuberculosis

Medicinal chemistry

ABSTRACT

Tuberculosis (TB) remains a global public health problem in recent years. TB originated mainly from various strains of *Mycobacterium tuberculosis*, is a highly infectious and chronic disease with high infection rate since ancient times. Since the last 50 years, the same long-duration, multidrug treatment plan is being followed for the treatment of tuberculosis. Due to the development of resistance to conventional antibiotics there is a need for new therapeutic strategies to combat *M. tuberculosis*. Subsequently, there is an urgent need for the development of new drug molecules with newer targets and with an alternative mechanism of action. Among heterocyclic compounds, quinoline compounds are important privileged structure in medicinal chemistry, are widely used as “parental” compounds to synthesize molecules with medical benefits, especially with anti-malarial and anti-microbial activities. Certain, quinoline-based compounds, also show effective anti-TB activity. This broad spectrum of biological and biochemical activities has been further facilitated by the synthetic versatility of quinoline, which allows the generation of a large number of structurally diverse derivatives. To pave the way for future research, there is a need to collect the latest information in this promising area. In the present review, we have collated published reports on this versatile core to provide an insight so that its full therapeutic potential can be utilized for the treatment tuberculosis. It is hoped that, this review will be helpful for new thoughts in the quest for rational designs of more active and less toxic quinoline-based anti-TB drugs.

© 2014 Elsevier Masson SAS. All rights reserved.

Abbreviations: AIDS, acquired immunodeficiency syndrome; CQ, 7-chloroquinoline; DOS, directly observed therapy short-course; DNA, deoxyribonucleic acid; EMB, ethambutol; ETH, ethylenediamine spacer; GI, growth inhibition; HIV, human immunodeficiency virus; HTS, high throughput screening; INH, isoniazid; LORA, low-oxygen-recovery assay; LRP, luciferase reporter phage; LTBI, latent tuberculosis infection; MABA, Micro plate Alamar Blue Assay; *M. africanum*, *Mycobacterium africanum*; *M. smegmatis* (MS), *Mycobacterium smegmatis*; *M. bovis* (MB), *Mycobacterium bovis*; *M. caprae*, *Mycobacterium caprae*; *M. fortuitum* (MF), *Mycobacterium fortuitum*; MDR, multidrug-resistant; MIC, minimal inhibitory concentration; MDR-TB, multi-drug resistant tuberculosis; NTM, non-tubercular mycobacterial; MTB, *Mycobacterium tuberculosis* (*M. tuberculosis*); NR-MTB, non-replicating *Mycobacterium tuberculosis*; PZA, prazinaamide; R-Mtb, replicating *Mycobacterium tuberculosis*; RIF, rifampicin; RTK, receptor tyrosine kinases; SAR, structure activity relationship; SI, Selectivity index; TB, Tubercle Bacillus; TMC, Tibotec Medicinal Compound; WHO, World Health Organization; XDR-TB, Extensively drug-resistant tuberculosis.

* Corresponding author. Organic and Medicinal Chemistry Laboratory, Centre for Nano and Material Sciences, Jain University, Jain Global Campus, Jakkasandra post, Kanakapura Road, Ramanagara District, Karnataka-562112, India.

Tel.: +918027577199, 919620667075.

E-mail addresses: keriphd@gmail.com, sk.rangappa@jainuniversity.ac.in (R.S. Keri).

1. Introduction

Tuberculosis (TB) is a lung infection caused mainly by *Mycobacterium tuberculosis* (*M. tuberculosis* [MTB]). It is considered to be one of the most contagious and deadly diseases and is a major threat for public health. In combination with the HIV-1 infection TB is today amongst the biggest threat to the mankind. A large proportion of these new cases and deaths occur mostly in developing countries and the number of HIV-positive patients coinfected with MTB is constantly rising [1]. As a result, the TB situation may become even worse with the spread of HIV-1 worldwide, emergence of multi-drug (isoniazid and rifampin) resistant (MDR-TB) and the extensively drug resistant (XDR-TB) strains. Tuberculosis, also known as TB and ‘white plaque’, is caused by infection with members of the MTB complex, which includes *Mycobacterium tuberculosis* itself, *Mycobacterium africanum*, *Mycobacterium bovis*, *Mycobacterium caprae*, *Mycobacterium microti*, *Mycobacterium pinnipedii* and *Mycobacterium canettii* [2,3]. Robert Koch was the first scientist who isolated the bacteria, MTB in 1882 and got Nobel Prize for this discovery [4].

TB has been one of the deadliest diseases over the past few decades affecting nearly one-third of the world's population [5] with new infection occurring at 1% of population each year [6]. According to WHO studies, in 2011, there were 8.7 million new cases of TB (13% co-infected with HIV) and 1.4 million people died from TB including one million HIV negative people [7]. The estimated 8.8 million new cases every year correspond to 52,000 deaths per week or more than 7000 each day [8,9]. These number shows ever, are only a partial depiction of the global TB threat. More than 80% of TB patients are in the economically productive age of 15–49 years, which results in tremendous economic and social problems. It was estimated that nearly 1 billion more people will be infected with TB in the next 20 years. About 15% of that group (150 million) will exhibit symptoms of the disease, and about 3.6% (36 million) will die from TB if new disease prevention and treatment measures are not developed [10]. In 2012, nearly nine million people around the world became sick with TB disease. There were around 1.3 million TB-related deaths worldwide. Over 95% of TB deaths occur in low- and middle-income countries, and it is among the top three causes of death for women aged 15 to 44. In 2012, an estimated 530,000 children became ill with TB and 74,000 HIV-negative children died of TB [11]. These data facilitated chemists and biologist to discovery of novel drug targets, assisted the understanding of the biological phenomenon of *MTB*. Currently, the six to nine month multidrug protocol used in the treatment of TB is highly effective with drug-susceptible TB, but poor patient compliance promotes development of drug resistance [12]. Although the existing method of curing is very effective against TB, the length of treatment, the toxicity and the potential for drug-drug interactions are factors that highlight the need for new anti-TB drugs [13,14]. In addition, *MTB* is resistant to some of the first and second line drugs [15]. Therefore, effective new drugs [16] and strategies [17] are essential to treat the TB bacilli.

It has been established that heterocyclic compounds play an important role in designing new class of structural entities for medicinal applications. Among pharmacologically important heterocyclic compounds, quinoline and its derivatives are significant because of their wide spectrum of biological activities and




Fig. 1. Chemical structure and numbering of quinoline.

their presence in naturally occurring compounds. Quinoline is a heterocyclic aromatic nitrogen compound characterized by a double-ring structure that contains a benzene ring fused to pyridine at two adjacent carbon atoms (Fig. 1) [18,19]. It can also be named as, benzopyridine, benzo[b]pyridine, 1-azanaphthalene, 1-benzazine and benzazine.

In the recent time, quinoline nucleus has gathered an immense attention among chemists as well as biologists as it is one of the key building elements for many naturally occurring compounds. In particular, quinoline alkaloids are found in many different plants including Berberidaceae, Fumariaceae, Papavaraceae and Rutaceae [20–24]. Quinoline and its derivatives are important class of bioactive molecules in the field of drugs and pharmaceuticals. They exhibit significant activity against several viruses including antimalarial [25–27], antibiotic [28,29], anticancer [30], anti-inflammatory [31], antihypertensive [32], tyrosinase PDGF-RTK inhibition [33] and anti-HIV [34,35] properties. To list a few quinoline derivatives quinine (antipyretic, antimalarial, analgesic, and anti-inflammatory properties), chloroquine (antimalarial), amodiaquine (antimalarial and anti-inflammatory agent), camptothecin (DNA enzyme topoisomerase I), and saquinavir (anti-retroviral drug), which are actively used in pharmacological field are given below (Fig. 2).

This broad spectrum of biological and biochemical activities has been further facilitated by the synthetic versatility of quinoline, which allows creating a large number of structurally diverse derivatives. Quinoline has been considered a pharmacophore for




Fig. 2. A few quinoline derivatives in clinical use.




Fig. 3. Chemical structures of quinoline derivatives as anti-TB drugs.

the design of anti-TB agents. Ciprofloxacin and moxifloxacin (Fig. 3) are promising agents for the treatment of TB [36] having quinoline moiety. Quinoline based mefloquine (Fig. 3) is known for anti-tubercular activity [37–40] and its analogs have displayed moderate [41] to submicromolar [42] anti-TB activity. Tibotec Medicinal Compound 207 (TMC207) has emerged as a lead molecule out of this work and currently this compound is under phase II clinical assessment (Fig. 3). Detailed mechanistic study revealed that oligomeric (F ATPase) and proteolipic (V ATPase) subunit of ATP synthase of mycobacteria is the target of this compound. TMC207 is effective for resistant and nonresistant strains of *MTB* at MIC 0.03 $\mu\text{g/mL}$. The results of its clinical trials show that TMC207 may shorten the treatment of TB and be effective in its treatment [43–45].

Therefore, the syntheses of quinoline and its derivatives have received an increasing attention to synthetic organic chemists and biologists. It is still a challenge for the pharmaceutical chemist to develop more effective and less toxic agents to treat signs and symptoms of TB disorders. A large amount of effort has been invested in the past decade to develop quinoline-based compounds as modulator of anti-TB, which is active on different clinically approved therapeutic targets showing excellent therapeutic potency. By looking into the importance of this therapeutic area we decided to collect the published anti-TB data on quinoline (from 2010 to date), the indispensable anchor in medicinal chemistry. In this review, we have attempted to shed light and compile published reports on quinoline derivatives along with

some opinion on different approaches to help the medicinal chemists in designing future generation potent yet safer anti-TB agents.

2. Quinoline: structural requirements for anti-TB activity

From collected published data, it is found that the quinoline nucleus substituted at all position with varied substituent's has produced potent anti-TB activity. However the 5-position of the nucleus is unsubstituted. The 1-position of quinoline may be unsubstituted or substituent's may vary from alkyl and aryl groups. Among them, quinoline with ethyl substituent's, showed excellent anti-TB activity. Similarly the 2-position may be substituted with alkyl or bulky lipophilic aryl/heteroaryl groups and shows a good activity. The 3 or 4-position of the nucleus may be substituent's are more because of the conjugation and substituent's may range from functional groups like halogens, alkene linker, hydrazones, butanamide derivatives heteroaryl groups. The 6 or 7-position of the nucleus may be unsubstituted or substituent's may range from functional groups like halogens, nitro, amino, 5-nitrofuranyl, dialkylamino, 4-fluorophenoxy, dimethylamino groups and quinoline with $\text{at-}\text{CF}_3$ at 8th position, shows good anti-TB activity (Fig. 4).

3. Quinoline derivatives for treatment of tuberculosis

A spectrum of pharmacological activities exhibited by quinoline and its derivatives has been reviewed by several authors [18,46]. But, no one review published on quinoline as anti-TB agents.

In 2010, Eswaran et al., reported the synthesis of quinoline-3-carbohydrazide derivatives and evaluated for their *in vitro* anti-TB activity against *MTB H₃₇Rv*, *Mycobacterium smegmatis* (MC2), and *Mycobacterium fortuitum* by broth micro dilution assay method. The presence of substituted hydrazones and (3R)-3-amino-N,N-dimethyl-4-(phenylthio)butanamide respectively, at positions 3 and 4 of quinoline skeleton has tremendously enhanced the TB activity. Compounds (2) and (3) showed good activity against both *M. fortuitum* and *MTB H₃₇Rv* and they are more potent than isoniazid (INH) and rifampin (RIF). SAR study revealed that, the presence of $-\text{CF}_3$ group at position-8 enhanced the activity, while the introduction of $-\text{F}$ at position-6 partially lowered the activity [47]. Same research group synthesized quinoline derivatives carrying oxazolidinone ring and tested for their both preliminary and second level *in vitro* antimycobacterial activity against *MTB H₃₇Rv*, *M. smegmatis* and *M. fortuitum*, results were compared with standard anti-TB drugs INH and RIF. Compounds (4–6) were active






Fig. 4. Structural requirement around quinoline nucleus for an anti-TB activity.

at 0.625 $\mu\text{g/mL}$ against *MTB H₃₇Rv* strains. These compounds are more potent than the reference compound INH (MIC = 0.7 $\mu\text{g/mL}$) against *MTB H₃₇Rv*, 50 $\mu\text{g/mL}$ against *M. smegmatis* and 12.5 $\mu\text{g/mL}$ against *M. fortuitum*. Also, the compounds (4) and (6) exhibited promising activity against *M. smegmatis* strain at 2.5 $\mu\text{g/mL}$ concentration whereas compound (5) displayed activity at 2.5 $\mu\text{g/mL}$ against *M. fortuitum* strain. The promising activities of these compounds are mainly due to the presence of alkylamines, methyl, ethyl and acetyl substituted piperazines are responsible for their improved activity [48].


showed very good anti-TB activity against both the TB strains. It may be attributed to the presence of electron donating $-\text{CH}_3$ group, which is responsible for stabilizing the pyrazole ring, thereby making the quinoline ring more active species [49]. Trifluoromethyl-quinoline derivatives synthesized and screened for their in vitro antimycobacterial activity against *MTB H₃₇Rv*, and non-tubercular mycobacterial (NTM) species like *M. smegmatis*, and *M. fortuitum* by Resazurin assay method. The standard drugs, viz. INH and RIF were used for comparison. Compounds (8–12) displayed significant activity against *MTB H₃₇Rv* strain and (9) and




MIC(*MTB*) = 1 $\mu\text{g/mL}$
MIC(*MS*) = 10 $\mu\text{g/mL}$
MIC(*MF*) = 1 $\mu\text{g/mL}$
% GI > 95 %
IC₅₀ > 62.5 $\mu\text{g/mL}$




MIC(*MTB*) = 1 $\mu\text{g/mL}$
MIC(*MS*) = 10 $\mu\text{g/mL}$
MIC(*MF*) = 1 $\mu\text{g/mL}$
% GI > 95 %
IC₅₀ > 62.5 $\mu\text{g/mL}$




MIC(*MTB*) = 0.625 $\mu\text{g/mL}$
MIC(*MS*) = 2.5 $\mu\text{g/mL}$
MIC(*MF*) = 10 $\mu\text{g/mL}$
% GI = 95 %



MIC(*MTB*) = 0.625 $\mu\text{g/mL}$
MIC(*MS*) = 2.5 $\mu\text{g/mL}$
MIC(*MF*) = 10 $\mu\text{g/mL}$
% GI = 95 %




MIC(*MTB*) = 0.625 $\mu\text{g/mL}$
MIC(*MS*) = 10 $\mu\text{g/mL}$
MIC(*MF*) = 2.5 $\mu\text{g/mL}$
% GI = 90 %




MIC(*MTB*) = 0.625 $\mu\text{g/mL}$
MIC(*MS*) = 2.5 $\mu\text{g/mL}$
MIC(*MF*) = 10 $\mu\text{g/mL}$
% GI = 95 %

Same researcher reported, the synthesis of quinoline derivatives carrying biologically active entities viz., hydrazones, ureas, thioureas and pyrazoles and evaluated anti-TB activity against *MTB H₃₇Rv* and *MDR-TB*. Most of synthesized compounds showed good activity against *MDR-TB* strain with MIC ranging from 6.25–25 $\mu\text{g/mL}$ and were found more active than INH and RIF. The good anti-TB activity is attributed to the presence of pharmacologically active hetero-aryl groups attached to the quinoline ring. Compound (7)


(10) displayed significant activity at 1.25 mg/mL against *M. smegmatis*, Compounds (8–11) showed substantial activity against the *MDR-TB* strain at 6.25 mg/mL. The activities of these compounds could be attributed to the incorporation of heterocyclic compounds viz., substituted indoles, pyrrole, imidazole, benzotriazole and aromatic compounds with methoxy and fluoro substituents to active 4-hydroxy-8-(trifluoromethyl)quinoline-3-carbohydrazide [50].




MIC(*MTB*) = 3.12 µg/mL
 MIC(*MS*) = 6.25 µg/mL
 MIC(*MF*) = 10 µg/mL
 MIC(*MDR-TB*) = 6.25 µg/mL
 % GI = 99 %




MIC(*MTB*) = 0.625 µg/mL
 MIC(*MS*) = 10 µg/mL
 MIC(*MF*) = 10 µg/mL
 MIC(*MDR-TB*) = 6.25 µg/mL
 % GI = 95 %




MIC(*MTB*) = 0.625 µg/mL
 MIC(*MS*) = 1.25 µg/mL
 MIC(*MF*) = 10 µg/mL
 MIC(*MDR-TB*) = 6.25 µg/mL
 % GI = 95 %



MIC(*MTB*) = 0.625 µg/mL
 MIC(*MS*) = 1.25 µg/mL
 MIC(*MF*) = 10 µg/mL
 MIC(*MDR-TB*) = 6.25 µg/mL
 % GI = 95 %




MIC(*MTB*) = 0.625 µg/mL
 MIC(*MS*) = 10 µg/mL
 MIC(*MF*) = 10 µg/mL
 MIC(*MDR-TB*) = 6.25 µg/mL
 % GI = 95 %




MIC(*MTB*) = 0.625 µg/mL
 MIC(*MS*) = 10 µg/mL
 MIC(*MF*) = 10 µg/mL
 MIC(*MDR-TB*) = 6.25 µg/mL
 % GI = 95 %

Quinoline-triazoles carrying amides, sulphonamides and amidopiperazine derivatives reported as antimycobacterial agents. Compounds **(13–15)** showed promising activity against *MTB H₃₇Rv* at 0.625 µg/mL. Compound **(16)** displayed promising activity at 0.625 µg/mL against *MTB H₃₇Rv* and active at 10 µg/mL against *M. smegmatis* (*MS*) and *M. fortuitum* (*MF*). Compound **(17)** showed substantial activity at 0.625 µg/mL against *MTB* and more potent than INH against *M. smegmatis*. Overall, compounds **(13–17)** were active against *MTB*, *MS* and *MF* mycobacterial strains. The


substantial activity of these compounds may be attributed to the important pharmacophoric groups present in the molecule, viz., acetyl, methoxy, trifluoromethyl and fluoro group [51]. Fused oxazoloquinoline derivatives showed good anti-TB activity against *MTB H₃₇Rv* and compounds **(18a–b)**, **(19a–b)**, and **(20a–c)** showed very good anti-TB activity. The activity is attributed to the presence of substituted aryl group at position-2 of quinoline ring. SAR study reveals that with the introduction of 1,3-oxazole ring has tremendously increased the activity of the molecules [52].




MIC(*MTB*) = 0.625 µg/mL
 MIC(*MS*) = 10 µg/mL
 MIC(*MF*) = 10 µg/mL
 % GI = 95 %




MIC(*MTB*) = 0.625 µg/mL
 MIC(*MS*) = 5 µg/mL
 MIC(*MF*) = 10 µg/mL
 % GI = 95 %




MIC(*MTB*) = 0.625 µg/mL
 MIC(*MS*) = 10 µg/mL
 MIC(*MF*) = 5 µg/mL
 % GI = 95 %




MIC(*MTB*) = 0.625 µg/mL
 MIC(*MS*) = 10 µg/mL
 MIC(*MF*) = 10 µg/mL
 % GI = 90 %




a) R = 2-methoxy phenyl; MIC(*MTB*) = 1 µg/mL
 % GI = 99 %
 b) R = Benzyl; MIC(*MTB*) = 1 µg/mL
 % GI = 99 %




a) R = 4-fluoro phenyl; MIC(*MTB*) = 1 µg/mL
 % GI = 99 %
 b) R = 2-methoxy phenyl; MIC(*MTB*) = 1 µg/mL
 % GI = 99 %
 c) R = 3-chloro phenyl; MIC(*MTB*) = 1 µg/mL
 % GI = 99 %




MIC(*MTB*) = 0.625 µg/mL
 MIC(*MS*) = 5 µg/mL
 MIC(*MF*) = 10 µg/mL
 % GI = 90 %



a) R₁ = F, R₂ = H; MIC(*MTB*) = 1 µg/mL
 % GI = 99 %
 b) R₁ = H, R₂ = F; MIC(*MTB*) = 1 µg/mL
 % GI = 99 %




a) R = 2-methoxy phenyl; MIC(*MTB*) = 1 µg/mL
 % GI = 99 %
 b) R = Benzyl; MIC(*MTB*) = 1 µg/mL
 % GI = 99 %



a) R = 4-fluoro phenyl; MIC(*MTB*) = 1 µg/mL
 % GI = 99 %
 b) R = 2-methoxy phenyl; MIC(*MTB*) = 1 µg/mL
 % GI = 99 %
 c) R = 3-chloro phenyl; MIC(*MTB*) = 1 µg/mL
 % GI = 99 %


Aryl and thiophenyl-quinolinones derivative reported as antimycobacterial agents against *MTB H₃₇Rv*. Compounds (**21a-b**) inhibited *MTB* with MIC of 3.13 µg/mL, compared to one of the first line anti-TB drug ethambutol (MIC 3.13 µg/mL), are found equally active. SAR study revealed that, introduction lipophobic hydroxymethyl groups profoundly decreased their MIC values (>25 µg/mL). In comparison with aryl tethered dihydro-6H-quinolin-5-ones, thiophenyl tethered dihydro-6H-quinolin-5-ones are better placed to show potent anti-TB activity. Dihydro-6H-quinolin-5-ones derived from dimedone are structurally better correlates and showed potent activity against *MTB* [53]. Thienoquinolines derivatives were synthesized using by the Friedlander annulations and screened for their *in vitro* antimycobacterial activity against *MTB* and *MDR-TB*. All the synthesized compounds showed promising *in vitro* activity against *MTB* with MIC in the range of 0.90–36.82 µM and against

MDR-TB with MIC ranging from 0.95 to >15.30 µM. Compound (**22**) is the most active compound with MIC of 0.90 µM against *MTB*. This compound is five and eight times more active than ciprofloxacin and ethambutol respectively, while three and eight times less active than INH and RIF respectively. Compounds (**23**) displayed maximum activity *in vitro* with MIC of 0.95 µM against *MDR-TB*, being 40, 12 and four times more potent than ciprofloxacin, INH and RIF respectively. The thienoquinolines with chlorine at the 7-position displayed significantly greater activity than the corresponding unsubstituted compounds. It is found that the thienoquinolines with electron withdrawing groups like halogen or nitro group in the phenyl ring of the thiophene ring of 3 showed better activity than that with electron donating groups like alkyl. Disubstitution in the phenyl ring enhances the activity many-fold relative to monosubstitution [54].




(21a-b)

a) R = Cl; MIC(*Mtb* H₃₇Rv) = 3.13 µg/mL
 b) R = Br; MIC(*Mtb* H₃₇Rv) = 3.13 µg/mL



(22)


MIC(*MTB*) = 0.90 µg/mL
 MIC(*MDR-TB*) = 1.76 µg/mL



(23)


MIC(*MTB*) = 1.86 µg/mL
 MIC(*MDR-TB*) = 0.95 µg/mL

Pyranoquinoline analogues synthesized using SnCl₂·2H₂O as a catalyst by one-pot Povarov reaction and evaluated for their in vitro antimycobacterial activity against *MTB*, they show activity MIC ranging from 3.13–25.0 µg/mL. Among them, compounds (24–26) inhibited *MTB* with MIC 3.13 µg/mL. When compared to one of the first line anti-TB drug ethambutol (MIC 3.13 µg/mL), these three compounds are found equally active as ethambutol. From SAR, halo substituent's is needed for hexahydro-2H-pyrano[3,2-c]quinolines to be active against *MTB* and among them fluoro analogues with trans geometry are structurally better correlates and showed better activity [55].




(24)

MIC(*MTB*) = 3.13 µg/mL



(25)

MIC(*MTB*) = 3.13 µg/mL




(26)

MIC(*MTB*) = 3.13 µg/mL


Quinoline bearing pyrimidine motifs synthesized by Desai et al., and evaluated for their in vitro antimycobacterial activity against *MTB* strain by using Lowenstein-Jensen slope method. Compound (27) having methoxy at 4th position, at the phenyl ring of pyrimidine substitution showed most promising anti-TB

activity with MIC of 0.20 µg/mL. SAR study reveals, substituent's with electron donating groups such as methoxy, methyl and hydroxy at *para* position of phenyl ring demonstrated high inhibitory activity against *MTB* as compared *meta* substituted derivatives, indicating that the electronic properties of the substituents have major influence on the antimycobacterial activity [56]. Zhou et al. reported the synthesis of dihydroartemisinin-fluoroquinolone conjugates as potential anti-TB agents. Compound (28a) exhibited the good inhibitory activity (MIC = 0.0625 µg/mL), which was comparable to the positive control moxifloxacin and even stronger than ofloxacin.



(27)

MIC(*MTB*) = 0.20 µg/mL
 % GI = 99 %




(28a-b)


a) n=2; MIC(*MTB*) = 0.0625 µg/mL
 b) n=3; MIC(*MTB*) = 0.125 µg/mL

The ferrocene-based hydrazones showed better antimycobacterial activity against *MTB*. Especially, quinoline-ferrocene hybrid (**29**) exhibited significant activity ($\text{MIC} = 2.5\text{--}5\text{ }\mu\text{g/mL}$) against *Mtb*, it is comparable to the one of EMB ($\text{MIC} = 2.5\text{ }\mu\text{g/mL}$). The good anti-TB activity of (**29**) may be attributed to the presence of the quinoline ring [58]. Azole-fused indeno[2,1-*c*]quinolines synthesized and tested against *MTB* $H_{37}\text{Rv}$. Compound (**30**) and (**31**) inhibited growth of *MTB* very effectively at $\text{MIC} < 0.39\text{ }\mu\text{g mL}^{-1}$ ($0.89\text{ }\mu\text{M}$ and $1\text{ }\mu\text{M}$ respectively, which is comparable to that of the existing front-line drug INH ($\text{MIC } 0.25\text{ }\mu\text{g/mL}$) [59].


Mefloquine-isoxazole carboxylic acid esters synthesized and screened for anti-TB activity against *MTB* $H_{37}\text{Rv}$ using the microplate MABA and low oxygen recovery assay (LORA). Compound with an alkene linker between the 4-position of the quinoline ring and the 5-position of the isoxazole, had markedly improved anti-TB activity and the trans isomer (**35**) (MABA $\text{MIC} = 0.2\text{ }\mu\text{M}$) was found to have a MIC only 2-fold higher than RIF, the most active anti-TB drug used today. SAR study reveals, an alkene linker is important for activity, and a transalkene is more favored. The position(s) of the substitution(s) on the quinoline ring



$\text{MIC}(\text{MTB}) = 2.5\text{--}5\text{ }\mu\text{g/mL}$




$\text{MIC}(\text{MTB}) < 0.39\text{ }\mu\text{g/mL}$
% GI = 97 %




$\text{MIC}(\text{MTB}) < 0.39\text{ }\mu\text{g/mL}$
% GI = 97 %

Gonec et al. have reported quinoline-2-carboxamides as anti-TB agents. *N*-(2-Phenylethyl) quinoline-2-carboxamide (**32a**), *N*-cycloheptylquinoline-2-carboxamide (**32b**) and *N*-cyclohexylquinoline-2-carboxamide (**32c**) showed high activity against *MTB* and 2-(pyrrolidin-1-ylcarbonyl)quinoline (**32d**) showed high activity against *M. kansasii* and *M. avium paratuberculosis* and these are comparable with or higher than the standards INH or pyrazinamide [60]. Quinoline-based azetidinone and thiazolidinone analogues were synthesized and screened against mycobacteria. Compound (**33**) with 2-amino 5-methyl thiazole moiety to the azetidinone class as well as (**34**) with electron-withdrawing and strong electronegative fluoro substituent within the thiazolidinone class exhibited good inhibitory potential at $12.5\text{ }\mu\text{g/mL}$ of MIC against *MTB* $H_{37}\text{Rv}$ strain. The inhibitory potential of the said derivatives was half-fold as compared to standard drugs. Azetidinones and thiazolidinones bearing various amines containing halogen(s) such as chloro or fluoro and nitro functional groups have showed high potency [61,62].


may also affect the activity to some extent, while the electron effect of the substitution(s) is less important [63]. Quinoline with different heterocyclic compounds like oxadiazoles, pyrazolines, and pyrazoles moieties and screened for antimycobacterial activity performed with *M. smegmatis* strain using growth inhibition assay by turbidimetry. Heteroaryl substitutions at 5th position of oxadiazole ring, in compounds (**36**), and (**37**) led to better antimycobacterial activity and the free pyrazole with quinolinyl (**38**) have shows the good anti-TB activity. The aryl ring substitutions, with fluoro, methyl, methoxy have showed good activity [64]. Tukulula et al. reported synthesis of quinoline-tetrazole derivatives and evaluated *in vitro* antimycobacterial activities using *MTB* $H_{37}\text{Rv}$ strain in MABA and LORA assays. Compound (**39**), inhibited 94% ($\text{MIC}_{90} = 123.2\text{ }\mu\text{M}$) of non-replicating bacteria and (**40**), which showed 98% inhibition ($\text{MIC}_{90} = 92.5\text{ }\mu\text{M}$) of replicating bacteria. SAR study reveals; the replacement of the hydroxyl group of 4-quinoline was accompanied by complete loss of activity [65].




$\text{MIC}(\text{MTB}) = 109\text{ }\mu\text{mol/L}$




$\text{MIC}(\text{MTB}) = 111\text{ }\mu\text{mol/L}$




$\text{MIC}(\text{MTB}) = 125\text{ }\mu\text{mol/L}$




$\text{MIC}(\text{MK}) = 111\text{ }\mu\text{mol/L}$
 $\text{MIC}(\text{MAP}) = 111\text{ }\mu\text{mol/L}$




$\text{MIC}(\text{MTB}) = 12.5\text{ }\mu\text{g/mL}$
% GI = 99 %




$\text{MIC}(\text{MTB}) = 12.5\text{ }\mu\text{g/mL}$
% GI = 99 %




MIC(*MTB*) = 0.2 μ M (MABA)
= 2.6 μ M (LORA)




MIC(*MS*) = 22.71 μ g/mL




MIC(*MS*) = 16.83 μ g/mL



MIC(*MS*) = 14.66 μ g/mL




MIC₉₀(*MTB*) = 123.2 μ M (LORA)
% GI = 94 %




MIC₉₀(*MTB*) = 92.5 μ M (MABA)
% GI = 98 %

3-Heteroarylthioquinolines derivatives were synthesized and screened for their *in vitro* antimycobacterial activity against *MTB*, some of the compounds showed good *in vitro* activity against *MTB* with MIC ranging from 3.2– 55.9 μ M. Compounds **(41a–c)** and **(42)** inhibited *MTB* with MIC less than 6.5 μ M and were more potent than the first line anti-TB drug, ethambutol (MIC = 7.6 μ M). When compared to ciprofloxacin (MIC = 4.7 μ M), two compounds **(41a)** (MIC = 3.5 μ M) and **(41b)** (MIC = 3.2 μ M) were found to be more potent against *MTB*. Quinolines with sulfur heterocyclic unit at position 3 are found to be more active than quinolines having a heterocycle with four nitrogen atoms at this position. The presence of halogens in the aryl ring enhances the antimycobacterial activity


as seen from the MIC values of compound **(41a)** and compound **(41b)** [66]. Quinoline with piperazinyl/piperidinyl-s-triazines derivatives reported as anti-TB agents by Patel and co-workers against *MTB H₃₇Rv*. Compound **(43)** with substitution of two methyl groups at the 3rd and 5th position of the piperidine ring, **(44)** and **(45)** bearing trifluoromethylation and trimethoxy substitution to the phenyl ring of piperazine base coupled to the nucleus exhibited highest inhibition (99%) at a constant concentration level (6.25 μ g/mL) against *MTB*. In Lowenstein Jensen MIC method, **(43)** was displayed inhibition of *MTB* completely (99%) at the MIC of 3.12 μ g/mL and this compound better efficacy, than the standard drug, pyrazinamide [67].




a) R = Cl; MIC(*MTB*) = 3.5 μ M
b) R = Br; MIC(*MTB*) = 3.2 μ M
c) R = Ph; MIC(*MTB*) = 6.4 μ M




MIC(*MTB*) = 5.9 μ M



% inhibition (*MTB*) = 99 %
(6.25 μ g/mL)




% inhibition (*MTB*) = 99 % (6.25 μ g/mL)




% inhibition (*MTB*) = 99 % (6.25 μ g/mL)

Benzofuro[2,3-b]quinoline derivatives synthesized for evaluated for their anti-TB against *MTB*. The less bulky methoxy derivative (**46**), 11-aminated derivatives in which the less bulky methylamino derivative (**47**), and a tertiary amine derivative (**48**) exhibited significant activities against the growth of *MT* (MIC values of $< 0.20 \mu\text{g/mL}$) [68]. Tetrazolo[1,5-a]quinoline based tetrasubstituted imidazole derivatives synthesized and evaluated for their activity against *MTB H₃₇Rv* strain. Compounds (**49a**) and (**49b**) showed best activity against *MTB*. The SAR study reveals, change in the substituent might also affect the anti-TB activity, compounds (**49a**) and (**49b**) carrying electron negative groups on quinoline ring displayed excellent anti-TB activity against tubercular strains [69].




(46)




(47)

MIC(*MTB*)
 $< 0.20 \mu\text{g/mL}$



(48)

MIC(*MTB*) $< 0.20 \mu\text{g/mL}$




(49a-b)

MIC(*MTB*) $< 0.20 \mu\text{g/mL}$


a) R = CH₃; % inhibition (*MTB*) = 99 % (25 $\mu\text{g/mL}$)
b) R = OCH₃; % inhibition (*MTB*) = 98 % (62.5 $\mu\text{g/mL}$)

Raj et al. reported synthesis of quinoline- β -lactam hybrids and tested for their anti-TB activities. Most of the compounds exhibited better anti-TB activity than standard drugs ethionamide (up to three times) and cephalexin (up to five times). Compounds (**50–52**) showed good anti-TB activity at MIC 5 $\mu\text{g/mL}$ [70]. Quinoline coupled 1,2,3-triazoles compounds synthesized by 'click chemistry' and screened for anti-TB activity against *MTB* by luciferase reporter phage (LRP) assay. Quinoline coupled triazole sugar hybrid, (**53**) is the exhibit potent anti-TB activity against *MTB* strain with 76.41% and 78.37% reduction calculated based on percentage reduction in relative light units at 5 and 25 $\mu\text{g/mL}$, respectively. SAR study reveals, introduction of the phenyl ring at the C-4 position of triazole induced reasonable inhibition against *MTB*. It was apparent that the introduction of bulky and lipophilic substituent such as benzene exhibits potent inhibition against *MTB*. However, introduction of small hydrophilic substituent such as hydroxyl linked to the C-4 of triazole via methylene showed dramatic loss in potent inhibition against *MTB*. Pentoses or hexoses in the chair conformation with 3,4-trans diacetyl orientation such as quinoline glycoconjugate of D-galactose (**53**) (3S,4R), were generally more potent inhibitor than those with corresponding 3,4-cis-diacetyl orientation (3S,4S) among the saccharide coupled compounds [71].




(50)

MIC(*MTB*) = 5 $\mu\text{g/mL}$




(51)

MIC(*MTB*) = 5 $\mu\text{g/mL}$



(52)

MIC(*MTB*) = 5 $\mu\text{g/mL}$




(53)


MIC(*MTB*) = 5 $\mu\text{g/mL}$

Gonçalves et al. reported synthesis of mefloquine-oxazolidine derivatives as anti-TB agents against *MTB H₃₇Rv*. The most active of the compounds was the dimethoxy derivative (**54**) and nitro derivative (**55**) with a MIC value of 11.9 μ M and 12.1 μ M respectively, showed 2.7 times more active than mefloquine (MIC = 33 μ M), with a better tuberculostatic activity than the first line tuberculostatic agent ethambutol (MIC = 15.9). SAR study reveals, the longer alkyl chain length scarcely affected the biological activity [72]. Tripartite hybrids from pharmacophores 7-chloroquinoline (CQ), ethylenediamine spacer (ETH) and phenylurea as thiourea bioisostere (ISO) synthesized and tested in vitro for their antimycobacterial activity against *MTB H₃₇Rv*. Compounds (**56**) and (**57**) exhibited high mycobactericidal activity, these structures were found to be the most potent compounds with MIC's of 4 and 2 μ g/mL, respectively. Compound (**57**) was two fold more potent than reference drugs EMB and isoxyl (MIC 4 μ g/mL), whereas compound (**56**) was as active as these two antimycobacterial drugs [73].


Aldehyde-arylhydrazone-oxoquinoline derivatives were synthesized and evaluated for their anti-TB activity against *MTB H₃₇Rv* strain. The 1-ethyl-N'-[(1E)-(5-nitro-2-furyl)methylene]-4-oxo-1,4-dihydroquinoline-3-carbohydrazide (**58**) was effective against *MTB*. The determination of the MIC of (**58**) showed a value (6.25 μ g/mL) better than current drugs in the market. The insertion of the ethyl substituent in the nitrogen atom of quinoline nucleus as possibly involved in the lack of anti-TB activity of most compounds probably due to steric hindrance effects. The ethyl-quinoline ring without any substitution and the 5-nitro-2-furyl group at Ar in the active compound (**58**) seem to be important for the anti-TB activity [74]. Substituted quinoliny chalcones, quinoliny pyrimidines, and pyridines were synthesized and evaluated for their anti-TB activity in vitro against *MTB H₃₇Rv*. The analogs (**59–61**) exhibit promising activity and inhibit the growth of mycobacterium to 99%, 97%, and 99%, respectively, at a concentration of 6.25 μ g/mL. Analogs (**62,63**) were most active of the series inhibiting the growth of drug-sensitive bacteria to 99% at




MIC(*MTB*) = 11.9 μ g/mL
MIC(*MDR-TB*) = 11.9 μ g/mL



MIC(*MTB*) = 12.1 μ g/mL
MIC(*MDR-TB*) = 12.1 μ g/mL




MIC(*MTB*) = 4 μ g/mL




MIC(*MTB*) = 2 μ g/mL


4.25 and 6.25 μ g/mL concentration, respectively. Compound with fluorine substituent's in the phenyl ring enhances the anti-TB activity against *MTB H₃₇RV* in case of chalcone and cyanopyridine derivatives [75].




(58)

MIC(*MTB*) = 6.25 μ g/mL


(59)

% inhibition (*MTB*) = 99 % (6.25 μ g/mL)


(60)

% inhibition (*MTB*) = 97 % (5 μ g/mL)

(61)

% inhibition (*MTB*) = 99 % (6.25 μ g/mL)

(62)


% inhibition (*MTB*) = 99 % (4.5 μ g/mL)

(63)


% inhibition (*MTB*) = 99 % (6.25 μ g/mL)

Fluorine substituted biquinoline derivatives synthesized and screened for their in vitro anti-TB activity against *MTB H₃₇Rv* strain was determined using Lowenstein-Jensen medium. Compound **(64)** with 91% inhibition displayed excellent activity against *MTB H₃₇Rv*. Authors reported, activity because of the combination effect of electron releasing -OCH₃ and, electron withdrawing -F and -CF₃ groups [76]. Quinoline-3-carbohydrazide derivatives synthesized by Garudachari et al., screened against *M. smegmatis*. Compounds, **(65a-b)** showed the lowest MIC value of 6.25 μ g/mL against *M. smegmatis* indicating these compounds can be possible future anti-TB agents. The activity increases with increase of electron withdrawing group on pyrazole carbohydrazide and electron donating groups on phenyl carbohydrazide at third position of quinoline [77].

Carmo et al. reported the synthesis of 4-aminoquinoline analogues and their platinum(II) complexes as new antileishmanial and antitubercular agents. Compounds **(66)**, and **(67)** inhibit promising activity against *MTB*, with MIC values ranging from 15.6–12.5 μ g/mL, comparable to the “first and second line” drugs used to treat tuberculosis. The platinum complexes not show good activity, when compared without metal. SAR study reveals; the addition of certain mono- and di-alkynes to the intermediate amino compounds increases their antitubercular activity [78]. Recently, Nazarenko et al. patented some chloroquinoline derivatives as anti-TB agents. The compound 5,6,7-substituted 1-(2-chloroquinolin-3-yl)-4-dimethylamino-2-(naphthalen-1-yl)-1-phenylbutan-2-ols **(68)** showed excellent anti-TB activity, when compared with standard anti-TB drugs [79].




(64)

% inhibition (*MTB*) = 91 % (25 μ g/mL)


(65a-b)

a) R = NO₂; MIC (*MS*) = 6.25 μ g/mLb) R = CH₃; MIC (*MS*) = 6.25 μ g/mL




(66)

MIC(MTB) = 15.6 µg/mL



(67)

MIC(MTB) = 12.5 µg/mL



(68)

R₁ = H, MeO; R₂, R₃ = H, MeO, O(CH₂)_nO

4. Conclusions and future aspects

Tubercular infections pose a continuous and serious threat to human health and life in recent years. There has been an increased use of antitubercular agents and has resulted in the development of resistance. This has given rise to search for molecules acting on a novel target or a multi targeted combination therapy. With the increase in the number of new compounds screened against mycobacteria, the opportunity exists to develop a novel drug for the cure and complete eradication of TB. Numerous outstanding achievements revealed that quinoline-based compounds possess extensively potential anti-TB agents. To further optimize the full potential of quinoline compounds, the SAR-based study will likely continue to play an important role. It is highly likely that optimized quinoline compounds with excellent potency and little side effects will continue to be created. Some of these quinoline compounds will undoubtedly be used as first and second line anti-TB therapeutic agents in the near future. Further research in this field will bring innovative pharmaceutical developments with a considerable spectrum of use.

Disclosure of interest

The authors declare that they have no conflicts of interest concerning this article.

Acknowledgements

Authors acknowledge Jain University, Bangalore for the financial support.

References

- [1] Brooks JT, Kaplan JE, Holmes KK, Benson C, Pau A, Masur H. HIV-associated opportunistic infections- going, going, but not gone: the continued need for prevention and treatment guidelines. *Clin Infect Dis* 2009;48:609–11.
- [2] Grange JM. In: Schaaf S, Zumla AL, editors. *Tuberculosis: a comprehensive clinical reference*. Saunders; 2009. p. 44–59.
- [3] Dye C, Scheele S, Dolin P, Pathania V, Ravigione MC. Consensus statement. Global burden of tuberculosis: estimated incidence, prevalence, and mortality by country. WHO global surveillance and monitoring project. *JAMA* 1999;282:677–86.
- [4] Nobel Foundation. The Nobel Prize in Physiology or Medicine; 1905.
- [5] World Health Organization (WHO). Tuberculosis fact sheet; 2009.
- [6] World Health Organisation. Global tuberculosis report; 2012.
- [7] World Health Organization (WHO). Tuberculosis fact sheet; 2010.
- [8] Okada M, Kobayashi K. Recent progress in mycobacteriology. *Kekkaku* 2007;82(10):783–99.
- [9] World Health Organization (WHO) on TB epidemic. Global TB Programme. WHO Geneva; 1997.
- [10] World Health Organisation. Tuberculosis, Fact Sheet No. 104; 2007.
- [11] World Health Organisation. Global tuberculosis report; 2013.
- [12] Ginsberg AM, Spigelman M. Challenges in tuberculosis drug research and development. *Nat Med* 2007;13:290–4.
- [13] Dye C, Williams BG. The population dynamics and control of tuberculosis. *Science* 2010;328:856–61.
- [14] Burman WJ. Development of tuberculosis treatment in the 21st century. *Clin Infect Dis* 2010;50:S165–72.
- [15] LoBue P. Extensively drug-resistant tuberculosis. *Curr Opin Infect Dis* 2009;22:167–73.
- [16] Zhang Y, Post-Martens K, Denkin S. New drug candidates and therapeutic targets for tuberculosis therapy. *Drug Discov Today* 2006;11:21–7.
- [17] Brown ED, Wright GD. New targets and screening approaches in antimicrobial drug discovery. *Chem Rev* 2005;105:759–74.
- [18] Solomon VR, Lee H. Quinoline as a privileged scaffold in cancer drug discovery. *Curr Med Chem* 2011;18:1488–508.
- [19] Manske RH. The chemistry of quinolines. *Chem Rev* 1942;30:113–44.
- [20] Prescott TAK, Sadler IH, Kiapranis R, Maciver SK. Lunacridine from *Lunasia amara* is a DNA intercalating topoisomerase II inhibitor. *J Ethnopharmacol* 2007;109:289–94.
- [21] Srivastava V, Negi AS, Kumar JK, Gupta MM, Khanuja SPS. Plant based anticancer molecules: a chemical and biological profile of some important leads. *Bioorg Med Chem* 2005;21:5892–908.
- [22] Canel C, Moraes RM, Dayan FE, Ferreira D. Molecules of interest: podophyllo-toxin. *Phytochemistry* 2000;54:115–20.
- [23] Du W. Towards new anticancer drugs: a decade of advances in synthesis of camptothecins and related alkaloids. *Tetrahedron* 2003;59:8649–87.
- [24] Byler KG, Wang C, Setzer WN. Quinoline alkaloids as intercalative topoisomerase inhibitors. *J Mol Model* 2009;15:1417–26.
- [25] LaMontagne MP, Markovac AMS, Sami Khan M. Antimalarials. 13. 5-Alkoxy analogs of 4-methylprimaquine. *J Med Chem* 1982;25:964–8.

- [26] LaMontagne MP, Blumbergs P, Strube RE. Antimalarials. 14. 5-(Aryloxy)-4-methylprimaquine analogs. A highly effective series of blood and tissue schizonticidal agents. *J Med Chem* 1982;25:1094–7.
- [27] Nasveld P, Kitchener S. Treatment of acute vivax malaria with tafenoquine. *Trans Royal Soc Trop Med Hyg* 2005;99:2–5.
- [28] Mahamoud A, Chevalier J, Davin-Regli A, Barbe J, Pages JM. Quinoline derivatives as promising inhibitors of antibiotic efflux pump in multidrug resistant *Enterobacter aerogenes* isolates. *Curr Drug Targets* 2006;7:843–7.
- [29] Eswaran S, Adhikari AV, Shetty NS. Synthesis and antimicrobial activities of novel quinoline derivatives carrying 1,2,4-triazole moiety. *Eur J Med Chem* 2009;44:4637–47.
- [30] Denny WA, Wilson WR, Ware DC, Atwell GJ, Milbank JB, Stevenson RJ. Anticancer 2,3-dihydro-1H-pyrrolo[3,2-f]quinoline complexes of cobalt and chromium; 2006, US Patent 7064117B2.
- [31] Leatham PA, Bird HA, Wright V, Seymour D, Gordon A. A double blind study of antrafenine, naproxen and placebo in osteoarthritis. *Eur J Rheumatol Inflamm* 1983;6:209–11.
- [32] Muruganathan N, Sivakumar R, Anbalagan N, Gunasekaran V, Leonard JT. Synthesis, anticonvulsant and antihypertensive activities of 8-substituted quinoline derivatives. *Biol Pharm Bull* 2004;27:1683–7.
- [33] Maguire MP, Sheets KR, McVety K, Spada AP, Zilberstein A. A new series of PDGF receptor tyrosine kinase inhibitors: 3-substituted quinoline derivatives. *J Med Chem* 1994;37:2129–37.
- [34] Wilson WD, Zhao M, Patterson SE, Wydra RL, Janda L, Strekowski L. Design of RNA interactive anti-HIV agents: unfused aromatic intercalators. *Med Chem Res* 1992;2:102–10.
- [35] Strekowski L, Mokrosz JL, Honkan VA, Czarny A, Cegla MT, Patterson SE, et al. Synthesis and quantitative structure-activity relationship analysis of 2-(aryl or heteroaryl)quinolin-4-amines, a new class of anti-HIV-1 agents. *J Med Chem* 1991;34:1739–46.
- [36] (a) Guidelines for the Management of Drug-Resistant Tuberculosis WHO/TB/96-210 (Rev.1). Geneva: World Health Organization; 1997;
(b) Ginsburg AS, Grosset JH, Bishai WR. Fluoroquinolones, tuberculosis, and resistance. *Lancet Infect Dis* 2003;3:432–42.
- [37] Jayaprakash S, Iso Y, Wan B, Franzblau SG, Kozikowski AP. Synthesis, and SAR studies of mefloquine-based ligands as potential antituberculosis agents. *ChemMed Chem* 2006;1:593–7.
- [38] Bermudez LE, Kolonoski P, Wu M, Aralar PA, Inderlied CB, Young LS. Mefloquine is active *in vitro* and *in vivo* against *Mycobacterium avium* complex. *Antimicrob Agents Chemother* 1999;43:1870–4.
- [39] Bermudez LE, Kolonoski P, Seitz LE, Petrofsky M, Reynolds R, Wu M, et al. Moxifloxacin for treatment of murine *mycobacterium avium* complex disease. *Antimicrob Agents Chemother* 2004;48:3556–8.
- [40] Bermudez LE, Kolonoski P, Petrofsky M, Wu M, Inderlied CB, Young LS. Mefloquine, moxifloxacin and ethambutol are triple-drug alternative to macrolide-containing regimens for treatment of *Mycobacterium avium* disease. *J Infect Dis* 2003;187:1977–80.
- [41] Mao J, Wang Y, Wan B, Kozikowski AP, Franzblau SG. Design, synthesis, and pharmacological evaluation of mefloquine-based ligands as novel antituberculosis agents. *ChemMed Chem* 2007;2:1624–30.
- [42] Pieroni M, Lilienkamp A, Wan B, Wang Y, Franzblau SG, Kozikowski AP. Synthesis, biological evaluation, and structure-activity relationships for 5-[(E)-2-arylethenyl]-3-isoxazolecarboxylic acid alkyl ester derivatives as valuable antitubercular chemotypes. *J Med Chem* 2009;52:6287–96.
- [43] Andries K, Verhasselt P, Guillemont J, Goehlmann HWH, Neefs JM, Winkler H, et al. A diarylquinoline drug active on the ATP synthase of *Mycobacterium tuberculosis*. *Science* 2005;307:223–7.
- [44] Koul A, Dendouga N, Vergauwen K, Molenberghs B, Vranckx L, Willebrords R, et al. Diarylquinolines target subunit c of mycobacterial ATP synthase. *Nat Chem Biol* 2007;3:323–4.
- [45] Rustomjee R, Diacon AH, Allen J, Venter A, Reddy C, Patientia RF, et al. Early bactericidal activity and pharmacokinetics of the diarylquinoline TMC207 in treatment of pulmonary tuberculosis. *Antimicrob Agents Chemother* 2008;52:2831–5.
- [46] (a) Kumar S, Bawa S, Gupta H. Biological activities of quinoline derivatives. *Mini Rev Med Chem* 2009;9(14):1648–54;
(b) Marella A, Tanwar OP, Saha R, Ali MR, Srivastava S, Akhter M, et al. A versatile heterocyclic. *Saudi Pharm J* 2012. <http://dx.doi.org/10.1016/j.jsps.2012.03.002> [in press].
- [47] Eswaran S, Adhikari AV, Pal NK, Chowdhury IH. Design and synthesis of some new quinoline-3-carbohydrazide derivatives as potential antimycobacterial agents. *Bioorg Med Chem Lett* 2010;20:1040–4.
- [48] Thomas KD, Adhikari AV, Chowdhury IH, Sandeep T, Mahmood R, Bhattacharya B, et al. Design, synthesis and docking studies of quinoline-oxazolidinone hybrid molecules and their antitubercular properties. *Eur J Med Chem* 2011;46:4834–45.
- [49] Eswaran S, Adhikari AV, Chowdhury IH, Pal NK, Thomas KD. New quinoline derivatives: synthesis and investigation of antibacterial and antituberculosis properties. *Eur J Med Chem* 2010;45:3374–83.
- [50] Thomas KD, Adhikari AV, Telkar S, Chowdhury IH, Mahmood R, Pal NK, et al. Design, synthesis and docking studies of new quinoline-3-carbohydrazide derivatives as antitubercular agents. *Eur J Med Chem* 2011;46:5283–92.
- [51] Thomas KD, Adhikari AV, Telkar S, Chowdhury IH, Sumesh E, Pal NK. New quinolin-4-yl-1,2,3-triazoles carrying amides, sulphonamides and amidopiperazines as potential antitubercular agents. *Eur J Med Chem* 2011;46:2503–12.
- [52] Eswaran S, Adhikari AV, Ajay Kumar R. New 1,3-oxazolo[4,5-c]quinoline derivatives: synthesis and evaluation of antibacterial and antituberculosis properties. *Eur J Med Chem* 2010;45:957–66.
- [53] Kantevari S, Patpi SR, Sridhar B, Yogeewari P, Sriram D. Synthesis and antitubercular evaluation of novel substituted aryl and thiophenyl tethered dihydro-6H-quinolin-5-ones. *Bioorg Med Chem Lett* 2011;21:1214–7.
- [54] Balamurugan K, Jeyachandran V, Perumal S, Manjashetty TH, Yogeewari P, Sriram D. A microwave-assisted, facile, regioselective Friedländer synthesis and antitubercular evaluation of 2,9-diaryl-2,3-dihydrothieno-[3,2-b]quinolines. *Eur J Med Chem* 2010;45:682–8.
- [55] Kantevari S, Yempala T, Surineni G, Sridhar B, Yogeewari P, Sriram D. Synthesis and antitubercular evaluation of novel dibenzo[b,d]furan and 9-methyl-9H-carbazole derived hexahydro-2H-pyrano[3,2-c]quinolines via Povarov reaction. *Eur J Med Chem* 2011;46:4827–33.
- [56] Desai NC, Kotadiya GM, Trivedi AR. Studies on molecular properties prediction, antitubercular and antimicrobial activities of novel quinoline based pyrimidine motifs. *Bioorg Med Chem Lett* 2014. <http://dx.doi.org/10.1016/j.bmcl.2014.05.002> [in press].
- [57] Zhou FW, Lei HS, Fan L, Jiang L, Liu J, Peng XM, et al. Design, synthesis, and biological evaluation of dihydroartemisinin-fluoroquinolone conjugates as a novel type of potential antitubercular agents. *Bioorg Med Chem Lett* 2014;24:1912–7.
- [58] Mahajan A, Kremer L, Louw S, Guéradel Y, Chibale K, Biot C. Synthesis and *in vitro* antitubercular activity of ferrocene-based hydrazones. *Bioorg Med Chem Lett* 2011;21:2866–8.
- [59] Upadhyaya RS, Shinde PD, Sayyed AY, Kadam SA, Bawane AN, Poddar A, et al. Synthesis and structure of azole-fused indeno[2,1-c]quinolines and their antimycobacterial properties. *Org Biomol Chem* 2010;8:5661–73.
- [60] Gonet T, Bobal P, Suján J, Pesko M, Guo J, Kralova K, et al. Investigating the spectrum of biological activity of substituted quinoline-2-carboxamides and their isosteres. *Molecules* 2012;17:613–44.
- [61] Mistry BM, Jauhari S. Quinoline-based azetidinone and thiazolidinone analogs as antimicrobial and antituberculosis agents. *Med Chem Res* 2013;22(2):647–58.
- [62] Mistry Bhupendra M, Jauhari Smita. Synthesis and *in vitro* antimicrobial and anti-tubercular evaluation of some quinoline-based azetidinone and thiazolidinone analogues. *Med Chem Res* 2013;22(2):635–46.
- [63] Mao J, Yuan H, Wang Y, Wan B, Pieroni M, Huang Q, et al. From serendipity to rational antituberculosis drug discovery of mefloquine-isoxazole carboxylic acid esters. *J Med Chem* 2009;52:6966–78.
- [64] Rachakonda V, Alla M, Kotipalli SS, Ummani R. Design, diversity-oriented synthesis and structure activity relationship studies of quinolinyl heterocycles as antimycobacterial agents. *Eur J Med Chem* 2013;70:536–47.
- [65] Tukulula M, Little S, Gut J, Rosenthal PJ, Wan B, Franzblau SG, et al. The design, synthesis, *in silico* ADME profiling, antiparasmodial and antimycobacterial evaluation of new arylamino quinoline derivatives. *Eur J Med Chem* 2012;57:259–67.
- [66] Chitra S, Paul N, Muthusubramanian S, Manisankar P, Yogeewari P, Sriram D. Synthesis of 3-heteroarylthioquinoline derivatives and their *in vitro* antituberculosis and cytotoxicity studies. *Eur J Med Chem* 2011;46:4897–903.
- [67] Patel RV, Kumari P, Rajani DP, Chikhaliya KH. Synthesis and studies of novel 2-(4-cyano-3-trifluoromethylphenyl amino)-4-(quinoline-4-yloxy)-6-(piperazinyl/piperidinyl)-s-triazines as potential antimicrobial, antimycobacterial and anticancer agents. *Eur J Med Chem* 2011;46:4354–65.
- [68] Yang CL, Tseng CH, Chen YL, Lu CM, Kao CL, Wu MH, et al. Identification of benzofuro[2,3-b]quinoline derivatives as a new class of antituberculosis agents. *Eur J Med Chem* 2010;45:602–7.
- [69] Mungra DC, Kathrotiya HG, Ladani NK, Patel MP, Patel RG. Molecular iodine catalyzed synthesis of tetrazolo[1,5-a]-quinoline based imidazoles as a new class of antimicrobial and antituberculosis agents. *Chinese Chem Lett* 2012;23:1367–70.
- [70] Raj R, Biot C, Carrere-Kremer S, Kremer L, Guerardel Y, Gut J, et al. 4-aminoquinoline-β-lactam conjugates: synthesis, antimalarial, and antitubercular evaluation. *Chem Biol Drug Des* 2014;83:191–7.
- [71] Karthik Kumar K, Seenivasan SP, Kumar V, Das TM. Synthesis of quinoline coupled [1,2,3]-triazoles as a promising class of anti-tuberculosis agents. *Carbohydr Res* 2011;346:2084–90.
- [72] Gonçalves RSB, Kaiser CR, Lourenço MCS, Bezerra FAFM, de Souza MVN, Wardell JL, et al. Mefloquine-oxazolidine derivatives, derived from mefloquine and arenecarbaldehydes: *in vitro* activity including against the multidrug-resistant tuberculosis strain T113. *Bioorg Med Chem* 2012;20:243–8.
- [73] Nava-Zuazo C, Estrada-Soto S, Guerrero-Álvarez J, León-Rivera I, Molina-Salinas GM, Said-Fernández S, et al. Design, synthesis, and *in vitro* antiprotozoal, antimycobacterial activities of N-[2-[(7-chloroquinolin-4-yl)amino]ethyl]ureas. *Bioorg Med Chem* 2010;18:6398–403.
- [74] Santos FC, Castro HC, Lourenço MCS, Abreu PA, Batalha PN, Cunha AC, et al. Tuberculosis: finding a new potential antimycobacterium derivative in a aldehyde-arylhydrazide-oxoquinoline series. *Curr Microbiol* 2012;65:455–60.
- [75] Khunt RC, Khedkar VM, Coutinho EC. Synthesis and 3D-QSAR analysis of 2-chloroquinoline derivatives as *H₃₇RV* MTB inhibitors. *Chem Biol Drug Des* 2013;82:669–84.
- [76] Kanani MB, Patel MP. Design and synthesis of new (bis)trifluoromethyl-promoted N-aryl biquinoline derivatives as antitubercular and antimicrobial agents. *Med Chem Res* 2014. <http://dx.doi.org/10.1007/s00044-014-1140-8> [in press].

- [77] Garudachari B, Isloor AM, Satyanaraya MN, Ananda K, Fund HK. Synthesis, characterization and antimicrobial studies of some new trifluoromethyl quinoline-3-carbohydrazide and 1,3,4-oxadiazoles. *RSC Adv* 2014;4:30864–75.
- [78] Carmo AML, Silva FMC, Machado PA, Fontes APS, Pavan FR, Leite CQF, et al. Synthesis of 4-aminoquinoline analogues and their platinum(II) complexes as new antileishmanial and antitubercular agents. *Biomed Pharmacother* 2011;65:204–9.
- [79] Nazarenko AB, Fedorov VE, Il'in VI, Omel'kov AV, Ruchko EA. Quinoline derivatives, in particular 5,6,7-substituted 1-(2-chloroquinolin-3-yl)-4-dimethylamino-2-(naphthalen-1-yl)-1-phenylbutan-2-ols, method for producing same and use of said compounds. *PCT Int Appl* 2013. WO 2013039428 A2 20130321.



Dr. Rangappa Keri was born in Hosur village of the Belgaum district, Karnataka, India in 1979. He did his B.Sc. degree from JSS College, Gokak, Karnatak University, Dharwad, Karnataka, India. He obtained his M.Sc., and Ph.D., degrees in Chemistry from Karnatak University Dharwad, India which is one of the prestigious universities in India. He joined the Centre for Nano and Material Science, Jain University as an Assistant Professor in October, 2013. Before joining Jain University, he did his postdoctoral work at Instituto Superior Técnico, Universidade de Lisboa, Lisboa, Portugal and Kyung Hee University, Seoul, South Korea. Dr. Keri has published more than 25 research articles in national/international journals and also he is a member

of national/international editorial board of many journals such as *Journal of Chemistry & Applied Biochemistry*. Dr. Keri is in the reviewer board of various international journals published by Bentham, sciencedirect, Springer, RSC etc. Dr. Keri has awarded prestigious fellowships like Fundação para a Ciência e Tecnologia (FCT) Portugal, BK21 Government of Korea, Postdoctoral Fellowship, Korea, and etc. His research interest includes synthesis of nitrogen, oxygen and sulfur containing heterocycles like benzothiazole, benzimidazole, coumarin moieties. Development of new drugs including acetylcholinesterase (AChE) inhibitors (Alzheimer drug), anti-cancers, neuroprotective agents and anti-virals by organic syntheses. Development of metal-, organo- and metalloorgano-catalyzed enantioselective methods for the synthesis of heterocyclic scaffolds and their potential application in medicinal chemistry and material science.



Dr. Siddappa Patil was born in Krishna Kittur village of district Belgaum, Karnataka, India in 1977. He did his B.Sc. degree from Karnatak Science College Dharwad, Karnataka, India. He obtained his M.Sc. and Ph.D. degrees in Chemistry from Karnatak University Dharwad, India which is one of the prestigious universities of the Karnataka state, India. He has been awarded several merit scholarships for pursuing higher studies at college as well as university level. He has vast postdoctoral research experience from around the world including: National Chung-Hsing University, Taiwan, University College Dublin, Ireland, Central Michigan University, Michigan, University of Texas at Arlington, Texas, and Northern Illinois University, Illinois, USA. He has 55 research publications in different well-know international journals, and several more are in the pipeline for publication. He is an active member of American Chemical Society since 2011. Dr. Siddappa Patil has awarded prestigious fellowships like National Science Foundation Postdoctoral Fellowship, USA, Government of Ireland IRCSET Postdoctoral Fellowship, Ireland, and Visiting Research Scientist, Northern Illinois University, USA, etc. Currently, he is working as Assistant Professor at Center for Nano and Material Sciences, Jain University, Bangalore, Karnataka, India. His research interest includes Organometallic and Medicinal Chemistry. His work has been cited over 655 times with h-index 15.