

Technical Note**CD3 Bright Lymphocyte Population Reveal $\gamma\delta$ T Cells**

Claude Lambert* and Christian Genin

Immunology Laboratory, University Hospital, St. Etienne, France

Background: In routine CD3/CD4/CD8 T-cell analysis, a CD3 bright population of lymphocytes is frequently observed. The aim of the present study was to identify the immunological significance of such CD3 bright lymphocytes.

Methods: We analyzed samples from 31 healthy adult volunteers, 78 human immunodeficiency virus (HIV)-positive, and 78 renal transplanted patients.

Results: A clearly distinct CD3 bright (frequently CD4⁺/CD8⁺) T-cell fraction was observed in 84% of donors and was directly correlated with the fraction of $\gamma\delta$ T cells ($r^2 = 0.64$). CD3 overexpression on $\gamma\delta$ T cells was confirmed by a combination of monoclonal antibody staining (CD3-ECD, $\gamma\delta$ TCR-FITC, and $\alpha\beta$ TCR-PE-Cy5) or immunomagnetic purification of $\gamma\delta$ T cells (i.e., MdFI 20 vs 8.86). The $\gamma\delta$ T cells expressed CD8 polypeptide chains (α and β) in all possible combinations. The largest proportion, surprisingly, were cells expressing CD8 $\beta\beta$ homodimers (43.8 \pm 16.5%). CD8 $\alpha\alpha$ homodimers were expressed on 14.2% (\pm 12.3) of total $\gamma\delta$ T cells, whereas CD8 $\alpha\beta$ heterodimers were expressed on 12.2% (\pm 7.5). We also observed a bimodal distribution of the intensity of CD3 fluorescence of $\gamma\delta$ T cells in immunocompromised patients with a threshold at 105 cell/ μ l. CD3 bright $\gamma\delta$ T cells were more frequently observed in HIV patients (29%) compared with renal transplant patients (11%) and healthy donors (3%; χ^2 test: $P = 0.0007$).

Conclusions: The simple observation of a CD3 bright T-cell subset on CD3/CD4/CD8 routine analysis suggests a high $\gamma\delta$ T-cell fraction and, in our opinion, should be followed by a complementary analysis to determine precisely the number of $\gamma\delta$ T cells and to identify their CD8 α/β phenotype. When CD3 bright T cells/ μ l were more than 40%, high $\gamma\delta$ T cells were detected in more than 87% of cases, with a specificity of 76%. Occasionally, the CD3 bright subset appeared to be strongly homogeneous, suggesting an oligoclonal proliferation that could possibly reveal a chronic localized stimulation or an early lymphoproliferative disorder. Because the $\gamma\delta$ T cells have interesting immunological peculiarities, the clinical significance of their quantitative abnormality should be clarified in diseases such as HIV, organ transplantation, autoimmunity and lymphoma. © 2004 Wiley-Liss, Inc.

Key terms: $\gamma\delta$ T cells; CD8 $\alpha\alpha$; intraepithelial lymphocyte

Quantitative analysis of CD4/CD8 T-cell subsets is one of the most commonly requested analyses for immunological monitoring in conditions, such as acquired immunodeficiency disease syndrome (AIDS), post-bone marrow grafting, and during immunosuppressive treatment. Quality assurance (QA) standard protocols using four-color immunolabeling allow very reproducible identification of both CD4 and CD8 T lymphocytes in a single run (1). However, this analysis provides much more information than is rarely considered.

In CD3/CD4/CD8/CD45 routine analysis of peripheral T cells, we repeatedly observed a bimodal distribution of CD3 expression that suggested the presence of two distinct T-cell subsets.

Besides CD3, the antigen-specific T-cell receptor (TCR) is another pan-T-cell marker. There are two distinct types of TCR— $\gamma\delta$ TCR and $\alpha\beta$ TCR—with different ontogeny

and functional properties. In the clinic, attention has been focused primarily on either CD4 or CD8 $\alpha\beta$ T cells, which constitute the vast majority (>95%) of circulating T cells. The remaining 5% of cells are $\gamma\delta$ T cells, dedicated to mucosal immunity (2–4) and develop only partially from thymus (5).

The aim of this report was to identify the immunological relevance of the CD3⁺ bimodal distribution, firstly observed in immunocompromised patients, and confirmed in healthy donors.

*Correspondence to: Claude Lambert, Immunology laboratory, Hôpital Bellevue, CHU St Etienne, F 42055 St. Etienne Cedex 2, France.

E-mail: claudelambert@uni-st-etienne.fr

Received 25 April 2003; Accepted 5 December 2003

Published online 6 August 2004 in Wiley InterScience (www.interscience.wiley.com).

DOI: 10.1002/cyto.b.20005




FIG. 1. Bimodal distribution of CD3 fluorescence intensity is frequently observed on CD3/CD4/CD8/CD45 routine lymphocyte analysis (one representative case). (a) CD3 bright T cells do not express CD4, but express low levels of CD8 (c). Conventional CD3⁺ T cells express CD4 or CD8 (b).

PATIENTS AND METHODS

Lymphocyte analysis was performed on 187 samples from 31 healthy adult volunteers, 78 human immunodeficiency virus (HIV) patients, and 78 renal transplant patients. Lymphocytes were immunolabeled with a combination of CD3-PE-cyanine 5 (PE-Cy5, Clone UCHT1), CD4-PE-Texas Red (PE-TR, clone SFC12T4D11), CD8 α -phycoerythrin (PE, clone SFC121thyD3), and CD45-fluorescein isothiocyanate (FITC, clone B3821F4A), (TetraChrome, Beckman-Coulter, Fullerton, CA). The TCR/CD8 chain expressions were analyzed using TCR α / β PE-Cy5 (clone BMA031), TCR γ / δ FITC (clone Immu 510), CD8 α PE-TR (clone SFC121thyD3), and CD8 β PE (clone 2ST8.5H7) all purchased from Immunotech/Beckman-Coulter. TCR/CD3 expression was analyzed simultaneously in a few samples, using TCR α / β PE-Cy5 (clone BMA031), TCR γ / δ FITC (clone Immu 510), CD3 PE-TR (clone UCHT1), and CD8 β PE (clone 2ST8.5H7) all purchased from Immunotech/Beckman-Coulter, USA).

The immunolabeling procedures were performed on whole blood according to the manufacturer's recommen-

dations. Briefly, 100 μ l of ethylenediaminetetraacetic acid (EDTA) anti-coagulated fresh peripheral blood was incubated with 10–20 μ l of antibody, gently mixed, and incubated for 20 min at room temperature in the dark. Red blood cells were then lysed, and the remaining leukocytes were fixed with formaldehyde, using Immunoprep reagent on a T-Q prep machine (Beckman-Coulter). Samples were then stored in the dark, at room temperature, and analyzed within 1 h on a four-color EPICS XL flowcytometer (Beckman-Coulter). Immediately before analysis, CD3/CD4/CD8/CD45 labeled samples were mixed with an equal volume of standardized fluorescent beads (Flow-count, Beckman-Coulter) for absolute quantification. The analysis was performed on 5,000 CD3⁺ T cells. Because the fluorescence intensity did not always have a normal distribution, we chose to consider the median (Mdf), instead of the mean (MFI), as calculated automatically by the machine software (System II, Beckman-Coulter).

The flowcytometer signal settings were checked every day using Flow Check[®] beads (Beckman-Coulter) and stabilized blood samples (ImmunoTrol[®], Beckman-Coulter)




FIG. 2. $\gamma\delta$ T lymphocytes are more CD3 bright than $\alpha\beta$ T cells. One patient had a very high level (23.6%) of $\gamma\delta$ T lymphocytes. The $\gamma\delta$ T cells were purified with magnetic beads (Materials and Methods) after staining with $\gamma\delta$ TCR-FITC and $\alpha\beta$ TCR-PE-CY5 (a). CD3 expression was higher in the $\gamma\delta$ + sorted fraction (c, MdFI = 20) than in the $\gamma\delta$ - remaining cells (b: MdFI = 8.86).

in order to check the day-to-day stability of the CD3, CD4, CD8 fluorescence signals. Signal amplification for each photomultiplier (PMT) and compensation were automatically corrected once a week, after every maintenance and each time the daily test failed, using Flow Set[®] standardized beads and CytoComp reagent and cell kit[®] (Beckman-Coulter) in accordance the manufacturer's recommendations. The resulting settings were analyzed periodically by Levey-Jennings graphs on System II software (Beckman-Coulter). The external quality controls comprised an international survey Interlaboratory Quality Assurance Program (Beckman-Coulter) and United Kingdom National External Quality Assessment Schemes (NEQUAS; Clinical Pathology Accreditation Ltd., Nequas Sheffield, UK).

$\gamma\delta$ T cells were purified from a $\gamma\delta$ T-cell-rich (23.6%) sample, using magnetic beads. The peripheral blood mononuclear cells (PBMC) were separated on density gradient from EDTA anti-coagulated blood, washed in sterile,

calcium free, phosphate buffered saline (Eurobio, Les Ulis, France) and immunolabeled with anti $\gamma\delta$ TCR monoclonal antibody (clone 11F2) conjugated with a hapten. The labeled cells were then sorted using anti-hapten antibody dually conjugated with the magnetic colloid and FITC (Miltenyi Biotech, GmbH, Bergisch Gladbach, Germany). Both sorted and negative fractions were analyzed using the same procedures previously described excluding the lysis step.

Statistical analysis was performed using the χ^2 test to compare frequencies in different groups, paired or unpaired Student's *t*-tests to compare means between or within groups and linear regression curve with coefficient of determination (r^2) to evaluate correlation of quantitative values. The results were expressed as mean \pm 1SD. The receiver operating characteristic (ROC) was calculated at the website of the Russell H. Morgan Department of Radiological Science (Johns Hopkins University, Baltimore, MD) with a Java program by John Eng.




FIG. 3. $\gamma\delta$ T cells express all possible combinations of the two CD8 polypeptide chains α and β . CD8 expression was analyzed on both $\alpha\beta$ and $\gamma\delta$ T-cell fractions using four-parameter labeling (CD8 β -PE; CD8 α PE-TR; $\gamma\delta$ TCR-FITC, $\alpha\beta$ TCR-PE-Cy5) of whole blood with red blood cell lysing and no washing (Materials and Methods). $\gamma\delta$ TCR $^{+}$ and $\alpha\beta$ TCR $^{+}$ T cells were electronically selected (a) for analysis of CD8 α /CD8 β proportion (b,c) one representative sample is shown). $\gamma\delta$ T cells expressed either CD8 β chain alone ($43.8 \pm 16.5\%$), both α and β chains of CD8 ($12.2 \pm 7.5\%$), or CD8 α only ($14.2 \pm 12.3\%$; c). CD8 fluorescence intensity was significantly lower on $\gamma\delta$ T cells compared with $\alpha\beta$ T cells, for CD8 α chain (MdFI = 5.4 ± 3.2 compared with 32.4 ± 7.6 , $P < 0.0001$) as well as CD8 β chain (MdFI = 2.2 ± 1.0 compared with 10.3 ± 3.1 , $P < 0.0001$; b). Less than 2% of cells had fluorescence intensity >1.0 with the isotype controls (not shown).

RESULTS

Evidence for a CD3 Bright T-Cell Subset in Healthy Donors

In the first set of experiments, we confirmed the presence of a CD3 bright fraction of T cells in 26 of 31 (84%) healthy donors (15 females, 16 males, aged 46.5 ± 13.3 years old). CD3 bright T cells constituted $3.15 \pm 2.16\%$ of the lymphocytes (Fig. 1a). The bimodal CD3 distribution could be confirmed with anti-CD3 monoclonal antibody obtained from another manufacturer (Dako; clone UCHT-1) and did not depend on the amount of antibody used (5, 10 -recommended-, 20 or μ l per 100 μ l of blood; data not shown).

Identification of the CD3 Bright T Cells in Healthy Donors

Of these CD3 bright T cells, 55.8% were both CD4 and CD8 negative (Fig. 1c). Therefore, we suspected that they could be $\gamma\delta$ T cells. In healthy donors, we observed that $2.4 \pm 1.4\%$ (46 ± 26 cells/ μ l, range: 2-105) of peripheral blood lymphocytes expressed the $\gamma\delta$ TCR. The size of the

CD3 bright and $\gamma\delta$ T-lymphocyte fractions were directly correlated ($0.73x + 0.370$; $R^2 = 0.644$, $P = 0.012$; not shown).

$\gamma\delta$ T Cells Expressed High Levels of CD3

To confirm that high levels of CD3 are indeed expressed on $\gamma\delta$ T cells, we used a CD3 PE-TR labeling combined with $\gamma\delta$ T FITC and $\alpha\beta$ T PE-Cy5, in five patients. We observed that electronically selected $\gamma\delta$ T cells expressed a higher CD3 fluorescence intensity than the $\alpha\beta$ T cells (e.g., MdFI = 15.2 vs 12.1). Similarly, sorted $\gamma\delta$ T cells, which contained only 1.2% of $\alpha\beta$ T cells and less than 1% of non-T cells, had a higher CD3 median fluorescence intensity (MdFI = 20) compared with the $\gamma\delta$ TCR-negative fraction (MdFI = 8.86; Fig. 2), which contained less than 0.25% of $\gamma\delta$ TCR (labeled by the fluorescent magnetic colloids or free FITC-antibody we added).




Fig. 4. Increased CD3-bright cells highlight the possibility of abnormal increase in $\gamma\delta$ T cells. $\gamma\delta$ T lymphocyte analysis is shown (a) in 31 healthy donors (ct), 78 HIV patients (HIV), and 78 renal transplant patients (RT). Immunosuppressed patients had elevated peripheral $\gamma\delta$ T cells ($>105 \text{ cell}/\mu\text{L}$) specially among HIV (29.3%) and RT (10.5%) patients compared with healthy donors (3%, χ^2 test: $P < 0.001$). The receiver operating characteristic (ROC) curve (b) shows that more than 87% of patients with high $\gamma\delta$ T cells could be identified when CD3-bright cells were higher than 40 cells/ μL (specificity 76%).

Phenotypes of $\gamma\delta$ T Cells in Healthy Donors

On CD3/CD4/CD8 labeling, the CD3 bright T cells never expressed CD4 but frequently ($39.5 \pm 20.7\%$) expressed the CD8 α (Fig. 1c), although at a lower level than the conventional CD8 $^+$ $\alpha\beta$ T cells (Fig. 1b). The frequency of $\gamma\delta$ T cells expressing CD8 was correlated neither to the age of the patients nor to the CD4 $^+$ T-cell or CD8 $^+$ T-cell absolute count (results not shown). On CD8 α /CD8 β / $\gamma\delta$ TCR labeling, a substantial fraction of the $\gamma\delta$ T cells did express CD8 α chain either alone (CD8 $\alpha\alpha$: $14.2 \pm 12.3\%$) or with the CD8 β chain (CD8 $\alpha\beta$: $12.2 \pm 7.5\%$). Moreover, most ($43.8 \pm 16.5\%$) of $\gamma\delta$ T cells expressed (low levels of) the CD8 β chain alone (Fig. 3). The CD8 MFI on $\gamma\delta$ T cells was much lower than on $\alpha\beta$ T cells for both α chain (5.4 ± 3.2 compared with 32.4 ± 7.6 , paired t -test: $P < 0.0001$) and β chain (2.2 ± 1.0 compared with 10.3 ± 3.1 , paired t -test: $P < 0.0001$) while most (98%) of lymphocytes labeled with the isotype controls or the CD4 $^+$ T cells had a CD8 fluorescence intensity below 1.0 (MFI = 0.15 and 0.13 respectively; not shown).

Possible Clinical Significance of $\gamma\delta$ T Cells in Diseases

To investigate whether the CD3 bright/ $\gamma\delta$ T-cell fraction has any pathological significance, we analyzed their distribution in 156 immunocompromised patients: 78 HIV patients (22F/56M: 40.3 ± 10.2 years) and 78 renal transplant patients (26 F/52M: 52.3 ± 14.1 years). All HIV patients were treated and had a wide range of CD4 $^+$ T-cell levels ($511 \pm 288 \text{ cells}/\mu\text{L}$). The renal patients had renal transplant for more than 6 months (1–8 years follow-up), were treated with Cyclosporin A, low doses of steroids (10–30 mg/day) and azathioprim and had normal levels of CD4 $^+$ T cells ($790 \pm 402 \text{ cells}/\mu\text{L}$).

The mean of the absolute number of $\gamma\delta$ T cells was significantly increased in HIV patients ($88 \pm 53 \text{ cells}/\mu\text{L}$) compared with renal transplant patients ($51 \pm 32/\mu\text{L}$; t -test: $P = 0.0056$) and healthy donors ($51 \pm 38/\mu\text{L}$; t -test: $P = 0.037$). We did not find any correlation with age or sex of patients (not shown). In fact, the graphic analysis (Fig. 4) shows a bimodal distribution. All but one (3%) healthy donors had less than 105 $\gamma\delta$ T cells/ μL . A signifi-




FIG. 5. $\gamma\delta$ T cells express more CD8 α chain in HIV ($41.7 \pm 19.4\%$) compared with renal transplant (RT) patients ($28.8 \pm 14.1\%$; $P < 0.0001$) and healthy donors ($26.4 \pm 14.6\%$; $P = 0.002$). 35% of HIV patients had more than 35% of $\gamma\delta$ T cells expressing CD8 compared with 8% of RT patients and 7% or donors (χ^2 : $P = 0.0033$).

cantly high number of HIV patients ($n = 22$ – 29% , χ^2 test: $P = 0.0007$) had raised $\gamma\delta$ T cells (105 – 498 cells/ μ l). The finding of increased $\gamma\delta$ T cells was less frequent in renal transplant patients ($n = 8$ – 10.5% ; from 115 – 405). The increased $\gamma\delta$ T cells in HIV patients may be related to the rise in CD8 T cells in HIV patients (1167 ± 980 cells/ μ l) compared with renal transplant patients (457 ± 25.1 cells/ μ l, t -test: $P < 0.0001$) and healthy donors (518 ± 217 cells/ μ l, t -test: $P = 0.0004$). Indeed, the $\gamma\delta$ T-cell counts were (poorly) correlated to the CD8 $^+$ $\alpha\beta$ T-cell counts ($r^2 < 0.497$), but not to the CD4 T cytopenia in HIV ($r^2 < 0.024$). This was similar in patients with renal transplants.

Moreover, the $\gamma\delta$ T cells expressed the CD8 α chain more frequently in HIV patients ($41.7 \pm 19.4\%$ of $\gamma\delta$ T) compared with renal transplant patients ($28.8 \pm 14.1\%$; t -test: $P < 0.0001$) and healthy donors ($26.4 \pm 14.4\%$; t -test: $P = 0.002$; Fig. 5). In some instance, a clear extra CD3bright CD8 $^+$ dot plot was observed (Fig. 6a). This was correlated with a raise of $\gamma\delta$ T cells (Fig. 6b). The extra T cell subset were $\gamma\delta$ T cell expressing heteroimeric ($\alpha\beta$) form of CD8 (Fig. 6c).

Predictive Value of Increased CD3 Bright $\gamma\delta$ T Cells

The presence of a CD3 bright subset of T cells (>40 cells/ μ l) allowed the detection of more than 87% of patients with high $\gamma\delta$ T cells (>105 cells/ μ l) without further testing (area under ROC curve was 0.851 with 77.6% accuracy; Fig. 4).

DISCUSSION

Knowledge of fundamental immunology has improved exponentially over the past few years [reviewed in refs. (6,7)]. In contrast, the information conveyed to clinicians after analysis in clinical immunology remains at a very basic level (i.e., numbers of CD4 $^+$ and CD8 $^+$ T cells). Improved technology and quality assurance of certified protocols provide us with high degree of confidence in the analysis of minor circulating cell subsets with unconventional phenotypes. Using the current clinical protocol for sample analysis, much more reliable information can be provided to clinicians with no extra cost. Until now, this information was not made available for clinical use. Even if the clinical significance of much of these new data has yet to be elucidated, our aim was to give clinicians the opportunity to get more detailed information, which could then be correlated with appropriate clinical circumstances.

The increase in CD3 bright T cells is an observation we make frequently. We propose that this information should be provided to the clinician with relevant interpretative comments pertaining to our current level of understanding of its immunological significance. In the present study, we have shown several lines of evidence suggesting that the CD3 bright fraction was $\gamma\delta$ T cells. Thus, the commonly used CD3/CD4/CD8 analysis allows the indirect detection of circulating $\gamma\delta$ T-cell variation that is rarely considered in clinical practice, despite the increasing availability of experimental evidence for their pathological involvement (5,6). By analyzing $\gamma\delta$ and $\alpha\beta$ TCR T-cell phenotypes, our data suggest that $\gamma\delta$ T cells can rise in a significant number of patients. A threshold of 105 cells/ μ l differentiates normal levels from a group of patients who show a large (up to 5 times) increase of peripheral $\gamma\delta$ T cells. This threshold is in accordance with published work (7). Because the frequency of raised $\gamma\delta$ T cells only concerned 1 of 5 patients and was rare in healthy donors, it would be unnecessarily expensive to routinely perform additional analysis for $\gamma\delta$ T-cell counting. This is why we believe that a rise in CD3 bright T cells could be a good screening criterion for further analysis specifically focused on $\gamma\delta$ T cells.

The threshold between normal and bright CD3 fluorescence intensities was not always easy to set because of the two peaks overlap. This could explain the relatively poor correlation we observed between CD3 bright and $\gamma\delta$ T-cell fractions. The CD3 bright fraction was more or less easily distinguished by its relative size and the fluorochrome conjugated to the anti-CD3 antibody. In some instances, when the $\gamma\delta$ T-cell subset was low, the CD3 bright fraction was almost indistinguishable from electronic noise. The PE-Cy fluorochrome appeared more discriminant for CD3 labeling than PE-Texas red. However, the bimodal distribution of CD3 was not dependent on the source of monoclonal antibodies, at least under saturating concentrations of the anti-CD3 antibodies used. Because CD3 bright T cells never expressed CD4, the dot plot of the CD3/CD4 $^+$ subset could be a very convenient tool to




FIG. 6. Extra CD3 bright/CD8 dim T cells (a) are enriched for $\gamma\delta$ T cells expressing CD8 $\alpha\beta$. A representative sample from one donor is shown. $\gamma\delta$ T cells were more frequent (b: 6.8% of lymphocytes) and expressed unusually higher levels (lower than $\alpha\beta$ T cells) of both CD8 β (MdfI : 4.8) and CD8 α (MdfI : 8.6) chains (c: 61.9%).

determine the confident interval of conventional CD3 distribution and the threshold for defining CD3 bright T cells.

$\gamma\delta$ T cells have been extensively studied relatively recently. They have very restricted diversity (6), poor MHC restriction, and recognize antigens that are widely distributed and are not usually recognized (nonpeptidic phosphoantigens) by $\alpha\beta$ T cells [reviewed in (6-7,9)]. The $\gamma\delta$ T cells can enhance the immune response by producing high amounts of interferon- γ (IFN- γ), tumor necrosis factor- α (TNF- α), or chemokines (7,10,11). Moreover, $\gamma\delta$ T cells have effector (cytotoxic) activity on their own (12). From an evolutionary standpoint, $\gamma\delta$ T cells have a unique place between the highly specific $\alpha\beta$ T cells and the innate immune system to fulfill the immune defense of the body against environmental pathogens. Experimental data have shown that $\gamma\delta$ T cells were essential in immunity against organisms such as *Mycobacterium*, *Borellia*, *Francisella tularensis*, *Salmonella* (7,11,13,14), or virus

(15,16), and the HIV virus (17-19). Moreover, $\gamma\delta$ T cells have anti-tumor activity (20-24) and an increase in circulating $\gamma\delta$ T cells has been described in patients with lymphoma (25). The $\gamma\delta$ T cells are also involved in epithelium repair and homeostasis (26-28). In contrast, $\gamma\delta$ T cells can be involved in immunopathologies such as diabetes (29) or autoimmune disorders (30-32) and asthma (25,33).

The $\gamma\delta$ T cells have very rapid (starting after 4-6 days), high ($\times 200$ times), and prolonged (> 7 months) proliferative capacity to antigen challenge (7). Moreover, because of their limited diversity, antigen-specific $\gamma\delta$ T cells can reach as much as 48-98% of circulating $\gamma\delta$ T cells during an immune response (7). Thus, the detection of an increased circulating $\gamma\delta$ T-cell subset may suggest a recent or ongoing chronic stimulation, particularly at mucosal or cutaneous sites. That information should alert clinicians for a possible slowly progressing infectious disease. It should be more particularly relevant in immunocompro-

mised patients and even more in HIV patients because their $\gamma\delta$ T cells should be reduced due to virus-induced lymphopenia (17). Knowing that the distribution of the clonotype is strongly related to the site of residence (i.e., $\delta 1$ isotype from gut residents and $\delta 2$ from peripheral pool: 6), a more precise clonotyping of the peripheral $\gamma\delta$ T-cell population may help to locate the origin of the proliferative stimulus.

In our study, we observed that a portion of $\gamma\delta$ T cells expressed CD8 $\alpha\beta$ as previously described in rats (34). The CD8 appeared to be functional and not restricted to oligoclonal $\gamma\delta$ T cells. Compared with $\alpha\beta$ T-cell ontogeny, this observation may suggest that $\gamma\delta$ T cells expressing CD8 $\alpha\beta$ could develop differently from other $\gamma\delta$ T cells, and may even be of thymic origin (4).

We have also confirmed that $\gamma\delta$ T cells could express homodimeric CD8 $\alpha\alpha$, similar to a small fraction of the $\alpha\beta$ T cells. Interestingly, both $\gamma\delta$ T cells and CD8 $\alpha\alpha$ $\alpha\beta$ T cells have shared tropism for mucosal barriers (2–4,35). Further characterization of the clonotype should determine if the CD8 $\alpha\alpha$ ⁺ $\gamma\delta$ T cells are preferentially derived from the gut intraepithelial T-cell pool. Thus, an increase of peripheral CD8 $\alpha\alpha$ ⁺ $\gamma\delta$ T cells could have clinical significance revealing an intestinal inflammatory or lymphoproliferative process.

Finally, we repeatedly observed that $\gamma\delta$ T cells frequently expressed the CD8 β chain alone? This has never been described before and needs to be confirmed by other investigators. However, the level of expression remains very low and may not be physiologically relevant.

During antigen recognition, both CD8 $\alpha\beta$ and CD8 $\alpha\alpha$ receptors strengthen the $\alpha\beta$ TCR binding to the MHC-peptide complex (36,37). It is generally accepted that $\gamma\delta$ T-cell interact with nonclassical MHC molecules (6,7,9,14,38) and there is some evidence that CD8 molecules are involved in the interaction with these nonclassical MHC molecules (e.g., CD1, MICA, MICB) (6,9,14,38). It is relevant in this context that NK-T cells that mainly interact with classical MHC molecules express CD8 (14,39,40).

ACKNOWLEDGMENTS

The authors thank Pascale Saby for her skillful technical assistance and A.A. Mohammad Ibrahim, King's College Hospital, London, for helpful reviewing and discussion of the paper.

LITERATURE CITED

1. The NIAID DAIDS New Technologies Evaluation Group. Reimann KA, O'Gorman MR, Spritzler J, Wilkening CL, Sabath DE, Helm K, Campbell DE. Multisite comparison of CD4 and CD8 T-lymphocyte counting by single-versus multiple-platform methodologies: evaluation of Beckman Coulter flow-count fluorospheres and the tetraONE system. *Clin Diagn Lab Immunol* 2000;7:344–351.
2. Ferrick DA, King DP, Jackson KA, Braun RK, Tam S, Hyde DM, Beaman BL. Intraepithelial gamma delta T lymphocytes: sentinel cells at mucosal barriers. *Springer Semin Immunopathol* 2000;22:283–296.
3. Beagley KW, Husband AJ. Intraepithelial lymphocytes: origins, distribution, and function. *Crit Rev Immunol* 1998;18:237–254.
4. Bagriacik EU, Okabe M, Klein JR. Origins of intestinal intraepithelial lymphocytes: direct evidence for a thymus-derived gamma delta T cell component. *Immunol Lett* 2000;75:77–83.
5. Helgeland L, Brandtzaeg P, Rolstad B, Vaage JT. Sequential development of intraepithelial gamma delta and alpha beta T lymphocytes expressing CD8 alpha beta in neonatal rat intestine: requirement for the thymus. *Immunology* 1997;92:447–455.
6. Carding SR, Egan PJ. Gammadelta T cells: functional plasticity and heterogeneity. *Nat Rev Immunol* 2002;2:336–345.
7. Chen ZW, Letvin NL. Adaptive immune response of Vgamma2Vdelta2 T cells: a new paradigm. *Trends Immunol* 2003;24:213–219.
8. Hviid L, Akanmori BD, Loizon S, Kurtzhals JA, Ricke CH, Lim A, Koram KA, Nkrumah FK, Mercereau-Puijalon O, Behr C. High frequency of circulating gamma delta T cells with dominance of the v(delta)1 subset in a healthy population. *Int Immunol* 2000;12:797–805.
9. Steele CR, Oppenheim DE, Hayday AC. Gamma(delta) T cells: non-classical ligands for non-classical cells. *Curr Biol* 2000;10:R282–285.
10. Lehner T, Mitchell E, Bergmeier L, Singh M, Spallek R, Cranage M, Hall G, Dennis M, Villinger F, Wang Y. The role of gammadelta T cells in generating antiviral factors and beta-chemokines in protection against mucosal simian immunodeficiency virus infection. *Eur J Immunol* 2000;30:2245–2255.
11. Moore TA, Moore BB, Newstead MW, Standiford TJ. Gamma delta-T cells are critical for survival and early proinflammatory cytokine gene expression during murine *Klebsiella pneumoniae*. *J Immunol* 2000;165:2643–2650.
12. Dieli F, Troye-Blomberg M, Ivanyi J, Fournier JJ, Krensky AM, Bonneville M, Peyrat MA, Caccamo N, Sireci G, Salerno A. Granulysin-dependent killing of intracellular and extracellular *Mycobacterium tuberculosis* by gamma9/Vdelta2 T lymphocytes. *J Infect Dis* 2001;184:1082–1085.
13. Mogues T, Goodrich ME, Ryan L, LaCourse R, North RJ. The relative importance of T cell subsets in immunity and immunopathology of airborne *Mycobacterium tuberculosis* infection in mice. *J Exp Med* 2001;193:271–280.
14. Kawashima T, Norose Y, Watanabe Y, Enomoto Y, Narazaki H, Watari E, Tanaka S, Takahashi H, Yano I, Brenner MB, Sugita M. Major CD8 T cell response to live bacillus Calmette-Guérin is mediated by CD1 molecules. *J Immunol* 2003;170:5345–5348.
15. Lafarge X, Merville P, Cazin MC, Berge F, Potaux L, Moreau JF, Dechanet-Merville J. Cytomegalovirus infection in transplant recipients resolves when circulating gammadelta T lymphocytes expand, suggesting a protective antiviral role. *J Infect Dis* 2001;184:533–541.
16. Selin LK, Santolucito PA, Pinto AK, Szomolanyi-Tsuda E, Welsh RM. Innate immunity to viruses: control of vaccinia virus infection by gamma delta T cells. *J Immunol* 2001;166:6784–6794.
17. Kabelitz D, Wesch D. Role of gamma delta T-lymphocytes in HIV infection. *Eur J Med Res* 2001;6:169–174.
18. Pellegrin JL, Taupin JL, Dupon M, Ragnaud JM, Maugein J, Bonneville M, Moreau JF. Gammadelta T cells increase with *Mycobacterium avium* complex infection but not with tuberculosis in AIDS patients. *Int Immunol* 1999;11:1475–1478.
19. Sindhu ST, Ahmad R, Morisset R, Ahmad A, Menezes J. Peripheral blood cytotoxic gammadelta T lymphocytes from patients with human immunodeficiency virus type 1 infection and AIDS lyse uninfected CD4⁺ T cells, and their cytotoxic potential correlates with viral load. *J Virol* 2003;77:1848–1855.
20. Girardi M, Oppenheim DE, Steele CR, Lewis JM, Glusac E, Filler R, Hobby P, Sutton B, Tigelaar RE, Hayday AC. Regulation of cutaneous malignancy by gammadelta T cells. *Science* 2001;294:605–609.
21. Kato Y, Tanaka Y, Miyagawa F, Yamashita S, Minato N. Targeting of tumor cells for human gammadelta T cells by nonpeptide antigens. *J Immunol* 2001;167:5092–5098.
22. Chen J, Niu H, He W, Ba D. Antitumor activity of expanded human tumor-infiltrating gammadelta T lymphocytes. *Int Arch Allergy Immunol* 2001;125:256–263.
23. Kobayashi H, Tanaka Y, Yagi J, Toma H, Uchiyama T. Gamma/delta T cells provide innate immunity against renal cell carcinoma. *Cancer Immunol Immunother* 2001;50:115–124.
24. Ferrarini M, Ferrero E, Dagna L, Poggi A, Zocch MR. Human gamma-delta T cells: a nonredundant system in the immune-surveillance against cancer. *Trends Immunol* 2002;23:14–18.
25. McClanahan J, Fukushima PI, Stetler-Stevenson M. Increased peripheral blood gamma delta T-cells in patients with lymphoid neoplasia: a diagnostic dilemma in flow cytometry. *Cytometry* 1999;38:280–285.
26. Lahn M, Kanehiro A, Takeda K, Joetham A, Schwarze J, Kohler G, O'Brien R, Gelfand EW, Born W, Kanehiro A. Negative regulation of airway responsiveness that is dependent on gammadelta T cells and independent of alphabeta T cells. *Nat Med* 1999;5:1150–1156.
27. Born WK, Lahn M, Takeda K, Kanehiro A, O'Brien RL, Gelfand EW. Role of gammadelta T cells in protecting normal airway function. *Respir Res* 2000;1:151–158.

28. Komano H, Fujiura Y, Kawaguchi M, Matsumoto S, Hashimoto Y, Obana S, Mombaerts P, Tonegawa S, Yamamoto H, Itoharu S, Nanno M, Ishikawa H. Homeostatic regulation of intestinal epithelia by intraepithelial gamma delta T cells. *Proc Natl Acad Sci USA* 1995;92: 6147-6151.
29. Kretowski A, Mysliwiec J, Szelachowska M, Turowski D, Wysocka J, Kowalska I, Kinalska I. Gammadelta T-cells alterations in the peripheral blood of high risk diabetes type 1 subjects with subclinical pancreatic B-cells impairment. *Immunol Lett* 1999;68:289-293.
30. Ichikawa Y, Shimizu H, Yoshida M, Takaya M, Arimori S. T cells bearing gamma/delta T cell receptor and their expression of activation antigen in peripheral blood from patients with Sjogren's syndrome. *Clin Exp Rheumatol* 1991;9:603-609.
31. Robak E, Blonski JZ, Bartkowiak J, Niewiadomska H, Sysa-Jedrzejowska A, Robak T. Circulating TCR gammadelta cells in the patients with systemic lupus erythematosus. *Mediators Inflamm* 1999;8:305-312.
32. Yin Z, Craft J. gamma delta T cells in autoimmunity. *Springer Semin Immunopathol* 2000;22:311-320.
33. Schramm CM, Puddington L, Yiamouyiannis CA, Lingenheld EG, Whiteley HE, Wolyniec WW, Noonan TC, Thrall RS. Proinflammatory roles of T-cell receptor (TCR)gammadelta and TCRalphabeta lymphocytes in a murine model of asthma. *Am J Respir Cell Mol Biol* 2000;22:218-225.
34. Straube F, Herrmann T. Expression of functional CD8alpha Beta heterodimer on rat gamma delta T cells does not correlate with the CDR3 length of the TCR delta chain predicted for MHC class I-restricted antigen recognition. *Eur J Immunol* 2000;30:3562-3568.
35. Miller C, Roberts SJ, Ramsburg E, Hayday AC. Gamma delta cells in gut infection, immunopathology, and organogenesis. *Springer Semin Immunopathol* 2000;22:297-310.
36. Garcia KC, Scott CA, Brunmark A, Carbone FR, Peterson PA, Wilson IA, Teyton L. CD8 enhances formation of stable T-cell receptor/MHC class I molecule complexes. *Nature* 1996;384:577-581.
37. Sun J, Kavathas PB. Comparison of the roles of CD8 alpha alpha and CD8 alpha beta in interaction with MHC class I. *J Immunol* 1997;159: 6077-6082.
38. Gao GF, Willcox BE, Wyer JR, Boulter JM, O'Callaghan CA, Maenaka K, Stuart DI, Jones EY, Van Der Merwe PA, Bell JL, Jakobsen BK. Classical and nonclassical class I major histocompatibility complex molecules exhibit subtle conformational differences that affect binding to CD8alphaalpha. *J Biol Chem* 2000;275:15232-15238.
39. Gapin L, Matsuda JL, Surh CD, Kronenberg M. NKT cells derive from double-positive thymocytes that are positively selected by CD1d. *Nat Immunol* 2001;2:971-978.
40. Heinzel AS, Grotzke JE, Lines RA, Lewinsohn DA, McNabb AL, Streblow DN, Braud VM, Grieser HJ, Belisle JT, Lewinsohn DM. HLA-E-dependent presentation of Mtb-derived antigen to human CD8⁺ T cells. *J Exp Med* 2002;196:1473-1481.
41. Kondo H, Uematsu M, Watanabe J, Takahashi Y, Hayashi K, Iwasaki H. CD3⁺, CD4⁻, CD8⁻, TCR alpha beta⁻, TCR gamma delta⁺ granular lymphocyte proliferative disorder without lymphocytosis and clinical symptoms. *Acta Haematol* 2000;104:54-56.
42. Toro JR, Beaty M, Sorbara L, Turner ML, White J, Kingma DW, Raffeld M, Jaffe ES. Gamma delta T-cell lymphoma of the skin: a clinical, microscopic, and molecular study. *Arch Dermatol* 2000;136:1024-1032.
43. Roncella S, Cutrona G, Truini M, Airolidi I, Pezzolo A, Valetto A, Di Martino D, Dadati P, De Rossi A, Ulivi M, Fontana I, Nocera A, Valente U, Ferrarini M, Pistoia V. Late Epstein-Barr virus infection of a hepatosplenic gamma delta T-cell lymphoma arising in a kidney transplant recipient. *Haematologica* 2000;85:256-262.