

***k*-Resonant Benzenoid Systems and *k*-Cycle Resonant Graphs[†]**Xiaofeng Guo^{*,‡} and Fuji Zhang[§]

Institute of Mathematics and Physics, Xinjiang University, Wulumuqi Xinjiang 830046,
 People's Republic of China, and Department of Mathematics, Xiamen University, Xiamen Fujian 361005,
 People's Republic of China

Received June 21, 2000

A benzenoid system (or hexagonal system) H is said to be k -resonant if, for $1 \leq t \leq k$, any t disjoint hexagons of H are mutually resonant; that is, there is a Kekule structure (or perfect matching) K of H such that each of the k hexagons is a K -alternating hexagon. A connected graph G is said to be k -cycle resonant if, for $1 \leq t \leq k$, any t disjoint cycles in G are mutually resonant. The concept of k -resonant benzenoid systems is closely related to Clar's aromatic sextet theory, and the concept of k -cycle resonant graphs is a natural generalization of k -resonant benzenoid systems. Some necessary and sufficient conditions for a benzenoid system (respectively a graph) to be k -resonant (respectively k -cycle resonant) have been established. In this paper, we will give a survey on investigations of k -resonant benzenoid systems and k -cycle resonant graphs.

1. INTRODUCTION

In the topological theory of benzenoid hydrocarbons, a benzenoid system (or hexagonal system) denotes the carbon atom skeleton graph of a benzenoid hydrocarbon, which is a two-connected plane graph whose every interior face is bounded by a regular hexagon. A Kekule structure K of a benzenoid system H corresponds to a perfect matching (1-factor) of H . An edge in H is said to be a K -double bond if it belongs to K , otherwise a K -single bond. An edge in H is said to be a fixed double (respectively, single) bond if it belongs to (respectively does not belong to) every Kekule structure of H . A benzenoid system is said to be normal if it contains no fixed bond. A cycle (or circuit) C in H is said to be *conjugated* or *resonant* if there is a Kekule structure K of H such that C is a K -alternating cycle. In conjugated circuit model,^{1–12} conjugated circuits with different sizes have different resonant energies. If the size of a conjugated circuit is equal to $4n + 2$, then the smaller n is, the larger the resonant energy is. So the conjugated hexagon has the largest energy. On the other hand, from a purely empirical standpoint, Clar found that various electronic properties of polycyclic aromatic hydrocarbons can be predicted by appropriately defining an aromatic sextet for their Kekule structures.^{13–23} According to Clar's aromatic sextet theory, a *Clar formula* of a benzenoid system is a set of mutually resonant sextets with the maximum cardinal number, where sextets mean resonant hexagons and a set of mutually resonant sextets means a set of disjoint hexagons for which there is a Kekule structure K so that all of the disjoint hexagons are K -alternating hexagons. The number of sextets in a Clar formula of G is called the Clar number of G . For




Figure 1. Benzenoid system H with Clar number 5 and with two Clar formulas.

a benzenoid system H with Clar number c , Clar formulas of H may not be unique, and, for $1 \leq k \leq c$, any k -disjoint hexagons of H are not certainly mutually resonant. Figure 1 shows a benzenoid system H with Clar number 5, which has two Clar formulas, and the hexagon s is not resonant.

An interesting problem arises: under what conditions are any k disjoint hexagons of a benzenoid system H mutually resonant?

If a benzenoid system H satisfies such a property, that is, for a positive integer k and $1 \leq t \leq k$, with any t disjoint hexagons of H being mutually resonant, it is said to be k -resonant or k -coverable.

1-Resonant benzenoid systems were first investigated by Zhang and Chen.²⁴ Some necessary and sufficient conditions for a benzenoid system to be 1-resonant were given. Later Zhang and Zheng²⁵ gave a similar characterization for 1-resonant generalized benzenoid systems, where a generalized benzenoid system is a 2-connected subgraph of a benzenoid system, which may have some holes (see Figure 2). Zheng²⁶ further gave some pretty results for k -resonant benzenoid systems with $k \geq 2$.

The concept of k -cycle resonant graphs was first introduced by Guo and Zhang, which is a natural generalization of the concept of k -resonant benzenoid systems. Some properties and necessary and sufficient conditions of k -cycle resonant graphs were also given.

[†] This project was supported by NSFC.

^{*} To whom correspondence should be addressed. Telephone: 86-9912879377. E-mail: xfguo@xju.edu.cn.

[‡] Xinjiang University.

[§] Xiamen University. Telephone: 86-9922182832, E-mail: fjzhang@jingxian.xmu.edu.cn.




Figure 2. k -coverable generalized benzenoid system with $k \geq 3$.

In this paper, we will give a survey and review on investigations of k -resonant benzenoid systems and k -cycle resonant graphs.

2. K-RESONANT (OR K-COVERABLE) BENZENOID SYSTEMS

A *cover* of H is a set of disjoint hexagons of H such that after deleting all the vertexes on these hexagons, the remainder of H has a Kekule structure or is empty. A maximum cover of H is a cover with maximum cardinal number, which is also called a Clar formula. In other words, a cover of H is a set of mutually resonant hexagons of H , and a maximum cover is a set of mutually resonant hexagons with the maximum cardinal number.

Definition 1. For a positive integer k , a benzenoid system H is said to be k -resonant or k -coverable if, for $1 \leq t \leq k$, any t disjoint hexagons of H form a cover of H .

For 1-coverable benzenoid systems, Zhang and Chen gave the following theorem.

Theorem 1.²⁴ Let H be an hexagonal system. Then each hexagon of H covers H iff either (1) H contains no fixed bond or (2) there exists a perfect matching M of H such that the contour of H is an M -alternating cycle.

A generalized benzenoid system G is said to be complete if each edge of G is contained on a hexagon. For a complete generalized benzenoid system H , Zhang and Zheng²⁵ gave a similar necessary and sufficient condition for H to be k -resonant.

Theorem 2.²⁵ Every hexagon of a completely generalized hexagonal system H is resonant if and only if the boundaries of the infinite face and nonhexagon faces of H are resonant.

A sufficient condition for a benzenoid system to be 2-resonant was given by Zheng.²⁶

Theorem 3.²⁶ A benzenoid system H is 2-resonant if H is 1-resonant and any pair of two disjoint hexagons of H with at least one boundary hexagon are mutually resonant.

In a draft of a book,²⁷ the above sufficient condition is improved as the following theorem.

Theorem 4.²⁷ A benzenoid system H is 2-coverable if and only if H is 1-coverable and any two disjoint side hexagons of H form a cover of H .

A complete characterization of $k(\geq 3)$ -resonant benzenoid systems was given by Zheng.²⁶

Theorem 5.²⁶ A benzenoid system is $k(\geq 3)$ -resonant if and only if it is 3-resonant.

Zheng defined three k -resonant bricks as shown in Figure 3. If a benzenoid system H can be constructed from the three bricks by affixing them in heavy edges, H is said to have a k - r -brick decomposition (see Figure 3). A series of lemmas in ref 26 imply the following theorem.




Figure 3. Three k -resonant bricks and the k -resonant-brick decomposition of a k -resonant benzenoid system.

Theorem 6.²⁶ A benzenoid system H is $k(\geq 3)$ -resonant if and only if H has a $(k - r)$ -brick decomposition.

Chen and Guo²⁸ proved that the results of theorems 5 and 6 are also valid for generalized benzenoid systems (see Figure 2).

3. K-CYCLE RESONANT GRAPHS

After investigations of k -resonant benzenoid systems got great advance, Guo and Zhang began to consider whether the concept of k -resonant benzenoid systems can be generalized to general cases. Hence a new concept of k -cycle resonant graphs was introduced in ref 29.

Definition 2.²⁹ A graph G is said to be k -cycle resonant or k -cycle extendable if G contains at least $k (\geq 1)$ disjoint cycles and, for $1 \leq t \leq k$, any t disjoint cycles in G are mutually resonant; that is, there is a Kekule structure K of G such that the t disjoint cycles are K -alternating cycles.

The following theorems were given in ref 29.

Theorem 7.²⁹ Let G be a k -cycle resonant graph. Then (1) G is bipartite.

(2) For $1 \leq t \leq k$ and any t disjoint cycles C_1, C_2, \dots, C_t in G , $G - \cup_{i=1}^t C_i$ contains no odd component.

(3) Any two 2-connected components in G have no common vertex.

Theorem 8.²⁹ Let G be a k -cycle resonant graph. Then G is elementary or 1-extendable if and only if G is 2-connected.

Theorem 7 conditions 1 and 2 give some necessary conditions for a graph to be k -cycle resonant. It was proved that the necessary conditions are also sufficient.

Theorem 9.²⁹ A connected graph with at least k disjoint cycles is k -cycle resonant if and only if G is bipartite and, for $1 \leq t \leq k$ and any t disjoint cycles C_1, C_2, \dots, C_t in G , $G - \cup_{i=1}^t C_i$ contains no odd component.

From theorem 7 condition 3 and theorem 8, it is not difficult to see that the following theorem 10 holds.

Theorem 10.³⁰ Let G be a k -cycle resonant graph. Then, (i) for a 2-connected component G' of G with the maximum number k^* of disjoint cycles, if $k^* \leq k$, G' is k^* -cycle resonant, otherwise G' is k -cycle resonant; (ii) the forest induced by all the vertexes of G not in any 2-connected component of G has a perfect matching.

The above theorems imply that a non-2-connected k -cycle resonant graph with $k \geq 3$ can be constructed from some disjoint 2-connected $k^*(\text{or } k)$ -cycle resonant graphs and a forest with perfect matching by adding some edges between




Figure 4. (1) 1-Cycle resonant hexagonal system with two chains of even length. (2) k^* -Cycle resonant benzenoid system.

the 2-connected graphs and the forest so that the resultant graph is connected and the added edges are cut edges. Hence we need only to consider 2-connected k -cycle resonant graphs. However, in general cases, the construction of 2-connected k -cycle resonant graphs is still an open problem.

4. PLANAR K -CYCLE RESONANT GRAPHS

The necessary and sufficient conditions in theorem 9 is simple and formally graceful. However, when it is used to determine whether or not a graph is k -cycle resonant, one needs to check any t disjoint cycles. It is obviously tedious. This is why we need to find new, simpler, necessary, and sufficient conditions for planar graphs to be k -cycle resonant.

For a class of planar graphs, k -cycle resonant hexagonal systems, we obtained the following theorems.

A path P in a graph G is said to be a **chain** if all internal vertexes of P are of degree 2 in G and the degree of any end vertex of P is not equal to 2 in G . A hexagonal system is said to a catacondensed hexagonal system if any vertex of it lies on the boundary.

Theorem 11.²⁹ A hexagonal system H is 1-cycle resonant if and only if H is a catacondensed hexagonal system.

Theorem 12.²⁹ A hexagonal system H is 2-cycle resonant if and only if (1) H contains at least two disjoint cycles and (2) H is a catacondensed hexagonal system with no chain of even length.

Theorem 13.²⁹ Let H be a 2-cycle resonant hexagonal system, and let k^* be the maximum number of disjoint cycles in H . Then H is k^* -cycle resonant.

Theorem 14.²⁹ A hexagonal system H with $k^* \geq 2$ is k^* -cycle resonant if and only if H is a catacondensed hexagonal system with no chain of even length, where k^* is the maximum number of disjoint cycles in H .

Figure 4(1) shows a 1-cycle resonant hexagonal system, which is not 2-cycle resonant because it has two chains of length 2. Figure 4(2) shows a $k(\geq 2)$ -cycle resonant hexagonal system. It was pointed out in ref 29 that, in the hexagonal systems with h hexagons obtained from a same parent hexagonal system with $h - 1$ hexagons, k^* -cycle resonant hexagonal systems have greater resonance energies than 1-cycle resonant hexagonal systems; also 1-cycle resonant hexagonal systems have greater resonance energies than hexagonal systems not being 1-cycle resonant, where k^* is the maximum number of disjoint hexagons of a hexagonal system.

For general planar k -cycle resonant graphs, their characterization is more difficult. Recently, we just finished a manuscript on planar k -cycle resonant graphs with $k = 1, 2$, which has been submitted to *Discrete Applied Mathematics*.³⁰ In that paper, some new necessary and sufficient conditions

for a graph to be planar 1-cycle resonant graphs or planar 2-cycle resonant graphs are established.

Before stating these results, we need to give some terminology and notations

Let G be a connected graph, and H a subgraph of G . A vertex in H is said to be an *attachment vertex* of H if it is incident with an edge in $G - E(H)$. The set of all attachment vertexes of H is denoted by $V_A(H)$. A bridge B of H in G is either an edge in $G - E(H)$ with two end vertexes being in H , or a subgraph of G induced by all the edges in a connected component B' of $G - V(H)$ together with all the edges with an end vertex in B' and the other in H . The vertexes in $V(B) \cap V(H)$ are also attachment vertexes of B to H . A bridge with k attachment vertexes is called a k -bridge.

The attachment vertexes of a k -bridge B of a cycle C in G divide C into k edge-disjoint paths, called the segments of B . Two bridges of C avoid one another if all the attachment vertexes of one bridge lie in a single segment of the other bridge; otherwise they overlap. Two bridges B and B^* of C are skewed if there are four distinct vertexes on C , in the cyclic order u, u^*, v, v^* , such that u and v are attachment vertexes of B and u^* and v^* are attachment vertexes of B^* .

For a bipartite graph G , we always color vertexes of G white and black so that any two adjacent vertexes have different colors.

We first gave several equivalent propositions.

Theorem 15.³⁰ Let G be a 2-connected bipartite planar graph. Then the following statements are equivalent:

- (i) G is 1-cycle resonant.
- (ii) For any cycle C in G , $G - V(C)$ has no odd component.
- (iii) For any cycle C in G , any bridge of C has exactly two attachment vertexes which have different colors.
- (iv) For any cycle C in G , any two bridges of C avoid one another. Moreover, for any 2-connected subgraph B of G with exactly two attachment vertexes, the attachment vertexes of B have different colors.

From the above theorem, we can give the following necessary and sufficient conditions for a graph to be planar 1-cycle resonant.

Theorem 16.³⁰ A 2-connected graph G is planar 1-cycle resonant if and only if G is bipartite and either (i) for any cycle C in G , any bridge of C has exactly two attachment vertexes which have different colors or (ii) for any cycle C in G , any two bridges of C avoid one another; moreover, for any 2-connected subgraph B of G with exactly two attachment vertexes, the attachment vertexes of B have different colors.

A vertex u of a graph G is said to be *cycle-related* to another vertex v of G if u is contained in a 2-connected block of G , and any cycle containing u must also contain v . If v is also cycle-related to u , then u and v are *mutually cycle-related*.

Property 1.³⁰ If a vertex u of a connected graph G is cycle-related to another vertex v of G , then u and v belong to the same 2-connected block B in G and all the edges in $B - v$ incident with u are cut edges of $G - v$.

For a chain P in a graph G , let $V_i(P)$ denote the set of internal vertexes of P . For a subgraph B of G , let \bar{B} denote the subgraph of G induced by $E(G) \setminus E(B)$. The necessary and sufficient conditions for a planar graph to be 2-cycle resonant were also given in ref 30.

Theorem 17.³⁰ A 2-connected graph G is planar 2-cycle resonant if and only if, (i) G is planar 1-cycle resonant; (ii) for a chain P with even length and end vertexes v_1 and v_2 , $G - V_1(P)$ has exactly two blocks, each of which is 2-connected, and v_1 and v_2 are cycle-related to the common vertex of the two blocks; (iii) for a chain P with odd length and end vertexes v_1 and v_2 such that $G - V_1(P)$ is not 2-connected, either (a) $G - V_1(P)$ has exactly three blocks, each of which is 2-connected, and each of v_1 and v_2 is cycle-related to the other attachment vertex of the block containing it, and the attachment vertexes of the third block are mutually cycle-related in the third block, or (b) any two 2-connected blocks of $G - V_1(P)$ are disjoint; (iv) for a 2-connected subgraph B_1 of G with exactly two attachment vertexes, if \bar{B}_1 is not 2-connected and every block of \bar{B}_1 is 2-connected, then \bar{B}_1 has exactly three blocks, say B_2, B_3, B_4 , and the attachment vertexes of each of B_1, B_2, B_3, B_4 are mutually cycle-related in the block.

Based on the above necessary and sufficient conditions, an efficient algorithm for determining whether a 2-connected graph to be planar 1-cycle resonant or 2-cycle resonant can be developed.

k -Cycle resonant hexagonal systems are a special class of planar k -cycle resonant graphs, the construction of which was completely characterized in ref 29. For general planar k -cycle resonant graphs, their construction is more complex. Further investigations are needed.

5. CONCLUSION

Investigations of k -resonant benzenoid systems and k -cycle resonant graphs have obtained great advance. The above many results are very interesting. For example, the coefficients of the sextet polynomial of a $k(\geq 3)$ -coverable benzenoid system H can be obtained by enumerations of any $i(\geq 1)$ disjoint hexagons in H . The two classes of graphs not only have a strong chemistry background but are also important topics in matching theorem. In the investigation of matching theory, Lovasz et al.³¹⁻³⁹ introduced and investigated elementary graphs, 1-extendable graphs, and n -extendable graphs, etc. A graph G is said to be n -extendable if any n independent edges of G are contained in a perfect matching of G . We can similarly call k -cycle resonant graphs as k -cycle extendable graphs and call k -coverable benzenoid systems as k -hexagon extendable benzenoid systems. The above investigations are also a new advance of matching theory research. There are still some open problems for further investigations.

REFERENCES AND NOTES

- (1) Randic, M. *Chem. Phys. Lett.* **1976**, 38, 68.
- (2) Gutman, I.; Randic, M. A correlation between Kekule valence structure and conjugated circuits. *Chem. Phys.* **1979**, 41, 265.
- (3) Randic, M. *J. Am. Chem. Soc.* **1977**, 99, 444.
- (4) Randic, M. *Tetrahedron* **1977**, 33, 1905.
- (5) Nikolic, S.; Randic, M.; Klein, D. J.; Plavsic, D.; Trinajstic, N. *J. Mol. Struct.* **1989**, 198, 223.
- (6) Vogler, H.; Trinajstic, N. *Theor. Chim. Acta* **1988**, 73, 437.
- (7) Nikolic, S.; Trinajstic, N.; Klein, D. J. *Comput. Chem.* **1990**, 14, 313.
- (8) Plavsic, D.; Nikolic, S.; Trinajstic, N. *J. Math. Chem.* **1991**, 8, 113.
- (9) Plavsic, D.; Nikolic, S.; Trinajstic, N. *Croat. Chem. Acta* **1990**, 63, 683.
- (10) Trinajstic, N.; Nikolic, S. *J. Mol. Struct. (THEOCHEM)* **1991**, 229, 63.
- (11) Guo, X.; Randic, M. Recursive Method for Enumeration of Linearly Independent and Minimal Conjugated Circuits of Benzenoid Hydrocarbons. *J. Chem. Inf. Comput. Sci.* **1994**, 34, 339.
- (12) Guo, X.; Randic, M.; Klein, D. J. Analytical Expressions for the Count of LM-Conjugated Circuits of Benzenoid Hydrocarbons. *Int. J. Quantum Chem.* **1996**, 60, 943.
- (13) Clar, E. *The Aromatic Sextet*; Wiley: London, 1972.
- (14) Hosoya, H.; Yamaguchi, T. *Tetrahedron Lett.* **1975**, 4669.
- (15) Ohkami, N.; Motoyama, A.; Yamaguchi, T.; Hosoya, H.; Gutman, I. *Tetrahedron* **1981**, 37, 1113.
- (16) Gutman, I. *Math. Chem.* **1981**, 11, 127.
- (17) Gutman, I. *Bull. Soc. Chim., Beograd* **1982**, 47, 453.
- (18) Zhang, F.; Guo, X. Directed Tree Structure of the Set of Kekule Patterns of Generalized Polyhex Graphs, *Discrete Applied Mathematics*. **1991**, 32, 295.
- (19) Guo, X.; Zhang, F. An Improvement of the Canonical P-V Path Elimination for Recognizing Kekulean Benzenoid systems. *J. Math. Chem.* **1992**, 9, 11.
- (20) Guo, X.; Zhang, F. Mathematical Properties and Structures of Sets of Sextet Patterns of Generalized Polyhexes. *J. Math. Chem.* **1992**, 9, 279.
- (21) Chen, R.; Cyvin, S. J.; Cyvin, B. N. *Math. Chem.* **1990**, 25, 71.
- (22) Li, X.; Zhang, F. *Math. Chem.* **1990**, 25, 153.
- (23) Zhang, F.; Li, X. *Math. Chem.* **1990**, 25, 251.
- (24) Zhang, F.; Chen, R. When each hexagon of a hexagonal system covers it. *Discrete Appl. Math.* **1991**, 30, 63-75.
- (25) Zhang, F.; Zheng, M. Generalized hexagonal systems with each hexagon being resonant. *Discrete Appl. Math.* **1992**, 36, 67.
- (26) Zheng, M. k -Resonant Benzenoid Systems. *J. Mol. Struct. (THEOCHEM)* **1991**, 231, 321.
- (27) Zhang, F.; Chen, R.; Guo, X. Hexagonal System Theory. Manuscript in preparation.
- (28) Chen, R.; Guo, X. k -Coverable Coronoid Systems. *J. Math. Chem.* **1993**, 12, 147.
- (29) Guo, X.; Zhang, F. k -Cycle Resonance Graphs. *Discrete Math.* **1994**, 135, 113.
- (30) Guo, X.; Zhang, F. Planar k -Cycle Resonant Graphs with $k = 1, 2$, Submitted for publication in *Discrete Appl. Math.*
- (31) Lovasz, L.; Plummer, M. D. Matching Theory. *Ann. Discrete Math.* **1986**, 29.
- (32) Heteyi, G. Rectangular configurations which can be covered by 2×1 rectangles. *Pecsi Tan. Foisok. Kozl.* **1964**, 8, 351.
- (33) Lovasz, L. A note on factor-critical graphs. *Stud. Sci. Math. Hung.* **1972**, 7, 279.
- (34) Lovasz, L. Ear-decompositions of matching-covered graphs. *Combinatorica* **1983**, 2, 395.
- (35) Lovasz, L.; Plummer, M. D. On Bicritical Graphs, Infinite and Finite Sets (Colloq. Keszthely, Hungary, 1973), II, *Colloq. Math. Soc. Janos Bolyai* **1975**, 10, 1051.
- (36) Lovasz, L.; Plummer, M. D. On a family of planar bicritical graphs. *Proc. London Math. Soc. Ser.* **1975**, 30 (3), 160.
- (37) Lovasz, L.; Plummer, M. D. On minimal elementary bipartite graphs. *J. Comb. Theory, Ser. B* **1977**, 23, 127.
- (38) Naddef, D.; Pulleyblank, W. R. Ear decompositions of elementary graphs and GF(2)-rank of perfect matchings. *Am. Discrete Math.* **1982**, 16, 241.
- (39) Plummer, M. D. On n -extendable graphs. *Discrete Math.* **1980**, 31, 201.

CI000080Q