

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231462141>

Decacarbonyl(methyl)hydrotriosmium: NMR evidence for a carbon.cntdot..cntdot.hydrogen.cntdot..cntdot.osmium interaction

ARTICLE *in* JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · NOVEMBER 1978

Impact Factor: 12.11 · DOI: 10.1021/ja00492a047

CITATIONS

157

READS

24

2 AUTHORS, INCLUDING:



John R Shapley

University of Illinois, Urbana-Champaign

251 PUBLICATIONS 5,715 CITATIONS


SEE PROFILE

Communications to the Editor

$\text{HOs}_3(\text{CO})_{10}\text{CH}_3$: NMR Evidence for a $\text{C}\cdots\text{H}\cdots\text{Os}$ Interaction¹

Sir:

Compounds involving methyl groups apparently bridging two transition metals are rare, but are becoming less so.² Definitive structural data are very scarce,³ but structural proposals commonly have assumed a symmetrically bridging methyl group, as in I. This structural element is now accepted for $\text{Al}_2(\text{CH}_3)_6$,⁴ in contrast to the briefly controversial, unsymmetrical alternative II.⁵ For lack of evidence to the con-



trary, we^{2a} previously suggested a symmetrically bridging position for the methyl group in the cluster compound $\text{HOs}_3(\text{CO})_{10}\text{CH}_3$. However, we now report novel NMR observations that support a significant $\text{C}\cdots\text{H}\cdots\text{Os}$ interaction for this compound.

Treatment of $\text{D}_2\text{Os}_3(\text{CO})_{10}$ with CH_2N_2 at room temperature provides " $\text{Os}_3(\text{CO})_{10}\text{CH}_2\text{D}_2$ ", which in solution is an equilibrium mixture of partially deuterated methyl and methylene tautomers.^{2a,6} We^{2a} previously noted that the methyl ^1H NMR signal occurs at unusually high field. However, the spectrum of the deuterated material shows separate CH_2D and CHD_2 signals⁷ displaced significantly to even higher field from the CH_3 signal (see Figure 1). The separations $\Delta_1 = \tau(\text{CH}_2\text{D}) - \tau(\text{CH}_3)$ and $\Delta_2 = \tau(\text{CHD}_2) - \tau(\text{CH}_2\text{D})$ vary strongly with temperature, increasing from 0.34 and 0.39 ppm at 35 °C to 0.55 and 0.68 ppm at -76 °C. These large, temperature-dependent values of Δ_1 and Δ_2 are inconsistent with the relatively small effect commonly observed upon geminal substitution of H by D (~ 0.01 ppm)⁸ and therefore require a different explanation.

A model involving a $\text{C}\cdots\text{H}\cdots\text{Os}$ interaction rationalizes the observed effect. For the case in which the methyl group is CH_2D , three structures are possible as shown ($\text{X} = \text{H}$ or D). An isotope effect on this equilibrium is to be expected, since the interacting (bridging) C-H or C-D bond should have a lower stretching force constant than the analogous noninteracting (terminal) bond. This implies a lower frequency,⁹ and hence a lower zero-point energy, leading to a preference for the lighter nucleus in the bridging site.⁶ Thus, each of the H-bridged forms (III-H) will be slightly more abundant than the D-bridged form (III-D). Since the bridging hydrogen atom



also should resonate at higher field than the terminal hydrogen atom, the nonrandom distribution results in a net upfield shift for the CH_2D signal relative to the CH_3 signal. As the temperature is lowered, the equilibrium shifts toward the lower energy H-bridged form, increasing the net shift.

The positions of the CH_3 , CH_2D , and CHD_2 resonances can




Figure 1. A portion of the ^1H NMR spectrum (35 °C) for a sample of " $\text{Os}_3(\text{CO})_{10}\text{CH}_2\text{D}_2$ ", with some " $\text{Os}_3(\text{CO})_{10}\text{CH}_4$ " added as a reference.

be expressed quantitatively in terms of three parameters: τ_b and τ_t , the chemical shifts for the bridging and terminal methyl hydrogen atoms, respectively, and ΔE , the energy difference between the D-bridged and H-bridged forms.¹⁰ Defining $A = \exp(-\Delta E/RT)$, the expressions resulting are the following:

$$\tau(\text{CH}_3) = (2\tau_t + \tau_b)/3 \quad (1)$$

$$\tau(\text{CH}_2\text{D}) = (\tau_t + A\tau_t + \tau_b)/(A + 2) \quad (2)$$


$$\tau(\text{CHD}_2) = (2A\tau_t + \tau_b)/(2A + 1) \quad (3)$$

Equations 1–3 may be solved at each temperature for the three parameters, the mean results of which are $\tau_b = 25 \pm 1$, $\tau_t = 8 \pm 1$, and $\Delta E = 130 \pm 10$ cal/mol.

An abnormally large and temperature-dependent isotope effect is also apparent for the methyl C-H coupling constant. The average value of $^1J(^{13}\text{C}-^1\text{H})$ observed for each type of methyl group at 27 °C is 121.1 (CH_3), 118.9 (CH_2D), and 116.4 Hz (CHD_2), whereas at -80 °C it is 121.1 (CH_3), 116.7 (CH_2D), and 112.3 Hz (CHD_2).¹³ These trends are qualitatively in accord with the interaction model as illustrated by $\text{III-H} \leftrightarrow \text{III-D}$. Quantitative analysis with a set of equations analogous to eq 1–3 leads to the values $J_t = 150 \pm 10$ Hz and $J_b = 60 \pm 20$ Hz.


Certain other models can be eliminated by further consideration of the NMR data. Thus, both the methyl and the methylene ^{13}C NMR signals show normal chemical shift isotope effects, i.e., ~ 0.2 ppm upfield per D atom.⁷ This result

rules out the possibility that the methyl signal could arise from two different carbon sites in rapid equilibrium,¹⁴ since an order of magnitude larger effect would be predicted. Furthermore, eq 1–3 predict that $\Delta_2 > \Delta_1$, as observed. A model involving two interactions per methyl group, as in IV, leads to closely related equations, which however predict that $\Delta_2 < \Delta_1$. Thus, the Δ_2/Δ_1 ratio allows configurations II and IV to be differentiated.



Cotton and co-workers¹⁵ have shown that [EtB(pz)₂]-Mo(CO)₂(η^3 -2-phenylallyl) has a C-H...Mo interaction involving an α -C-H bond of one ethyl group. The strength of the interaction was estimated to be ~ 19 kcal/mol from DNMR studies, but the barrier to exchange between the interacting and noninteracting methylene hydrogens was somewhat lower, ~ 14 kcal/mol.^{15a} We have attempted to determine the barrier to bridge-terminal exchange in $\text{HOs}_3(\text{CO})_{10}\text{CH}_3$, but the methyl ¹H NMR signal does not broaden relative to internal SiMe₄ down to -100°C . From this result an upper limit to ΔG^\ddagger of ~ 5 kcal/mol can be estimated.¹⁶ However, since bridge-terminal exchange could proceed via a doubly bridged configuration such as IV, the exchange barrier may represent only a lower bound for the strength of the C-H...Os interaction.

The general significance of our conclusion that the methyl ligand in $\text{HOs}_3(\text{CO})_{10}\text{CH}_3$ adopts configuration II instead of I is not clear at this time. Neither the Os-Os separation, expected to be $\sim 2.8 \text{ \AA}$ in comparison with 2.61 \AA for Al_2Me_6 ,^{4a} 2.72 \AA for $(\text{MgMe}_2)_n$,¹⁷ 3.06 \AA for $(\eta^5\text{-C}_5\text{H}_5)_2\text{YAlMe}_4$,¹⁸ and $\sim 3.5 \text{ \AA}$ for $[(\eta^5\text{-C}_5\text{H}_5)_2\text{YbMe}]_2$,^{2c} nor steric crowding, which was proposed as determining the formation of Si-H...W bridges in $\text{W}_2(\text{CO})_8(\text{SiHEt}_2)_2$,¹⁹ appears to be a rational basis for the preference. Nevertheless, the new structural model for $\text{HOs}_3(\text{CO})_{10}\text{CH}_3$ leads to a revised and clarified picture (shown) of the hydrogen transfer from carbon to osmium involved in the formation of $\text{H}_2\text{Os}_3(\text{CO})_{10}\text{CH}_2$. Spin saturation transfer experiments confirm the selective exchange between the methyl group in $\text{HOs}_3(\text{CO})_{10}\text{CH}_3$ and one hydride site in $\text{H}_2\text{Os}_3(\text{CO})_{10}\text{CH}_2$.²⁰



Partial deuteration should be a useful probe for C-H...M interactions in other cases.^{21,22} It is noteworthy that the number of H bridges per ligand can be determined even with rapid bridge-terminal exchange. Extension of the method to borohydride-metal complexes should complement the IR technique²³ for determining whether the BH₄ ligand is mono-, bi-, or tridentate.

References and Notes

- (1) This research was supported by National Science Foundation Grant No. CHE 75-14460. We acknowledge a loan of OsO₄ from Engelhard Industries.
- (2) (a) R. B. Calvert and J. R. Shapley, *J. Am. Chem. Soc.*, **99**, 5225 (1977); (b) A. F. Masters, K. Mertis, J. F. Gibson, and G. Wilkinson, *Nouveau J. Chim.*, **1**, 389 (1977); (c) J. Holton, M. F. Lappert, D. G. H. Ballard, R. Pearce, J. L. Atwood, and W. E. Hunter, *J. Chem. Soc., Chem. Commun.*, 480 (1976); (d) K. Fischer, K. Jonas, P. Misbach, R. Stabba, and G. Wilke, *Angew. Chem., Int. Ed. Engl.*, **12**, 943 (1973); B. Bogdanovic, H. Bonnemann, and G. Wilke, *ibid.*, **5**, 582 (1966); (e) P. C. Wailes, H. Weigold, and A. P. Bell, *J. Organomet. Chem.*, **34**, 155 (1972).
- (3) For structurally characterized compounds with $\mu\text{-CH}_2\text{SiMe}_3$ and related groups see the following: (a) R. A. Andersen, R. A. Jones, G. Wilkinson, K. M. A. Malik, and M. B. Hursthouse, *J. Chem. Soc., Chem. Commun.*, 283 (1977); (b) R. A. Andersen, E. Carmona-Guzman, J. F. Gibson, and G. Wilkinson, *J. Chem. Soc., Dalton Trans.*, 2204 (1976); (c) J. A. J. Jarvis, B. T. Kilbourn, R. Pearce, and M. F. Lappert, *J. Chem. Soc., Chem. Commun.*, 475 (1973).
- (4) (a) J. C. Huffman and W. E. Streib, *Chem. Commun.*, 911 (1971); (b) A. Almenningen, S. Salvorsen, and A. Haaland, *Acta Chem. Scand.*, **25**, 1937 (1971); (c) R. G. Vranka and E. L. Amma, *J. Am. Chem. Soc.*, **89**, 3121 (1967).
- (5) (a) S. K. Byram, J. K. Fawcett, S. C. Nyburg, and R. J. O'Brien, *Chem. Commun.*, 16 (1970); (b) M. J. S. Dewar and D. B. Patterson, *ibid.*, 544 (1970); (c) F. A. Cotton, *Inorg. Chem.*, **9**, 2804 (1970).
- (6) A combined neutron-diffraction/NMR study on $\text{HDOs}_3(\text{CO})_{10}(\text{CHD})$ has shown the operation of an equilibrium isotope effect favoring H in the osmium-bound sites and D in the carbon-bound sites: R. B. Calvert, J. R. Shapley, A. J. Schultz, J. M. Williams, S. L. Suib, and G. D. Stucky, *J. Am. Chem. Soc.*, **100**, 6240 (1978). The effect is readily rationalized in terms of the zero-point energy difference between C-H and Os-H-Os vibrational modes.
- (7) Evidence for the assignment of these signals is the following. (a) Samples with a lower level of deuteration show the expected alteration in intensities; i.e., the CH₃ signal increases at the expense of the CHD₂ signal. (b) Near room temperature the line widths are in the order CH₃ < CH₂D < CHD₂, consistent with partially relaxed H-D coupling. As the temperature is lowered, quadrupolar relaxation apparently becomes more effective and by -60°C all three signals have the same width. (c) The ¹³C NMR spectrum of "¹³Os₃(CO)₁₀¹³CH₂D₂" shows overlapping triplet and quintet patterns due to ¹³C-H coupling.
- (8) J. B. Lambert and L. G. Greifenstein, *J. Am. Chem. Soc.*, **96**, 5120 (1974), and references therein.
- (9) We have observed IR bands at 2985 (m), 2930 (s), 2840 (w), and 2520 (s, br) cm⁻¹ for a carefully purified sample of the $\text{HOs}_3(\text{CO})_{10}\text{CH}_3\text{-H}_2\text{Os}_3(\text{CO})_{10}\text{CH}_2$ mixture in CCl₄ solution. The broad band near 2520 cm⁻¹ appears to be a $\nu_{\text{CO}} + \nu_{\text{Os-C}}$ combination band.^{2a} The 2840-cm⁻¹ band may be due to the C-H₂ moiety. However, we note that a similar band has been observed for $\text{Mn}(\text{CO})_5\text{CH}_3$ and assigned as an overtone: A. B. Dempster, D. B. Powell, and N. Sheppard, *J. Chem. Soc. A*, 1129 (1970). We are attempting to assign the bending modes for "¹³Os₃(CO)₁₀CH₄" and "¹³Os₃(CO)₁₀CH₂D₂" in order to identify those associated with the perturbed C-H bond.
- (10) Using only one chemical shift parameter for the terminal positions is an approximation, since the two sites are diastereotopic. However, the chemical shift difference between the two analogous sites in $\text{HOs}_3(\text{CO})_{10}(\text{CH}_2\text{CO}_2\text{Et})^{11}$ is only 0.2 ppm, which is insignificant in comparison with the 17-ppm overall bridge-terminal shift difference in $\text{HOs}_3(\text{CO})_{10}\text{CH}_3$. Furthermore, although ΔE and $\Delta \tau$ could be slightly different for the CH₂D compound than for the CHD₂ compound, separate analysis of the two sets of shifts as a function of temperature gave identical results within experimental error. A closely similar, though not identical, analysis has been applied in studying the effect of deuterium substitution on equilibrating carbonium ions.¹²
- (11) J. B. Keister and J. R. Shapley, *J. Am. Chem. Soc.*, **98**, 1056 (1976).
- (12) M. Saunders, L. Telkowski, and M. R. Kates, *J. Am. Chem. Soc.*, **99**, 8070 (1977); M. Saunders, M. H. Jaffe, and P. Vogel, *ibid.*, **93**, 2558 (1971).
- (13) The observed values of $\Delta^1\text{J}(\text{CH})_{\text{av}}$ are significantly larger than the secondary isotope effects sometimes observed; e.g., $^1\text{J}(\text{CH})$ is estimated to be 0.6 ± 0.4 Hz lower for $\text{C}_6\text{H}_5^{13}\text{CH}_2\text{D}$ than for $\text{C}_6\text{H}_5^{13}\text{CH}_3$; M. Murray, *J. Magn. Reson.*, **9**, 326 (1973).
- (14) One possible model for the two sites would be a symmetrically bridging methyl and a terminal methylene.
- (15) (a) F. A. Cotton and A. G. Stanislawski, *J. Am. Chem. Soc.*, **96**, 5074 (1974); (b) F. A. Cotton, T. LaCour, and A. G. Stanislawski, *ibid.*, **96**, 754 (1974).
- (16) Approximating by use of two-site exchange expressions, for fast exchange $k_1 = (\pi/2)(1/\Delta W) \times (\Delta\nu)^2$ and at coalescence $k_c = (\pi/\sqrt{2})(\Delta\nu)$. Assuming $\Delta\nu = 1700$ Hz and that $\Delta W \leq 1$ Hz at 163 K, the former expression leads to ΔG^\ddagger (163 K) ≤ 4.4 kcal/mol. The coalescence expression with $T_c \leq 143$ K leads to ΔG^\ddagger (143 K) ≤ 5.8 kcal/mol. An exchange barrier of 3.6 kcal/mol has been estimated from ¹³C T₁ data: J. R. Norton and R. Jordan, personal communication.
- (17) E. Weiss, *J. Organomet. Chem.*, **2**, 314 (1964).
- (18) J. Holton, M. F. Lappert, G. F. Scollary, D. G. H. Ballard, R. Pearce, J. L. Atwood, and W. E. Hunter, *J. Chem. Soc., Chem. Commun.*, 425 (1976).
- (19) M. J. Bennett and K. A. Simpson, *J. Am. Chem. Soc.*, **93**, 7156 (1971). See also M. Cowie and M. J. Bennett, *Inorg. Chem.*, **16**, 2325 (1977).
- (20) The hydride site assignments for $\text{H}_2\text{Os}_3(\text{CO})_{10}\text{CH}_2$ are based on chemical shift and coupling constant comparisons. These arguments together with the spin saturation transfer data will be presented in a full paper.
- (21) M. Brookhart, T. H. Whitesides, and J. M. Crockett, *Inorg. Chem.*, **15**, 1550 (1976).
- (22) S. D. Ittel, F. A. Van-Catledge, C. A. Tolman, and J. P. Jesson, *J. Am. Chem. Soc.*, **100**, 1317 (1978). Similar effects have been observed upon D⁺ incorporation into the complexes described (S. D. Ittel, personal communication).
- (23) T. J. Marks and J. R. Kolb, *Chem. Rev.*, **77**, 263 (1977).

R. Bruce Calvert, John R. Shapley*

Department of Chemistry, University of Illinois
Urbana, Illinois 61801

Received July 28, 1978