

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231624941>

Asymmetric Addition of Trimethylsilyl Cyanide to Benzaldehydes Catalyzed by Samarium(III) Chloride and Chiral Phosphorus(V) Reagents

ARTICLE *in* THE JOURNAL OF ORGANIC CHEMISTRY · JANUARY 1998

Impact Factor: 4.72 · DOI: 10.1021/jo971868j

CITATIONS

53

READS

26

2 AUTHORS, INCLUDING:



Wen-Bin Yang

Academia Sinica

63 PUBLICATIONS 1,059 CITATIONS

SEE PROFILE

Asymmetric Addition of Trimethylsilyl Cyanide to Benzaldehydes Catalyzed by Samarium(III) Chloride and Chiral Phosphorus(V) Reagents

Wen-Bin Yang and Jim-Min Fang*

Department of Chemistry, National Taiwan University,
Taipei 106, Taiwan, Republic of China

Received October 9, 1997

Chiral phosphorus(V) reagents have been used as resolving agents,¹ auxiliaries² or ligands³ to promote various asymmetric reactions. In general, those P(V)-reagents prepared from C2-symmetric diols or diamines do not contain asymmetric phosphorus centers. We speculate that a bis-phosphoramidate reagent, such as **3a**, not only exhibits the advantageous effect of C2-symmetry, but also exerts better asymmetric induction for it contains phosphorus stereocenters closer to reactive sites.⁴ In this paper, we focus on the combined use of Lewis acid and bis-phosphoramidate reagent in promotion of asymmetric cyanosilylation of benzaldehydes.

Cyanohydrins are important synthons,⁵ such as in preparation of α -hydroxy acids and β -amino alcohols. Optically active cyanohydrins can be obtained by biological⁶ or chemical⁷ methods. For example, racemic cyanohydrins can be resolved by lipase-catalyzed acetylation.^{5a,6d}

A variety of aromatic aldehydes and some aliphatic aldehydes can be converted to optically active cyanohydrins by the catalysis of oxynitrilases.^{6a} Catalytic asymmetric addition of HCN or Me₃SiCN onto aldehydes can be achieved by using chiral ligands, such as dipeptides (especially diketopiperazines derived from histidine and phenylalanine),^{7f} bisoxazolines,^{7j} salens,⁷ⁿ Schiff's bases,^{7k} sulfoximines,^{7m} cinchonine,^{7g} and binaphthol,^{7q} together with Lewis acids containing Ti(IV) (most frequently used),^{7a} Al(III),^{7h} B(III),^{7c} Y(III),^{7p} Sn(II),^{7g} or Mg(II)^{7j} ions. However, most of reports indicate that the reaction must be conducted at low temperature (e.g. -78 °C) in order to obtain high enantioselectivity.

The bis-phosphoramidate **3a** was prepared in 77% overall yield by treatment of (1*R*,2*S*)-(-)-ephedrine subsequently with POCl₃ and ethylenediamine (Scheme 1).^{8a} The (2*S*,2'*S*,4*S*,4'*S*,5*R*,5'*R*)-configuration of **3a** was unambiguously determined by NMR and X-ray analyses. A minor diastereomer **3b** exhibiting the (2*R*,2'*R*,4*S*,4'*S*,5*R*,5'*R*)-configuration was also prepared. By a similar procedure, compound **3c** (the antipode of **3a**) was prepared from (1*S*,2*R*)-(+)-ephedrine. A preliminary survey of the reaction of Me₃SiCN with benzaldehyde in the presence of the chiral ligand **3a** indicated that Lewis acids SmCl₃, SmI₃, LaCl₃, and Sc(OTf)₃ were effective catalysts, whereas Sm(O*i*-Pr)₃ or Ti(O*i*-Pr)₄ were ineffective, in terms of conversion and enantioselectivity. We report herein mainly the SmCl₃-catalyzed cyanosilylation of various benzaldehydes.⁹ Comparison experiments showed that the SmCl₃-catalyzed cyanosilylation was accelerated by addition of the bis-phosphoramidate reagent **3a**. However, use of **3a** alone (as a base) did not promote the cyanosilylation.

(1) Chiral P(V)-reagents as resolving agents: (a) Jacques, J.; Fouquey, C.; Vitero, R. *Tetrahedron Lett.* **1971**, 4617. (b) Mikes, F.; Boshart, G. *J. Chem. Soc., Chem. Commun.* **1978**, 173. (c) Arnold, W.; Daly, J. J.; Imhof, R.; Kyburz, E. *Tetrahedron Lett.* **1983**, 24, 343. (d) Anderson, R. C.; Shapiro, M. J. *J. Org. Chem.* **1984**, 49, 1304. (e) Johnson, C. R.; Elliott, R. C.; Penning, T. D. *J. Am. Chem. Soc.* **1984**, 106, 5019. (f) Alexakis, A.; Mutti, S.; Normant, J. F.; Mangeney, P. *Tetrahedron Asymmetry* **1990**, 1, 437.

(2) Representative examples of chiral P(V)-auxiliaries: (a) Hua, D. H.; Chan-Yu-King, R.; McKie, J. A.; Myer, L. *J. Am. Chem. Soc.* **1987**, 109, 5026. (b) Sting, M.; Steglich, W. *Synthesis* **1990**, 132. (c) Denmark, S. E.; Chen, C.-T. *J. Am. Chem. Soc.* **1992**, 114, 10674. (d) Narasaka, K.; Hidai, E.; Hayashi, Y.; Gras, J.-L. *J. Chem. Soc., Chem. Commun.* **1993**, 102. (f) Hanessian, S.; Andreotti, D.; Gomtsyan, A. *J. Am. Chem. Soc.* **1995**, 117, 10393. (g) Schrader, T. *Angew. Chem., Int. Ed. Engl.* **1995**, 34, 917.

(3) For reviews: (a) Hall, C. R.; Inch, T. D. *Tetrahedron* **1980**, 21, 2059. (b) Katti, K. V.; Reddy, V. S.; Singh, P. R. *Chem. Soc. Rev.* **1995**, 97. Representative examples for using chiral P(V)-reagents as ligands: (c) Soai, K.; Hirose, Y.; Ohno, Y. *Tetrahedron Asymmetry* **1993**, 4, 1473. (d) Hulst, R.; Heres, H.; Fitzpatrick, K.; Peper, N. C. M. W.; Kellogg, R. M. *Tetrahedron Asymmetry* **1996**, 7, 2755. Representative examples for using chiral P(V)-reagents as Lewis bases: (e) Chiodi, O.; Fotiadu, F.; Sylvestre, M.; Buono, G. *Tetrahedron Lett.* **1996**, 37, 39. (f) Denmark, S. E.; Wong, K.-T.; Stavenger, R. A. *J. Am. Chem. Soc.* **1997**, 119, 2333. (g) Burns, B.; Gamble, M. P.; Simm, A. R. C.; Studley, J. R.; Alcock, N. W.; Wills, M. *Tetrahedron Asymmetry* **1997**, 8, 73.

(4) For a chiral bis-azaphosphole: Brenchley, G.; Fedouloff, M.; Merifield, E.; Wills, M. *Tetrahedron Asymmetry* **1996**, 7, 2809.

(5) Examples for transformation of optically active cyanohydrins: (a) Ohta, H.; Kimura, Y.; Sugano, Y. *Tetrahedron Lett.* **1988**, 29, 6957. (b) Effenberger, F.; Stelzer, U. *Angew. Chem., Int. Ed. Engl.* **1991**, 30, 873. (c) Warmerdam, E. G. J. C.; Brusse, J.; Kruse, C. G.; Van der Gen, A. *Tetrahedron* **1993**, 49, 1063.

(6) A review for preparation of optically active cyanohydrins by using biological catalysts: (a) Effenberger, F.; *Angew. Chem., Int. Ed. Engl.* **1994**, 33, 1555. Representative examples: (b) Brusse, J.; Roos, E. C.; Van der Gen, A. *Tetrahedron Lett.* **1988**, 29, 4485. (c) Brusse, J.; Loos, W. T.; Kruse, C. G.; Van der Gen, A. *Tetrahedron* **1990**, 46, 979. (d) Inagaki, M.; Hiratake, J.; Nishioka, T.; Oda, J. *J. Am. Chem. Soc.* **1991**, 113, 9360. (e) Ognyanov, V. I.; Datcheva, V. K.; Kyler, K. S. *J. Am. Chem. Soc.* **1991**, 113, 6992. (f) Klempier, N.; Griengl, H.; Hayn, M. *Tetrahedron Lett.* **1993**, 34, 4769. (g) Schmidt, M.; Herve, S.; Klempier, N.; Griengl, H. *Tetrahedron* **1996**, 52, 7833.

(7) Reviews for preparation of optically active cyanohydrins and their silyl ether derivatives by using chemical catalysts: (a) North, M. *Synlett* **1993**, 807. (b) Hayashi, M.; Oguni, N. *J. Synth. Org. Chem.* **1994**, 52, 488. Representative examples: (c) Reetz, M. T.; Kunisch, R.; Heitmann, P. *Tetrahedron Lett.* **1986**, 39, 4721. (d) Matthews, B. R.; Jackson, W. R.; Jayatilake, G. S.; Wilshire, C.; Jacobs, H. A. *Aust. J. Chem.* **1988**, 41, 1697. (e) Minamikawa, H.; Hayakawa, S.; Yamada, N.; Iwasawa, N.; Narasaka, K. *Bull. Chem. Soc. Jpn.* **1988**, 61, 4379. (f) Tanaka, K.; Mori, A.; Inoue, S. *J. Org. Chem.* **1990**, 55, 181. (g) Kobayashi, S.; Tsuchiya, Y.; Mukaiyama, T. *Chem. Lett.* **1991**, 541. (h) Ohno, H.; Nitta, H.; Tanaka, K.; Mori, A.; Inoue, S. *J. Org. Chem.* **1992**, 57, 6778. (i) Nitta, H.; Yu, D.; Kudo, M.; Mori, A.; Inoue, S. *J. Am. Chem. Soc.* **1992**, 114, 7969. (j) Corey, E. J.; Wang, Z. *Tetrahedron Lett.* **1993**, 34, 4001. (k) Hayashi, M.; Miyamoto, Y.; Inoue, T.; Oguni, N. *J. Org. Chem.* **1993**, 58, 1515. (l) Kim, H. J.; Jackson, W. R. *Tetrahedron Asymmetry* **1994**, 5, 1541. (m) Bolm, C.; Müller, P. *Tetrahedron Lett.* **1995**, 36, 1625. (n) Pan, W.; Feng, X.; Gong, L.; Hu, W.; Li, Z.; Mi, A.; Jiang, Y. *Synlett* **1996**, 337. (o) Shvo, Y.; Gal, M.; Becker, Y.; Elgovi, A. *Tetrahedron Asymmetry* **1996**, 7, 911. (p) Abiko, A.; Wang, G.-q. *J. Org. Chem.* **1996**, 61, 2264. (q) Mori, M.; Imma, H.; Nakai, T. *Tetrahedron Lett.* **1997**, 38, 6229.


(8) Examples of ephedrine-derived P(V)-reagents: (a) Cooper, D. F.; Hall, C. R.; Harrison, J. M.; Inch, T. D. *J. Chem. Soc., Perkin Trans. 1* **1977**, 1969. (b) Welch, S. C.; Levine, J. A.; Bernal, I.; Cetrullo, J. *J. Org. Chem.* **1990**, 55, 5991. (c) Sum, V.; Davies, A. J.; Kee, T. P. *J. Chem. Soc., Chem. Commun.* **1992**, 1771. (d) Jommi, G.; Miglierini, G.; Pagliarini, R.; Sello, G.; Sisti, M. *Tetrahedron* **1992**, 48, 7275. (e) Neidlein, R.; Keller, H.; Boese, R. *Heterocycles* **1993**, 35, 1185. (f) Sheehan, S. K.; Jiang, M.; McKinsty, L.; Livinghouse, T.; Garton, D. *Tetrahedron* **1994**, 50, 6155.

(9) Examples for lanthanide ion-promoted asymmetric reactions: (a) Evans, D. A.; Hoveyda, A. H. *J. Am. Chem. Soc.* **1990**, 112, 6447. (b) Shibasaki, M.; Sasai, H. *J. Synth. Org. Chem. Jpn.* **1993**, 51, 972. (c) Giardello, M. A.; Conticello, V. P.; Brard, L.; Gagné, M. R.; Marks, T. J. *J. Am. Chem. Soc.* **1994**, 116, 10241. (d) Sasai, H.; Arai, S.; Tahara, Y.; Shibasaki, M. *J. Org. Chem.* **1995**, 60, 6656. (e) Sasai, H.; Arai, T.; Satow, Y.; Houk, K. N.; Shibasaki, M. *J. Am. Chem. Soc.* **1995**, 117, 6194.

Table 1. Addition Reaction of Benzaldehyde with Trimethylsilyl Cyanide in the Presence of Lewis Acid SmCl₃ and Chiral Ligand **3a**, Giving the Silyl Ether **4a**^a


entry	SmCl ₃ /equiv	3a /equiv	solvent	reaction temp/°C	reaction time/h	product 4a , ee/% ^b	config ^c
1	0.1	0.1	CH ₂ Cl ₂	25 ± 2	5	65	<i>R</i>
2	0.01	0.01	PhCH ₃	25 ± 2	4.5	72	<i>R</i>
3	0.01	0.01	C ₆ H ₁₄	25 ± 2	6	12	<i>R</i>
4	0.01	0.01	PhH	25 ± 2	6	51	<i>R</i>
5	0.01	0.01	THF	25 ± 2	4.5	8	<i>R</i>
6	0.01	0.01	CH ₃ CN	25 ± 2	4.5	12	<i>S</i>
7	0.01	0.01	EtOH	25 ± 2	6	0	
8	0.01	0.01	CH ₂ Cl ₂ ^d	25 ± 2	6	42	<i>R</i>
9	0.1	0.1	CH ₂ Cl ₂	-15 ± 5	15	75	<i>R</i>
10	0.01	0.01	CH ₂ Cl ₂ ^e	-15 ± 5	16	67	<i>R</i>
12	0.1	0.1 ^f	CH ₂ Cl ₂	-15 ± 5	24	63	<i>R</i>
11	0.1	0.1 ^g	CH ₂ Cl ₂	25 ± 2	4	62	<i>R</i>
13	0.01	0.01	PhCH ₃	-15 ± 5	16	80	<i>R</i>
14	0.001	0.003	PhCH ₃	-15 ± 5	15	84	<i>R</i>
15	0.001	0.003 ^g	PhCH ₃	-15 ± 5	15	76	<i>R</i>

^a Molar ratio: PhCHO/Me₃SiCN = 1:2. The reaction completed in the indicated time (>97% conversion) according to the ¹H NMR analysis. ^b The ee value was determined by comparison of optical rotation with the reported value^{6g} of the cyanohydrin **4b**, [α]_D²⁰ +45 (CHCl₃, *c* 1). Occasionally, **4b** was converted to the corresponding MTPA ester, of which diastereomeric ratio was determined by the ¹⁹F NMR analysis. In entry 13, the silyl ether **4a** was analyzed by HPLC on a Chiralcel OD column. ^c The configuration of major enantiomer. ^d EtOH (0.01 equiv) was added. ^e 2,6-Lutidine (0.05 equiv) was added. ^f The crude ligand **3a**, without recrystallization, was used. ^g The recovered ligand **3a** was reused.



Reagents and conditions: (i) POCl₃, Et₃N, PhH, -10 °C, 6 h; **2a** (85%), **2b** (8%), **2c** (87%). (ii) H₂NCH₂CH₂NH₂, Et₃N, THF, 27 °C, 6 h; **3a** (90%), **3b** (93%), **3c** (90%).

A combined use of SmCl₃ and bis-phosphoramidate **3a** (with phosphorus stereocenters) showed significant asymmetric induction in the cyanosilylation of benzaldehyde (Table 1). Thus, treatment of PhCHO with Me₃SiCN (1.5–2 equiv), SmCl₃ (1 mol %), and ligand **3a** (1 mol %) in toluene (5 mL) at room temperature for 5 h gave a quantitative yield of cyanohydrin silyl ether **4a** with 72% enantiomeric excess (ee). Toluene was the solvent of choice; CH₂Cl₂ and benzene were also suitable, but not hexane, THF, CH₃CN, or EtOH (entries 1–7). The ee value of **4a** was determined by HPLC analysis on a Chiralcel OD column. Furthermore, hydrolysis of **4a** yielded the corresponding dextrorotatory cyanohydrin **4b**.

**Figure 1.** The major enantiomers obtained from the cyanosilylation of benzaldehydes (Tables 1 and 2).

By comparison with the reported optical rotation,^{6g} [α]_D²⁰ +45 (CHCl₃, *c* 1, >99% *R*-isomer), the *R*-configuration for the major enantiomers in **4a** and **4b** was deduced. Cyanohydrin **4b** was also converted to the corresponding MTPA esters **4c**, of which diastereomeric ratio was determined by the ¹H and ¹⁹F NMR or HPLC analyses to double check the optical purity of **4a**. The MTPA ester was generally prepared by treatment of a cyanohydrin with (*R*)-(+)- α -methoxy- α -(trifluoromethyl)phenylacetyl chloride and Et₃N at room temperature for 1 h. Use of a stronger base, 4-(dimethylamino)pyridine, or prolonged reaction period might cause some degree of racemization.^{7d}

The presumed SmCl₃–ligand complex was insoluble or sparingly soluble in toluene or CH₂Cl₂ at the outset of cyanosilylation. However, the heterogeneous mixture became homogeneous solution at the end of reaction. After the reaction, the ligand **3a** was easily separated by filtration. The recovered ligand (>80% yield) could be reused in the cyanosilylation with slightly decreased enantioselectivity (compared entries 1 vs 11, and 14 vs 15). Modest enantioselectivity was also procured by using crude ligand **3a** (entry 12). However, the Lewis acid SmCl₃ must be kept from moisture, otherwise, the enantioselectivity decreased dramatically. The turnover of this cyanosilylation was very high; thus, up to 84% ee of (*R*)-**4a** was obtained from the reaction at -15 °C (entry 14) using very small amounts of SmCl₃ (0.1 mol %) and ligand **3a** (0.3 mol %).

Addition of protic solvent, such as EtOH in entry 8, greatly deteriorated the asymmetric cyanosilylation; however, addition of lutidine only caused a slight change of enantioselectivity (entry 10). Beside solvent effect, the reaction temperature appeared to be another key factor for achieving asymmetric induction. The reaction at -15 °C generally resulted in higher enantioselectivity than

Table 2. Cyanosilylation of Aldehydes Catalyzed by SmCl₃ and Ligand **3a**^a

entry	aldehyde	SmCl ₃ /equiv	3a /equiv	reaction temp/°C ^a	product	[α] _D (CHCl ₃)	ee/%	config ^b
1 ^c	<i>p</i> -MeOC ₆ H ₄ CHO	0.1	0.1	25 ± 2	5a		61 ^{d,e}	<i>R</i>
2	<i>p</i> -MeOC ₆ H ₄ CHO	0.002	0.006	−15 ± 5	5a	+20.14	90 ^{d,e,f}	<i>R</i>
3	<i>p</i> -MeOC ₆ H ₄ CHO	0.001	0.003	−15 ± 5	5a	+18.13	81 ^{d,e,f}	<i>R</i>
4	<i>m</i> -MeOC ₆ H ₄ CHO	0.002	0.006	−15 ± 5	6a	+17.62	83 ^{d,e,f}	<i>R</i>
5	<i>o</i> -MeOC ₆ H ₄ CHO	0.002	0.006	−15 ± 5	7a	+7.55	80 ^{d,f}	<i>R</i>
6 ^c	<i>p</i> -CH ₃ C ₆ H ₄ CHO	0.1	0.11	25 ± 2	8a		56 ^{d,e}	<i>R</i>
7	<i>p</i> -CH ₃ C ₆ H ₄ CHO	0.002	0.006	−15 ± 5	8a	+18.54	66 ^{d,e,f}	<i>R</i>
8	<i>p</i> -CH ₃ C ₆ H ₄ CHO	0.001	0.01	−15 ± 5	8a	+20.29	73 ^{d,e,f}	<i>R</i>
9	<i>p</i> -PhC ₆ H ₄ CHO	0.002	0.006	−15 ± 5	9a	+7.35	49 ^{d,f}	<i>R</i>
10	<i>m</i> -PhOC ₆ H ₄ CHO	0.002	0.006	−15 ± 5	10a	+6.37	45 ^{d,f}	<i>R</i>
12	<i>p</i> -FC ₆ H ₄ CHO	0.002	0.006	−15 ± 5	11a	+17.25	77 ^{d,f}	<i>R</i>
11 ^c	<i>p</i> -NCC ₆ H ₄ CHO	0.1	0.11	25 ± 2	12a		20 ^d	<i>R</i>
13	<i>p</i> -NCC ₆ H ₄ CHO	0.002	0.006	−15 ± 5	12a	+7.43	35 ^{d,f}	<i>R</i>
14	<i>p</i> -O ₂ NC ₆ H ₄ CHO	0.001	0.003	−15 ± 5	13a		12 ^f	<i>R</i>
15	<i>p</i> -O ₂ NC ₆ H ₄ CHO	0.002	0.006	−70 ± 5	13a	+4.40	29 ^f	<i>R</i>

^a The reactions were generally conducted in toluene with a molar ratio of PhCHO/Me₃SiCN = 1:2. ^b The configuration of major enantiomer. ^c The reactions were conducted in CH₂Cl₂. ^d The value was deduced from the optical rotation of the corresponding cyanohydrin. ^e The value was determined by NMR analysis of the corresponding MTPA ester. ^f The value was determined by HPLC analysis of silyl ether on a Chiralcel OD column.

that performed at room temperature. The reaction at lower temperature (such as −40 or −78 °C) was very slow (impractical), though the enantioselectivity might be improved.

As one can expect that (*S*)-**4a** was obtained from the cyanosilylation of benzaldehyde by using the antipode **3c** and SmCl₃ as the combined catalyst. However, the reaction using ligand **3b** (a diastereomer of **3a**) resulted in a low enantioselectivity.

The asymmetric cyanosilylation of substituted benzaldehydes were also carried out by the catalysis of SmCl₃ and **3a** (Table 2). Benzaldehydes bearing electron-donating groups tended to give higher enantioselectivity than those bearing electron-withdrawing groups. The cyanosilylation of 4-methoxybenzaldehyde at −15 °C by using SmCl₃ (0.2 mol %) and **3a** (0.6 mol %) yielded the silyl ether **5a** with predominance of (*R*)-enantiomer (90% ee).

The real nature of the active catalyst awaits further investigation. It is not a trivial problem especially in dealing with the SmCl₃–ligand complexes of tiny and hygroscopic powder forms. Two preliminary experiments were carried out as follows. First, SmCl₃ (0.01 mmol) and **3a** (0.01 mmol) were mixed in CH₂Cl₂ (5 mL). The “clear” solution was taken by filtration through fritted glass (10–16 mm porosity) or by centrifuge. The cyanosilylation of benzaldehyde (1 mmol) was then conducted in this “clear” CH₂Cl₂ solution at room temperature to give the product **4a** with 56% ee (somewhat lower than 65% ee listed in the entry 1 of Table 1). On the other hand, SmCl₃ (0.01 mmol) and **3a** (0.01 mmol) were mixed in toluene (5 mL). The “clear” solution and the solid mass were separated. No cyanosilylation was effected in the “clear” toluene solution. On addition of “fresh” toluene to the solid mass, the suspension then functioned as an active catalyst to effect the cyanosilylation with comparable rate and enantioselectivity as shown in the entry 2 of Table 1. At this moment, we cannot conclude whether the solid mass or the minute soluble species is the real catalyst for asymmetric cyanosilylations.

In summary, bis-phosphoramidate reagents **3a** or **3c** bearing phosphorus stereocenters were readily prepared from inexpensive ephedrine, POCl₃, and ethylenediamine. Asymmetric cyanotrimethylsilylation of various benzaldehydes was realized by using very small amounts

of SmCl₃ and the C₂-symmetric chiral ligand (**3a** or **3c**) at −15 °C or room temperature. Although the enantioselectivity of cyanosilylation products is not extremely high, our method still exhibits several promising features: (i) the cyanosilylation occurs with reasonable enantioselectivity at room temperature, (ii) the bis-phosphoramidate ligand can be used in minute amounts to effect the asymmetric cyanosilylation, and (iii) the ligand is easily separated by simple filtration from the reaction mixture (after trituration with Et₂O) and reused in the cyanosilylation.

Experimental Section

Melting points are uncorrected. Optical rotations were measured on a digital polarimeter with a cuvette of 1 dm length. ¹H NMR spectra were recorded at 200, 300, or 400 MHz; ¹³C NMR spectra were recorded at 50, 75, or 100 MHz; ¹⁹F NMR spectra were recorded at 376 MHz; ³¹P NMR spectra were recorded at 81, 121, or 162 MHz. Tetramethylsilane (δ = 0 ppm) was used as internal standard in ¹H NMR spectra; phosphoric acid (δ = 0 ppm) or triphenylphosphine (δ = −17.16 ppm) were used as external standards in ³¹P NMR spectra; trifluorotoluene (δ = −67.73 ppm) was used as external standard in ¹⁹F NMR spectra. Mass spectra were recorded at an ionizing voltage of 70 or 20 eV. Merck silica gel 60F sheets were used for analytical thin-layer chromatography. Column chromatography was performed on SiO₂ (70–230 mesh); gradients of EtOAc and *n*-hexane were used as eluents. High-pressure liquid chromatography was carried out on a liquid chromatograph equipped with UV (254 nm) and refractive index detectors. Enantiomeric excess was determined by HPLC using a Chiralcel OD column (0.46 cm ID × 25 cm).

Preparation of Chiral Phosphorus(V) Ligands. A benzene solution (100 mL) of (1*R*,2*S*)-(−)-ephedrine hydrochloric salt (4.1 g, 20 mmol) and Et₃N (10 mL, 72 mmol) was cooled to −10 °C in an ice–salt bath. POCl₃ (2.0 mL, 20 mmol) was added dropwise, and the mixture was stirred for 6 h. The resulting precipitates were filtered, and the filtrate was concentrated and subjected to chromatography (silica gel, EtOAc/hexane (1:2)) to give (2*R*,4*S*,5*R*)-2-chloro-3,4-dimethyl-5-phenyl-1,3,2-oxazaphospholidin-2-one (**2a**, 4.17 g, 85%) and the (2*S*,4*S*,5*R*)-isomer **2b** (0.42 g, 8%).

A THF solution (10 mL) of ethylenediamine (0.7 mL, 11 mmol) and Et₃N (2 mL, 14 mmol) was added dropwise to a THF solution (30 mL) of **2a** (2.6 g, 11 mmol) at room temperature (27 °C). The mixture was stirred for 6 h, and the resulting precipitates were filtered. The filtrate was concentrated to give crude **3a** as solids, which were recrystallized from CH₂Cl₂/EtOAc (1:1) to afford pure product (2.4 g, 90%). (2*S*,2'*S*,4*S*,4'*S*,5*R*,5'*R*)-*N,N*-Bis(3,4-dimethyl-2-oxo-5-phenyl-1,3,2-oxazaphospholan-2-yl)-

ethane-1,2-diamine (**3a**): Solid, mp 199–201 °C; $[\alpha]_D^{19}$ –115.9 (CHCl₃, *c* 4); TLC (MeOH/CH₂Cl₂ (1:4)) R_f = 0.41; IR (KBr) 3230, 1453, 1328, 1242 cm⁻¹; ¹H NMR (CDCl₃, 200 MHz) δ 0.69 (6 H, d, *J* = 6.6 Hz, two CH₃), 2.70 (6 H, d, *J* = 9.5 Hz, two CH₃), 3.08–3.24 (4 H, m, CH₂), 3.50–3.68 (2 H, m, H-4), 3.66–3.80 (2 H, m, NH, shiftable), 5.64 (2 H, d, *J* = 6.6 Hz, H-5, H-5'), 7.15–7.38 (10 H, m); ³¹P NMR (CDCl₃, 121 MHz) δ 25.18; ¹³C NMR (CDCl₃, 50 MHz) δ 14.7, 29.0 (d, *J*_{P-C} 5.9 Hz), 43.3 (d, *J*_{P-C} 5.7 Hz), 59.2 (d, *J*_{P-C} 12.2 Hz), 78.3, 125.6, 127.9, 128.3, 136.4 (d, *J*_{P-C} 10.4 Hz); FAB-MS *m/z* 479.2 (*M*⁺ + 1); HRMS calcd for [*M*]⁺ 478.1879, found *m/z* 478.1913. Anal. Calcd for C₂₂H₃₂N₄O₄P₂: C, 55.23; H, 6.74; N, 11.71. Found: C, 55.23; H, 6.58; N, 11.31.

By a procedure similar to that for **3a**, the minor chloro compound **2b** reacted with ethylenediamine to give **3b** (2*R*,2'*R*,4*S*,4'*S*,5*R*,5'*R*) in 93% yield after recrystallization. Solid, mp 193–195 °C (from CH₂Cl₂/EtOAc (1:1)); $[\alpha]_D^{28}$ –43.01 (CHCl₃, *c* 0.6). (1*S*,2*R*)-(+)-Ephedrine hydrochloric acid reacted with POCl₃, followed by treatment with ethylenediamine to give **3c** (2*R*,2'*R*,4*R*,4'*R*,5*S*,5'*S*) in 78% total yield. Solid, mp 200–201 °C (from CH₂Cl₂/EtOAc (1:1)); $[\alpha]_D^{28}$ +115.5 (CHCl₃, *c* 0.6).

General Procedure for Asymmetric Reaction of Aldehydes and Trimethylsilyl Cyanide. Under an atmosphere of nitrogen, a suspension of SmCl₃ (0.5 mg, 0.002 mmol) and ligand **3a** (ground to fine powders, 3.0 mg, 0.006 mmol) in toluene (5 mL) was placed in a flame-dried flask and cooled to –78 °C. Benzaldehyde (freshly distilled, 106 μ L, 1.0 mmol) and Me₃SiCN (200–266 μ L, 1.5–2.0 mmol) were added sequentially at –78 °C. The heterogeneous mixture was stirred for 24 h while the temperature was kept between –10 and –20 °C in a freezer. The mixture appeared as a homogeneous, transparent solution at this stage. Toluene and excess of Me₃SiCN were removed in vacuo, and the residue was triturated with Et₂O (3 mL). The precipitates were filtered through a pad of anhydrous Na₂SO₄ and rinsed with Et₂O (3 mL). The filtrate was concentrated in vacuo to give a cyanohydrin silyl ether **4a** (201 mg, 99%) enriched in (*R*)-enantiomer. The ratio of enantiomers was determined by HPLC analysis on a Chiralcel OD column (*i*-PrOH/hexane (0.25:100), 2.5 mL/min flow rate). The mixture of precipitates and Na₂SO₄ was taken up with CH₂Cl₂ (3 mL), and the soluble part was concentrated in vacuo to recover the ligand **3a** (80–92%).

General Procedure for Hydrolysis of Silyl Ethers. The cyanohydrin trimethylsilyl ether **4a** (103 mg, 0.5 mmol) was treated with HCl (1 N, 2 mL) in EtOAc (20 mL) at room temperature (27 °C) for 4–6 h. After which, the organic phase was added dropwise into a saturated NaHCO₃ solution (30 mL) and stirred for 5–10 min. The organic phase was separated, washed with NaHCO₃ (30 mL) and brine, dried (Na₂SO₄), and concentrated in vacuo to give the cyanohydrin **4b** (63 mg, 95%). The enantiomeric excess of **4b** was determined by comparison of optical rotation with the reported value, lit.^{6g} $[\alpha]_D^{20}$ +45 (CHCl₃, *c* 1).

General Procedure for Preparation of MTPA Esters. A solution of cyanohydrin **4b** (ca. 10 mg) in CH₂Cl₂ (1 mL) was treated with (*R*)-(+)- α -methoxy- α -(trifluoromethyl)phenylacetyl chloride (ca. 20 mg) and Et₃N (or pyridine, ca. 0.05 mL) at room temperature (27 °C) for 1 h. After which, the mixture was concentrated, and the ratio of resulting MTPA-esters was determined by ¹⁹F NMR analysis. Furthermore, the mixture was taken up with EtOAc (10 mL) and H₂O (10 mL). The aqueous phase was separated and extracted with EtOAc (10 mL \times 2). The combined organic phase was washed with brine, dried (Na₂SO₄), and concentrated. The residue was chromatographed on a silica gel column by elution with EtOAc/hexane (1:10) to give the MTPA esters of **4b**. The ratio of diastereomers was double checked by ¹H NMR and HPLC analyses. It was noticed that a significant degree of racemization might occur when the reaction was conducted with a stronger base such as 4-(dimethylamino)pyridine.

Acknowledgment. We thank the National Science Council for financial support (NSC 87-2113-M-002-042).

Supporting Information Available: Physical and spectral data of compounds **2–13b**, ORTEP drawing and crystal data of compound **3a**, and ¹H NMR spectra of **3a** and **3b** (11 pages). This material is contained in libraries on microfiche, immediately follows this article in the microfilm version of the journal, and can be ordered from the ACS; see any current masthead page for ordering information.

JO971868J