

# Correction to An On-the-Fly Surface-Hopping Program JADE for Nonadiabatic Molecular Dynamics of Polyatomic Systems: Implementation and Applications

Likai Du and Zhenggang Lan\*

*J. Chem. Theory Comput.* **2015**, *11*, 1360–1374, DOI: 10.1021/ct501106d

In eq 13 of page 1363, there is a typographical mistake:

$$\sigma_{ji}\left(t + \frac{\Delta t}{2}\right) = \frac{\langle \varphi_i^*(t) | \varphi_j(t + \Delta t) \rangle - \langle \varphi_i^*(t + \Delta t) | \varphi_j(t) \rangle}{2\Delta t} \quad (13)$$

It should instead read as follows:

$$\sigma_{ji}\left(t + \frac{\Delta t}{2}\right) = \frac{\langle \varphi_j(t) | \varphi_i(t + \Delta t) \rangle - \langle \varphi_j(t + \Delta t) | \varphi_i(t) \rangle}{2\Delta t} \quad (13)$$

Also, in the Supporting Information, eq S1

$$\sigma_{ji} = \langle \Psi_i^*(t) | \frac{\partial}{\partial t} | \Psi_j(t) \rangle \quad (S1)$$

should be

$$\sigma_{ji} = \langle \Psi_j(t) | \frac{\partial}{\partial t} | \Psi_i(t) \rangle \quad (S1)$$

and Supporting Information eq S2

$$\sigma_{ji}\left(t + \frac{\Delta t}{2}\right) = \frac{\langle \Psi_i^*(t) | \Psi_j(t + \Delta t) \rangle - \langle \Psi_i^*(t + \Delta t) | \Psi_j(t) \rangle}{2\Delta t} \quad (S2)$$

should be

$$\sigma_{ji}\left(t + \frac{\Delta t}{2}\right) = \frac{\langle \Psi_j(t) | \Psi_i(t + \Delta t) \rangle - \langle \Psi_j(t + \Delta t) | \Psi_i(t) \rangle}{2\Delta t} \quad (S2)$$

Furthermore, in Supporting Information eq S6, one should note that the superscripts  $\alpha$  and  $\beta$  are not referred to a specific spin state, and they are just used to distinguish two Slater determinants.

$$\langle \Phi^\alpha(\mathbf{r}; \mathbf{R}(t)) | \Phi^\beta(\mathbf{r}; \mathbf{R}(t + \Delta t)) \rangle = \det\{\langle \phi_k(t) | \phi'_{k'}(t + \Delta t) \rangle\} \quad (S6)$$

It should be more readable, if other superscripts were adopted; for example,

$$\langle \Phi^a(\mathbf{r}; \mathbf{R}(t)) | \Phi^b(\mathbf{r}; \mathbf{R}(t + \Delta t)) \rangle = \det\{\langle \phi_k(t) | \phi'_{k'}(t + \Delta t) \rangle\}, \quad a, b \in \alpha, \beta \quad (S6)$$

Equation 15 of page 1363 is only one approximated option to obtain the expansion coefficients for the excited state  $K$ :

$$c_{i,a}^K = \sqrt{\frac{\epsilon_a - \epsilon_i}{\omega_K}} (\mathbf{X} + \mathbf{Y})_{i,a}^K \quad (15)$$

And eq 15 should be corrected to

$$c_{i,a}^K = \sqrt{\frac{\epsilon_a - \epsilon_i}{\omega_K}} \mathbf{Z}_{i,a}^K \quad (15)$$

$\mathbf{Z}_{i,a}^K$  represents the solution of the TDDFT pseudoeigenvalue problem.<sup>1,2</sup> In eq 15,  $\epsilon_a, \epsilon_i$  are the energies of virtual and occupied molecular orbitals, respectively. And  $\omega_K$  is the corresponding excitation energy. A few variations to approximate the expansion coefficients ( $c_{i,a}^K$ ) have been reported;<sup>3–10</sup> i.e.,

$$c_{i,a}^K \approx \sqrt{\frac{1}{\epsilon_a - \epsilon_i}} (\mathbf{X} + \mathbf{Y})_{i,a}^K$$

$$c_{i,a}^K \approx (\mathbf{X} + \mathbf{Y})_{i,a}^K$$

$$c_{i,a}^K \approx (\mathbf{X})_{i,a}^K$$

$\mathbf{X}$  and  $\mathbf{Y}$  represent the solutions of the TDDFT/RPA or TDHF eigenvalue problem.<sup>1</sup> Our JADE code may select any of preceding assignments by switching via a particular keyword. Please note that some previous work also included the additional prefactor that does not change the preceding assignment due to normalization. For the case of TDDFT/TDA and CIS, only the  $\mathbf{X}$  vector is available, and one can simply take the  $\mathbf{X}$  vector. Recently, we also noted further discussions of the pseudo wavefunction description of time-dependent Hartree–Fock (TDHF) states by Subotnik and co-workers.<sup>11,12</sup> In practice, although different approximated assignment ways were made, the numerical test gave very similar and consistent results, possibly owing to the small  $\mathbf{Y}$  amplitudes. And some numerical test examples with different assignment ways can be found in the Supporting Information (SI).

## REFERENCES

- (1) Casida, M. E. Time-Dependent Density Functional Response Theory for Molecules. *Recent Advances in Density Functional Methods*; World Scientific: Singapore, 1995; Vol. 1, Chapter 5, pp 155–192, DOI: 10.1142/9789812830586\_0005.
- (2) Dreuw, A.; Head-Gordon, M. Single-Reference Ab Initio Methods for the Calculation of Excited States of Large Molecules. *Chem. Rev.* **2005**, *105*, 4009–4037.
- (3) Tapavicza, E.; Tavernelli, I.; Rothlisberger, U. *Phys. Rev. Lett.* **2007**, *98*, 023001.
- (4) Werner, U.; Mitrić, R.; Suzuki, T.; Bonačić-Koutecký, V. *Chem. Phys.* **2008**, *349*, 319–324.
- (5) Tavernelli, I.; Tapavicza, E.; Rothlisberger, U. *J. Mol. Struct.: THEOCHEM* **2009**, *914*, 22–29.

Published: July 23, 2015



- (6) Tapavicza, E.; Tavernelli, I.; Rothlisberger, U.; Filippi, C.; Casida, M. E. *J. Chem. Phys.* **2008**, *129*, 124108.
- (7) Tavernelli, I.; Tapavicza, E.; Rothlisberger, U. *J. Chem. Phys.* **2009**, *130*, 124107.
- (8) Wohlgemuth, M.; Bonacic-Koutecky, V.; Mitric, R. *J. Chem. Phys.* **2011**, *135*, 054105.
- (9) Plasser, F.; Crespo-Otero, R.; Pederzoli, M.; Pittner, J.; Lischka, H.; Barbatti, M. *J. Chem. Theory Comput.* **2014**, *10*, 1395–1405.
- (10) Zhang, X.; Li, Z.; Lu, G. *Phys. Rev. B: Condens. Matter Mater. Phys.* **2011**, *84*, 235208.
- (11) Alguire, E. C.; Ou, Q.; Subotnik, J. E. *J. Phys. Chem. B* **2015**, *119*, 7140–7149.
- (12) Ou, Q.; Alguire, E. C.; Subotnik, J. E. *J. Phys. Chem. B* **2015**, *119*, 7150–7161.