

Are Thermodynamic and Kinetic Stabilities Correlated? A Topological Index of Reactivity toward Electrophiles Used as a Criterion of Aromaticity of Polycyclic Benzenoid Hydrocarbons[†]

Arkadiusz Ciesielski,^{*,†,‡} Tadeusz M. Krygowski,[‡] Michał K. Cyrański,^{*,‡}
Michał A. Dobrowolski,[‡] and Alexandru T. Balaban[§]

Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Pawińskiego 5a, 02-106 Warsaw, Poland, Department of Chemistry, Warsaw University, Pasteura 1, 02-093 Warsaw, Poland, and Texas A&M University at Galveston, 5007 Avenue U, Galveston, Texas 77551

Received October 29, 2008

A topological index of reactivity (*TIR*) of benzenoid hydrocarbons is defined basing on an approximate value of the bicentric localization energies. *TIR* values correlate with all known (24) Hammett–Streitwieser position constants, based on kinetic data for electrophilic substitution in benzenoid hydrocarbons. The maximum value of the index, denoted by *TIR(max)*, defines the stability of a molecule toward electrophiles. For all 35 nonisoarithmetic molecules of benzenoid hydrocarbons for which Hess and Schaad data are known, *TIR(max)* values correlate with classical numerical characteristics of aromaticity: resonance energy per π -electron (*REPE*), HOMO–LUMO gap, and geometry based aromaticity index *HOMA*. Correlation between *TIR(max)* and exaltation of magnetic susceptibility is also found for cata-condensed benzenoid hydrocarbons, whereas if the peri-condensed ones are included, no correlation is observed. This can be ascribed to the presence of both paratropic and diatropic rings in perifusenes.


INTRODUCTION

Aromaticity, electronegativity, ionicity, van der Waals radii of atoms, substituent effects, and many other terms belong to the group of notions named by C. A. Coulson *Primitive Patterns of Understanding*.¹ To some extent they play in chemistry a similar role to that of axioms and postulates in mathematics. They are frequently used in the everyday research language of chemistry and related fields. The common feature of these terms is that in spite of their frequent use, most of them are not uniquely defined. To justify using these imprecise terms, let us quote E. D. Bergmann's opinion:² "Classification and theory are not ends in themselves. If they generate new experimental work, new compounds, new processes, new methods - they are good; if they are sterile - they are bad". A substantial number of papers devoted to the problems of how aromaticity is manifested and how to define this notion qualitatively or quantitatively encouraged us to undertake an attempt to present one more view on aromaticity. Most researchers in this field have agreed that classically,³ aromaticity is characterized by a collection of properties proper for cyclic π -electron systems.^{4–8} This is independent whether the cause of aromatic stability is considered as due to the π -electron or σ -electron structure.^{9,10} Earlier it was generally accepted that aromaticity was strongly associated with the planarity

of the system.⁴ In the last decades it was shown theoretically¹¹ and experimentally¹² that planarity is not a *conditio sine qua non* for aromatic properties.¹³ Benzene itself in crystalline state at 20 K is nonplanar.¹⁴ This has been attributed to intermolecular interactions in the crystal lattice. Recent definitions of aromaticity¹⁵ related this phenomenon to four characteristic features which are components of the whole definition. These are as follows: energy, geometry, magnetism, and "reactivity-based criteria", which are outlined briefly below.

"ENERGY-BASED CRITERIA" OF AROMATICITY

The earliest chemical observation for aromatic systems—first found for benzene—was associated with their resistance to chemical reactions.^{16,17} Then it was found that aromatics are thermodynamically more stable than their olefinic analogues, as documented for benzene by pioneer papers by Pauling et al.¹⁸ and Kistiakowsky et al.,¹⁹ and then extended by Wheland for a wider group of π -electron systems.²⁰ Their concept of resonance energy (*RE*) led to estimations of Aromatic Stabilization Energy (*ASE*), which were based on comparing the energy of a system having cyclic π -electron delocalization to appropriate reference π -electron systems in which no such delocalization is present. This is usually schematically formulated in a form of a reaction. An example for benzenoid hydrocarbons is presented by eq 1:²¹



Numerous papers have been published about these kinds of treatments, which differ in details but preserve the general idea of eq 1. For recent reviews see ref 22.

* Corresponding author fax: (+48) 22 8222892 (A.C.), (+48) 22 8222892 (M.K.C.); e-mail: arcad@chem.uw.edu.pl (A.C.). Corresponding author address: Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Pawińskiego 5a, 02-106 Warsaw, Poland (A.C.), Department of Chemistry, Warsaw University, Pasteura 1, 02-093 Warsaw, Poland (M.K.C.).

[†] This paper is dedicated to Prof. Zbigniew Galus (University of Warsaw) on the occasion of his 70th birthday.

[‡] Polish Academy of Sciences.

[§] Warsaw University.

[§] Texas A&M University at Galveston.

“GEOMETRY-BASED CRITERIA” OF AROMATICITY

Since the beginning of structural organic chemistry, bond lengths have been considered to be very important parameters related to chemical reactivity.¹⁷ Aromaticity was recognized early as typical for cyclic systems with bonds of intermediate length between those for single and double ones. Quantitatively, Julg et al.²³ were the first who used this view and introduced a quantitative measure of aromaticity, named sometimes as an aromaticity index, A_J . The A_J index was defined as a function of variance of CC bond lengths in the perimeter of molecules. Schematically this approach takes the form of eq 2:

$$A_J = 1 - \frac{225}{m} \sum \left(1 - \frac{R_r}{R} \right)^2 \quad (2)$$

where m is the number of peripheral bonds with individual bond lengths R_r , and R is their mean value.

Replacement of the mean value in the formula 2 for the variance of bond lengths in a ring or in the whole π -electron system by so-called “optimal bond lengths”, R_{opt} , led to the geometry-based index of aromaticity, $HOMA$,²⁴ which is calculated from geometry by use of formula 3:

$$HOMA = 1 - \frac{1}{m} \sum_{i=1}^m \alpha (R_{opt} - R_i)^2 \quad (3)$$

where m is the number of bonds taken into the summation; α is a normalization constant (for C–C, $R_{C-C} = 257.7$) fixed to give $HOMA = 0$ for a model nonaromatic system, e.g., one cyclohexatrienic Kekulé structure of benzene, and $HOMA = 1$ for the system with all bonds equal to the optimal value R_{opt} . For C–C bonds, $R_{opt,C-C} = 1.388$ Å; R_i stands for a running bond length, whereas the R_{opt} was defined as a bond for fully delocalized π -electron system. For a review on structural aspects of aromaticity see ref 25. The higher the $HOMA$ value, the more “aromatic” the ring in question and, hence, the more delocalized the π -electrons of the system.

“MAGNETISM-BASED CRITERIA” OF AROMATICITY

Measurements of diamagnetic anisotropy and exaltation of the magnetic susceptibility were used²⁶ at a very early stage for evaluating aromaticity numerically.²⁷ The exaltation needs additional assumptions (as in the case of RE) on some reference molecules. For critical reviews of these methods see refs 6 and/or 28. Soon afterward, measurements of proton NMR spectra followed, for which the term “aromatic chemical shift” was introduced and applied in many different situations.²⁹ Along with experimental approaches, in the last decades several theoretical indices of aromaticity have appeared—the best known is the *nucleus independent chemical shift* ($NICS$),³⁰ which is defined as the absolute value of shielding estimated at a given site of a cyclic π -electron system (most frequently in the center or at 1 Å above it or a perpendicular component of the tensor of $NICS(I)$). The negative values of $NICS$ characterize aromatic systems. The more aromatic system is, the more negative value of $NICS$ is expected.^{15b}

For all three criteria mentioned above, there are known numerical parameters, “aromaticity indices”, which quantify the “aromatic character” of a given molecule or its particular fragment. Even if they are not equivalent to one another,⁵

they allow us to find numerically how much they differ in “measuring” the aromaticity.

Unfortunately, the fourth criterion of aromaticity^{15a}—the **reactivity-based one**, cannot be unambiguously defined in a numerical way. If a given system follows the rule of maintaining its π -electron structure after undergoing a chemical reaction (e.g., in an “aromatic electrophilic substitution reaction”),³¹ then the criterion is fulfilled, but a more subtle differentiation is very difficult, because of the large variety of possible media, reagents, and reaction conditions used for this purpose. A topological approach for a different type of reactivity, namely a [4 + 2] cycloaddition for benzenoid hydrocarbons with maleic anhydride was suggested for the interpretation of experimental rate constants.³²

The purpose of this work is to present a topology-based method which allows us to distinguish reactivity in electrophilic substitutions between positions in molecules of benzenoid hydrocarbons, depending on how favorable is it energetically in comparison with all other possible positions. This index of reactivity is then compared with traditional energy-, magnetism-, and geometry-based indices of aromaticity.

CONCEPT AND METHODOLOGY

Consider a benzenoid hydrocarbon in which one of carbon atoms, say C_i , is a potential site of reaction, whereas all other carbons are labeled C_k . When the C_i atom is already involved in the reaction (i.e., it is excluded from the π -electron distribution), the rest of the molecule has a different π -electron structure compared to the initial state. In order to assign a canonical structure for the remaining part of the molecule another carbon C_j (alternant in respect to C_i) has to be excluded from the distribution. Then the remaining part of the molecule (i.e., the carbon atoms C_k ; $k \neq i, j$) may be described by different sets of canonical structures.³³ For each position i we have n independent sets of canonical structures in which bicentric π -electron localization occurs that for the i -th carbon atom corresponds to the number of so-called Pauling bond orders generalized to long bonds.³⁴ Consider anthracene as an illustrative example. First we divide the carbon atoms into two sets of n atoms as starred and unstarred (see Figure 1a). We localize first one electron in a starred position $i = 1$, and then we obtain 7 sets of canonical structures in which the second localization j is in unstarred positions 2, 4, 5, 7, 10, 12, and 13. Thus we have 7 sets of canonical structures for each of $n = 7$ sets of bicentric reactions involving the i -th carbon atom (see Figure 1c). The n value is always equal the half of the number N of π -electrons of the system: $n = N/2$.

Each of these sets represents a bicentric reaction. Each of these canonical structures is a realization of a virtual state in which electrons are localized in positions i and j , whereas the remainder of the molecule is described by a collection of single and double CC bonds. It is well-known that the larger the number of canonical structures, the more stable the π -electron system.¹⁷ This is represented by formula $RE \sim \ln K$, where K is the number of canonical structures.³⁵ On the other hand, K may be considered as a number of states characterized by different π -electron distributions, i.e. canonical structures. This quantity in the case of anthracene is $K = 4$ (see Figure 1b). By analogy, this notion may be compared to Boltzmann’s definition of entropy S in which




Figure 1. a) The molecule of anthracene with starred and unstarred carbon atoms. b) All unexcited canonical resonance structures of anthracene. c) Canonical resonance structures for all seven positions for $i = 1$ carbon atom for anthracene (in which the bicentric reaction occurs). K_{ij} stands for the number of canonical structures of the remaining part of the molecule. d) The TIR values for four independent positions in anthracene.

$S \sim \ln W$, where W denotes a thermodynamic probability associated with a number of ways in which a given system may be realized. Hence for a given benzenoid hydrocarbon we can define a “ π -electron delocalization entropy” in the meaning presented above: $S = \ln K$.

In our case, for each molecule of benzenoid hydrocarbon there are defined bicentric reactions in which for a fixed i -th atom there is a set of localizations labeled by j . For each set we can describe a “partial entropy” denoted as $S_{ij} = \ln K_{ij}$.

Now we define a topological index of reactivity, TIR_i of the i -th carbon atom, as a ratio of the mean partial entropy of the system (in which the bicentric reaction occurs) to the entropy of an initial state.

$$TIR_i = \frac{\bar{S}_{ij}}{S} \quad (4)$$

Hence

$$TIR_i = \frac{\frac{1}{n} \sum_{j=1}^n S_{ij}}{S} = \frac{\frac{1}{n} \sum_{j=1}^n \ln K_{ij}}{\ln K} \quad (5)$$

The summation is over n , which is the number of sets of bicentric reactions $n = N/2$, where N is total number of π -electrons

Thus

$$TIR_i = \frac{\sum_j^{N/2} \ln K_{ij}}{\frac{N}{2} \ln K} \quad (6)$$

For the archetypal aromatic compound, benzene, $TIR_i = 0$, since each bicentric reaction leads to a fully localized π -electron structure, with $K_{ij} = 1$ and hence $\ln K_{ij} = 0$. In this way the reactivity of benzene is the lowest possible among polycyclic benzenoid hydrocarbons, in line with chemical experience.




































Let us consider the example for anthracene in position $i = 1$. Application of (6) leads to

$$TIR_1 = \frac{(\ln K_{1,2} + \ln K_{1,4} + \ln K_{1,5} + \ln K_{1,7} + \ln K_{1,10} + \ln K_{1,12} + \ln K_{1,13})}{7 \cdot \ln K}$$

$$TIR_1 = \frac{(1.099 + 1.099 + 0 + 0 + 0.693 + 0 + 0)}{7 \cdot 1.386} = 0.297$$

The TIR values of four independent positions for anthracene are presented in Figure 1d. As a consequence of much higher TIR values for C_1 , C_2 , and C_9 the reaction should be most effective for position 9; therefore, position 9 is decisive about reactivity of a system in question. The TIR value for C_{11} and all equivalent positions (C_{12} , C_{13} , C_{14}) is dramatically lower than for other ones. For this reason the

Table 1. Values of $TIR(max)$, $HOMA$, $REPE$ ($\times 10^3$ [β]), Exaltation of Magnetic Susceptibility, Λ [cgs-ppm], and the Values of the HOMO–LUMO Gap $\Delta\epsilon$ in au^a



No	Structure	$TIR(max)$	$HOMA$	$REPE$	Λ	$\Delta\epsilon$
1	 benzene	0.000	0.991	65	14.5	0.247
2	 naphthalene	0.252	0.811	55	29.6	0.176
3	 anthracene	0.571	0.718	47	45.5	0.132
4	 phenanthrene	0.318	0.742	55	41.4	0.173
5	 pyrene	0.585	0.742	51	59.2	0.141
6	 naphthacene	0.638	0.670	42	62.2	0.102
7	 benz[a]anthracene	0.568	0.696	50	55.2	0.138
8	 chrysene	0.466	0.709	53	55.5	0.156
9	 triphenylene	0.338	0.691	56	49.3	0.180
10	 perylene	0.636	0.656	48	42.8	0.111
11	 benzo[e]pyrene	0.589	0.690	53	66.9	0.147
12	 benzo[a]pyrene	0.702	0.700	49	72.2	0.124
13	 pentacene	0.739	0.644	38	79.9	0.081
14	 benzo[a]naphthacene	0.631	0.660	45	70.3	0.108
15	 dibenz[a,h]anthracene	0.558	0.683	51	66.6	0.142
16	 benzo[b]chrysene	0.590	0.680	49	70.5	0.128
17	 picene	0.497	0.697	53	68.6	0.155
18	 benzo[ghi]perylene	0.629	0.707	51	79.8	0.129
19	 anthanthrene	0.712	0.691	45	89.3	0.106
20	 naphtho[2,1,8-gra]naphthacene	0.707	0.665	45	85.7	0.105
21	 benzo[a]perylene	0.761	0.639	45	62.3	0.094
22	 benzo[b]perylene	0.656	0.642	49	56.0	0.113
23	 coronene	0.557	0.742	53	123.9	0.148
24	 zethrene	0.759	0.623	41	45.5	0.082
25	 benzo[a]pentacene	0.716	0.638	42	86.7	0.086
26	 dibenzo[b,k]chrysene	0.579	0.661	46	85.9	0.115
27	 naphtho[2,3-g]chrysene	0.616	0.657	51	76.2	0.131
28	 naphtho[8,1,2-bcd]perylene	0.702	0.661	47	71.9	0.101
29	 dibenzo[cd,lm]perylene	0.720	0.690	48	105.1	0.104
30	 dibenzo[a,f]perylene	0.837	0.630	43	81.1	0.076
31	 phenanthro[1,10,9,8-opqra]perylene	0.879	0.600	42	58.2	0.067
32	 dibenzo[de,op]pentacene	0.856	0.620	38	48.4	0.066
33	 dibenzo[a,l]pentacene	0.699	0.637	44	94.5	0.092
34	 benzo[2,1-a:3,4-a']dianthracene	0.563	0.618	47	88.6	0.125
35	 naphtho[2,1,8-ya]hexacene	0.790	0.636	40	116.5	0.073

^a Dots represent carbon atoms with the maximal values of TIR_r .




Table 2. Juxtaposition of Experimental Position Constants σ_r^{+56} and TIR_r Values for 24 Positions of 14 Benzenoid Hydrocarbons^a

	σ_r^{+}	TIR_r
benzene	0.00	0.000*
1-naphthalene	0.35	0.252*
2-naphthalene	0.25	0.126
1-anthracene	0.41	0.297
2-anthracene	0.36	0.185
9-anthracene	0.72	0.571*
5-naphthalene	0.80	0.638*
3-perylene	0.74	0.636*
1-pyrene	0.67	0.585*
2-pyrene	0.22	0.202
4-pyrene	0.36	0.314
7,12-benz[a]anthracene	0.64	0.568*
coronene	0.44	0.557*
1-triphenylene	0.32	0.338*
2-triphenylene	0.26	0.327
6-anthanthrene	0.81	0.712*
1-phenanthrene	0.34	0.318*
2-phenanthrene	0.25	0.221
3-phenanthrene	0.29	0.282
4-phenanthrene	0.33	0.257
9-phenanthrene	0.365	0.308
6-chrysene	0.46	0.466*
6-benzo[a]pyrene	0.86	0.702*
7-dibenz[a,h]anthracene	0.65	0.429*

^a The r -value precedes the hydrocarbon name. An asterisk indicates $TIR(max)$.

**Figure 2.** Scatter plot of position constants σ_r^{+} and TIR values for 24 positions of 14 benzenoid hydrocarbons. Correlation coefficients $R = 0.95$ (for all points) and $R = 0.94$ (benzene not included).**Figure 3.** The scatter plot of $REPE^{51}$ on $TIR(max)$. Correlation coefficients $R = -0.88$ (for all points) and $R = -0.83$ (benzene not included).

maximal TIR value (TIR_{max}) may be used as a reactivity-based descriptor of aromaticity. Importantly the minimal value of TIR (TIR_{min}) characterizes the inner carbon atoms of anthracene (i.e., atoms not bound to hydrogen, for which

**Figure 4.** The scatter plot of the HOMO–LUMO gap in au on $TIR(max)$. Correlation coefficients $R = -0.96$ (for all points) and $R = -0.94$ (benzene not included).**Figure 5.** The scatter plot between Λ per number of π -electrons vs $TIR(max)$. Correlation coefficient $R = 0.00$.**Figure 6.** The scatter plot between Λ per number of π -electrons vs $TIR(max)$ for cata-condensed benzenoids. Correlation coefficients $R = 0.83$ (for all points) and $R = 0.70$ (benzene not included).

the electrophilic substitution does not occur). This is general for all benzenoids hydrocarbons considered (data not shown). Therefore TIR_{min} should not be taken into consideration as a descriptor of reactivity.

CALCULATIONS

The magnetic susceptibility exaltations (Λ) for systems (1)–(35), see Table 1, were evaluated by using eq 1. The magnetic susceptibility computations were made at the CSGT/B3LYP/6–311G** level of theory³⁶ by using optimized geometries at the B3LYP/6–311G** DFT level.³⁷ All molecules in eq 1 corresponded to real minima on the potential energy surface, with no imaginary vibrational frequencies. This method reproduces reasonably well the




Figure 7. The scatter plot of *HOMA* vs *TIR(max)*. Correlation coefficients $R = -0.86$ (for all points) and $R = -0.78$ (benzene not included).

experimental data of magnetic susceptibilities for benzenoid hydrocarbons. For example the calculated and experimental values are as follows: benzene: -53.0 cgs-ppm (calc.) and -54.8 cgs-ppm^{38,39} (exp.); naphthalene: -90.3 (calc.) and -89.8 cgs-ppm,^{39,40} or -91.6 cgs-ppm,^{39,41–43} or -93.6 cgs-ppm^{39,44} (exp.); anthracene: -128.4 cgs-ppm (calc.) and -123.7 ^{39,40,45} cgs-ppm (exp.); pyrene: -151.5 cgs-ppm (calc.) and -149.4 ^{39,41,46} or -154.9 ^{39,46,47} (exp.); chrysene: -160.7 (calc.) and -160.7 cgs-ppm^{39,44,47,48} (exp.); perylene: -157.4 cgs-ppm (calc.) and -159.6 cgs-ppm^{39,49} (exp.). The calculations were performed using the Gaussian 03 program.⁵⁰

The values of *HOMA*, the HOMO–LUMO gap, and exaltation of magnetic susceptibility, Λ (Table 1), were calculated from optimized structures at the same (B3LYP/6–311G**) level of theory. The values of *REPE* were taken from the paper by Hess and Schaad.⁵¹ They are given in Table 1.

RESULTS AND DISCUSSION

Reactivity is associated with a particular site in a molecule—the reaction site. Hence, in cases when there are a few possible reaction sites—positions in benzenoid hydrocarbons—it has to be decided which position should be taken as a criterion of aromaticity. The most reactive position should be decisive about the overall (kinetic) stability of a molecule. Thus, the problem is to find this kind of position in polycyclic benzenoid hydrocarbons. The classical works on electrophilic aromatic substitution in benzenoid hydrocarbons came from the mid-twentieth century,⁵² was summarized in a monograph by Streitwieser,⁵³ and then reviewed extensively several decades ago.³¹ Streitwieser defined reactivity constants for aromatic substitution (mostly the rate constants of nitration in acetic anhydride), σ_r^+ , where r is the position in the benzenoid hydrocarbon molecule.⁵⁴ These constants described well various kinds of reactivity of a given position.⁵⁴ It was also shown that physicochemical properties of the functional groups attached to position r in a hydrocarbon follow in a regular way the changes in σ_r^+ values.^{55,56}

The following question arises: do the topological indices *TIR_r* of reactivity follow any relationship with the experimental indices σ_r^+ . The data of Table 2 and Figure 2 answer this question.

The scatter plot in Figure 2 illustrates the fact that the purely topological index of reactivity for given positions,

TIR_r, correlates with the experimentally described reactivity in these positions given by σ_r^+ . Thus, at least approximately, the maximum values, *TIR(max)*, of *TIR_r* for a given position in a given molecule of a benzenoid hydrocarbon can be used as a reactivity-based index of aromaticity. Figures 3 and 4 show how this approach works in comparison with global aromaticity indices such as *REPE*⁵¹ and the HOMO–LUMO gap computed by use of Gaussian 03 at the B3LYP/6–311G** DFT level of theory,⁵⁰ which are also considered as measures of stability.⁵⁷ Figures 3 and 4 present scatter plots of *REPE* and the HOMO–LUMO gap versus *TIR(max)*. Taking into account the very approximate way of estimation the reactivity index *TIR* as well as values of *REPE* and the HOMO–LUMO gap, the observed correlation should be taken as a convincing argument that there is some equivalency between the maximal reactivity of a given benzenoid hydrocarbon and its energetic characteristic.

A completely different picture results from a correlation of the exaltation of magnetic susceptibility data divided by the number of π -electrons with the *TIR(max)* values for the whole data set of all molecules of benzenoid hydrocarbons for which data are available (35 nonisoarithmetic^{58,59} molecules listed by Hess and Schaad).⁵¹ As is apparent from Figure 5, there is no correlation at all. However, if only molecules of cata-condensed benzenoid hydrocarbons (catafusenes) are taken into account, as in Figure 6, a correlation becomes evident, qualitatively even acceptable.

The disagreement between scatter plots in Figures 5 and 6 may result from the fact that, in peri-condensed benzenoid hydrocarbons (perifusenes), the overall magnetic susceptibility is a sum of ring currents, which for some rings are clockwise and for other ones are anticlockwise. Coronene is the best illustration.⁶⁰ Moreover, some support comes also from recent studies showing that there is no direct relationship between resonance energy and magnetic response properties.⁶¹ Since *TIR(max)* is related to *REPE* (Figure 3), the lack of correlation observed in Figure 5 seems to be understandable.

Finally, the relationship between *HOMA* and *TIR(max)* shown by Figure 7 illustrates also an acceptable dependence between these indices. Again the correlation is not as good as for the previous relationships (Figures 2 and 3), but note that (when taking all points into account) $100 \times R^2 = 74.0$ means that 74.0% of the total variance of *HOMA* values is explained by the linear regression presented in Figure 7.

In conclusion it may be said that for a homogeneous set of 35 benzenoid hydrocarbons, all four criteria of aromaticity, represented by numerical indices describing energetic, magnetic (for catafusenes), geometric, and reactivity properties, of the whole molecule are in a qualitative agreement. It is important to mention that all compared quantities are intensive in nature, whereas the extensive quantities such as exaltation of magnetic susceptibilities and resonance energies are calculated per π -electron.

ACKNOWLEDGMENT

The Interdisciplinary Centre for Mathematical and Computational Modelling (Warsaw, Poland) provided computational facilities.

Supporting Information Available: Cartesian coordinates, absolute electronic energies at B3LYP/6–311G** zero

point vibrational energies (ZPE) at B3LYP/6-311G** and magnetic susceptibilities calculated at CSGT/B3LYP/6-311G** for molecules presented in Table 1. This material is available free of charge via the Internet at <http://pubs.acs.org>.

REFERENCES AND NOTES

- (1) McWeeny, R. *Coulson's Valence*; Oxford University Press: Oxford, U.K., 1979; Preface.
- (2) Concluding remarks by the following: Bergmann E. D. In *Aromaticity, Pseudoaromaticity, Antiaromaticity, Proceedings of an International Symposium held in Jerusalem 1970*; Bergmann, E. D., Pullman, B., Eds.; Israel Academy of Science and Humanities: Jerusalem, 1971.
- (3) For other, wider use of the term, see: (a) Minkin, V. I.; Glukhovtsev, M. N.; Simkin, B. Ya. *Aromaticity and Antiaromaticity*; Wiley: New York, 1994. (b) Special issue devoted to aromaticity. *Chem. Rev.* **2001**, *101*, 1115–1566.
- (4) (a) Elvidge, J. A.; Jackman, L. M. Studies of Aromaticity by Nuclear Magnetic Resonance Spectroscopy. Part I. 2-Pyridones and Related Systems. *J. Chem. Soc.* **1961**, 859–866. (b) Sondheimer, F. Recent Advances in the Chemistry of Large-Ring Conjugated Systems. *Pure Appl. Chem.* **1963**, *7*, 363–388. (c) Dewar, M. J. S. A Molecular Orbital Theory of Organic Chemistry - VIII. Aromaticity and Electrocyclic Reactions. *Tetrahedron Suppl.* **1966**, *8*, 75–92.
- (5) (a) Katritzky, A. R.; Barczyński, B.; Musumma, G.; Pisano, D.; Szafran, M. Aromaticity as a Quantitative Concept. 1. A Statistical Demonstration of the Orthogonality of Classical and Magnetic Aromaticity in Five- and Six-Membered Heterocycles. *J. Am. Chem. Soc.* **1989**, *111*, 7–15. (b) Jug, K.; Köster, A. Aromaticity as a Multi-Dimensional Phenomenon. *J. Phys. Org. Chem.* **1991**, *4*, 163–169. (c) Krygowski, T. M.; Ciesielski, A.; Bird, C. W.; Kotschy, A. Aromatic Character of the Benzene Ring Present in Various Topological Environments in Benzenoid Hydrocarbons. Nonequivalence of Indices of Aromaticity. *J. Chem. Inf. Comput. Sci.* **1995**, *35*, 203–210. (d) Katritzky, A. R.; Karelson, M.; Sild, S.; Krygowski, T. M.; Jug, K. Aromaticity as a Quantitative Concept. 7. Aromaticity Reaffirmed as a Multidimensional Characteristic. *J. Org. Chem.* **1998**, *63*, 5228–5231. (e) Cyrański, M. K.; Krygowski, T. M.; Katritzky, A. R.; Schleyer, P. v. R. To What Extent Can Aromaticity Be Defined Uniquely. *J. Org. Chem.* **2002**, *67*, 1333–1338. (f) Cyrański, M. K.; Schleyer, P. v. R.; Krygowski, T. M.; Jiao, H. J.; Hohlneicher, G. Facts and Artifacts about Aromatic Stability Estimation. *Tetrahedron* **2003**, *59*, 1657–1665. (g) Sadlej-Sosnowska, N. Mutual Relationships Between Magnetic Aromaticity Indices of Heterocyclic Compounds. *J. Phys. Org. Chem.* **2004**, *17*, 303–311.
- (6) Lazaretti, P. Ring Currents. *Prog. Nucl. Magn. Reson. Spectrosc.* **2000**, *36*, 1–88.
- (7) (a) Katritzky, A. R.; Jug, K.; Onciu, D. C. Quantitative Measures of Aromaticity for Mono-, Bi-, and Tricyclic Penta- and Hexatomic Heteroaromatic Ring Systems and Their Interrelationships. *Chem. Rev.* **2001**, *101*, 1421–1450. (b) Balaban, A. T.; Onciu, D.; Katritzky, A. R. Aromaticity as a Cornerstone of Heterocyclic Chemistry. *Chem. Rev.* **2004**, *104*, 2777–2812.
- (8) Schleyer, P. v. R.; Jiao, H. What Is Aromaticity. *Pure Appl. Chem.* **1996**, *68*, 209–218.
- (9) (a) Shaik, S. S.; Hiberty, P. C.; Ohanessian, G.; Lefour, J. M. Allylic Resonance is Forced Upon the π -System by the σ -Framework. *Nouv. J. Chim.* **1985**, *9*, 385–388. (b) Shaik, S. S.; Hiberty, P. C.; Ohanessian, G.; Lefour, J. M. Is the Delocalized π -System of Benzene a Stable Electronic System. *J. Org. Chem.* **1985**, *50*, 4657–4659. (c) Shaik, S. S.; Hiberty, P. C.; Ohanessian, G.; Lefour, J. M. When Does Electronic Delocalization Become a Driving Force of Chemical Bonding. *J. Phys. Chem.* **1988**, *92*, 5086–5094. (d) Jug, K.; Köster, A. M. Influence of σ and π Electrons on Aromaticity. *J. Am. Chem. Soc.* **1990**, *112*, 6772–6777. (e) Shaik, S. S.; Shurki, A.; Danovich, D.; Hiberty, P. C. A Different Story of π -Delocalization - The Distortivity of π -Electrons and Its Chemical Manifestations. *Chem. Rev.* **2001**, *101*, 1501–1539. (f) Maksić, Z. B.; Muller, T. Clar's Sextet Rule Is a Consequence of the σ -Electron Framework. *J. Phys. Chem. A* **2006**, *110*, 10135–10147. (g) Barić, D.; Kovacević, B.; Maksić, Z. B.; Muller, T. A Novel Approach in Analyzing Aromaticity by Homo- and Isostructural Reactions: An ab Initio Study of Fluorobenzenes. *J. Phys. Chem. A* **2005**, *109*, 10594–10606.
- (10) Jug, K.; Hiberty, P. C.; Shaik, S. σ - π Energy Separation in Modern Electronic Theory for Ground States of Conjugated Systems. *Chem. Rev.* **2001**, *101*, 1477–1500.
- (11) (a) Anet, F. A. L.; Yavari, I. Force-field Calculations for some Unsaturated Cyclic Hydrocarbons. *Tetrahedron* **1978**, *34*, 2879–2886. (b) Lipkowitz, K. B.; Peterson, M. A. Benzene Is Not Very Rigid. *J. Comput. Chem.* **1993**, *14*, 121–125.
- (12) (a) Bodwell, G. J.; Miller, D. O.; Vermeij, R. Nonplanar Aromatic Compounds. 6. [2]Paracyclo[2](2,7)pyrenophane. A Novel Strained Cyclophane and a First Step on the Road to a "Vögtle" Belt. *Org. Lett.* **2001**, *3*, 2093–2096. (b) Bickelhaupt, F. Small Cyclophanes: The Bent Benzene Business. *Pure Appl. Chem.* **1990**, *62*, 373–382. (c) Pascal, R. A., Jr. Twisted Acenes. *Chem. Rev.* **2006**, *106*, 4809–4819.
- (13) (a) Jenneskens, W.; Klammer, J. C.; de Boer, H. J. R.; de Wolf, W. H.; Bickelhaupt, F.; Stam, C. H. Molekülstruktur von 8,11-Dichlor[5]metacyclophan: Ein stark verbogener Benzolring. (Molecular structure of 8,11-dichlor[5]metacyclophane: a strongly bent benzene ring). *Angew. Chem.* **1984**, *96*, 236–237. (b) Dijkstra, F.; van Lenthe, J. H. Aromaticity of bent benzene rings: A VBSCF study. *Int. J. Quantum Chem.* **1999**, *74*, 213–221. (c) Ma, B.; Sulzbach, H. M.; Remington, R. B.; Schaefer, H. F., III. Structure, Strain Energy, and Magnetic Susceptibility of [4]Paracyclophane and the Activation Energy for Its Interconversion with 1,4-Tetramethylene Dewar Benzene. *J. Am. Chem. Soc.* **1995**, *117*, 8392–8400. (d) Bodwell, G. J.; Fleming, J. J.; Mannion, M. R.; Miller, D. O. Nonplanar Aromatic Compounds. 3. A Proposed New Strategy for the Synthesis of Buckybowls. Synthesis, Structure and Reactions of [7-], [8-], and [9](2,7)Pyrenophanes. *J. Org. Chem.* **2000**, *65*, 5360–5370. (e) Bodwell, G. J.; Bridson, J. N.; Cyrański, M. K.; Kennedy, J. W. J.; Krygowski, T. M.; Mannion, M. R.; Miller, D. O. Nonplanar Aromatic Compounds. 8. Synthesis, Crystal Structures, and Aromaticity Investigations of the 1,n-Dioxan[2](7)pyrenophanes. How Does Bending Affect the Cyclic π -Electron Delocalization of the Pyrene System. *J. Org. Chem.* **2003**, *68*, 2089–2098. (f) Dobrowolski, M. A.; Cyrański, M. K.; Merner, B. L.; Bodwell, G. J.; Wu, J. I.; Schleyer, P. v. R. Interplay of π -Electron Delocalization and Strain in [n](2,7)Pyrenophanes. *J. Org. Chem.* **2008**, *73*, 8001–8009.
- (14) Jeffrey, G. A.; Ruble, J. R.; McMullan, R. K.; Pople, J. A. The Crystal Structure of Deuterated Benzene. *Proc. R. Soc., London* **1987**, *414*, 47–57.
- (15) (a) Krygowski, T. M.; Cyrański, M. K.; Häfelfinger, G.; Katritzky, A. R. Aromaticity: a Theoretical Concept of Immense Practical Importance. *Tetrahedron* **2000**, *56*, 1783–1796. (b) Chen, Z.; Wannere, C. S.; Corminboeuf, C.; Puchta, R.; Schleyer, P. v. R. Nucleus-Independent Chemical Shifts (NICS) as an Aromaticity Criterion. *Chem. Rev.* **2005**, *105*, 3842–3888.
- (16) Ingold, C. K. *Structure and Mechanism in Organic Chemistry*; Cornell University Press: Ithaca, U.S.A., 1953; Chapter 4.
- (17) Müller, E. *Neuere Anschauungen der organischen Chemie (New views in organic chemistry)*; Springer Verlag: Berlin, Germany, 1957; Chapter 3.
- (18) (a) Pauling, L.; Wheland, G. W. The Nature of the Chemical Bond. V. The Quantum-Mechanical Calculation of the Resonance Energy of Benzene and Naphthalene and the Hydrocarbon Free Radicals. *J. Chem. Phys.* **1933**, *1*, 362–374. (b) Pauling, L.; Sherman, J. The Nature of the Chemical Bond. VI. The Calculation from Thermochemical Data of the Energy of Resonance of Molecules Among Several Electronic Structures. *J. Chem. Phys.* **1933**, *1*, 606–617.
- (19) Kistiakowsky, G. B.; Ruhoff, J. R.; Smith, H. A.; Vaughan, W. E. Heats of Organic Reactions. IV. Hydrogenation of Some Dienes and of Benzene. *J. Am. Chem. Soc.* **1936**, *58*, 146–153.
- (20) Wheland, G. W. *The Theory of Resonance and its Application to Organic Chemistry*; Wiley: New York, U.S.A., 1944.
- (21) (a) George, P.; Trachtman, M.; Bock, C. W.; Brett, A. M. Homodesmotic Reactions for the Assessment of Stabilization Energies in Benzenoid and Other Conjugated Cyclic Hydrocarbons. *J. Chem. Soc., Perkin Trans. 2* **1976**, 1222–1227. (b) George, P.; Trachtman, M.; Bock, C. W.; Brett, A. M. The use of 90°-1,3-butadiene as a reference structure for the evaluation of stabilization energies for benzene and other conjugated cyclic hydrocarbons. *Tetrahedron* **1976**, *32*, 1357–1362.
- (22) (a) Schaad, L. J.; Hess, B. A. Dewar Resonance Energy. *Chem. Rev.* **2001**, *101*, 1465–1476. (b) Slayden, W. S.; Liebman, J. F. The Energetics of Aromatic Hydrocarbons: An Experimental Thermochemical Perspective. *Chem. Rev.* **2001**, *101*, 1541–1566. (c) Cyrański, M. K. Energetic Aspects of Cyclic π -Electron Delocalization: Evaluation of the Methods of Estimating Aromatic Stabilization Energies. *Chem. Rev.* **2005**, *105*, 3773–3811.
- (23) Jug, A.; Françoise, P. Recherches sur la Géométrie de Quelques Hydrocarbures Non-alternants: Son Influence sur les Énergies de Transition, une Nouvelle Définition de L'aromaticité (Research on geometry of Some Non-alternant Hydrocarbons Its Influence on Transition Energy, a New Definition of Aromaticity). *Theor. Chim. Acta* **1967**, *8*, 249–259.
- (24) (a) Kruszewski, J.; Krygowski, T. M. Definition of aromaticity basing on the harmonic oscillator model. *Tetrahedron Lett.* **1972**, *13*, 3839–3842. (b) Krygowski, T. M. Crystallographic Studies of Inter- and Intramolecular Interactions Reflected in Aromatic Character of π -Electron Systems. *J. Chem. Inf. Comput. Sci.* **1993**, *33*, 70–78.

- (25) Krygowski, T. M.; Cyrański, M. K. Structural Aspects of Aromaticity. *Chem. Rev.* **2001**, *101*, 1385–1419.
- (26) Dauben, H. J.; Wilson, J. D.; Laity, J. L. Diamagnetic Susceptibility Exaltation as a Criterion of Aromaticity. *J. Am. Chem. Soc.* **1968**, *90*, 811–813.
- (27) Flygare, W. H. Magnetic Interactions in Molecules and an Analysis of Molecular Electronic Charge Distribution from Magnetic Parameters. *Chem. Rev.* **1974**, *74*, 653–687.
- (28) (a) Gomes, J. A. N. F.; Mallion, R. B. Aromaticity and Ring Currents. *Chem. Rev.* **2001**, *101*, 1349–1384. (b) Mallion, R. B. Topological Ring-Currents in Condensed Benzenoid Hydrocarbons. *Croat. Chem. Acta* **2008**, *81*, 227–246.
- (29) Mitchell, R. H. Measuring Aromaticity by NMR. *Chem. Rev.* **2001**, *101*, 1301–1315.
- (30) Schleyer, P. v. R.; Maerker, C.; Dransfeld, H.; Jiao, H.; van Eikemma Hommes, N. J. R. Nucleus-Independent Chemical Shifts: A Simple and Efficient Aromaticity Probe. *J. Am. Chem. Soc.* **1996**, *118*, 6317–6318.
- (31) (a) Taylor, R. *Electrophilic Aromatic Substitution*; Wiley: New York, U.S.A., 1990. (b) De la Mare, P. B. D.; Ridd, J. H. *Aromatic Substitution, Nitration and Halogenation*; Butterworths: London, U.K., 1959.
- (32) Balaban, A. T.; Biermann, D.; Schmidt, W. Chemical Graphs. 42. Dualist Graph Approach for Correlating Diels-Alder Reactivities of Polycyclic Aromatic Hydrocarbons. *Nouv. J. Chim.* **1985**, *9*, 443–449.
- (33) Our approach is based on chemical graph theory; therefore, we do not consider systems (or their fragments) in which the canonical structures involve ions. Note that the number of canonical structures K for a given benzenoid hydrocarbon corresponds to all valence states, and it can be derived in a simple way from the connectivity matrix of the molecule as the square root of the absolute value of the matrix determinant. Even though the ionic structures are included (which needs a completely different approach than presented in this paper) their contribution to stability of a given benzenoid hydrocarbon is small and should not significantly influence our findings.
- (34) Morikawa, T.; Narita, S.; Klein, D. J. Molecular Electric Conductance and Long-bond Structure Counting for Conjugated-carbon Nanostructures. *Chem. Phys. Lett.* **2005**, *402*, 554–558.
- (35) Randić, M. Aromaticity of Polycyclic Conjugated Hydrocarbons. *Chem. Rev.* **2003**, *103*, 3449–3605.
- (36) (a) Keith, T. A.; Bader, R. F. W. Calculation of Magnetic Response Properties using a Continuous Set of Gauge Transformations. *Chem. Phys. Lett.* **1992**, *194*, 1–8. (b) Keith, T. A.; Bader, R. F. W. Calculation of Magnetic Response Properties using a Continuous Set of Gauge Transformations. *Chem. Phys. Lett.* **1993**, *210*, 223–231. (c) Cheeseman, J. R.; Trucks, G. W.; Keith, T. A.; Frisch, M. J. A comparison of Models for Calculating Nuclear Magnetic Resonance Shielding Tensors. *J. Chem. Phys.* **1996**, *104*, 5497–5509.
- (37) (a) Becke, A. D. Density-Functional Thermochemistry. III. The Role of Exact Exchange. *J. Chem. Phys.* **1993**, *98*, 5648–5652. (b) Lee, C.; Yang, W.; Parr, R. G. Development of the Colle-Salvetti Correlation-energy Formula into a Functional of the Electron Density. *Phys. Rev. B* **1988**, *37*, 785–789.
- (38) Hoarau, J.; Jousot-Dubien, J.; Lemaire, B.; Lumbroso, N.; Pacault, A. *Cah. Phys.* **1956**, *74*, 34.
- (39) Gupta, R. R. *Landolt-Börnstein, Numerical Data and Functional Relationship in Science and Technology, New Series, II/16, Diamagnetic Susceptibility*; Springer: Berlin, Germany, 1986.
- (40) Krishnan, K. S.; Guha, B. C.; Banerjee, S. Investigations on Magnetic Crystalline Action. Part I: Diamagnetism. *Phil. Trans. R. Soc. London A* **1933**, *231*, 235–262.
- (41) Poquet, E. *J. Chim. Phys.* **1963**, *60*, 566.
- (42) Abrahams, R. J.; Robertson, J. M.; White, J. G. The Crystal and Molecular Structure of Naphthalene. I. X-ray Measurements. *Acta Crystallogr.* **1949**, *2*, 233–238.
- (43) Abrahams, R. J.; Robertson, J. M.; White, J. G. The Crystal and Molecular Structure of Naphthalene. II. Structure Investigation by the Triple Fourier Series Method. *Acta Crystallogr.* **1949**, *2*, 238–244.
- (44) Lonsdale, K.; Krishnan, K. S. Diamagnetic Anisotropy of Crystals in Relation to Their Molecular Structure. *Proc. R. Soc. London A* **1936**, *156*, 597–613.
- (45) Robertson, J. M. The Crystalline Structure of Anthracene. A Quantitative X-Ray Investigation. *Proc. R. Soc. London A* **1933**, *140*, 79–98.
- (46) Robertson, J. M.; White, J. G. The Crystal Structure of Pyrene. A Quantitative X-ray Investigation. *J. Chem. Soc.* **1947**, 358–368.
- (47) (a) Krishnan, K. S.; Banerjee, S. Investigations on Magnetic Crystalline Action. III. Further Studies on Organic Crystals. *Phil. Trans. R. Soc. London A* **1935**, *234*, 265–298. (b) Dhar, J.; Guha, A. Z. *Kristallogr.* **1935**, *91*, 123.
- (48) Lonsdale, K. Molecular Anisotropy of Urea, $\text{CO}(\text{NH}_2)_2$, and of Related Compounds. *Proc. R. Soc. London A* **1941**, *177*, 272–280.
- (49) (a) Banerjee, S. Z. *Kristallogr.* **1939**, *100*, 316–355. (b) Hertel, E.; Romer, G. H. Der Strukturelle Feinbau der Strukturisomeren Kohlenwasserstoffe Quaterphenyl und Triphenylbenzol. (The fine structure of the structural isomers of hydrocarbons: quaterphenyl and triphenylbenzene). *Z. Phys. Chem. B* **1933**, *23*, 226–234.
- (50) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Montgomery, J. A., Jr.; Vreven, T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; Pople, J. A. *GAUSSIAN03*; Gaussian Inc.: Wallingford, CT, 2004.
- (51) Hess, B. A.; Schaad, L. J. Hückel Molecular Orbital π Resonance Energies. The Benzenoid Hydrocarbons. *J. Am. Chem. Soc.* **1971**, *93*, 2413–2416.
- (52) Dewar, M. J. S.; Mole, T.; Urch, D. S.; Warford, E. W. T. 689. Electrophilic Substitution. Part IV. The Nitration of Diphenyl, Chrysene, Benzo[a]pyrene, and Anthanthrene. *J. Chem. Soc.* **1956**, 3572–3575.
- (53) Streitwieser, A., Jr. *Molecular Orbital Theory for Organic Chemists*; Wiley: New York, U.S.A., 1961; p 322 ff.
- (54) Table 11.1 in ref 53.
- (55) (a) Krygowski, T. M.; Stencel, M.; Galus, Z. Polarographic and Voltametric Study of Mono-nitro Derivatives of Benzenoid Hydrocarbons in DMF - Interpolation within Hammett-Streitwieser Equation and HMO-Theory. *J. Electroanal. Chem.* **1972**, *39*, 395–405. (b) Kamiński, B.; Krygowski, T. M. Application of the Hammett-Streitwieser Equation to Interpretation of the Chemical Shifts of Nonaromatic Protons in Substituted Arenes. *Tetrahedron Lett.* **1972**, 681–684. (c) Krygowski, T. M.; Kruszewski, J. Application of the Hammett-Streitwieser Equation to Predictions of Para-band Location in UV and VIS Spectra of Benzologues of Perylene and Fluoranthrene. *Bull. Acad. Polon. Sci., Ser. Sci. Chim.* **1974**, *22*, 1059–1064.
- (56) Krygowski, T. M. Towards the Unification of the Substituent (Position) Constants in Hammett-Streitwieser Equation. *Tetrahedron* **1972**, *28*, 4981–4987.
- (57) (a) Zhou, Z.; Parr, R. G.; Garst, J. F. Absolute Hardness as a Measure of Aromaticity. **1988**, *29*, 4843–4846. (b) De Proft, F.; Geerlings, P. Relative Hardness as a Measure of Aromaticity. *Phys. Chem. Chem. Phys.* **2004**, *6*, 242–248.
- (58) Isoarithmic catafusenes (cata-condensed benzenoids) have the same numbers of Kekule structures, which are in 1-to-1 correspondence; they differ only in the direction of the “kink” condensation, and in their 3-digit code this difference is reflected in permutations between digits 1 and 2 of the code. See: (a) Balaban, A. T.; Harary, F. Chemical graphs. V. Enumeration and proposed nomenclature of benzenoid cata-condensed polycyclic aromatic hydrocarbons. *Tetrahedron* **1968**, *24*, 2505–2516. (b) Balaban, A. T. Chemical graphs. VII. Proposed nomenclature of branched cata-condensed benzenoid polycyclic hydrocarbons. *Tetrahedron* **1969**, *25*, 2949–2956. (c) Balaban, A. T. Chemical graphs. Part 50. Symmetry and enumeration of fibonacenes (unbranched catacondensed benzenoids isarithmic with helixenes and zigzag catafusenes). *Math. Chem. (MATCH)* **1989**, *24*, 29–38.
- (59) Balaban, A. T.; Tomescu, I. Algebraic Expressions for the Number of Kekule Structures of Isoarithmic Cata-Condensed Benzenoid Polycyclic Hydrocarbons. *Match Commun. Math. Comput. Chem.* **1983**, *14*, 155–182.
- (60) Steiner, E.; Fowler, P. W.; Jenneskens, I. W. Counter-Rotating Ring Currents in Coronene and Corannulene. *Angew. Chem., Int. Ed.* **2001**, *40*, 362–366.
- (61) Havenith, R. W. A. Electric and Magnetic Properties Computed for Valence Bond Structures: Is There a Link between Pauling Resonance Energy and Ring Current. *J. Org. Chem.* **2006**, *71*, 3559–3563.