

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51213117>

Synthesis and Biological Evaluation of N-Fluoroalkyl and 2-Fluoroalkoxy Substituted Aporphines: Potential PET Ligands for Dopamine D(2) Receptors.

ARTICLE in ACS MEDICINAL CHEMISTRY LETTERS · MARCH 2011

Impact Factor: 3.12 · DOI: 10.1021/ml1001689 · Source: PubMed

CITATIONS

7

READS

28

6 AUTHORS, INCLUDING:



[Anna Sromek](#)

McLean Hospital

22 PUBLICATIONS 1,417 CITATIONS

SEE PROFILE



[Philip Seeman](#)

University of Toronto

496 PUBLICATIONS 32,638 CITATIONS

SEE PROFILE



[John L Neumeyer](#)

McLean Hospital

253 PUBLICATIONS 5,250 CITATIONS

SEE PROFILE

Published in final edited form as:

ACS Med Chem Lett. 2011 March 10; 2(3): 189–194. doi:10.1021/ml1001689.

Synthesis and Biological Evaluation of *N*-Fluoroalkyl and 2-Fluoroalkoxy Substituted Aporphines: Potential PET Ligands for Dopamine D₂ Receptors

Anna W. Sromek[†], Yu-Gui Si[†], Tangzhi Zhang[†], Susan R. George[‡], Philip Seeman[‡], and John L. Neumeyer^{*,†}

[†]Alcohol & Drug Abuse Research Center, McLean Hospital, Harvard Medical School, 115 Mill Street, Belmont, MA 02478-9106 USA

[‡]Department of Pharmacology, University of Toronto, Toronto, Ontario, Canada

Abstract

The 2-fluoroalkoxy substituted catechol-aporphines **6**, **8a-f** and 11-monohydroxyaporphines **11a-e** were synthesized and found to have high in vitro affinity and selectivity for the dopamine D₂ receptors. The catechol aporphines, **8b** and **8d**, and the monohydroxy aporphines, **11a-d**, were identified as candidates for development as potential PET ligands.

Keywords

aporphine; D₂ agonist; neurological disorders; positron emission tomography; dopamine receptors

Dopamine is unarguably one of the most important neurotransmitters in the brain. Disturbances in the dopaminergic system, and especially irregularities in dopamine D₂ receptor function, have been implicated in many different neurological and psychiatric disorders, including Parkinson's disease, Huntington's chorea, schizophrenia, attention deficit-hyperactivity disorder, Tourette's syndrome, restless leg syndrome, and addiction.¹ Early diagnosis of these disorders is desirable, as early treatment would allow for a better outcome for the patient, either by slowing the progression of the disease or lessening the severity of the symptoms or future episodes. Physical symptoms tend to manifest themselves much later, after significant changes occur in the brain. Thus, identification of subtle changes in the brain early in the course of the disease, before a clinical diagnosis from physical symptoms can be made, would offer the best opportunity for early treatment.

Noninvasive imaging of molecular and biological processes in living subjects with positron emission tomography (PET)² and single photon emission computed tomography (SPECT)³ are invaluable tools for the investigation of human neurochemistry and neuropharmacology in vivo.⁴ Thus, extensive research efforts have been directed toward the development of PET radioligands suitable for probing the dopaminergic system. For example, the PET ligands [¹⁸F]-DOPA and the dopamine transporter ligand, [¹¹C]-PE2I, have been used to quantify the presynaptic dopamine levels in patients suffering from Parkinson's disease.² However, these radioligands do not elucidate the postsynaptic dopamine functions in neurological disorders. In order to gain more insight into dopamine D₂ receptor function,

*To whom correspondence should be addressed. Tel: 617-855-3388; Fax: 617-855-2519; jneumeyer@mclean.harvard.edu.

Supporting Information Available: Experimental details and characterization of all compounds, biological methods. This material is available free of charge via the Internet at <http://pubs.acs.org>.

several different D₂ receptor radioligands have been developed to date. These include radioligands for striatal D₂ binding, [¹¹C]methylspiperone, [¹¹C]raclopride; and high affinity ligands for extrastriatal binding, [¹¹C]FLB457, [¹¹C]cyclopropyl-FLB457, and [¹⁸F]fallypride (see Figure 1 for structures).⁵

Although these radioligands are invaluable tools for studying or diagnosing diseases, they have certain limitations. As mentioned above, the presynaptic radioligands do not directly provide information on postsynaptic dopamine function. Of the D₂ radioligands discussed above, [¹¹C]raclopride binding is reduced when synaptic dopamine concentration is high. For others, selectivity may be an issue. For example, [¹¹C]methylspiperone also has 5HT₂ affinity, while the higher affinity ligands [¹¹C]FLB457, and [¹⁸F]fallypride do not discriminate between D₂ and D₃ sites.⁶ It has been hypothesized that, in schizophrenia and other DA-dependent neurological disorders, more D₂ receptors exist in the D₂^{high} state,^{7,8} and that D₂^{high} is the primary and common target for the antiparkinson action of dopamine agonists.⁹ However, all of the D₂ radioligands discussed above are based on benzamide D₂ antagonists, which do not discriminate between high affinity (D₂^{high}) and low affinity (D₂^{low}) states of the D₂ receptor. Therefore, it is anticipated that an agonist tracer would be more sensitive to endogenous DA concentration changes than that of an antagonist tracer, and thus, will serve as a superior probe for quantifying endogenous DA concentration.¹⁰ There have been very few attempts to develop D₂ agonist radiotracers to date. The agonist tracer [¹⁸F]F-PHNO was recently reported to have high binding affinities to D₂ and D₃ receptors *in vitro* as well as good brain penetration; however, in contrast to [³H]-(+)-PHNO (Figure 1), it did not perform well *ex vivo*.^{11a} However, [¹¹C](+)-PHNO proved to be a nonselective D₂/D₃ receptor agonist tracer with good brain uptake and favorable kinetics for PET in humans.^{11b,c} As an alternative, aporphines exhibiting D₂ receptor activity in the brain have been considered as potential agonist PET tracers. Several ¹¹C and ¹⁸F labeled apomorphine analogues have been investigated for their *in vivo* binding potency to the D₂ receptors and their distribution in brain and peripheral tissues of rats or monkey, such as *N*-[¹¹C]methylnorapomorphine,¹² *N*-*n*-3-[¹⁸F]fluoropropylnorapomorphine,^{13a,b} *N*-*n*-2-[¹⁸F]fluoroethylnorapomorphine,^{13b} *N*-[¹¹C]-propylnorapomorphine (NPA),^{13c} 2-[¹¹C]methoxy-NPA,^{13d,e} and 2-chloro-*N*-[¹¹C]-propylnorapomorphine.^{13f}

A need for even more potent dopamine ligands, with higher selectivity and oral availability, has led to the development of many novel aporphines.¹⁴ Furthermore, substituents in the 2-position of aporphines have been demonstrated to modulate the dopaminergic receptor potency and D₂/D₁ selectivity.¹⁵ In this report, several novel 2-fluoroalkoxy aporphines were synthesized and tested for dopamine receptor affinities and selectivities.

The synthesis of the twelve target molecules (**6**, **8a-f** and **11a-e**) is shown in Scheme 1. 3-Deoxynormorphine **4** was prepared from morphine according to our published procedure in 4 steps.¹⁶ *N*-Alkylation of **4** with 1-bromo-3-fluoropropane led to the *N*-substituted-3-deoxynormorphine **5**. Acid-catalyzed rearrangement of **5** with methanesulfonic acid at 90-100 °C yielded the target compound 11-hydroxy-*N*-(3-fluoropropyl)aporphine **6**. Starting from thebaine, *N*-ethyl and -propyl nororipavines **7b,c** were prepared in 4 steps using our previously reported procedure.¹⁷ Acid-catalyzed rearrangement of oripavine **7a** or nororipavines **7b,c** with methanesulfonic acid at 90-95 °C¹⁸ in the presence of 2-fluoroethanol or 3-fluoropropanol yielded the corresponding fluorinated compounds **8a-f**. *N*-*n*-Alkyl-3-*O*-[(trifluoromethyl)sulfonyl]nororipavines **9b-c** were prepared in 5 steps from thebaine according the published procedure.¹⁹ 3-*O*-[(trifluoromethyl)sulfonyl] oripavine **9a** was obtained in one step from oripavine.¹⁹ Acid-catalyzed rearrangement of **9a-c** with methanesulfonic acid at 90-100 °C in the presence of either fluoroethanol or fluoropropanol yielded **10a-e**. Pd/C catalyzed reduction of the latter with Mg metal in MeOH at rt in the presence of NH₄OAc¹⁹ provided the target compounds 2-(fluoroalkoxy)-11-hydroxy-*N*-

alkylnoraporphines **11a-e**. Alternatively, triflates **10a-e** could be reduced using a Pd-triethylhydrosilane system to furnish 11-hydroxy aporphine derivatives **11a-e**.²⁰

The receptor affinities of the twelve novel compounds **6**, **8a-f** and **11a-e** at D₁ and D₂ dopamine receptors were assessed using competitive radioreceptor binding assays with membrane-containing homogenates of rat corpus striatum tissue. Affinities to the D₃ receptor were assessed using human D₃ clones following procedures previously reported in detail^{9b} (see Supporting Information for details). However, the receptor affinities at D₃ dopamine receptors for compounds **11a-d** were assessed using rat clones following procedures reported in detail²¹ (see Supporting Information for details). The results are summarized in Table 1.

From the binding data shown in Table 1, we observed that the cold compounds **6**, **8a-f** and **11a-e** showed good to high affinity at D₂^{high} site, high selectivity of D₂ versus D₁ and, in contrast to [¹¹C]-(+)-PHNO¹¹ and [¹⁸F]fallypride⁶, exhibited low affinity or no affinity at all to the D₃ site. N-fluoropropyl aporphine **6** retained a similar binding affinity as N-propyl analog **3b** to D₂^{high} (6.9 and 4.9 nM, respectively). However, it has been previously shown that increasing the length of the N-substituent beyond three carbons causes D₂ binding affinity to drop, thereby limiting the potential for improving binding affinity and selectivity by varying the labeled N-substituent.²³ Placing a fluoroalkoxy group at position 2 would allow for more flexibility with respect to ligand design. Thus, a series of different N-*n*-propyl, ethyl, and methyl aporphines were systematically synthesized and evaluated, while varying the fluoropropanoxy and fluoroethoxy chains at position 2 (see Table 1). Likewise, the corresponding 10,11-dihydroxy (**8a-f**) and 11-hydroxy analogs (**11a-e**), all aimed at achieving the best combination of binding affinity, selectivity, and lipophilicity, were also evaluated.

We began by focusing on N-*n*-propyl catechol-aporphines, since these have been shown to have consistently higher D₂ binding affinities and selectivities over their N-ethyl and N-methyl counterparts.²³ Unfortunately, the 2-fluoropropanoxy analog **8a** exhibited a loss of D₂^{high} affinity compared to NPA (**2a**), 2-MeO-NPA (**2b**),²² and 2-F-NPA (**2c**) (27 nM vs. 5.1 to 2.7 nM range). However, in comparison, we found that by removing one carbon from the 2-substituent, 2-fluoroethoxy analog (**8b**) restored D₂^{high} affinity (3.7 nM) without compromising the remaining DA receptor binding affinity profile.

We next focused our attention on N-ethyl analogs, with the expectation that we may be able to improve D₂^{high} binding affinities. In comparison to the N-propyl catechol-aporphine analog **8a**, the N-ethyl-2-fluoropropanoxy catechol-aporphine **8c** afforded about a 4-fold increase in D₂^{high} affinity while simultaneously showing higher selectivity against D₃. The 2-fluoroethoxy analog (**8d**) afforded a more than 2-fold improvement in D₂^{high} binding affinity while retaining a similar binding affinity profile among the other dopamine receptors tested.

Encouraged by these findings, we investigated the N-methyl series. It was found that the 2-fluoropropanoxy catechol aporphine **8e** exhibited a D₂^{high} affinity consistent with **8a**, although, unlike **8a** or even **8c**, it did not exhibit any D₃^{high} affinity. The 2-fluoroethoxy analog **8f** afforded further improvement in D₂^{high} binding over the N-propyl and N-ethyl analogs **8b** and **8d**, again with no appreciable affinity to D₃.

Finally, we investigated the series of 11-monohydroxy aporphines in order to determine the effect of the absence of the 10-hydroxy group on D₂^{high} binding affinities. It was reasoned that, although 11-monohydroxyaporphines exhibited reduced D₂^{high} binding affinities compared to the 10,11-dihydroxy analogs (see **3a**, **3b**, and **6** compared to **1**, **2a**, **2b**, and **2c**, Table 1), they were also far less prone to oxidation than the catechol-aporphines, and may

thus be a viable consideration for development of imaging agents. Since catechol-aporphine **8f** exhibited the highest D_2^{high} binding affinity among the catechol-aporphine derivatives, we began by synthesizing and testing its 11-monohydroxy analog, **11e**. We expected to obtain a more stable analog, hopefully without significantly sacrificing D_2^{high} binding. We were pleased to find that the binding profile of analog **11e** exceeded our expectations, having a binding affinity to the D_2^{high} receptor in the same range as **8f**. Encouraged by this promising result, we proceeded to synthesize and test other fluorinated 11-monohydroxy aporphines, which we hypothesized might also have analogous relationships in binding affinities to their parent catechol-aporphines as the **8f** – **11e** pair. Another advantage to developing N-alkyl aporphines is that they are predicted to be more lipophilic than N-methyl derivatives, which may make them better candidates as potential PET ligands. Next, we tested N-propyl-2-fluoroethoxy-11-monohydroxy aporphine **11b**, the analog of catechol-aporphine **8b**, and found that, indeed, as in the **8f** – **11e** pair, the D_2^{high} binding affinity was about the same. We were pleased to find that the N-ethyl 11-monohydroxy analog **11d** was found to have binding affinity on the order of 1 nM and exhibiting an improved binding affinity than its 10,11-dihydroxy analog **8d**. Next we tested N-ethyl-2-fluoropropanoxy-11-monohydroxy aporphine **11c**. As expected, it also exhibited an overall improved binding affinity to D_2^{high} compared to its 10,11-dihydroxy analog **8c**. Unexpectedly, the last analog, N-propyl-2-fluoropropanoxy-11-monohydroxy aporphine **11a**, had the highest D_2^{high} binding affinity of any of the aporphines tested in this study. It was measured to have an average K_i value of 0.54 nM, which is at minimum an order of magnitude higher than its dihydroxy analog **8a**. Our goal of finding a fluorinated aporphine with high binding affinity and high selectivity to D_2^{high} was achieved. Having a series of fluorinated aporphines derivatives in hand which exhibit binding affinities in the 0-2 nM range (**8b,d,f**; **11a-e**), we can proceed with radiolabelling and *in vivo* PET studies.

Conclusion

A series of aporphines containing different N-substituents, substituents at the 2-position, 10,11-dihydroxy-, and 11-monohydroxy- groups have been systematically synthesized and tested for dopamine receptor binding affinity. Some conclusions could be made from the obtained binding data. It was found that 2-fluoroethoxy catechol-aporphines generally tended to have higher affinity to D_2^{high} than the 2-fluoropropanoxy analogs. In contrast to the generally accepted trend for nitrogen substituents in the aporphine series, smaller substituents on nitrogen (Me>Et>*n*Pr) allowed for improved D_2^{high} binding and selectivity. Contrary to our expectations, removal of the 10-hydroxy group generally *enhanced* the binding affinity to the D_2^{high} site. This is an especially attractive characteristic since the 11-hydroxy aporphines are more resistant to oxidation and are orally active²⁴ compared to their 10,11-dihydroxy counterparts. Two of the 11-hydroxy aporphines, **11a** and **11d**, were found to have average subnanomolar affinity to the D_2^{high} binding site. Unlike the series of catechol-aporphines, no clear structure-activity relationships between the substituents and binding affinities could be concluded in the 11-monohydroxy aporphine series. Finally, this new class of fluorinated aporphines could potentially prove to be valuable as therapeutics or as PET ligands. Such studies are currently in progress.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by grant from the Branfman Family Foundation (JLN) and NIH NIDA Research Training Grant T-32-DA007252 (AWS). For compounds **11a-d**, D_3 K_i determinations were generously provided by the

National Institute of Mental Health's Psychoactive Drug Screening Program, Contract # HHSN-271-2008-00025-C (NIMH PDSP). The NIMH PDSP is directed by Bryan L. Roth MD, PhD at the University of North Carolina at Chapel Hill and Project Officer Jamie Driscoll at NIMH, Bethesda MD, USA. Experimental details are described in the Supporting Information. Thebaine, morphine, and oripavine were generously donated by Mallinkrodt Inc.

References

1. (a) Bozzi Y, Borrelli E. Dopamine in neurotoxicity and neuroprotection: what do D₂ receptors have to do with it? *TRENDS in Neurosciences*. 2006; 29:167–174. [PubMed: 16443286] (b) Marsden CA. Dopamine: the rewarding years. *British J Pharm*. 2006; 147:S136–S144.
2. Ametamey SM, Horner M, Schubiger PA. Molecular imaging with PET. *Chem Rev*. 2008; 108:1501–1516. [PubMed: 18426240]
3. (a) Innis RB, Seibyl JB, Scanley BE, Laruelle M, Abi-Dargham A, Wallace E, Baldwin RM, Zea-Ponce Y, Zoghbi S, Wang S, Gao Y, Neumeyer JL, Charney DS, Hoffer PB, Marek KL. Single photon emission computed imaging demonstrates loss of dopamine transporters in Parkinson's Disease. *Proc Natl Acad Sci USA*. 1993; 90:11965–11969. [PubMed: 8265656] (b) Seibyl JP, Marek K, Sheff K, Zoghbi S, Baldwin RM, Charney DS, van Dyck CH, Innis RB. Iodine-123-beta-CIT and iodine-123-FPCIT SPECT measurement of dopamine transporters in healthy subjects and Parkinson's patients. *J Nucl Med*. 1998; 39(9):1500–1508. [PubMed: 9744331]
4. Neumeyer JL, Wang S, Gao Y, Milius RA, Kula NS, Campbell A, Baldessarini RJ, Zea-Ponce Y, Baldwin RM, Innis RB. N- ω -Fluoroalkyl Analogs of (1R)-2 β -Carbomethoxy-3 β -(4-iodophenyl)-tropine (β -CIT): Radiotracers for Positron Emission Tomography and Single Photon Emission Computed Tomography Imaging of Dopamine Transporters. *J Med Chem*. 1994; 37:1558–1561. [PubMed: 8201589]
5. Airaksinen AJ, Nag S, Finnema SJ, Mukherjee J, Chattopadhyay S, Gulyas B, Farde L, Halldin C. [¹¹C]Cyclopropyl-FLB457: A PET radioligand for low densities of dopamine D₂ receptors. *Bioorg Med Chem*. 2008; 16:6467–6473. [PubMed: 18534857]
6. Elsinga PH, Hatano K, Ishiwata K. PET tracers for imaging of the dopaminergic system. *Curr Med Chem*. 2006; 13:2139–2153. [PubMed: 16918344]
7. (a) Seeman P. All psychotic roads lead to increased dopamine D^{2high} receptors: a perspective. *Clin Schizophrenia and related psychoses*. 2008:351–355. (b) Seeman P, Weinshenker D, Quirion R, Srivastava LK, Bhardwaj SK, Grandy DK, Premont RT, Sotnikova TD, Boksa P, El-Ghundi M, O'Dowd BF, George SR, Perreault ML, Männistö PT, Robinson S, Palmiter RD, Talerico T. Dopamine supersensitivity correlates with D^{2high} states, implying many paths to psychosis. *Proc Natl Acad Sci*. 2005; 102:3513–3518. [PubMed: 15716360] (c) Seeman P. Dopamine D₂^{high} Receptors Measured Ex Vivo are Elevated in Amphetamine-Sensitized Animals. *Synapse*. 2009; 63:186–192. [PubMed: 19086090]
8. Seeman P. Dopamine D₂^{high} Receptors on Intact Cells. *Synapse*. 2008; 62:314–318. [PubMed: 18241049]
9. (a) Seeman P. Antiparkinson therapeutic potencies correlate with their affinities at dopamine D₂^{high} receptors. *Synapse*. 2007; 61:1013–1018. [PubMed: 17853435] (b) Seeman P, Ko F, Willeit M, McCormick P, Ginovart N. Antiparkinson concentrations of pramipexole and PHNO occupy dopamine D₂^{High} and D₃^{High} receptors. *Synapse*. 2005; 58:122–128. [PubMed: 16088951]
10. (a) Laruelle M. Imaging synaptic neurotransmission with in vivo binding competition techniques: a critical review. *J Cereb Blood Flow Metab*. 2000; 20:423–451. [PubMed: 10724107] (b) Ross SB, Jackson DM. Kinetic properties of the in vivo accumulation of ³H-(–)-N-n-propylnorapomorphine in mouse brain. *Naunyn-Schmeideberg's Arch Pharmacol*. 1989; 340:13–20.
11. (a) Vasdev N, Seeman P, Garcia A, Stableford WT, Nobrega JN, Houle S, Wilson AA. Syntheses and in vitro evaluation of fluorinated naphthoxazines as dopamine D₂/D₃ receptor agonists: Radiosynthesis, ex vivo biodistribution and autoradiography of [¹⁸F]F-PHNO. *Nucl Med Biol*. 2007; 34:195–203. [PubMed: 17307127] (b) Narendran R, Slifstein M, Guillain O, Hwang Y, Hwang D-R, Scher E, Reeder S, Rabiner E, Laruelle M. Dopamine (D_{2/3}) Receptor Agonist Positron Emission Tomography Radiotracer [¹¹C]-(+)-PHNO is a D₃ Receptor Preferring Agonist In Vivo. *Synapse*. 2006; 60:485–495. [PubMed: 16952157] (c) Willeit M, Ginovart N, Kapur S, Houle S, Hussey D, Seeman P, Wilson AA. High-Affinity States of Human Brain Dopamine D_{2/3} Receptors Imaged by the Agonist [¹¹C]-(+)-PHNO. *Biol Psych*. 2006; 59:389–394.

12. Zijlstra S, Vanderworp H, Wiegman T, Visser GM, Korf J, Vaalburg W. Synthesis and in vivo distribution in the rat of a dopamine agonist: *N*-([¹¹C]methyl)norapomorphine. *Nucl Med Biol.* 1993; 20:7–12. [PubMed: 8096418]
13. (a) Zijlstra S, Visser GM, Korf J, Vaalburg W. Synthesis and in vivo distribution in the rat of several fluorine-18 labeled *N*-fluoroalkylaporphines. *Appl Radiat Isot.* 1993; 44:651–658. [PubMed: 8097131] (b) Morin I, Gao Y, Kula NS, Baldessarini RJ, Neumeyer JL. *N*-monofluoroalkylnoraporphines: synthesis and binding dopamine receptor studies. *Med Chem Res.* 1992; 2:354–360. (c) Finnema SJ, Seneca N, Farde L, Shchukin E, Sovago J, Gulyas B, Wikstrom HV, Innis RB, Neumeyer JL, Halldin C. A Preliminary PET Evaluation of the New Dopamine D₂ Receptor Agonist [¹¹C]MNPA in Cynomolgus Monkey. *Nucl Med Biol.* 2005; 32:353–360. [PubMed: 15878504] (d) Hwang D-R, Narendran R, Laruelle M. Positron-labeled dopamine agonists for probing the high affinity states of dopamine subtype 2 receptors. *Bioconjugate Chem.* 2005; 16:27–31. (e) Skinbjerg M, Namkung Y, Halldin C, Innis RB, Sibley DR. Pharmacological characterization of 2-methoxy-*N*-propylnorapomorphine's interactions with D₂ and D₃ dopamine receptors. *Synapse.* 2009; 63:462–475. [PubMed: 19217026] (f) Palner M, McCormick P, Gillings N, Begtrup M, Wilson AA, Knudsen GM. Radiosynthesis and ex vivo evaluation of (*R*)-(-)-2-chloro-*N*-[1-¹¹C-propyl]*n*-propylnorapomorphine. *Nucl Med Biol.* 2010; 37:35–40. [PubMed: 20122666]
14. Zhang A, Zhang Y, Branfman AR, Baldessarini RJ, Neumeyer JL. Advances in development of dopaminergic aporphinoids. *J Med Chem.* 2007; 50:171–181. [PubMed: 17228858]
15. (a) Ramsby S, Neumeyer JL, Grigoriadis D, Seeman P. 2-Haloaporphines as potent dopamine agonists. *J Med Chem.* 1989; 32:1198–1201. [PubMed: 2524592] (b) Neumeyer JL, Gao Y, Kula NS, Baldessarini RJ. Synthesis and dopamine receptor affinity of (*R*)-(-)-2-fluoro-*N*-*n*-propylnorapomorphine: a highly potent and selective dopamine D₂ agonist. *J Med Chem.* 1990; 33:3122–3124. [PubMed: 2147956] (c) Søndergaard K, Kristensen JL, Gillings N, Begtrup M. Synthesis of (*R*)-(-)-2-fluoro norapomorphine – a precursor for the synthesis of (*R*)-(-)-2-fluoro-*N*-[¹¹C]propylnorapomorphine for evaluation as a dopamine D₂ agonist ligand for PET investigations. *Eur J Org Chem.* 2005:4428–4433. (d) Zhang A, Csutoras C, Zong R, Neumeyer JL. Synthesis of 2-fluoro-11-hydroxy-*N*-propylnorapomorphine: a potential dopamine D₂ agonist. *Org Lett.* 2005; 7:3239–3242. [PubMed: 16018630] (e) Si Y-G, Gardner MP, Tarazi FI, Baldessarini RJ, Neumeyer JL. Synthesis and binding studies 2-*O*- and 11-*O*-substituted *N*-alkylnoraporphines. *Bioorg Med Chem Lett.* 2008; 18:3971–3973. [PubMed: 18585036] (f) Si Y-G, Gardner MP, Tarazi FI, Baldessarini RJ, Neumeyer JL. Synthesis and dopamine receptor affinities of *N*-alkyl-11-hydroxy-2-methoxynoraporphines: *N*-alkyl substituents determine D₁ versus D₂ receptor selectivity. *J Med Chem.* 2008; 51:983–987. [PubMed: 18251489]
16. Csutoras C, Zhang A, Bidlack JM, Neumeyer JL. An investigation of the *N*-demethylation of 3-deoxymorphine and the affinity of the alkylation products to μ, δ, and κ receptors. *Bioorg Med Chem Lett.* 2004; 17:2687–2690.
17. Si YG, Gardner MP, Tarazi FI, Baldessarini RJ, Neumeyer JL. Synthesis and Dopamine Receptor Affinities of *N*-Alkyl-11-hydroxy-2-methoxynoraporphines: *N*-Alkyl Substituents Determine D₁ versus D₂ Receptor Selectivity. *J Med Chem.* 2008; 51:983–987. [PubMed: 18251489]
18. Sipos A, Csutoras C, Berenyi S, Uustare A, Rinken A. Synthesis and neuropharmacological characterization of 2-*O*-substituted apomorphines. *Bioorg Med Chem.* 2008; 16:4563–4568. [PubMed: 18313931]
19. Si YG, Neumeyer JL. Facile Synthesis of 2-Methoxy-11-hydroxyaporphine: A Potential D₁ Receptor Ligand. *Synthesis.* 2007; 24:3787–3790.
20. Hupp CD, Neumeyer JL. Rapid Access to Morphinones: Removal of 4,5-Ether Bridge with Pd-catalyzed triflate reduction. *Tetrahedron Lett.* 2010; 51:2359–2361. [PubMed: 20495617]
21. [Nov 12, 2010]. <http://pdsp.med.unc.edu/UNC-CH/Protocol/Book.pdf>
22. Gao Y, Baldessarini RJ, Kula NS, Neumeyer JL. Synthesis and Dopamine Receptor Affinities of Enantiomers of 2-Substituted Apomorphines and Their *N*-*n*-Propyl Analogs. *J Med Chem.* 1990; 33:1800–1805. [PubMed: 1971309]
23. Gao Y, Ram VJ, Campbell A, Kula NS, Baldessarini RJ, Neumeyer JL. Synthesis and structural requirements on *N*-substituted norapomorphines for affinity and activity at dopamine D-1, D-2, and agonist receptor sites in rat brain. *J Med Chem.* 1990; 33:39–44. [PubMed: 2136919]

24. Campbell A, Baldessarini RJ, Gao Y, Ram VJ, Neumeyer JL. *R*(-) and *S*(+) stereoisomers of 11-hydroxy-*N-n*-propylnoraporphine: Central dopaminergic behavioral activity in the rat. *Neuropharmacology*. 1990; 29:527–536. [PubMed: 1974710]





Figure 1.
High affinity D₂ ligands used in PET imaging.





**Scheme 1.**





Synthesis of aporphine analogs **6**, **8a-f**, and **11a-e**





a) ref. 16. b) 1-bromo-3-fluoropropane, NaHCO_3 , EtOH/reflux; c) MeSO_3H , 95 °C; d) ref. 17. e) ROH, MeSO_3H , 95 °C; f) NaI, acetone, reflux; g) PhNTf_2 , Et_3N , CH_2Cl_2 ; h) Pd/c, Mg, NH_4OAc , MeOH.




Table 1

Affinities (K_i) at Dopamine D_2 and D_3 receptors^b

Compound	K_i (nM) ^{a,b}				
	D_2^{low} [³ H] domperidone	D_2^{high} [³ H] domperidone	D_3 [³ H] domperidone	CLogP ^c	
 1 (APO)	98±40 ^d	1.8±0.9 ^d	2.6 ^e	2.49	
 2a (NPA)	54±20 ^d	0.18±0.03 ^d	0.44 ^g	3.55	
 2b^f (2-OMe-NPA)	810±140	5.1±1.3	1.02 ^g	3.51	
 2c (2-F-NPA)	1800±340	2.7±1.3	>10μM ^h	3.77	

Compound	K_i (nM) ^{a,b}				CLogP ^c
	D ₂ ^{low} [³ H] domperidone	D ₂ ^{high} [³ H] domperidone	D ₃ [³ H] domperidone	D ₃ [³ H] domperidone	
 3a (MCL-509)	1400±370	20±6	>10μM ^h	4.10	
 3b (11-OH-NPa)	1410±220	4.9±1.2	1700±250 ^h	4.15	
 6 (MCL-517)	860±170	6.9±2.1	>10μM	3.57	
 8a (MCL-526)	3000	28±15	430±64 ^h	3.99	

Compound	K_i (nM) ^{a,b}			D_2^{low} [³ H] domperidone	D_2^{high} [³ H] domperidone	D_3 [³ H] domperidone	CLogP ^c
	8b (MCL-524)	990±35	3.7±1.2	2200±330 ^h		3.77	
	8c (MCL-527)	1600±780	6.1±3	>10μM ^h		3.47	
	8d (MCL-528)	2400±1500	2.5±0.8	>10μM ^h		3.24	
	8e (MCL-531)	>10,000	31 ± 9	>10μM		2.94	

Compound	K_i (nM) ^{a,b}				D ₂ ^{low} [³ H] domperidone	D ₂ ^{high} [³ H] domperidone	D ₃ [³ H] domperidone	CLogP ^c
	8f (MCL-530)	620±260	2.0±0.96	>10μM				2.71
	11a (MCL-536)	490 ±280	0.54±1.6	100±14 ⁱ				4.58
	11b (MCL-522)	56 ±37	3.5±2.0	410±62 ⁱ				4.35

The image displays three chemical structures. Structure 1 is a complex polycyclic molecule featuring a benzene ring fused to a cycloheptane ring, which is further fused to a pyridine ring. It has an ethyl group on the nitrogen atom, a 3-fluoropropoxy group, and a hydroxyl group. Structures 2 and 3 are precursors to structure 1, showing the same core but with different substituents.

^aSource and radioligands: D1: rat striatum [³H]SCH23390; D2: rat striatum [³H]domperidone; D3: human D3 clone [³H]domperidone; errors are expressed as standard deviations.

^bCompounds in Table 1 have been tested and found to have low affinity to D₁ (K_i >5000) except for the following: **1** (K_i =650±310 nM), **2a** (K_i =490±220 nM), **3b** (K_i =4300±250 nM), **8f** (K_i =3700±1200 nM) and **11e** (K_i =340±44 nM); and no affinity to D₁^{high}, except for the following: **1** (K_i =4.6±1.2 nM), **2a** (K_i =1±0.2 nM), and **2b** (K_i =8.1±0.7 nM); data for **1** and **2a** obtained from ref. 9a.

^cCalculated using the chemical properties feature in Cambridgesoft ChemDraw Ultra, version 12.0. d Data from ref. 9a.^eData from ref. 9b.

^fFor preparation see ref 22.

^gSee ref. 13e: HEK293T cell homogenate used with [3H]methy/isipiperone.

^h The following compounds were also found to have D₃^{high} affinity: **2c** ($K_i = 3.8 \pm 2 \text{ nM}$), **3a** ($K_i = 130 \pm 100 \text{ nM}$), **3b** ($K_i = 1.2 \pm 1 \text{ nM}$), **8a** ($K_i = 1.1 \pm 2 \text{ nM}$), **8b** ($K_i = 1.9 \pm 1.5 \text{ nM}$), **8c** ($K_i = 230 \pm 140 \text{ nM}$), and **8d** ($K_i = 250 \pm 19 \text{ nM}$).

ⁱ Source and radioligands: D₃ rat clone [³H] domperidone; data provided by PDSP