

Enantioselective Synthesis of 2-Aryl-4-piperidones via Rhodium/Phosphoramidite-Catalyzed Conjugate Addition of Arylboroxines


Richard B. C. Jagt, Johannes G. de Vries,* Ben L. Feringa,* and Adriaan J. Minnaard*

Department of Organic and Molecular Inorganic Chemistry, Stratingh Institute, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

j.g.de.vries@rug.nl; b.l.feringa@rug.nl; a.j.minnaard@rug.nl

Received April 6, 2005

ABSTRACT



The highly enantioselective synthesis of 2-aryl-4-piperidones by rhodium/phosphoramidite-catalyzed conjugate addition of arylboroxines to 2,3-dihydro-4-pyridones is described. Both enantiomers of a variety of products with sterically and electronically different R substituents were obtained in high isolated yield and with excellent enantioselectivity up to 99%.

The piperidine ring system is a frequently encountered heterocyclic unit in natural compounds and drug candidates.¹ Piperidine alkaloids exhibit a range of biological activities and as such represent important synthetic targets. Piperidones serve an important role as intermediates *en route* to substituted piperidines² and can be found as a part of more complex biologically active compounds.³ Therefore, the development of short, enantioselective routes to substituted piperidones is a major goal.⁴ An attractive catalytic route toward enantiopure piperidones is based on the enantioselective conjugate addition to readily accessible *N*-protected 2,3-dihydro-4-pyridones,⁵ which are frequently used in alkaloid

synthesis.⁶ Until recently,^{7–9} however, no suitable procedures had been developed.

Monodentate phosphoramidite ligands comprise a cheap and easily tunable class of ligands that has already proven to be successful in a variety of reactions, including rhodium-catalyzed asymmetric hydrogenations,¹⁰ rhodium-catalyzed conjugate additions of trifluoroborates¹¹ and boronic acids,¹² and copper-catalyzed asymmetric conjugate additions of

(1) For a general review on piperidines, see: Rubiralta, M.; Giralt, E.; Diez, A. *Piperidine: Structure, Preparation and Synthetic Applications of Piperidine and its Derivatives*; Elsevier: Amsterdam, 1991.

(2) For some recent examples of the use of 4-piperidones in the synthesis of biologically active 4-piperidines, see: (a) Tawara, J. N.; Lorenz, P.; Stermitz, F. R. *J. Nat. Prod.* **1999**, *62*, 321–323. (b) Watson, P. S.; Jiang, B.; Scott, B. *Org. Lett.* **2000**, *2*, 3679–3681. (c) Brooks, C. A.; Comins, D. L. *Tetrahedron Lett.* **2000**, *41*, 3551–3553.

(3) For an example, see: Mukhtar, T. A.; Wright, G. D. *Chem. Rev.* **2005**, *105*, 529–542.

(4) (a) For a review of the recent developments in asymmetric routes to substituted piperidines, see: Bailey, P. D.; Millwood, P. A.; Smith, P. D. *Chem. Commun.* **1998**, 633–640. (b) For a review on the recent advances in the synthesis of piperidines and piperidones, see: Weintraub, P. M.; Sabol, J. S.; Kane, J. M.; Borcharding, D. R. *Tetrahedron* **2003**, *59*, 2953–2989.

(5) For the preparation of *N*-protected 2,3-dihydropyridones, see: (a) Kozikowski, A. P.; Park, P.-U. *J. Org. Chem.* **1990**, *55*, 4668–4682. (b) Comins, D. L.; Chung, G.; Foley, M. A. *Heterocycles* **1994**, *37*, 1121–1140.

(6) For a review on the asymmetric preparation and synthetic utility of 2,3-dihydro-4-pyridones, see: Comins, D. L. *J. Heterocycl. Chem.* **1999**, *36*, 1491–1500.

(7) Shintani, R.; Tokunaga, N.; Doi, H.; Hayashi, T. *J. Am. Chem. Soc.* **2004**, *126*, 6240–6241.

(8) Šebesta, R.; Pizzuti, M. G.; Boersma, A. J.; Minnaard, A. J.; Feringa, B. L. *Chem. Commun.* **2005**, 1711–1713.




Figure 1. Phosphoramidite **L**, a highly efficient ligand in the rhodium-catalyzed conjugate addition of boronic acids.

diorganozinc reagents.¹³ Recently, we have reported the synthesis of 2-alkyl-4-piperidones with high enantiomeric excess using the copper/phosphoramidite-catalyzed conjugate addition of dialkylzinc reagents to *N*-protected 2,3-dihydro-4-pyridones.⁸ It was noted that this type of substrate is less reactive toward 1,4-addition than cyclic enones, *e.g.*, 2-cyclohexenone.

Although highly enantioselective 1,4-addition of diphenylzinc, using the same catalyst, has been reported for 2-cyclohexenone,¹⁴ the lack of readily available diarylzinc reagents severely limits this method. A more convenient method for the introduction of aryl and alkenyl moieties is the asymmetric rhodium-catalyzed conjugate addition of boronic acids pioneered by Hayashi and Miyaura.¹⁵ Excellent levels of enantioselectivity have been achieved for a broad range of enones using BINAP.¹⁵ Also phosphonites¹⁶ and amidophosphines¹⁷ were successfully applied as chiral ligands. It was shown by our group that phosphoramidites (*i.e.*, **L**, Figure 1) are exceptionally efficient ligands for this reaction in terms of reaction rate, chemoselectivity, and enantioselectivity.¹²

(9) For related 1,4-additions to 2-piperidones, see: Pineschi, M.; Del Moro, F.; Gini, F.; Minnaard, A. J.; Feringa, B. L. *Chem. Commun.* **2004**, 1244–1245.

(10) (a) Van den Berg, M.; Minnaard, A. J.; Schudde, E. P.; Van Esch, J.; de Vries, A. H. M.; de Vries, J. G.; Feringa, B. L. *J. Am. Chem. Soc.* **2000**, *122*, 11539–11540. (b) Peña, D.; Minnaard, A. J.; de Vries, J. G.; Feringa, B. L. *J. Am. Chem. Soc.* **2002**, *124*, 14552–12553. (c) Van den Berg, M.; Minnaard, A. J.; Haak, R. M.; Leeman, M.; Schudde, E. P.; Meetsma, A.; Feringa, B. L.; de Vries, A. H. M.; Maljaars, C. E. P.; Willans, C. E.; Hyett, D.; Boogers, J. A. F.; Henderickx, H. J. W.; de Vries, J. G. *Adv. Synth. Catal.* **2003**, *345*, 308–323.

(11) (a) Duursma, A.; Boiteau, J.-G.; Lefort, L.; Boogers, J. A. F.; de Vries, A. H. M.; de Vries, J. G.; Minnaard, A. J.; Feringa, B. L. *J. Org. Chem.* **2004**, *69*, 8045–8052. (b) Duursma, A.; Lefort, L.; Boogers, J. A. F.; de Vries, A. H. M.; de Vries, J. G.; Minnaard, A. J.; Feringa, B. L. *Org. Biomol. Chem.* **2004**, *2*, 1682–1684.

(12) (a) Boiteau, J.-G.; Imbos, R.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2003**, *5*, 681–684. See also: Boiteau, J.-G.; Imbos, R.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2003**, *5*, 1385. (b) Boiteau, J.-G.; Minnaard, A. J.; Feringa, B. L. *J. Org. Chem.* **2003**, *68*, 9481–9484. (c) Duursma, A.; Hoen, R.; Schuppan, J.; Hulst, R.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2003**, *5*, 3111–3113.

(13) (a) Feringa, B. L. *Acc. Chem. Res.* **2000**, *33*, 346–353. (b) Naasz, R.; Arnold, L. A.; Minnaard, A. J.; Feringa, B. L. *Angew. Chem., Int. Ed.* **2001**, *40*, 927–930. (c) Arnold, L. A.; Naasz, R.; Minnaard, A. J.; Feringa, B. L. *J. Am. Chem. Soc.* **2003**, *125*, 3700–3701.


(14) Peña, D.; Lopez, F.; Harutyunyan, A. R.; Minnaard, A. J.; Feringa, B. L. *Chem. Commun.* **2004**, 1836–1837.

(15) (a) Sakai, M.; Hayashi, H.; Miyaura, N. *Organometallics* **1997**, *16*, 4229–4231. (b) Takaya, Y.; Ogasawara, M.; Hayashi, T.; Sakai, M.; Miyaura, M. *J. Am. Chem. Soc.* **1998**, *120*, 5579–5581. (c) For a recent review, see: Hayashi, T.; Yamasaki, K. *Chem. Rev.* **2003**, *103*, 2829–2844.

(16) Reetz, M. T.; Moulin, D.; Gosberg, A. *Org. Lett.* **2001**, *3*, 4083–4085.

(17) Kuriyama, M.; Nagai, K.; Yamada, K.; Miwa, Y.; Taga, T.; Tomioka, K. *J. Am. Chem. Soc.* **2002**, *124*, 8932–8939.

Scheme 1



We envisioned that introduction of aryl groups to **1** (Scheme 1), using the rhodium/phosphoramidite-catalyzed conjugate addition of arylboronic acids, could provide a pathway to 2-substituted 4-piperidones that is complementary to our work with dialkylzinc reagents. During our studies, Hayashi *et al.* reported the enantioselective addition of arylzinc chlorides to 2,3-dihydro-4-pyridones.⁷ In that study, it was noted also that this type of substrate is less reactive toward 1,4-addition compared to other enones. The rhodium/BINAP-catalyzed conjugate addition of phenylboronic acid failed to proceed to full conversion, although the enantioselectivity was excellent.

Initial screening of our catalyst system on substrate **1** was performed under standard conditions in a mixture of dioxane/water (10/1) at 100 °C with a catalyst generated from 3 mol % Rh(acac)(C₂H₄)₂ and 7.5 mol % **L**. As in the report of Hayashi, with 3 equiv of phenylboronic acid, the reaction did not go to completion according to ¹H NMR (entry 1, Table 1). The enantioselectivity was, however, excellent

Table 1. Optimization of the Reaction Conditions for the Rhodium-Catalyzed Conjugate Addition to **1**

entry	"PhB" (equiv)	conditions ^a	conversion % ^b	ee (%) ^c
1	PhB(OH) ₂ (3.0)	A	80	96
2	(PhBO) ₃ (1.0)	B	60	99
3	(PhBO) ₃ (3.0)	B	75	99
4	(PhBO) ₃ (1.0)	C	84	99
5	(PhBO) ₃ (2.0)	C	92	99
6	(PhBO) ₃ (3.0)	C	100	99

^a All reactions were performed on a 0.2 mmol scale with 3 mol % Rh(acac)(C₂H₄)₂ and 7.5 mol % (*R*)-**L** at 100 °C for 2 h. Conditions A: 0.55 mL of dioxane/H₂O (10/1). Conditions B: 0.5 mL of dioxane, 1 equiv of H₂O with respect to boron. Conditions C: 0.5 mL of dioxane, slow addition of water by syringe pump, 100 °C, 1 h. ^b Determined by ¹H NMR. ^c Determined by chiral HPLC.

(96% ee). We then decided to generate phenylboronic acid *in situ* from phenylboroxine ((PhBO)₃)¹⁸ and water (one

(18) Arylboroxines are obtained by dehydration of arylboronic acids by azeotropic removal of water from their xylene solution or heating at 300 °C *in vacuo*. For a review, see: Lappert, M. F. *Chem. Rev.* **1956**, *56*, 959–1064.

equivalent with respect to boron), providing mild reaction conditions (entries 2–5).¹⁹ The use of this reagent did not improve the conversion but did improve the enantioselectivity to an excellent ee of 99%. Upon slow addition of water, thereby preventing premature hydrolysis of the boroxine, the reaction could be driven to 84% conversion using 1 equiv of boroxine (entry 4) and to full conversion with retention of 99% ee using 3 equiv of the reagent (entry 6).

To show the applicability of this reaction for synthesis on a laboratory scale, it was performed on a 0.5 g (2.2 mmol) scale. After flash chromatography, the product was isolated in 86% yield with 99% ee.

With these optimized conditions in hand, the scope of the asymmetric conjugate addition of arylboroxines to **1** was investigated. High ee values could be obtained with a variety of sterically and electronically diverse arylboroxines (entries 1–8, Table 2). *meta*- and *para*-tolyl groups can be introduced with high enantioselectivity and high yield (entries 3 and 4). A dramatic drop in enantioselectivity was observed for the more sterically demanding *ortho*-tolyl group (entry 2), illustrating a possible limitation of the catalytic method. Products with one or two electron-donating substituents on the aryl were obtained in high yield with high enantioselectivity (entries 5 and 6). However, electron-withdrawing groups such as chloride or fluoride slow the reaction, leading to incomplete conversions (entries 7 and 8). Despite this observation, the enantioselectivity is largely independent of the electronic properties of the substituents, and all *para*- or *meta*-substituted products are obtained with excellent ee values between 94 and 98%.

In summary, we have shown that conjugate addition of arylboroxines with a rhodium/phosphoramidite catalyst can be used to prepare 2-aryl-4-piperidones in high isolated yield (82–92%) and with excellent enantioselectivity (up to 99%). We are currently directing our efforts toward enhancing the

(19) Use of arylboroxines was previously reported to have a beneficial effect on both conversion and enantioselectivity in the conjugate addition to highly deactivated 1-alkenylphosphonates; see: Hayashi, T.; Senda, T.; Takaya, Y.; Ogasawara, M. *J. Am. Chem. Soc.* **1999**, *121*, 11591–11592.

Table 2. Scope of Arylboroxines in the Rhodium-Catalyzed Asymmetric 1,4-Addition to **1**^a

entry	Ar	product	yield (%) ^b	ee (%) ^c
1	Ph	2a	86 ^d	99
2	<i>o</i> -MeC ₆ H ₄	2b	82 ^d	24
3	<i>m</i> -MeC ₆ H ₄	2c	92 ^d	98
4	<i>p</i> -MeC ₆ H ₄	2d	86 ^d	95
5	<i>p</i> -MeOC ₆ H ₄	2e	85 ^d	96
6	<i>m,p</i> -(MeO) ₂ C ₆ H ₃	2f	86 ^d	98
7	<i>p</i> -FC ₆ H ₄	2g	71	94
8	<i>p</i> -ClC ₆ H ₄	2h	55	96

^a All reactions were performed in duplicate with both enantiomers of the ligand on a 0.2 mmol scale with 3 mol % Rh(acac)(C₂H₄)₂ and 7.5 mol % **L** at 100 °C for 2 h. (*R*)-**L** gave the (*R*)-enantiomer of the product in all cases; see Supporting Information. ^b Isolated yield. ^c Determined by chiral HPLC. ^d Thin-layer chromatography shows a spot to spot conversion in 2 h.

scope and applications of this method in the synthesis of more complex heterocycles.

Acknowledgment. We thank Mrs. T. D. Tiemersma-Wegman for the technical support. Financial support from DSM, the Ministry of Economic Affairs, and NWO/CW, administered through the CW/Combichem program, is gratefully acknowledged.

Supporting Information Available: Experimental details and chromatographic and spectroscopic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL050734M