

Möbius Aromaticity in [12]Annulene: Cis–Trans Isomerization via Twist-Coupled Bond Shifting

Claire Castro,* William L. Karney,* Miguel A. Valencia, Catherine M. H. Vu, and Ryan P. Pemberton

Department of Chemistry, University of San Francisco, 2130 Fulton Street, San Francisco, California 94117

Received April 14, 2005; E-mail: castroc@usfca.edu; karney@usfca.edu

The study of annulenes and the dynamic processes that they undergo has led to many fascinating discoveries.¹ Bond shifting in cyclobutadiene is now believed to proceed via heavy-atom tunneling.² Ring inversion and bond shifting in cyclooctatetraene (COT) have been extensively investigated;³ the latter process occurs via a singlet, diradical transition state that violates Hund's rule.^{3b} Comparisons of theory and experiment for [10]annulene provided critical insights regarding the validity of different methods for computation of annulenes.^{4,5} Recent work by Wannere et al. revealed that [14]- and [18]annulene are *not* bond-equalized; rather, both exhibit pronounced C–C bond-length alternation, with the implication being that bond shifting *is* relevant to these $[4n + 2]$ -annulenes.⁵ In all of these cases ([4]-, [8]-, [10]-, [14]-, and [18]-annulene), bond shifting is believed to occur via a planar or nearly planar structure.

In contrast with the planar annulenes mentioned above, there is a resurgence of interest in potential aromaticity in Möbius $[4n]$ annulenes,^{6–10} exemplified by the synthesis of a Möbius [16]-annulene derivative,⁸ though later work argues it is nonaromatic.⁹ The preparation of a stable, neutral, clearly aromatic Möbius annulene remains elusive. Here, we present the first evidence that neutral, highly aromatic, Möbius annulenes play a critical role in the known chemistry of $[4n]$ annulenes. We show, using density functional and coupled cluster calculations,^{11–13} that bond shifting in [12]annulene can take place via such species, and that this provides a mechanism for the thermal cis–trans isomerization proposed by Oth and Schröder over 30 years ago.¹⁸ The work described herein unites three fundamental concepts: thermal cis–trans isomerization, bond shifting in annulenes, and Möbius aromaticity.

After characterizing tri-*trans*-[12]annulene **1a** by low-temperature NMR, Oth and Schröder found that upon warming to -40°C , the bicyclic product **3** was formed, presumably via electrocyclization of the transient di-*trans* isomer **2** (Scheme 1).^{18a} Because **2** was never observed, the determined barrier of 17.4 ± 0.1 kcal/mol effectively pertains to the isomerization **1a** \rightarrow **2**. For comparison, similar configuration change in [16]annulene was found to have a barrier of ca. 10 kcal/mol.^{18b} Both of these barriers are much lower than those for thermal cis–trans isomerization (via diradicals) in acyclic systems of similar size ($E_a > 28$ kcal/mol),¹⁹ suggesting that cis–trans isomerization in annulenes occurs via a different mechanism than in acyclic systems. We find that this is indeed the case for [12]annulene.

Given an appropriate starting conformation of **1**, bond shifting can effectively result in cis–trans isomerization to **2**.²⁰ Assuming that bond shifting proceeds via a bond-equalized transition state, the solution to finding the appropriate conformation lies in Heilbronner's idea that for $[4n]$ annulenes, distortions from planar to Möbius topology¹⁰ could occur without loss of π -electron energy, assuming that angle strain is not prohibitive.^{6a} On the basis of our prior computational work on conformational automerization in **1**,²¹





Figure 1. BH&HLYP/6-311+G** optimized geometries of three Möbius conformations of tri-*trans*-[12]annulene **1** (left), transition states for bond shifting from each (middle), and bond-shifting products (right). Selected C–C distances (Å) and CCCC dihedral angles ($^\circ$) are shown.

Scheme 1



the lowest-energy conformer of **1** (**1a**, Scheme 1, Figure S1)²² can undergo slight bond rotation with minimal energy cost (< 1 kcal/mol) to form the Möbius, *nonaromatic*, conformation **1b** (Figure 1).

From this latter conformation, twisting of the C4–C5, C5–C6, and C6–C7 bonds, coupled with bond-length equalization of all C–C bonds, leads to a highly delocalized transition state for bond shifting (**TS1b**, Figure 1). We refer to this process as “twist-coupled bond shifting”. IRC calculations indicate that **TS1b** connects **1b** to di-*trans* isomer **2a**, which can readily undergo conformational change to **2b** (Figure S2), followed by thermal electrocyclization to **3** ($E_a = 13.8$ kcal/mol).²² Figure 2 summarizes the energetics of the entire process at the CCSD(T)/cc-pVDZ//BH&HLYP/6-311+G** level. The computed overall barrier of 18.0 kcal/mol for the **1a** \rightarrow **3** process (bond shifting being the rate-determining step) agrees well with the experimental barrier of ca. 17.4 kcal/mol.^{18a}




Figure 2. CCSD(T)/cc-pVDZ//BH&HLYP/6-311+G** potential energy curve for the conversion of tri-*trans*-[12]annulene **1a** to di-*trans* isomer **2a** and ultimately to *cis*-bicyclo[6.4.0]dodeca-2,4,6,9,11-pentaene (**3**).

Table 1. Relative Energies (kcal/mol) and Aromaticity Indicators for [12]Annulene Stationary Points and Related Species^a

species	sym	config	BHH rel E	CC rel E	Δr	NICS	Λ
1a	C ₁	CTCTCT	0.0	0.0	0.147	1.3	10.6
1b	C ₁	CTCTCT	0.8	0.7	0.156	-1.6	0.4
TS1b	C ₁		20.2	18.0	0.031	-13.9	-43.8
2a	C ₁	CCCTCT	5.4	5.7	0.150		
TS2a2b	C ₁	CCCTCT	13.9	13.7			
2b	C _s	CCCTCT	2.8	4.0			
TS2b3	C _s		19.3	17.8			
3	C ₁		-15.2	-16.5			
1c	C ₂	CTCTCT	4.7	4.5	0.157	-0.3	6.0
TS1c	C ₂		27.8	25.5	0.035	-13.4	-43.8
4a	C ₂	CCTCCT	0.5	0.9	0.152		
1d	C ₂	CTCTCT	8.3	7.9	0.154	0.5	5.9
TS1d	C ₂		26.8	25.8	0.027	-15.3	-46.7
4b	C ₂	CCTCCT	5.7	6.3	0.135		

^a Config = *cis-trans* configuration. BHH = BH&HLYP/6-311+G**. CC = CCSD(T)/cc-pVDZ//BH&HLYP/6-311+G**. Relative energies (kcal/mol) are corrected for differences in ZPE. Δr = difference, in Å, between the longest and shortest C–C bonds in the annulene ring. NICS and Λ computed at the B3LYP/6-311+G**/BH&HLYP/6-311+G** level. NICS = nucleus-independent chemical shift, in ppm, computed at the ring center using the GIAO method. Λ = magnetic susceptibility exaltation (MSE, in cgs-ppm), computed using the CSGT method and the increment method. Details can be found in Supporting Information.

TS1b is a clear example of a Möbius aromatic species. Moreover, it is the first example of a Möbius aromatic transition state that does not involve breaking or forming σ bonds.^{6b,23} As expected for a bond-shifting transition state, **TS1b** exhibits a small Δr value of 0.031 Å. Its large negative NICS value²⁴ (–13.9 ppm) and magnetic susceptibility exaltation²⁵ (Λ) (–43.8 cgs-ppm) reveal that **TS1b** is highly aromatic (Table 1). Thus, the transformation of **1** \rightarrow **2** via **TS1b** suggests that Oth and Schröder prepared a Möbius aromatic [4*n*]annulene, albeit in the form of a transition state, over 30 years ago.²⁶

Two other conformations of **1** (**1c** and **1d**, Figure 1)²⁷ can undergo twist-coupled bond shifting via Möbius aromatic transition states, though these barriers are higher than that for **1b** (Table 1). Analogous to **TS1b**, the transition states **TS1c** and **TS1d** (Figure 1) exhibit strong aromatic character, based on the large negative NICS and Λ values (Table 1). Bond shifting via **TS1c** or **TS1d** leads to two different conformations of di-*trans*-[12]annulene **4** (**4a** and **4b**, respectively, Figure 1), an isomer previously studied.^{7c,28,29}

Whereas planar bond shifting interconverts degenerate species, twist-coupled bond shifting produces configuration change. This new mechanism (i) suggests that neutral, highly aromatic, Möbius geometries serve as the transition states for *cis-trans* isomerization

in the case of [12]annulene, and (ii) probably explains the numerous known examples of facile *cis-trans* isomerization in [4*n*]annulenes ($n \geq 3$).^{1,18b}

Acknowledgment. Dedicated to the memory of Orville Chapman. This work was supported in part by a grant from the University of San Francisco Faculty Development Fund.

Supporting Information Available: Computational details, Figures S1–S3, absolute energies, thermal corrections, and Cartesian coordinates for all optimized structures, and complete ref 11. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Reviews: (a) Sondheimer, F. *Acc. Chem. Res.* **1972**, *5*, 81. (b) Kennedy, R. D.; Lloyd, D.; McNab, H. J. *Chem. Soc., Perkin Trans. 1* **2002**, 1601.
- (2) Carpenter, B. K. *J. Am. Chem. Soc.* **1983**, *105*, 1700.
- (3) (a) Paquette, L. A. *Acc. Chem. Res.* **1993**, *26*, 57. (b) Wenthold, P. G.; Hrovat, D. A.; Borden, W. T.; Lineberger, W. C. *Science* **1996**, *272*, 1456.
- (4) (a) King, R. A.; Crawford, T. D.; Stanton, J. F.; Schaefer, H. F. *J. Am. Chem. Soc.* **1999**, *121*, 10788. (b) Price, D. R.; Stanton, J. F. *Org. Lett.* **2002**, *4*, 2809.
- (5) Wannere, C. S.; Sattelmeyer, K. W.; Schaefer, H. F.; Schleyer, P. v. R. *Angew. Chem., Int. Ed.* **2004**, *43*, 4200.
- (6) (a) Heilbronner, E. *Tetrahedron Lett.* **1964**, 1923. (b) See also: Zimmerman, H. E. *Acc. Chem. Res.* **1971**, *4*, 272 and references therein.
- (7) (a) Mauksch, M.; Gogonea, V.; Jiao, H.; Schleyer, P. v. R. *Angew. Chem., Int. Ed.* **1998**, *37*, 2395. (b) Martín-Santamaría, S.; Lavan, B.; Rzepa, H. S. *J. Chem. Soc., Perkin Trans. 2* **2000**, 1415. (c) Castro, C.; Isborn, C. M.; Karney, W. L.; Mauksch, M.; Schleyer, P. v. R. *Org. Lett.* **2002**, *4*, 3431.
- (8) Ajami, D.; Oeckler, O.; Simon, A.; Herges, R. *Nature* **2003**, *426*, 819.
- (9) Castro, C.; Chen, Z.; Wannere, C. S.; Jiao, H.; Karney, W. L.; Mauksch, M.; Puchta, R.; Hommes, N. J. R. v. E.; Schleyer, P. v. R. *J. Am. Chem. Soc.* **2005**, *127*, 2425.
- (10) In tracing the π system along one face of the ring, a return to the starting point on the same face indicates Hückel topology, whereas a return on the opposite face indicates Möbius topology.
- (11) All calculations were performed with Gaussian 98: Frisch, M. J. et al. *Gaussian 98*, revision A.11.3; Gaussian, Inc.: Pittsburgh, PA, 2002.
- (12) Geometry optimizations and vibrational analyses were performed at the BH&HLYP/6-311+G** level.^{14,15} BH&HLYP performs much better than B3LYP in computing the relative energy ordering of bond-equalized versus bond-alternating isomers of [10]-, [14]-, [18]-, and [22]annulene.⁵
- (13) Single-point energies were computed at the CCSD(T)/cc-pVDZ//BH&HLYP/6-311+G** level^{16,17} and were corrected for differences in unscaled zero-point energies. The success of the CCSD(T) method for the problematic case of [10]annulene is described in ref 4.
- (14) (a) Becke, A. D. *J. Chem. Phys.* **1992**, *98*, 1372. (b) Miehlisch, B.; Savin, A.; Stoll, H.; Preuss, H. *Chem. Phys. Lett.* **1989**, *157*, 200.
- (15) Hariharan P. C.; Pople, J. A. *Theor. Chim. Acta* **1973**, *28*, 213.
- (16) (a) Bartlett, R. J. *J. Phys. Chem.* **1989**, *93*, 1697. (b) Scuseria, G. E. *Chem. Phys. Lett.* **1991**, *176*, 27.
- (17) Dunning, T. H., Jr. *J. Chem. Phys.* **1989**, *90*, 1007.
- (18) (a) Oth, J. F. M.; Röttele, H.; Schröder, G. *Tetrahedron Lett.* **1970**, 61. (b) Oth, J. F. M. *Pure Appl. Chem.* **1971**, *25*, 573.
- (19) Doering, W. v. E.; Kitagawa, T. *J. Am. Chem. Soc.* **1991**, *113*, 4288.
- (20) Oth and Schröder found no NMR evidence for degenerate bond shifting.¹⁸
- (21) Castro, C.; Karney, W. L.; Vu, C. M. H.; Burkhardt, S. E.; Valencia, M. A. *J. Org. Chem.* **2005**, *70*, 3602.
- (22) Details of all stationary points can be found in the Supporting Information.
- (23) Jiao, H.; Schleyer, P. v. R. *J. Chem. Soc., Perkin Trans. 2* **1994**, 407.
- (24) Schleyer, P. v. R.; Maerker, C.; Dransfeld, A.; Jiao, H.; Hommes, N. J. R. v. E. *J. Am. Chem. Soc.* **1996**, *118*, 6317.
- (25) Dauben, H. J.; Wilson, J. D.; Laity, J. L. *J. Am. Chem. Soc.* **1969**, *91*, 1991.
- (26) Despite extensive searching, we failed to locate a diradical mechanism for *cis-trans* isomerization of **1** to **2**.
- (27) These conformations have been described in ref 21. Despite appearances, species **1c** and **1d** are neither equivalent nor enantiomeric. H1 and H10 are syn to each other in **1c** and **TS1c**, whereas they are anti in **1d** and **TS1d**.
- (28) Gard, M. N.; Reiter, R. C.; Stevenson, C. D. *Org. Lett.* **2004**, *6*, 393.
- (29) We have also found a Möbius bond-shifting transition state connecting **4a** to a mono-*trans* isomer via a 12.7 kcal/mol barrier (Figure S3, Table S2).

JA052447J