

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/244438874>

Elimination of Chlorine (Radical) or Tosylate (Anion) from C2' of Nucleoside C3' Free Radicals as Model Reactions Postulated To Occur at the Active Site of Ribonucleotide Reductase...

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · APRIL 1997

Impact Factor: 12.11 · DOI: 10.1021/ja970171c

CITATIONS

23

READS

19

3 AUTHORS, INCLUDING:



Stanislaw F Wnuk

Florida International University

162 PUBLICATIONS 2,096 CITATIONS

SEE PROFILE

Elimination of Chlorine (Radical) or Tosylate (Anion) from C2' of Nucleoside C3' Free Radicals as Model Reactions Postulated To Occur at the Active Site of Ribonucleotide Reductases¹

Morris J. Robins,* Zhiqiang Guo, and Stanislaw F. Wnuk

Department of Chemistry and Biochemistry
Brigham Young University
Provo, Utah 84602-5700


Received January 21, 1997

Ribonucleotide reductases (RNRs) catalyze the conversion of ribonucleoside 5'-di- or -triphosphates to 2'-deoxynucleotides that are required for DNA biosynthesis.² The ribonucleoside diphosphate reductase (RDPR) from *Escherichia coli* (EC 1.17.4.1) has two nonidentical subunits (R1 and R2) whose structures have been determined by X-ray crystallography.³ The R1 subunit contains allosteric control sites and five cysteine residues that participate in catalytic turnover and/or as redox dithiol/disulfide pairs. The R2 subunit contains a diiron chelate and a tyrosine-centered free radical that is responsible for generation of a proximate thiyl radical⁴ on R1 via coupled electron and proton transfer reactions. The thiyl radical has been proposed to initiate nucleotide reduction by abstraction of H3' from the substrate ribonucleotide.^{2c} Water (O2') is then lost from C2' of the resulting C3' radical.^{2a,5} Interaction of a carboxylate group (glutamate) with OH3' has recently been invoked to assist with the heterolytic release of water.^{2c}

Abstraction of H3' from 2'-chloro-2'-deoxynucleoside 5'-diphosphates to generate C3' radicals was proposed^{2a,c} to initiate reactions leading to inactivation of RDPR.⁶ Spontaneous loss of chloride and transfer of the OH3' proton to glutamate would give 2'-deoxy-3'-ketonucleotide intermediates without involvement of a cysteine pair on R1.^{2a,c} Successive β -eliminations (H2'/base and H4'/pyrophosphate) would give the Michael acceptor 2-methylene-3(2H)-furanone, which could effect covalent inactivation of the enzyme.⁷

We recently demonstrated a mechanistic alternative for potential generation of the Michael acceptor that involved loss of a radical, rather than an anionic, species from C2' of model 2'-substituted nucleosides.^{8,9} Thus, treatment of 2'-(azido, bromo, chloro, iodo, or methylthio)nucleoside 3'-thionocarbonates with tributylstannane/AIBN resulted in loss of the 2'-substituents, as presumed radicals, to give 2',3'-didehydro-2',3'-dideoxy derivatives upon generation of C3' radicals (without O3'); whereas 3'-thionocarbonates with 2'-fluoro or 2'-O-(mesyl or tosyl) substituents underwent radical-mediated hydrogen transfer to C3' to give the 3'-deoxy-2'-[fluoro or O-(mesyl or

Scheme 1^a



^a (a) (i) (Bu₃Sn)₂O/CHCl₃/Δ; (ii) Br₂. (b) TBDMSCl/pyridine. (c) (i) TsNHNH₂/MeOH; (ii) NaBH₄/MeOH/Δ. (d) (i) CrO₃/pyridine/Ac₂O; (ii) NaBH₄/EtOH. (e) TBAF/THF. (f) BzCl/pyridine. (g) (i) TFA/H₂O; (ii) Ac₂O/pyridine. (h) (i) Adenine/SnCl₄/CH₃CN; (ii) NH₃/MeOH. (i) (i) Bu₃SnO/MeOH; (ii) TsCl/Et₃N. (j) (i) HCl/MeOH; (ii) Me₂CO/Me₂C(OMe)₂/Δ. (k) HNO₃/Ac₂O/−60 °C. (l) (i) Amberlite IR-120 (H⁺)/MeOH; (ii) Ac₂O/DMAP.

tosyl)] products.⁸ We also demonstrated that 6'-oxy radicals (e.g., **20**, produced¹⁰ from the 6'-O-nitro derivative **19**) generated OH3'-containing C3' radicals that underwent chlorine loss and β -elimination (H/base) to provide the first model simulation of the initiation/elimination cascade that occurs during mechanism-based inactivation of RNRs with 2'-substituted nucleotides.⁹ We now describe synthesis of 6'-O-nitro-2'-O-tosylhomoadenosine (**13**) and its treatment with Bu₃SnD/AIBN. Generation¹⁰ of the 6'-oxy radical, relay abstraction of H3' (by [1,5]-hydrogen shift via a six-membered transition state¹¹) to produce the C3' radical, loss of tosylate, and elimination (H/base) gave partially deuterated 2(R)-(2-hydroxyethyl)-3-(2H)-furanone (**18**).

Regioselective oxidation of 1,2-isopropylidene- α -D-glucofuranose (**1**) gave the 5-ulose **2**^{12,13} (91%, Scheme 1). Silylation (O6) and deoxygenation (C5, via its tosylhydrazone¹⁴) of **3** gave the 5-deoxy sugar **4** (~60% from **1**). Oxidation (C3) of **4** and stereoselective reduction¹⁵ gave **5** which was desilylated to give ribohexofuranose **6** (77%). Homoadenosine¹⁶ (**7**) was obtained by benzoylation (O3 and O6) of **6**, acetal hydrolysis, acetylation, coupling¹⁷ of the anomeric acetates (adenine, SnCl₄), and deacylation. However, glycosyl cleavage occurred upon attempted nitration¹⁸ of derivatives of **7**.

Methanolysis¹⁹ of **5** and one-pot treatment with acetone gave **9**, which was nitrated¹⁸ to give **10**. Acetal hydrolysis [Amberlite (H⁺)] and acetylation gave **11** (81%) which was coupled

(1) Nucleic Acid Related Compounds. 95. Paper 94: Robins, M. J.; Sarker, S.; Samano, V.; Wnuk, S. F. *Tetrahedron* **1997**, *53*, 447–456.

(2) For recent reviews, see: (a) Stubbe, J. *Adv. Enzymol. Relat. Areas Mol. Biol.* **1989**, *63*, 349–419. (b) Reichard, P. *Science* **1993**, *260*, 1773–1777. (c) Stubbe, J.; van der Donk, W. A. *Chem. Biol.* **1995**, *2*, 793–801. (d) Robins, M. J.; Samano, M. C.; Samano, V. *Nucleosides Nucleotides* **1995**, *14*, 485–493. (e) Sjöberg, B.-M. In *Nucleic Acids and Molecular Biology*; Eckstein, F.; Lilley, D. M. J., Eds.; Springer-Verlag: Berlin, 1995; Vol. 9, pp 192–221.

(3) (a) Nordlund, P.; Sjöberg, B.-M.; Eklund, H. *Nature* **1990**, *345*, 593–598. (b) Uhlin, U.; Eklund, H. *Nature* **1994**, *370*, 533–539.

(4) Mao, S. S.; Yu G. X.; Chalfoun, D.; Stubbe, J. *Biochemistry* **1992**, *31*, 9752–9759.

(5) Stubbe, J.; Ator, M.; Krenitsky, T. J. *Biol. Chem.* **1983**, *258*, 1625–1631.

(6) Thelander, L.; Larsson, B.; Hobbs, J.; Eckstein, F. *J. Biol. Chem.* **1976**, *251*, 1398–1405.

(7) (a) Harris, G.; Ator, M.; Stubbe, J. *Biochemistry* **1984**, *23*, 5214–5225. (b) Ator, M. A.; Stubbe, J. *Biochemistry* **1985**, *24*, 7214–7221.

(8) Robins, M. J.; Wnuk, S. F.; Henández-Thirring, A. E.; Samano, M. C. *J. Am. Chem. Soc.* **1996**, *118*, 11341–11348.

(9) Robins, M. J.; Guo, Z.; Samano, M. C.; Wnuk, S. F. *J. Am. Chem. Soc.* **1996**, *118*, 11317–11318.

(10) (a) Vite, G. D.; Fraser-Reid, B. *Synth. Commun.* **1988**, *18*, 1339–1342. (b) Lopez, J. C.; Alonso, R.; Fraser-Reid, B. *J. Am. Chem. Soc.* **1989**, *111*, 6471–6473.

(11) (a) Barton, D. H. R.; Beaton, J. M.; Geller, L. E.; Pechet, M. M. *J. Am. Chem. Soc.* **1961**, *83*, 4076–4083. (b) Kabasakalian, P.; Townley, E. R.; Yudis, M. D. *J. Am. Chem. Soc.* **1962**, *84*, 2716–2718.

(12) Tsuda, Y.; Hanajima, M.; Matsuhira, N.; Okuno, Y.; Kanemitsu, K. *Chem. Pharm. Bull.* **1989**, *37*, 2344–2350.

(13) New compounds had satisfactory ¹H and ¹³C NMR spectra and elemental analyses (C,H,N) and/or HRMS.

(14) Caglioti, L. *Organic Syntheses*; Wiley: New York, 1988; Collect. Vol. VI, pp 62–63.


(15) (a) Garegg, P. J.; Samuelsson, B. *Carbohydr. Res.* **1978**, *67*, 267–270. (b) Sowa, W.; Thomas, G. H. S. *Can. J. Chem.* **1966**, *44*, 836–838.

(16) Ryan, K. J.; Arzoumanian, H.; Acton, E. M.; Goodman, L. J. *Am. Chem. Soc.* **1964**, *86*, 2503–2508.

(17) Saneyoshi, M.; Satoh, E. *Chem. Pharm. Bull.* **1979**, *27*, 2518–2521. (18) Lichtenhaler, F. W.; Müller, H. J. *Synthesis* **1974**, 199–201.

(19) Lerner, L. M. *J. Org. Chem.* **1978**, *43*, 2469–2473.



(20) Wagner, D.; Verheyden, J. P. H.; Moffatt, J. G. *J. Org. Chem.* **1974**, *39*, 24–30.

Scheme 2^a^a (a) Bu₃SnD/AIBN/benzene/Δ.

(adenine, SnCl₄)¹⁷ and deprotected to give 6'-O-nitrohomoadenosine (12). Regioselective 2'-O-tosylation²⁰ of 7 and 12 gave 8 (62%) and 13 (63%), respectively.

Treatment of 13 (Scheme 2) with Bu₃SnD/AIBN/benzene/Δ/2 h¹⁰ resulted in its total conversion to adenine, 2(R)-(2-hydroxyethyl)-3-(2H)-furanone⁹ (18, 62%), and 8 [28%, deuterium transfer to the 6'-oxy radical; no observed ²H exchange at C3' (¹H NMR)]. ¹H NMR spectra of our homologated⁹ furanone⁷ 18 had ~30% integrated reduction in the signal at δ 5.71 (H4), and HRMS peaks at *m/z* 129.0545 (100, MH⁺ [C₆H₉O₃] = 129.0552) and 130.0619 (41, MH⁺ [C₆H₈DO₃] = 130.0614) confirmed the incorporation of deuterium. A mechanism for conversion of 13 into 18 is in harmony with generation of 6'-oxy radical 14 followed by a [1,5]-shift of H3' to give the C3' radical 15. Departure of the 2'-tosylate anion, induced by a [1,2]-electron shift, would produce the C2'-radical intermediate 16. Deuterium transfer (Bu₃SnD to 16) should occur selectively from the less-hindered α-face²¹ to give the unstable 2'-deoxy-2'-deuterio-3'-ketohomoadenosines 17 [C2' (*R/S*) ≈ 30:70] which would undergo β-elimination to give 18 (with ~30% deuterium incorporation at C4). Loss of tosylate from 15 to give 16 is analogous to the [1,2]-hydride shift rearrangement^{22,23} observed during conversion of 2'-O-tosyladenosine into 9-(2-deoxy-β-D-threo-pentofuranosyl)adenine (LiEt₃BH/DMSO).²² In that case, a 2'-deoxy-3'-keto intermediate is formed by a [1,2]-hydride shift (H3' from C3' to C2') with tosylate expulsion. The present [1,2]-electron shift with generation of a carbonyl group (at C3') and electron charge repulsion at C2' would provide the driving force for expulsion of the 2'-tosylate (15 → 16).

Deuterium incorporation into 18 [C4 (D/H) ≈ 30:70] occurred in contrast with our parallel treatment of 2'-chloro-2'-deoxy-6'-O-nitrohomouridine (19) (no ²H in 18).^{9,24} Departure of a chlorine atom, rather than a chloride anion,⁷ from 20 followed by elimination (H/base) gives enol 21.⁹ Deuterium transfer from Bu₃SnD to chlorine would propagate radical chains (Scheme

Scheme 3^a^a (a) Bu₃SnD/AIBN/benzene/Δ.Scheme 4^a^a (a) TBDPSCl/pyridine. (b) Dess–Martin periodinane/CH₂Cl₂. (c) Bu₃SnD/AIBN/benzene/Δ.

3, no ²H incorporation into 18); whereas tosylate loss from 15 produces radical 16 which would undergo deuterium transfer from Bu₃SnD to propagate the 13 to 16 radical chains (Scheme 2).

Support for the latter mechanism was provided by parallel treatment of the 2'-deoxy-3'-keto²⁵ derivatives 24 (Scheme 4). Silylation of 2'-deoxy-2'-deuterioadenosine²¹ [22; C2' (*R/S*) ≈ 85:15] gave 23, and Dess–Martin oxidation²⁶ (C3') gave 24. Downfield shifts of the H2', 2'' signals, reduction in their intensities, and simplification in splittings in ¹H NMR spectra were in harmony with 24. Subjection of 24 to Bu₃SnD/AIBN/benzene/Δ gave 2(R)-{[(*tert*-butyldiphenylsilyl)oxy]methyl}-3-(2H)-furanone (25), a dehomologated analogue of 18. Integrated reduction (~15%) of the ¹H NMR signal at δ 5.75 (H4) was in harmony with trans stereoselective β-elimination (D/adenine) from 24 (~85% *S*-[²H]). Spontaneous decomposition of such 2'-deoxy-3'-ketonucleosides with elimination of the base is well-known.^{23a,27}

In summary, we prepared 6'-O-nitro-2'-O-tosylhomoadenosine (13) and demonstrated its radical-induced decomposition (Bu₃SnD/AIBN) to adenine and 2(R)-(2-hydroxyethyl)-3-(2H)-furanone (18) with ~30% deuterium at C4. Parallel treatment of 2'-chloro-2'-deoxy-6'-O-nitrohomouridine (19) gave 18 without deuterium. This provides the first biomimetic models for differentiation between radical (no ²H in 18) and anionic (~30% ²H in 18) departure of 2'-substituents upon generation of O3'-containing C3' radicals at active sites of RNRs.

Acknowledgment. We thank the American Cancer Society (DHP-34) and Brigham Young University development funds for support and Mrs. Jeanny K. Gordon for assistance with the manuscript.

Supporting Information Available: Experimental details and characterization/spectral data for compounds 2–13, 18, and 23–25 (10 pages). See any current masthead page for ordering and Internet access instruction.

JA970171C

(21) Robins, M. J.; Wilson, J. S.; Hansske, F. J. *Am. Chem. Soc.* **1983**, 105, 4059–4065.

(22) Hansske, F.; Robins, M. J. *J. Am. Chem. Soc.* **1983**, 105, 6736–6737.

(23) (a) Kawana, M.; Takeuchi, K.; Ohba, T.; Kuzuhara, H. *Nucleic Acids Res., Symp. Ser.* **1986**, (17), 37–40. (b) Kawana, M.; Kuzuhara, H. *Tetrahedron Lett.* **1987**, 28, 4075–4078. (c) Kawana, M.; Kuzuhara, H. *J. Chem. Soc., Perkin Trans. I* **1992**, 469–478.

(24) Careful repetition of the 19/Bu₃SnD/AIBN/benzene/Δ experiment⁹ showed <2% ²H incorporation at C4 of 18 (¹H NMR integration limit; no [²H]-containing ion peaks with HRMS).

(25) Samano, V.; Robins, M. J. *J. Org. Chem.* **1990**, 55, 5186–5188.

(26) (a) Dess, D. B.; Martin, J. C. *J. Org. Chem.* **1983**, 48, 4155–4156. (b) Ireland, R. E.; Liu, L. *J. Org. Chem.* **1993**, 58, 2899–2899.

(27) (a) Binkley, R. W.; Hehemann, D. G.; Binkley, W. W. *J. Org. Chem.* **1978**, 43, 2573–2575. (b) Hansske, F.; Madej, D.; Robins, M. J. *Tetrahedron* **1984**, 40, 125–135.