

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/11087671>

Asymmetric Total Synthesis of (–)-Azaspirene, a Novel Angiogenesis Inhibitor

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · NOVEMBER 2002

Impact Factor: 12.11 · DOI: 10.1021/ja0276826 · Source: PubMed

CITATIONS

35

READS

54

10 AUTHORS, INCLUDING:



Yujiro Hayashi

Tohoku University

260 PUBLICATIONS 7,414 CITATIONS

SEE PROFILE



Mitsuru Shoji

Keio University

132 PUBLICATIONS 3,937 CITATIONS

SEE PROFILE



Junichiro Yamaguchi

Nagoya University

97 PUBLICATIONS 3,078 CITATIONS

SEE PROFILE



Hiroyuki Osada

RIKEN

482 PUBLICATIONS 10,224 CITATIONS

SEE PROFILE


Asymmetric Total Synthesis of (–)-Azaspirene, a Novel Angiogenesis Inhibitor

Yujiro Hayashi,^{*,†} Mitsuru Shoji,[†] Junichiro Yamaguchi,[†] Kenji Sato,[†] Shinpei Yamaguchi,[†] Takasuke Mukaiyama,[†] Ken Sakai,[§] Yukihiro Asami,^{||} Hideaki Kakeya,^{||} and Hiroyuki Osada^{||}

Department of Industrial Chemistry, Faculty of Engineering, and Department of Applied Chemistry, Faculty of Science, Tokyo University of Science, Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan, and Antibiotics Laboratory, RIKEN, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan


Received July 12, 2002

The inhibition of angiogenesis is a promising method of treating angiogenesis-related diseases such as cancer and rheumatoid arthritis.¹ We have recently completed an asymmetric total synthesis of epoxyquinols A and B,² which we have isolated and identified as novel, unique angiogenesis inhibitors.³ Our continuing search for new angiogenesis inhibitors from natural sources led us to azaspirene (**1**), isolated from the fungus *Neosartorya* sp.⁴ Structurally, azaspirene (**1**) contains a highly oxygenated 1-oxa-7-azaspiro-[4.4]non-2-ene-4,6-dione skeleton with benzyl and hexadiene substituents, and a core structure also found in the pseurotins⁵ and synerazol.⁶ Although several synthetic studies on the pseurotins have been reported,^{5e–h} their total synthesis has not yet been accomplished. Because of its interesting biological properties and rare structure, we have investigated the total synthesis of azaspirene (**1**), aiming to determine its absolute stereochemistry.



Our synthesis (see Scheme 1) started with the Sharpless asymmetric dihydroxylation of methyl 2-pentenoate (**2**) using (DHQD)₂PHAL as the chiral ligand.⁷ This gave (2R,3S)-diol **3** in 88% yield. This was treated with dimethoxypropane in the presence of a catalytic amount of TsOH·H₂O, to afford acetal **4** in 95% ee⁸ and 79% yield. The next step, an aldol condensation of **4** with phenylpropargyl aldehyde (**5**),⁹ proved troublesome. Although the lithium enolate of **4** reacted with **5** smoothly at –78 °C, all four possible aldol products **6** were obtained with very poor diastereoselectivity (57:20:13:8), albeit in good yield (total 98%). The desired *syn*-aldol **6a** was obtained in only a small amount (13%), while the major isomer was *anti*-aldol **6b** (57%). As screening of solvent, base, and additives did not significantly improve this result, we examined the alternative, Mukaiyama aldol reaction.

After conversion of ester **4** to its ketene silyl acetal **7**,¹⁰ the aldol reaction of this was investigated in the presence of various Lewis acids (eq 1),



* To whom correspondence should be addressed. E-mail: hayashi@ci.kagu.tus.ac.jp.

[†] Department of Industrial Chemistry, Faculty of Engineering, Tokyo University of Science.

[§] Department of Applied Chemistry, Faculty of Science, Tokyo University of Science.

^{||} Antibiotics Laboratory, RIKEN.

Table 1. Effect of Lewis Acid in Mukaiyama Aldol Reaction of **5** and **7**^a

Lewis acid	yield/% ^b	isomer ratio/%		other isomers
		6a	6b	
SnCl ₄	67	61	33	6
BF ₃ ·OEt ₂	41	63	34	3
ZnBr ₂	62	68	31	1
Ti(O- <i>i</i> -Pr) ₄	49	91	9	0
MgBr ₂ ·OEt ₂	72	94	6	0


^a Reaction conditions; **7**:**5**:Lewis acid = 1:1.5:2.5, CH₂Cl₂, –78 °C.

^b Isolated yield based on ester **4**. ^c Isomer ratio was determined by 400 MHz ¹H NMR analysis.

with the results summarized in Table 1. Although the desired *syn*-aldol **6a** was obtained as the major isomer using SnCl₄, BF₃·OEt₂, and ZnBr₂, the diastereoselectivity was insufficient. Ti(O-*i*-Pr)₄¹¹ afforded the desired isomer **6a** with high selectivity but in poor yield. MgBr₂·OEt₂,¹² on the other hand, was found to be a suitable promoter, affording **6a** with both high diastereoselectivity and in good yield, without loss of enantioselectivity as checked by chiral-HPLC analysis.⁸ The stereochemistry of **6a** was established unambiguously by X-ray crystallographic analysis of a crystalline amide,¹³ synthesized by the following sequence: protection of the hydroxy group of **6a** as its benzyl ether, ester hydrolysis, and amide formation. The stereochemistry of **6b** was determined by oxidation of **6a**, followed by reduction, affording **6a** and **6b**.

Protection of **6a** with TIPSOt and 2,6-lutidine, afforded silyl ether **8** in 98% yield, which was hydrolyzed to provide carboxylic acid **9**. This was next treated with oxalyl chloride and NEt₃, providing the acid chloride, which was reacted with NH₃ to give amide **10** in 45% yield over two steps, along with recovered carboxylic acid **9** in 34% yield. Amide **10** was crystalline, and a single recrystallization gave optically pure compound.⁸ When this amide **10** was reacted with NaH in DMF at room temperature for 1 h, the *Z*-benzylidene γ -lactam **11**¹⁴ and its *E*-isomer¹⁴ were obtained in 92 and 7% yield, respectively.¹⁵ Treatment of **11** with CF₃CO₂H in the presence of MeOH cleaved the acetal, affording diol **12** in 92% yield without affecting the benzylidene moiety. Oxidation of **12** with the Dess–Martin periodinane (DMP)¹⁶ in the presence of water according to Schreiber's modified conditions¹⁷ gave α -hydroxy ketone **13** in 82% yield. Aldol condensation of the lithium enolate of **13** and heptadienal in the presence of HMPA afforded a mixture of diastereomeric aldols **14** in 50% yield. Ketone **13** was recovered in 47% yield, and thus a high conversion yield (94%) had been achieved. It should be noted that it is not necessary to protect the *tert*-alcohol or amide functionality during this aldol reaction. Oxidation of **14** with DMP gave 1,3-diketone **15**,¹⁸ which was partially converted to the azaspiro[4.4]nonenedione bicycle **16** when purified on thin-layer chromatography (**15** + **16**, 94%, **15**:

Scheme 1. Total Synthesis of (–)-azaspirene (1)



16 = 1:1). When a mixture of **15** and **16** was treated with a catalytic amount of TsOH·H₂O for 2.8 h, complete formation of the azaspiro[4.4]nonenedione bicyclic and hydration of the benzylidene group occurred concurrently to afford **17** as a single isomer¹⁹ in 91% yield. Deprotection of the TIPS group with NH₄F in MeOH afforded azaspirene (**1**) in 35% yield. The order of the last two reactions is very important: When the benzylidene derivative **16**²⁰ was first deprotected with NH₄F, and then hydrated with TsOH·H₂O, racemic azaspirene was formed, probably because of a retro-aldol reaction. During the sequence of hydration and deprotection, the presence of an hydroxy group at C8 of the azaspiro[4.4]nonenedione seems to prevent the racemization. Synthetic azaspirene exhibited properties identical to those of the natural product (¹H NMR, ¹³C NMR, IR, mp, *R_f* value, and chiral HPLC analysis). Comparison of the optical rotation (synthetic **1**; [α]_D²⁵ −207 (*c* = 0.13, MeOH), natural **1**¹⁴; [α]_D²⁵ −204.4 (*c* = 0.158, MeOH), established the absolute stereochemistry of the natural product to be (5*S*,8*R*,9*R*).

In summary, the first asymmetric total synthesis of (–)-azaspirene (**1**) has been achieved, and its absolute stereochemistry has been determined. There are several noteworthy features to this total synthesis: The MgBr₂·OEt₂-mediated, diastereoselective Mukaiyama aldol reaction of **5** and **7**, the NaH-promoted intramolecular cyclization of the alkynylamide **10** to form selectively the *Z*-benzylidene γ-lactam **11**, the aldol reaction of **13** containing functionalized γ-lactam moiety without protection of *tert*-alcohol and amide functionalities, and the importance of the order of the last two reactions.

Acknowledgment. This work was partially supported by the Sumitomo Foundation.

Supporting Information Available: Detailed experimental procedures, full characterization, copies of ¹H, ¹³C NMR, and IR spectra of all new compounds (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Folkman, J. *J. Natl. Cancer Inst.* **1990**, *82*, 4. (b) Risau, W. *Nature* **1997**, *386*, 671. (c) Klagsbrum, M.; Moses, M. A. *Chem. Biol.* **1999**, *6*, R217. (d) Gasparini, G. *Drugs* **1998**, *58*, 17.
- (2) Shoji, M.; Yamaguchi, J.; Kakeya, H.; Osada, H.; Hayashi, Y. *Angew. Chem., Int. Ed.* **2002**, *41*, 3192.
- (3) Kakeya, H.; Onose, R.; Koshino, H.; Yoshida, A.; Kobayashi, K.; Kageyama, S.-I.; Osada, H. *J. Am. Chem. Soc.* **2002**, *124*, 3496.
- (4) Asami, Y.; Kakeya, H.; Onose, R.; Yoshida, A.; Matsuzaki, H.; Osada, H. *Org. Lett.* **2002**, *4*, 2845.
- (5) (a) Bloch, P.; Tamm, C.; Bollinger, P.; Petcher, T. J.; Weber, H. P. *Helv. Chim. Acta* **1976**, *59*, 133. (b) Bloch, P.; Tamm, C. *Helv. Chim. Acta* **1981**, *64*, 304. (c) Breitenstein, W.; Cheval, K. K.; Mohr, P.; Tamm, C. *Helv. Chim. Acta* **1981**, *64*, 379. (d) Mohr, P.; Tamm, C. *Tetrahedron* **1981**, *37*, 201. (e) Dolder, M.; Shao, X.; Tamm, C. *Helv. Chim. Acta* **1990**, *73*, 63. (f) Shao, X.; Dolder, M.; Tamm, C. *Helv. Chim. Acta* **1990**, *73*, 483. (g) Su, Z.; Tamm, C. *Helv. Chim. Acta* **1995**, *78*, 1278. (h) Su, Z.; Tamm, C. *Tetrahedron* **1995**, *51*, 11177. (i) Komagata, D.; Fujita, S.; Yamashita, N.; Saito, S.; Morino, T. *J. Antibiot.* **1996**, *49*, 958.
- (6) Ando, O.; Satake, H.; Nakajima, M.; Sato, A.; Nakamura, T.; Kinoshita, T.; Furuya, K.; Haneishi, T. *J. Antibiot.* **1991**, *44*, 382.
- (7) Sharpless, K. B.; Amberg, W.; Bennani, Y. L.; Crispino, G. A.; Hartung, J.; Jeong, K.-S.; Kwong, H.-L.; Morikawa, K.; Wang, Z.-M.; Xu, D.; Zhang, X.-L. *J. Org. Chem.* **1992**, *57*, 2768.
- (8) For the determination of ee, see the Supporting Information.
- (9) Allen, C. F. H.; Enders, C. O., Jr. *Organic Syntheses*; Wiley & Sons: New York, 1973; Collect. Vol. 3, p 731.
- (10) Ketene silyl acetal **7** was obtained as a single isomer, the geometry of which has not been determined.
- (11) In the elegant total syntheses of zaragozic acid and cinatrin by Evans et al., Ti(O-*i*-Pr)₄ is an effective Lewis acid in Mukaiyama aldol reaction of a silylketene acetal derived from di-*tert*-butyl-(2*S*,3*S*)-1,4-dioxaspiro[4.4]nonane-2,3-dicarboxylate. Evans, D. A.; Barrow, J. C.; Leighton, J. L.; Robichaud, A. J.; Sefkow, M. *J. Am. Chem. Soc.* **1994**, *116*, 12111; Evans, D. A.; Trotter, B. W.; Barrow, J. C. *Tetrahedron* **1997**, *53*, 8779.
- (12) For recent elegant use of MgBr₂·OEt₂ as an effective Lewis acid, see: (a) Fujisawa, H.; Sasaki, Y.; Mukaiyama, T. *Chem. Lett.* **2001**, 190. (b) Evans, D. A.; Tedrow, J. S.; Shaw, J. T.; Downey, C. W. *J. Am. Chem. Soc.* **2002**, *124*, 392. (c) Sibi, M. P.; Sausker, J. B. *J. Am. Chem. Soc.* **2002**, *124*, 984.
- (13) Crystal data: monoclinic, space group *P*2₁, *a* = 11.404(3) Å, *b* = 7.941(2) Å, *c* = 12.843(3) Å, β = 105.643(5)°, *V* = 1120.0(5) Å³, *Z* = 2, *R*₁ = 0.0698 for *I* > 2.0σ(*I*), *wR*₂ = 0.1713 for all data (5002 reflections). The details are supplied as Supporting Information (Figure S1).
- (14) The geometry of the olefins was determined by NOESY experiment.
- (15) (a) Koseki, Y.; Kusano, S.; Ichi, D.; Yoshida, K.; Nagasaka, T. *Tetrahedron* **2000**, *56*, 8855. (b) Cid, M. M.; Dominguez, D. D.; Castedo, L.; Vazquez-Lopez, E. M. *Tetrahedron* **1999**, *55*, 5599.
- (16) (a) Dess, D. B.; Martin, J. C. *J. Org. Chem.* **1983**, *48*, 4155. (b) Dess, D. B.; Martin, J. C. *J. Am. Chem. Soc.* **1991**, *113*, 7277. (c) Ireland, R. E.; Liu, L. *J. Org. Chem.* **1993**, *58*, 2899.
- (17) Meyer, S. D.; Schreiber, S. L. *J. Org. Chem.* **1994**, *59*, 7549.
- (18) ¹H and ¹³C NMR analyses show that **15** exists in the 1,3-diketone form rather than the keto–enol form.
- (19) The stereochemistry at C8 was confirmed by NOESY and difference NOE experiments, see Supporting Information.
- (20) **16** was obtained in 94% yield from **15** by repeated treatment with thin-layer chromatography (TLC).

JA0276826