

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/228440163>

Synthesis and cytotoxic activity of some 17-picolyl and 17-picolinylidene androstane derivatives

ARTICLE in EUROPEAN JOURNAL OF MEDICINAL CHEMISTRY · JUNE 2012

Impact Factor: 3.45 · DOI: 10.1016/j.ejmech.2012.06.030 · Source: PubMed

CITATIONS

9

READS

58

7 AUTHORS, INCLUDING:



Jovana Ajdukovic

University of Novi Sad

22 PUBLICATIONS 88 CITATIONS

SEE PROFILE



Marija N. Sakač

University of Novi Sad

84 PUBLICATIONS 347 CITATIONS

SEE PROFILE



Gordana Bogdanovic

148 PUBLICATIONS 1,047 CITATIONS

SEE PROFILE



Katarina Penov Gaši

University of Novi Sad

128 PUBLICATIONS 440 CITATIONS

SEE PROFILE



Original article

Synthesis and cytotoxic activity of some 17-picolyl and 17-picolinyldene androstane derivatives

Evgenija A. Djurendić^a, Jovana J. Ajduković^{a,*}, Marija N. Sakač^a, János J. Csanádi^a, Vesna V. Kojić^b, Gordana M. Bogdanović^b, Katarina M. Penov Gaši^a

^a Department of Chemistry, Faculty of Sciences, University of Novi Sad, Trg Dositeja Obradovića 3, 21000 Novi Sad, Serbia

^b Oncology Institute of Vojvodina, Institutski put 4, 21204 Sremska Kamenica, Serbia

ARTICLE INFO

Article history:

Received 21 September 2011

Received in revised form

14 June 2012

Accepted 15 June 2012

Available online 23 June 2012

Keywords:

Androst-5-ene derivatives

17 α -Picolyl

17-Picolinyldene derivatives

Cytotoxicity

ABSTRACT

New 17-picolyl and 17-picolinyldene androstane derivatives, **3–10**, **15**, **18**, **19**, **22** and **23**, were synthesized starting from 17 α -picolyl-androst-5-en-3 β ,17 β -diol (**1**) and 17(Z)-picolinyldene-androst-5-en-3 β -ol (**2**). Reaction of **1** with *m*-chloroperoxybenzoic acid gives 5 α ,6 α -epoxy N-oxide derivative **3**, or, with Jones reagent, 3,6-dione derivative **4**; while 17 α -picolyl-androst-5-en-3 β ,4 α ,17 β -triol (**5**) or 3 β ,4 β ,17 β -triol (**6**) derivatives are obtainable from **1** using SeO₂ in dioxane. Base-catalyzed tosyl group elimination from **7** or **9** affords AB conjugated derivatives **8** and **10**. Oppenauer oxidation of **1** and **2** yields 4-en-3-one derivatives **11** and **12**, which, with H₂O₂ in 4 M NaOH, affords 4 α ,5 α and 4 β ,5 β -epoxides **13**, **14**, **16** and **17**. New 4-methoxy-3-keto derivatives **15** and **18** were obtained from **13** and **14**, or, with methanol in 4 M NaOH, from **16** and **17**. Reduction of **11** with NaBH₄ gives **22**, which was then acetylated to obtain **23**. All new derivatives were screened for antitumor activity against human breast adenocarcinoma ER+, MCF-7; human breast adenocarcinoma ER-, MDA-MB-231; prostate cancer AR-, PC-3; human cervix carcinoma, HeLa; and colon cancer, HT-29 cells; as well as one human non-tumor cell line, MRC-5. Compounds **3**, **5**, **6**, **8**, **10**, **18**, **19** and **22** exhibited significant antitumor activity against MDA-MB-231 breast cancer cells; while **5**, **6** and **10** also showed strong cytotoxicity against HT-29. Only compound **19** exhibited significant activity against MCF-7 breast cancer cells. No compounds displayed cytotoxicity against non-tumor MRC-5 cells.

© 2012 Elsevier Masson SAS. All rights reserved.

1. Introduction

Steroids are a fundamental class of natural biological signaling molecules, which have demonstrated potential for development into potent drugs for the treatment of a large number of diseases, including: brain tumors, breast cancer, prostate cancer, cardiovascular disease or autoimmune disorders [1,2].

Androgens are male steroid hormones which have a wide-range of important physiological functions, including central involvement in the development of male sex characteristics and other phenotypes [3–5]. Anti-androgens are a diverse class of drugs which can compete with circulating androgens for binding sites on androgen receptors within prostate cells, leading to targeted apoptosis and inhibition of androgen-dependent prostate cancer growth [5].

Several steroidal derivatives with substitutions at the C-17 position have been previously described in the literature as potent

biologically active compounds [6,7]. Similarly, in our own work [8–10], we described the synthesis and characterization of several 17 α -picolyl and 17-picolinyldene-androst-5-ene derivatives, with anti-aromatase (CYP19A1) and/or anti-tumor activity against a range of tumor cell lines. Several of these tested androstane derivatives displayed significant aromatase inhibitory activity, and/or strong activity against 3 common tumor cell lines (human cervix carcinoma, HeLa; human melanoma, FemX; and human myelogenous leukemia, K562), with IC₅₀ values between 4 and 10 μ M [8]. Because it is well known [11] that the presence of nitrogen heteroatoms can interfere with steroidal hydroxylation by directly interacting with the cytochrome P450 heme iron in aromatase; these androstane compounds were specifically designed to incorporate nitrogen in the steroidal structure. In the present study, we continue our investigation of C-17 substitutions, and report a simple synthetic route for the preparation of 17-picolyl and 17-picolinyldene androst-5-ene derivatives. The effect of these new androstane derivatives on cell proliferation was tested against a wide-range of common tumor cell lines, including: human breast

* Corresponding author. Tel.: +381 21 485 27 33; fax: +381 21 454 065.

E-mail address: jovana.ajdukovic@dh.uns.ac.rs (J.J. Ajduković).

adenocarcinoma ER+, MCF-7; human breast adenocarcinoma ER-, MDA-MB-231; prostate cancer AR-, PC-3; human cervix carcinoma, HeLa; and colon cancer, HT-29.

2. Results and discussion

2.1. Chemistry

The initial steps in the preparation of 17-picolyl and 17-picolinylidene androstane derivatives consist of a series of different reactions of 17 α -picolyl-androst-5-en-3 β -ol (**1**) [12,13] and 17-picolinylidene-androst-5-en-3 β -ol (**2**) [12,13] (see Fig. 1). Specifically, compound **3** (5 α ,6 α -epoxy N-oxide) was obtained from compound **1** by reaction with *m*-chloroperoxybenzoic acid (MCPBA) at 0 °C, 1 h \rightarrow 10 °C, 3 h \rightarrow and room temperature, 24 h, at a molar ratio of **1** : MCPBA = 1:3 (Scheme 1). The presence of 5 α ,6 α -epoxy function in compound **3** was confirmed by ¹H and ¹³C NMR spectroscopy: the doublet at 2.93 ppm was attributed to H-6 β and ¹³C NMR signals for C-5 and C-6 atoms were identified at 65.98 and 59.15 ppm, respectively. In addition, a signal for the 17 β -OH proton was visible at 6.65 ppm for compound **3** (but not compound **1**), suggesting a lack of rapid proton exchange; while significant chemical shift changes were also observed for pyridine C-atoms in compound **3** (but not compound **1**). Together, these results strongly suggest the presence of an intramolecular hydrogen bond between 17 β -OH and the N-oxide function on the pyridine core in compound **3**, which is absent in compound **1**.

In contrast with the above described synthesis, reaction of compound **1** with Jones reagent [14] at 0 °C for 1.5 h affords 17 β -hydroxy-17 α -picolyl-androst-4-en-3,6-dione (**4**). Consistent with this product, ¹H NMR data for compound **4** shows the expected singlet for the H-4 proton at 6.19 ppm, in place of the H-6 doublet at 5.36 ppm visible in ¹H NMR spectra of compound **1**. In addition, signals for C-3 and C-6 carbonyl carbons are visible in ¹³C NMR spectra of compound **4** at 199.46 and 201.95 ppm, respectively.

Interestingly, reaction of compound **1** with selenium-dioxide in dioxane was found to give a mixture of 4 α and 4 β hydroxylated derivatives (**5** and **6**). Although a 4 β -hydroxylation mechanism was previously reported for steroidal 5-en-3-ols via selenium-dioxide oxidation by Ma and Choi [15], these authors obtained only a 4 β -hydroxy group. After separation by column chromatography, the 17 α -picolyl-androst-5-ene-3 β ,4 α -diol (**5**) product was identified by the appearance of a new ¹H NMR triplet signal at 3.89 ppm (*J* = 3.8 Hz) corresponding to the H-4 proton. In addition, a C-4 carbon signal was also visible for compound **5** at 65.06 ppm by ¹³C NMR. Analogously, for compound **6**, a new ¹H NMR doublet appeared at 4.16 ppm (*J* = 3.2 Hz), corresponding to the H-4 proton, and a C-4 carbon signal was observed at 77.24 ppm by ¹³C NMR. The β -configuration of the 4-hydroxy group in compound **6** was assigned by ¹H-NOE NMR: irradiation of the H-4 proton at 4.16 ppm

resulted in NOE enhancements at H-3 α (3.57 ppm) and H-6 (5.71 ppm). In contrast, for compound **5**, selective irradiation of the H-4 proton signal at 3.89 ppm yielded no such NOE enhancement of the H-3 signal, indicating that the C4-OH proton is directed toward the α -position, consistent with our assignment of an α -configuration for the 4-hydroxy group in compound **5**.

Base-catalyzed tosyl group elimination from **7** or **9** was used to obtain AB conjugated derivatives **8** and **10**. Specifically, compound **8**, which contains a 3,5-diene system, was obtained from compound **1** via **7** (in which the 3 β -hydroxy group is tosylated). The tosyl group was then eliminated by reaction of **7** with bis(2-aminoethyl)amine under reflux for 1.5 h, yielding only compound **8**. A similar scheme for formation of a Δ^3 double bond was previously reported by King and Bigelow [16], by reaction of epicholesteryl *p*-toluenesulfonate with pyridine; and by Numazawa et al. [17], via elimination of the tosyl group at C-3 in basic media with *N*-methylpyrrolidone. The structure of compound **8** was identified by the presence of an ¹H NMR multiplet at 5.63 ppm, and a doublet at 5.94 ppm (*J* = 9.9 Hz), corresponding to H-3 and H-4 protons, respectively. Consistent with this, ¹³C NMR signals were also recorded for C-3 (128.90 ppm) and C-4 (125.14 ppm).

Similar reactions were performed on the 17-picolinylidene-substituted derivative **2** (Scheme 1), to obtain compound **10** via 3 β tosyl derivative **9**. In this case, the tosyloxy group was also eliminated with bis(2-aminoethyl)amine at reflux for 1 h, and compound **10** was isolated from the resulting complex reaction mixture. The identity of compound **10** was also confirmed by NMR. ¹H NMR spectra of **10** shows a multiplet at 5.42 ppm for the H-3 proton, and a doublet at 5.95 ppm (*J* = 9.7 Hz) for the H-4 proton; while another multiplet signal at 5.61 ppm was assigned to the H-6 proton. Similarly, ¹³C NMR spectra contain characteristic signals for C-3, C-4, C-5 and C-6 atoms, at 128.89, 125.16, 141.57 and 124.72 ppm, respectively.

In addition to the above, we also investigated the effects of applying Oppenauer oxidation [8] to our starting compounds **1** and **2**. Oppenauer oxidation of **1** and **2** yields 4-en-3-one derivatives **11** and **12**, which, with H₂O₂ in 4 M NaOH, affords 4 α ,5 α and 4 β ,5 β -epoxides **13**, **14**, **16** and **17** (Scheme 2), respectively. Compounds **15** and **18**, which contain a 4-methoxy group, were then obtained from 17-picolyl epoxy derivatives **13** and **14** [9] and 17-picolinylidene epoxy derivatives **16** and **17** [9], in reaction with methanol and sodium-hydroxide at reflux for 3 h. Analysis of compounds **15** and **18** by ¹H NMR confirmed the expected 4-methoxy group signal at 3.60 and 3.59 ppm, respectively. Similarly, ¹³C NMR signals were observed for C-4 atoms at 147.23 ppm (**15**) and at 154.96 ppm (**18**), respectively.

We next attempted a series of synthetic schemes involving compounds **11** and **12**. Reaction of compounds **11** and **12** with potassium-*tert*-butoxide in *tert*-butanol at 45 °C for 3 h was found to afford 4,17 β -dihydroxy-17 α -picolyl-androsta-4,6-dien-





Fig. 1. 17 α -Picolyl-androst-5-en-3 β ,17 β -diol (**1**) and 17(Z)-picolinylidene-androst-5-en-3 β -ol (**2**).



Scheme 1. Reagents and reaction conditions: a) MCPBA, CH₂Cl₂, 5% NaHCO₃, 0 °C, 1 h → 10 °C, 3 h → r.t., 24 h; b) Jones reagent, acetone, 0 °C, 1.5 h; c) SeO₂, dioxane, reflux, 25.5 h; d) *p*-TsCl, Py, r.t., 45 h (for 1) or 19 h (for 2); e) bis(2-aminoethyl)amine, reflux, 1 h.

3-one (**19**), accompanied by compound **4**, and 4-hydroxy-17-picolynilidene-androsta-4,6-dien-3-one (**21**) [10] accompanied by compound **20** (Scheme 2). ¹H NMR analysis of compound **19** showed a double doublet at 6.06 ppm (*J*_{6,7} = 9.4 Hz, *J*_{7,8} = 2.0 Hz) and 6.65 ppm (*J*_{6,7} = 9.4 Hz, *J*_{6,8} = 2.8 Hz) for the H-7 and H-6 protons, as expected. Similarly, ¹³C NMR spectral signals were assigned for double bond carbons at 124.71 (C-6), 134.70 (C-5), 137.10 (C-7) and 140.21 (C-4) ppm. The molecular formula of compound **19** was confirmed by high resolution mass spectra (HRMS) (TOF); and by comparison with spectroscopic data previously reported by our group for compound **21** [10].

Based on the above, we further investigated possible synthetic routes based on compounds **11** and **12**. Interestingly, reduction of compound **11** with NaBH₄ in ethanol at room temperature for 2.5 h was found to yield a 3β-hydroxy derivative, compound **22**. Mechanistically, attack of the hydride anion in the axial direction of the carbonyl group could result in generation of the 3β-hydroxy configuration observed for compound **22**. Analogously, reduction of compound **12** with NaBH₄ in absolute ethanol at room temperature for 2 h, affords compound **24** [10]. The ¹H NMR spectrum of compound **22** was characterized by a multiplet at 4.15 ppm, attributed to the H-3α proton; while ¹³C NMR spectroscopy revealed the corresponding C-3 atom signal at 67.93 ppm. The stereochemistry of compound **22** was confirmed using selective NOE difference experiments. Because selective irradiation of the C-10 methyl protons did not result in NOE enhancement of the H-3 proton signal, the H-3 and C-10 angular methyl group are assumed to be in *trans*; and the OH group at C-3 is thus directed toward the β position.


Finally, NaBH₄ reduction of compound **11** to **22**, followed by acetylation with acetic anhydride in absolute pyridine at room temperature for 24 h was found to give compound **23**.

2.2. Cytotoxicity

As a first step toward testing the potential of the above synthesized compounds for the development of novel steroidal therapeutics, we performed a series of cytotoxicity assays. The cytotoxic activity of selected synthesized compounds was evaluated against several common human tumor cell lines, originating from 4 different solid tumor types. As a control, the selected compounds were also tested against normal fetal lung fibroblasts, MRC-5. Cytotoxic activity was evaluated *in vitro* after 48 h treatment using the SRB assay, as previously described [18].

Interestingly, all of the synthesized compounds with a 17α-picolyl substituent (**3**, **5**, **6**, **8**, **19** and **22**) appear to have stronger cytotoxicity against ER- human breast adenocarcinoma cells (MDA-MB-231) (see Table 1). Of these, compound **19**, which contains a 4-hydroxy-4,6-dien-3-on system, displayed the strongest cytotoxicity (IC₅₀ 1.2 μM); while compound **8**, which contains a 3,5-dien system, also displayed significant cytotoxicity (IC₅₀ 2.3 μM). In comparison, the cytotoxicity for the 17-picolynilidene compound **21** (IC₅₀ 9.3 μM) [10] was approximately 8-fold less than that of the 17α-picolyl derivative **19** (IC₅₀ 1.2 μM); while compound **10** (17-picolynilidene derivative) was 2-fold less cytotoxic (IC₅₀ 4.6 μM) than compound **8** (IC₅₀ 2.3 μM). One possibly interesting exception to this rule was compound **18**, a 17-picolynilidene derivative with a 4-methoxy group: compound **18** was measured to have 10-fold stronger cytotoxicity (IC₅₀ 4.1 μM) than compound **15** (IC₅₀ 42.6 μM), which also has a 4-methoxy group, but contains a 17α-picolyl substituent.

Compounds **5** and **6**, which contain an additional 4-hydroxy group, both displayed significantly stronger cytotoxicity against ER- human breast adenocarcinoma cells (MDA-MB-231) (17-fold



Scheme 2. Reagents and reaction conditions: a) 30% H_2O_2 , 4M NaOH, MeOH, 0 °C, 1 h \rightarrow 10 °C, 24 h; b) 4M NaOH, MeOH, reflux, 3.5 h (or 3 h); c) 1M t-BuOK, t-BuOH, 45 °C, 3 h (or r.t., 45 h); d) NaBH_4 , EtOH, r.t., 24 h (or 2 h); e) Ac_2O , Py, r.t., 24 h.

and 10-fold, respectively) vs. compound **1** [9] (IC_{50} 96.1 μM). Similarly, compound **3**, which contains a 5 α ,6 α -epoxy and N-oxide function, also had significantly stronger cytotoxicity (IC_{50} 7.5 μM) than compound **1** against MDA-MB-231 cells.

Intriguingly, replacement of the 3 β -hydroxy group in compound **22** with a 3 β -acetoxy group (compound **23**), resulted in a reduction

in cytotoxicity (IC_{50} 10.2 μM vs. IC_{50} 3.8 μM for **23** vs. **22**, respectively). Similarly, compound **25**, which also contains a 3 β -acetoxy group [10], showed low cytotoxicity (IC_{50} 39.3 μM) against the same cell line. However, compound **24** [10] (17-picolinylidene), with a 3 β -hydroxy group, appears to be inactive (IC_{50} > 100 μM) against MDA-MB-231 cells.

In contrast with our results from ER- breast adenocarcinoma cell lines, only compound **19** strong cytotoxicity against ER + human breast adenocarcinoma cells (MCF-7 cells) (IC_{50} 3.6 μM); although compound **8** did display moderate MCF-7 cytotoxicity (IC_{50} 17.4 μM). Interestingly, compound **19** also displayed the strongest cytotoxicity against ER- breast cancer cells, followed by compound **8**.

With respect to prostate cancer, PC-3 cells were the most sensitive to compound **1** [9] (IC_{50} 6.3 μM), although the majority of the tested compounds (**3**, **5**, **8**, **10**, **12** [9], **17** [9], **18**, **19**, **22** and **24**) showed moderate PC-3 cytotoxicity: ranging from a maximum IC_{50} of 10.1 μM (**24**) to a minimum IC_{50} of 34.1 μM (**18**). Of these, **5**, **8**, **10**, **18** and **22** also displayed moderate cytotoxicity against HeLa cancer cells (along with **6**). Unlike our results from ER- breast cancer cells, compound **24** [10] had 2-fold higher cytotoxicity (IC_{50} 10.1 μM) than compound **22** (IC_{50} 22.3 μM) against PC-3 cells.

Cytotoxicity results against the colon cancer-derived cell line (HT-29) revealed a different set a compounds with significantly

Table 1
In vitro cytotoxic activity of the tested compounds – IC_{50} values.

Compounds	IC_{50} (μM)					
	MCF-7	MDA-MB-231	PC-3	HeLa	HT-29	MRC-5
1 [9]	>100	96.1	6.3	—	—	>100
3	>100	7.5	33.6	33.5	25.8	>100
4	>100	>100	4.3	>100	>100	>100
5	>100	5.6	21.3	21.3	2.6	>100
6	>100	9.0	52.1	26.6	8.0	>100
8	17.4	2.3	24.7	27.5	11.7	>100
10	55.8	4.6	19.9	23.3	1.8	>100
15	>100	42.6	54.8	19.1	>100	>100
18	>100	4.1	34.1	15.3	56.0	>100
19	3.6	1.2	25.6	36.6	>100	>100
21 [10]	>100	9.3	>100	—	—	>100
22	>100	3.8	22.3	26.9	>100	>100
23	>100	10.2	31.3	>100	10.6	>100
24 [10]	>100	>100	10.1	—	—	>100
Doxorubicin	0.75	0.12	95.61	1.17	0.32	0.12
Formestane	>100	55.5	48.36	5.55	>100	>100

strong cytotoxic activity: according to IC_{50} values, **5** and **10** showed strong cytotoxicity (IC_{50} 2.6 μ M and 1.8 μ M, respectively) against HT-29 cells; followed by compound **6** (IC_{50} 8.0 μ M).

Overall, the ER- breast cancer cell line (MDA-MB-231) appears to be the most sensitive to the majority of tested compounds (Fig. 2).

In fact, all of the tested compounds induced a dose-dependent cytotoxic response against MDA-MB-231 cancer cells: with some compounds displaying more toxicity than others. For example, compounds **5**, **18** and **19** showed stronger cytotoxic activity for the whole range of concentrations vs. Formetan (a CYP19A1 aromatase inhibitor in clinical use); whereas compounds **8** and **22** only induced higher cytotoxicity at concentrations > 0.1 μ M; and compounds **3**, **4**, **6**, **15**, and **23** showed either similar or less cytotoxicity than Formetan.

As an indication of their potential potency, the cytotoxic activities of compounds **19**, **8** and **5** were found to be similar to that of Doxorubicin at concentrations ≥ 0.1 μ M. However, only compound **18** displayed higher cytotoxicity than Doxorubicin (up to 10% at concentrations < 1 μ M against MDA-MB-231 cells).

As a control, all of the newly synthesized compounds were confirmed to be non-toxic to normal MRC-5 cells, whereas Doxorubicin was very toxic to these cells.

3. Conclusion

Novel androstane derivatives **3–10**, **15**, **18**, **19**, **22** and **23** were synthesized from 17 α -picolyl-androst-5-en-3 β ,17 β -diol (**1**) and 17(Z)-picolinylidene-androst-5-en-3 β -ol (**2**), respectively. These compounds were evaluated for cytotoxic activity against five common tumor cell lines, and one normal cell line. Synthesized compounds with a 17-picoly substituent (**3**, **5**, **6**, **8**, **10**, **18**, **19** and **22**) showed markedly strong cytotoxic activity against ER- breast cancer cells (MDA-MB-231 cells); while compound **19** (followed by **8**) also displayed the strongest cytotoxicity against ER + breast cancer cells (MCF-7); and compounds **5**, **6** and **10** were the most cytotoxic to HT-29 colon cancer cells.

4. Experimental

4.1. Chemistry

All melting points were determined using a Büchi SMP 20 apparatus and are reported uncorrected. IR spectra were recorded on a NEXUS 670 SP-IR spectrometer (wave numbers in cm^{-1}). NMR spectra were recorded on a Bruker AC 250E spectrometer




Fig. 2. Cytotoxic activity of selected synthesized compounds against MD-MB-231 cells. For- formetan; DOX- doxorubicin.

operating at 250 MHz (^1H) and 62.5 MHz (^{13}C), and are reported in ppm (symbol 100 \f “Symbol” \s 11d-scale) downfield from a tetramethylsilane internal standard; coupling constants (J) are given in Hz. GC/MS analyses were performed on an Agilent Technologies GC 6890N instrument with Mass Selective Detector 5973; the first number denotes the m/z value, while ion abundances are given in parentheses. High resolution mass spectra (TOF) were recorded on a 6210 Time-of-Flight LC/MS Agilent Technologies (ESI+) instrument. Chromatographic separations were performed on silica gel columns (Kieselgel 60, 0.063–0.20 mm, Merck). All reagents used were of analytical reagent grade. All solutions were dried over anhydrous sodium sulfate.

4.1.1. 5 α ,6 α -Epoxy-17 α -picolyl-androstane-N-oxide-3 β ,17 β -diol (**3**)

A 5% solution of NaHCO_3 (2.5 ml) was added to a solution of compound **1** (0.305 g, 0.799 mmol) in dichloromethane (10 ml) at 0 °C, to pH 8. Then, MCPBA (0.414 g, 2.38 mmol) was added and the reaction mixture was stirred for 1 h at 0 °C, 3 h at 10 °C and 24 h at room temperature in the dark. The resulting organic layer was then washed successively with a solution of 10% $\text{Na}_2\text{S}_2\text{O}_3$ (2 \times 10 ml), 5% NaHCO_3 (2 \times 10 ml) and water (2 \times 10 ml). After drying and removal of dichloromethane, the mixture of reaction products was separated by column chromatography (15 g silica gel, chloroform-acetone 1:1), yielding pure compound **3** (0.138 g, 42%, m.p. 248–250 °C after recrystallization from dichloromethane-hexane).

IR (KBr, cm^{-1}): 3416, 2940, 2868, 1637, 1560, 1489, 1438, 1377, 1215, 1060, 1034, 771; ^1H NMR (250 MHz, CDCl_3): δ 0.87 (s, 3H, H-18), 1.09 (s, 3H, H-19), 2.56 (d, 1H, $J_{\text{gem}} = 13.4$ Hz, CH_2Py), 2.93 (d, 1H, $J_{6,7} = 4.3$ Hz, H-6 β), 3.79 (d, 1H, $J_{\text{gem}} = 13.4$ Hz, CH_2Py), 3.89 (m, 1H, H-3), 6.65 (s, 1H, 17 β -OH \cdots O \leftarrow N-Py), 7.22–7.39 (m, 3H, H-3', H-4', H-5', Py), 8.28 (d, 1H, $J_{5',6'} = 6.1$ Hz, H-6', Py); ^{13}C NMR (62.5 MHz, CDCl_3): δ 14.12 (C-18), 15.96 (C-19), 20.42, 23.04, 28.47, 30.82 (CH), 31.05, 32.52, 33.74, 34.93, 38.44, 39.74, 42.87 (CH), 47.02, 51.45 (CH), 59.15 (CH), 65.98 (C-5), 68.42 (C-6), 69.16 (C-3), 85.18 (C-17), 124.28 (CH), 127.74 (CH), 128.97 (CH), 139.95 (CH), 150.76 (C-2', Py); GC/MS (m/e , %): 304 ([M- $\text{CH}_2\text{Py-O}$] $^+$, 100), 286 ([M- $\text{CH}_2\text{Py-O-H}_2\text{O}$] $^+$, 29), 271 (24), 253 (28), 227 (16); HRMS (TOF), m/z : for $\text{C}_{25}\text{H}_{35}\text{NO}_4$ [M + H] $^+$ calcd. 414.26389, found 414.26441.

4.1.2. 17 β -Hydroxy-17 α -picolyl-androst-4-ene-3,6-dione (**4**)

Jones reagent [14] (2 ml) was added dropwise (\sim 1 drop/10 s) to a solution of compound **1** (0.352 g, 0.923 mmol) in acetone (35 ml) at 0 °C, with vigorous stirring, and the reaction mixture was stirred for 1.5 h. The resulting reaction mixture was then quenched with methanol (15 ml, to eliminate any excess reagent) and solvent was removed *in vacuo*. Water (20 ml) and NaHCO_3 (to pH 8) were then added, and crude product was extracted with dichloromethane (5 \times 10 ml). Finally, purification by column chromatography (15 g silica gel, toluene-ethyl acetate 7:1) afforded pure compound **4** (0.044 g, 12%, m.p. 243–245 °C after recrystallization from dichloromethane-hexane).

IR (KBr, cm^{-1}): 3299, 3063, 2923, 2857, 1726, 1693, 1596, 1569, 1440, 1378, 1264, 1119, 1055, 1022, 869; ^1H NMR (250 MHz, CDCl_3): δ 1.01 (s, 3H, H-18), 1.20 (s, 3H, H-19), 2.80 (d, 1H, $J_{\text{gem}} = 14.5$ Hz, CH_2Py), 3.09 (d, 1H, $J_{\text{gem}} = 14.5$ Hz, CH_2Py), 6.19 (s, 1H, H-4), 7.18 (m, 2H, H-3' and H-5', Py), 7.65 (td, 1H, $J_{3',4'} = J_{4',5'} = 7.7$ Hz, $J_{4',6'} = 1.7$ Hz, H-4', Py), 8.47 (d, 1H, $J_{6',5'} = 4.4$ Hz, H-6', Py); ^{13}C NMR (62.5 MHz, CDCl_3): δ 14.08 (C-18), 17.50 (C-19), 18.72, 20.56, 23.58, 31.64, 33.95, 34.84, 35.55, 35.73, 39.75, 42.94, 46.55, 50.94, 50.99, 83.06 (C-17), 121.53 (C-5', Py), 124.68 (C-3', Py), 125.51 (C-4), 136.93 (C-4', Py), 148.02 (C-6', Py), 160.27 (C-2', Py), 160.82 (C-5), 199.46 (C-3), 201.95 (C-6); HRMS (TOF), m/z : for $\text{C}_{25}\text{H}_{32}\text{NO}_3$ [M + H] $^+$ calcd. 394.23767, found 394.23581.

4.1.3. 17 α -Picolyl-androst-5-en-3 β ,4 α ,17 β -triol (**5**) and 17 α -picolyl-androst-5-en-3 β ,4 β ,17 β -triol (**6**)

Selene-dioxide (0.089 g, 0.81 mmol) was added to a solution of compound **1** (0.200 g, 0.52 mmol) dissolved in dioxane (4 ml), and the reaction mixture was stirred under reflux for 25.5 h. The resulting mixture was then filtered, and the precipitate washed with dichloromethane (10 ml) and ethyl acetate (10 ml). Water (10 ml) was then added, and the mother liquor was extracted with dichloromethane (5 \times 10 ml). After drying and solvent removal, crude products were separated by column chromatography (10 g silica gel, hexane-ethyl acetate 2:1 and 1:2), affording pure compound **6** (0.045 g, 22%) and compound **5** (0.058 g, 28%), both in the form of a colorless oil.

Compound **5**: IR (film, cm^{-1}): 3353, 2926, 2854, 1665, 1596, 1570, 1438, 1377, 1260, 1216, 1053, 1080, 955, 757; ^1H NMR (250 MHz, CDCl_3): δ 0.98 (s, 3H, H-18), 1.04 (s, 3H, H-19), 2.88 (d, 1H, $J_{\text{gem}} = 14.5$ Hz, CH_2Py), 3.10 (d, 1H, $J_{\text{gem}} = 14.5$ Hz, CH_2Py), 3.60 (m, 1H, H-3), 3.89 (t, 1H, $J = 3.8$ Hz, H-4), 5.64 (m, 1H, H-6), 7.18 (m, 2H, H-3' and H-5', Py), 7.63 (td, 1H, $J_{3',4'} = J_{4',5'} = 7.7$ Hz, $J_{4',6'} = 1.8$ Hz, H-4', Py), 8.46 (d, 1H, $J_{5',6'} = 4.1$ Hz, H-6', Py); ^{13}C NMR (62.5 MHz, CDCl_3): δ 13.90 (C-18), 18.27 (C-19), 20.47, 21.47, 23.99, 29.69, 31.38, 31.78, 36.07, 37.09, 38.45, 42.00, 42.47, 43.01, 46.09, 65.06 (C-4), 71.35 (C-3), 83.52 (C-17), 121.35 (C-5', Py), 123.81 (C-3', Py), 124.86 (C-6), 136.79 (C-4', Py), 146.44 (C-5), 147.90 (C-6', Py), 160.83 (C-2', Py); GC/MS (m/e , %): 271 ([M- $\text{CH}_2\text{Py-2OH}$] $^+$, 41), 260 (100), 253 (58); HRMS (TOF), m/z : for $\text{C}_{25}\text{H}_{36}\text{NO}_3$ [M + H] $^+$ calcd. 398.26897, found 398.26828.

Compound **6**: IR (film, cm^{-1}): 3362, 2926, 2855, 1661, 1597, 1438, 1379, 1250, 1052, 1027, 967, 757, 666; ^1H NMR (250 MHz, CDCl_3): δ 0.98 (s, 3H, H-18), 1.22 (s, 3H, H-19), 2.80 (d, 1H, $J_{\text{gem}} = 14.5$ Hz, CH_2Py), 3.08 (d, 1H, $J_{\text{gem}} = 14.5$ Hz, CH_2Py), 3.57 (m, 1H, H-3), 4.16 (d, 1H, $J = 3.2$ Hz, H-4), 5.71 (m, 1H, H-6), 7.16 (m, 2H, H-3' and H-5', Py), 7.64 (td, 1H, $J_{3',4'} = J_{4',5'} = 7.7$ Hz, $J_{4',6'} = 1.8$ Hz, H-4', Py), 8.46 (m, 1H, H-6', Py); ^{13}C NMR (62.5 MHz, CDCl_3): δ 14.11 (C-18), 20.27, 21.05 (C-19), 23.96, 25.41, 29.69, 32.13, 32.64, 36.13, 37.01, 38.14, 43.19, 46.34, 50.33, 51.31, 72.48 (C-3), 77.24 (C-4), 83.48 (C-17), 121.35 (C-5', Py), 124.74 (C-3', Py), 128.46 (C-6), 136.75 (C-4', Py), 142.94 (C-5), 148.01 (C-6', Py), 160.80 (C-2', Py); GC/MS (m/e , %): 304 ([M- CH_3Py] $^+$, 23), 286 ([M- $\text{CH}_3\text{Py-H}_2\text{O}$] $^+$, 45), 260 (100), 247 (75); HRMS (TOF), m/z : for $\text{C}_{25}\text{H}_{36}\text{NO}_3$ [M + H] $^+$ calcd. 398.26897, found 398.26769.

4.1.4. General procedure for 17 β -hydroxy-17 α -picolyl-androst-5-en-3 β -yl *p*-toluenesulfonylate (**7**) and 17-picolinylden-androst-5-en-3 β -yl *p*-toluenesulfonylate (**9**)

A solution of *p*-toluenesulfonylchloride (1.028 g, 5.41 mmol for compound **1** or 0.511 g, 2.69 mmol for compound **2**) was added to **1** (0.506 g, 1.33 mmol) or **2** (0.200 g, 0.55 mmol) dissolved in absolute pyridine (8 ml or 4 ml) at 0 °C; and the resulting mixture was stirred 45 h (for **1**) or 19 h (for **2**) at room temperature. Reactions were stopped by addition of ice, and poured into water (20 or 40 ml). Crude product (**7**) was precipitated and purified by column chromatography (30 g silica gel, toluene-ethyl acetate 20:1 and 15:1), giving pure compound **7** (0.140 g, 20%, m.p. 163–165 °C after recrystallization from dichloromethane-hexane). The reaction mixture for **9** was extracted with dichloromethane (8 \times 10 ml), and, after drying and solvent removal, crude product (**9**) was recrystallized from dichloromethane-hexane, giving pure compound **9** (0.145 g, 51%, m.p. 176–178 °C).

Compound **7**: IR (KBr, cm^{-1}): 3323, 2951, 2870, 1597, 1569, 1473, 1438, 1361, 1188, 1175, 1098, 938, 866, 755, 667; ^1H NMR (250 MHz, CDCl_3): δ 0.94 (s, 3H, H-18), 1.00 (s, 3H, H-19), 2.45 (s, 3H, CH_3 from OTs group), 2.79 (d, 1H, $J_{\text{gem}} = 14.5$ Hz, CH_2Py), 3.06 (d, 1H, $J_{\text{gem}} = 14.5$ Hz, CH_2Py), 4.36 (m, 1H, H-3), 5.32 (d, 1H, $J = 4.9$ Hz, H-6), 7.16 (m, 2H, H-3' and H-5', Py), 7.34 (d, 2H from OTs group), 7.63 (td,

^1H , $J_{3',4'} = J_{4',5'} = 7.8$ Hz, $J_{4',6'} = 1.7$ Hz, H-4', Py), 7.80 (d, 2H from OTs group), 8.46 (d, 1H, $J_{5',6'} = 5.6$ Hz, H-6', Py); ^{13}C NMR (62.5 MHz, CDCl_3): δ 14.05 (C-18), 19.13 (C-19), 20.67, 21.62 (CH_3 , OTs), 23.93, 28.59, 31.75, 32.05, 32.55, 35.94, 36.45, 36.93, 38.84, 43.11, 46.26, 49.99, 50.94, 82.27 (C-3), 83.42 (C-17), 121.34 (C-5', Py), 123.23 (C-6), 124.73 (C-3', Py), 127.57 (C-2' and C-6', OTs), 129.72 (C-3' and C-5', OTs), 134.71 (C-4', OTs), 136.76 (C-4', Py), 138.95 (C-5), 144.38 (C-1', OTs), 147.95 (C-6', Py), 160.72 (C-2', Py); HRMS (TOF), m/z : for $\text{C}_{32}\text{H}_{42}\text{NSO}_4$ $[\text{M} + \text{H}]^+$ calcd. 536.28291, found 536.28358.

Compound **9**: IR (film, cm^{-1}): 2945, 1653, 1586, 1562, 1467, 1432, 1362, 1175, 1098, 938, 865, 753. ^1H NMR (250 MHz, CDCl_3): δ 0.88 (s, 3H, H-18), 1.01 (s, 3H, H-19), 2.44 (s, 3H, CH_3 from OTs), 4.32 (m, 1H, H-3), 5.32 (d, 1H, $J = 5.0$ Hz, H-6), 6.26 (s, 1H, H-20), 7.07 (m, 2H, H-3' and H-5', Py), 7.33 (d, 2H from OTs group), 7.65 (t, 1H, $J_{4',3'} = J_{4',5'} = 9.2$ Hz, H-4', Py), 7.79 (d, 2H from OTs group), 8.58 (m, 1H, H-6', Py); ^{13}C NMR (62.5 MHz, CDCl_3): δ 18.50 (C-18), 19.14 (C-19), 20.94, 21.58 (CH_3 , OTs), 24.94, 28.51, 29.81, 31.48, 31.61, 35.54, 36.40, 36.81 (C-10), 38.81, 45.79 (C-13), 50.09, 53.74 (C-9), 82.17 (C-3), 117.33 (C-20), 120.39 (C-5', Py), 123.16 (C-6), 126.01 (C-3', Py), 127.57 (C-2' and C-6', OTs), 129.70 (C-3' and C-5', OTs), 134.53 (C-4', OTs), 136.55 (C-4', Py), 138.93 (C-5), 144.38 (C-1', OTs), 148.39 (C-6', Py), 156.93 (C-17), 161.24 (C-2', Py).

4.1.5. General procedure for 17 α -picolyl-androsta-3,5-dien-17 β -ol (**8**) and 17-picolinylden-androsta-3,5-diene (**10**)

Compound **7** (0.080 g, 0.149 mmol) or compound **9** (0.379 g, 0.733 mmol) were stirred with bis(2-aminoethyl)amine (4.5 ml, or 20 ml) under reflux for 1 h, cooled to room temperature, poured into water, and extracted with dichloromethane (5×10 ml). The organic phase was dried, filtered and evaporated. Crude product was then purified by column chromatography (1 g silica gel, toluene-ethyl acetate 30:1 for **8**, or 6 g silica gel, toluene-ethyl acetate 30:1 for **10**), affording pure compound **8** (0.008 g, 15%) and pure compound **10** (0.018 g, 7%), both in the form of a colorless oil.

Compound **8**: IR (film, cm^{-1}): 3328, 3017, 2926, 2855, 1652, 1596, 1570, 1438, 1377, 1024, 846, 753; ^1H NMR (250 MHz, CDCl_3): δ 0.99 (s, 3H, H-18), 1.26 (s, 3H, H-19), 2.82 (d, 1H, $J_{\text{gem}} = 14.6$ Hz, CH_2Py), 3.09 (d, 1H, $J_{\text{gem}} = 14.6$ Hz, CH_2Py), 5.41 (m, 1H, H-3), 5.63 (m, 1H, H-6), 5.94 (d, 1H, $J = 9.9$ Hz, H-4), 7.16 (m, 2H, H-3' and H-5', Py), 7.63 (td, 1H, $J_{4',5'} = J_{4',3'} = 7.7$ Hz, $J_{4',6'} = 1.8$ Hz, H-4', Py), 8.47 (m, 1H, H-6', Py); ^{13}C NMR (62.5 MHz, CDCl_3): δ 14.21 (C-18), 18.77 (C-19), 20.66, 23.03, 23.89, 29.68, 31.68, 32.20, 32.56, 33.83, 36.04, 43.22, 46.46, 48.53, 51.33, 83.48 (C-17), 121.31 (C-5', Py), 122.80 (C-6), 124.74 (C-3', Py), 125.14 (C-4), 128.90 (C-3), 136.72 (C-4', Py), 141.56 (C-5), 147.99 (C-6', Py), 160.88 (C-2', Py); HRMS (TOF), m/z : for $\text{C}_{25}\text{H}_{34}\text{NO}$ $[\text{M} + \text{H}]^+$ calcd. 364.26349, found 364.26255.

Compound **10**: IR (film, cm^{-1}): 3016, 2963, 2851, 1652, 1586, 1560, 1470, 1428, 1372, 1149, 857, 756; ^1H NMR (250 MHz, CDCl_3): δ 0.94 (s, 3H, H-18), 1.01 (s, 3H, H-19), 5.42 (m, 1H, H-3), 5.61 (m, 1H, H-6), 5.95 (d, 1H, $J_{4,3} = 9.7$ Hz, H-4), 6.25 (t, 1H, $J = 2.4$ Hz, H-20), 7.03 (m, 1H, H-5', Py), 7.29 (m, 1H, H-3', Py), 7.61 (td, 1H, $J_{4',5'} = J_{4',3'} = 7.8$ Hz, $J_{4',6'} = 1.8$ Hz, H-4', Py), 8.57 (d, 1H, $J_{6',5'} = 4.8$ Hz, H-6', Py); ^{13}C NMR (62.5 MHz, CDCl_3): δ 18.79 (C-18), 18.83 (C-19), 21.01, 23.03, 24.98, 29.87, 31.62, 31.66, 33.77, 35.32, 35.82, 45.90, 48.67, 54.17, 118.02 (C-20), 120.17 (C-5', Py), 122.77 (C-3', Py), 124.72 (C-6), 125.16 (C-4), 128.89 (C-3), 135.80 (C-4', Py), 141.57 (C-5), 149.16 (C-6', Py), 157.65 (C-17), 160.44 (C-2', Py); HRMS (TOF), m/z : for $\text{C}_{25}\text{H}_{32}\text{N}$ $[\text{M} + \text{H}]^+$ calcd. 346.25293, found 346.25184.

4.1.6. General procedure for 17 β -hydroxy-4-methoxy-17 α -picolyl-androst-4-en-3-one (**15**) and 4-methoxy-17-picolinylden-androst-4-en-3-one (**18**)

A solution of 4 M NaOH (2 ml or 0.6 ml) was added to a solution of a mixture of compounds **13** and **14** [9] (0.175 g,

0.44 mmol) or compounds **16** and **17** [9] (0.056, 0.14 mmol) in methanol (22 ml or 7 ml), and the reaction mixture was stirred under reflux for 3.5 h (or 3 h). After reaction completion, the resulting reaction mixture was extracted with dichloromethane (5×10 ml or 5×5 ml), solvent was dried and removed, and crude product was separated by column chromatography (5 g or 1 g silica gel). Elution with hexane-ethyl acetate 3:1 (for **15**) or toluene-ethyl acetate 9:1 (for **18**) afforded pure compound **15** (0.007 g, 4%), or pure compound **18** (0.006 g, 10%), both in the form of a colorless oil.

Compound **15**: IR (film, cm^{-1}): 3391, 2929, 1681, 1597, 1570, 1439, 1379, 1317, 1260, 1214, 1086, 1022, 927, 891, 800, 756; ^1H NMR (250 MHz, CDCl_3): δ 1.00 (s, 3H, H-18), 1.22 (s, 3H, H-19), 2.80 (d, 1H, $J_{\text{gem}} = 14.3$ Hz, CH_2Py), 3.07 (d, 1H, $J_{\text{gem}} = 14.3$ Hz, CH_2Py), 3.60 (s, 3H, from 4-OCH₃), 4.72 (bs, 1H, 17 β -OH), 7.18 (m, 2H, H-3' and H-5', Py), 7.63 (td, 1H, $J_{4',5'} = J_{4',3'} = 7.6$ Hz, $J_{4',6'} = 1.7$ Hz, H-4', Py), 8.46 (d, 1H, $J_{6',5'} = 5.1$ Hz, H-6', Py); ^{13}C NMR (62.5 MHz, CDCl_3): δ 14.15 (C-18), 17.50 (C-19), 20.69, 23.41, 23.79, 29.66, 31.90, 34.25, 34.84, 35.91 (C-10), 36.01, 38.78, 43.02, 44.23 (C-13), 50.20 (C-14), 54.33 (C-9), 60.23 (O-CH₃), 83.34 (C-17), 121.43 (C-5', Py), 124.75 (C-3', Py), 136.88 (C-4', Py), 146.21 (C-5), 147.23 (C-4), 147.91 (C-6', Py), 160.57 (C-2', Py), 191.39 (C-3); GC/MS (m/e , %): 316 ($[\text{M}-\text{CH}_3\text{Py}]^+$, 59), 284 ($[\text{M}-\text{CH}_3\text{Py}-\text{CH}_3\text{OH}]^+$, 26), 273 (100); HRMS (TOF), m/z : for $\text{C}_{26}\text{H}_{36}\text{NO}_3$ $[\text{M} + \text{H}]^+$ calcd. 410.26897, found 410.26718.

Compound **18**: IR (film, cm^{-1}): 2924, 1682, 1607, 1586, 1561, 1470, 1435, 1359, 1334, 1314, 1207, 1150, 1095, 1066, 1035, 776, 732. ^1H NMR (250 MHz, CDCl_3): δ 0.93 (s, 3H, H-18), 1.25 (s, 3H, H-19), 3.59 (s, 3H, from 4-OCH₃), 6.23 (s, 1H, H-20), 7.04 (m, 1H, H-5', Py), 7.27 (m, 1H, H-3', Py), 7.62 (td, 1H, $J_{4',5'} = J_{4',3'} = 7.8$ Hz, $J_{4',6'} = 1.8$ Hz, H-4', Py), 8.57 (d, 1H, $J_{6',5'} = 4.7$ Hz, H-6', Py); ^{13}C NMR (62.5 MHz, CDCl_3): δ 17.52 (C-18), 18.80 (C-19), 21.06, 23.40, 24.92, 29.74, 31.40, 34.24, 34.81, 35.14, 35.70 (C-10), 38.80, 45.72 (C-13), 53.15 (C-14), 54.43 (C-9), 60.32 (O-CH₃), 118.14 (C-20), 120.30 (C-5', Py), 122.82 (C-3', Py), 135.89 (C-4', Py), 146.66 (C-5), 149.16 (C-6', Py), 154.96 (C-4), 157.46 (C-17), 159.82 (C-2', Py), 194.14 (C-3); GC/MS (m/e , %): 391 (M^+ , 100), 377 (59), 348 (6), 170 (7), 130 (20), 93 (15); HRMS (TOF), m/z : for $\text{C}_{26}\text{H}_{34}\text{NO}_2$ $[\text{M} + \text{H}]^+$ calcd. 392.25841, found 392.25724.

4.1.7. 4,17 β -Dihydroxy-17 α -picolyl-androsta-4,6-dien-3-one (**19**) and 17 β -hydroxy-17 α -picolyl-androst-4-en-3,6-dione (**4**)

1 M potassium *tert*-butoxide (2.5 ml) was added dropwise to a solution of compound **11** (0.202 g, 0.54 mmol) dissolved in *tert*-butyl alcohol (6 ml) at room temperature, and the reaction mixture was stirred at 45 °C for 3 h (protected from light). When the reaction was complete, the resulting mixture was poured into water (10 ml) and extracted with ethyl acetate (5×10 ml), followed by dichloromethane (5×10 ml). Combined extracts were then dried and solvent removed. The resulting crude product was separated and purified by column chromatography (20 g silica gel, toluene-ethyl acetate 8:1 and 5:1), giving compound **19** (0.052 g, 25%, m.p. 147–149 °C after recrystallization from methanol) and compound **4** (0.028 g, 13%, m.p. 243–245 °C after recrystallization from methanol).

Compound **19**: IR (KBr, cm^{-1}): 3423, 3054, 2941, 2872, 1655, 1616, 1597, 1570, 1380, 1265, 894; ^1H NMR (250 MHz, CDCl_3): δ 1.04 (s, 3H, H-18), 1.12 (s, 3H, H-19), 2.80 (d, 1H, $J_{\text{gem}} = 14.5$ Hz, CH_2Py), 3.10 (d, 1H, $J_{\text{gem}} = 14.5$ Hz, CH_2Py), 6.06 (dd, 1H, $J_{6,7} = 9.4$ Hz, $J_{7,8} = 2.0$ Hz, H-7), 6.65 (dd, 1H, $J_{6,7} = 9.4$ Hz, $J_{6,8} = 2.8$ Hz, H-6), 7.18 (m, 2H, H-3' and H-5', Py), 7.66 (td, 1H, $J_{3',4'} = J_{4',5'} = 7.6$ Hz, $J_{4',6'} = 1.8$ Hz, H-4', Py), 8.47 (d, 1H, $J_{5',6'} = 4.0$ Hz, H-6', Py); ^{13}C NMR (62.5 MHz, CDCl_3): δ 14.08 (C-18), 16.44 (C-19), 20.27, 23.43, 29.70, 31.93, 33.29, 35.75, 36.48, 37.93, 42.98, 46.87, 48.04, 51.44, 83.17 (C-17), 121.55 (C-5', Py), 123.94 (C-3', Py), 124.71 (C-6), 134.70 (C-5),

136.76 (C-4', Py), 137.10 (C-7), 140.21 (C-4), 147.83 (C-6', Py), 160.39 (C-2', Py), 193.77 (C-3); HRMS (TOF), m/z : for $C_{25}H_{32}NO_3$ $[M + H]^+$ calcd. 394.23767, found 394.24972.

4.1.8. 17 α -Picolyl-androst-4-en-3 β ,17 β -diol (**22**)

$NaBH_4$ (0.142 g, 3.74 mmol) was added to a solution of compound **11** (0.203 g, 0.535 mmol) dissolved in absolute ethanol (15 ml), and the reaction mixture was stirred at room temperature for 2.5 h. When the reaction was complete, the resulting mixture was poured into water (20 ml), and crude product was filtered and recrystallized from dichloromethane-methanol, to give pure compound **22** (0.191 g, 94%, m.p. 96–98 °C) as white crystals.

Compound **22**: IR (KBr, cm^{-1}): 3417, 2934, 2853, 1634, 1597, 1570, 1440, 1380, 1327, 1230, 1193, 1101, 1056, 1031, 860, 760, 657, 588; 1H NMR (250 MHz, $CDCl_3$): δ 0.97 (s, 3H, H-18), 1.08 (s, 3H, H-19), 2.78 (d, 1H, $J_{gem} = 14.5$ Hz, CH_2Py), 3.07 (d, 1H, $J_{gem} = 14.5$ Hz, CH_2Py), 4.15 (m, 1H, H-3), 5.30 (s, 1H, H-4), 7.17 (m, 2H, H-3' and H-5', Py), 7.64 (td, 1H, $J_{4,3'} = J_{4,5'} = 7.7$ Hz, $J_{4,6'} = 1.8$ Hz, H-4', Py), 8.45 (d, 1H, $J_{6,5'} = 4.4$ Hz, H-6', Py); ^{13}C NMR (62.5 MHz, $CDCl_3$): δ 14.16 (C-18), 18.92 (C-19), 20.68 (C-11), 23.84, 29.50 (C-2), 32.14 (C-7), 32.17, 32.92 (C-1), 35.43, 35.88, 36.65 (C-8), 37.42 (C-10), 43.11 (C-20), 46.28 (C-13), 50.37 (C-14), 54.60 (C-9), 67.93 (C-3), 83.41 (C-17), 121.32 (C-5', Py), 123.41 (C-4), 124.71 (C-3', Py), 136.74 (C-4', Py), 147.90 (C-5), 147.96 (C-6', Py), 160.77 (C-2', Py); HRMS (TOF), m/z : for $C_{25}H_{36}NO_2$ $[M + H]^+$ calcd. 382.27406, found 382.27769.

4.1.9. 17 β -Hydroxy-17 α -picolyl-androst-4-en-3 β -yl acetate (**23**)

Acetic anhydride (1.5 ml) was added to a solution of compound **22** (0.049 g, 0.13 mmol) in absolute pyridine (1.5 ml) and the reaction mixture was stirred at room temperature for 24 h. The resulting reaction mixture was slowly poured into cold water (10 ml), yielding a precipitate, which was filtered and recrystallized from dichloromethane-hexane, affording pure compound **23** (0.031 g, 56%, m.p. 179–181 °C) as white crystals.

Compound **23**: IR (KBr, cm^{-1}): 3437, 2942, 2852, 1730, 1598, 1570, 1441, 1371, 1240, 1097, 1023, 970, 927, 901, 760, 675, 612; 1H NMR (250 MHz, $CDCl_3$): δ 0.97 (s, 3H, H-18), 1.09 (s, 3H, H-19), 2.06 (s, 3H, CH_3 from OAc), 2.78 (d, 1H, $J_{gem} = 14.6$ Hz, CH_2Py), 3.05 (d, 1H, $J_{gem} = 14.6$ Hz, CH_2Py), 5.24 (m, 1H, H-3), 5.30 (m, 1H, H-4), 6.65 (bs, 1H, 17 β -OH \cdots N-Py), 7.15 (m, 2H, H-3' and H-5', Py), 7.63 (td, 1H, $J_{4,3'} = J_{4,5'} = 7.8$ Hz, $J_{4,6'} = 1.6$ Hz, H-4', Py), 8.45 (d, 1H, $J_{6,5'} = 5.2$ Hz, H-6', Py); ^{13}C NMR (62.5 MHz, $CDCl_3$): δ 14.15 (C-18), 18.84 (C-19), 20.65, 21.45 (CH_3 from OAc), 23.86, 25.08, 32.18, 32.77, 35.04, 35.94, 36.62, 37.42, 43.13, 46.43, 50.32, 54.38, 70.88 (C-3), 83.37 (C-17), 119.04 (C-4), 121.32 (C-5', Py), 124.73 (C-3', Py), 136.75 (C-4', Py), 147.98 (C-5), 149.48 (C-6', Py), 160.82 (C-2', Py), 171.04 (C=O from OAc); HRMS (TOF), m/z : for $C_{27}H_{38}NO_3$ $[M + H]^+$ calcd. 424.28462, found 424.28428.

4.2. Cytotoxicity

Cell lines. Five human tumor cell lines and one human non-tumor cell line were used in the present study: human breast adenocarcinoma ER+, MCF-7; human breast adenocarcinoma ER-, MDA-MB-231; prostate cancer AR-, PC-3; human cervix carcinoma, HeLa; colon cancer, HT-29; and normal fetal lung fibroblasts, MRC-5.

Cells were grown in Dulbecco's modified Eagle's medium (DMEM) with addition of 4.5% glucose. Media were supplemented with 10% fetal calf serum (FCS, NIVNS) and the following antibiotics: 100 IU/ml of penicillin and 100 μ g/ml of streptomycin (ICN Galenika). All cell lines were cultured in flasks (Costar, 25 cm^2) at 37 °C in a 100% humidity atmosphere with 5% CO_2 . Only viable cells

were used in the assay. Viability was determined by the dye exclusion assay with trypan blue.

Cytotoxicity assay. Cytotoxic activity was evaluated using the colorimetric sulforhodamine B (SRB) assay, following the method of Skehan et al. [18]. Briefly, a single cell suspension was plated onto 96-well microtiter plates (Costar, flat bottom): 5×10^3 cells/well in 180 μ l of medium. Plates were pre-incubated 24 h at 37 °C, 5% CO_2 . Tested substances were then added to all wells (except controls), at concentrations ranging from 10^{-8} to 10^{-4} M. After incubation (48 h/37 °C/5% CO_2), the SRB assay was carried out as follows: 50 μ l of 50% trichloroacetic acid (TCA) was added to all wells and plates were incubated for 1 h; plates were washed with distilled water, 75 μ l of 0.4% SRB was then added to all wells and plates were incubated for 30 min; plates were then washed with citric acid (1%) and dried at room temperature. Finally, 200 μ l of 10 mmol TRIS (pH 10.5) was added to all wells, and absorbance (A) was measured at 540/690 nm on a microplate reader (Multiscan MCC340, Labsystems). Wells without cells, but containing complete medium only, acted as blanks. Cytotoxic activity was calculated according to the formula:

$$1 - A_{TEST}/A_{CONTROL} \times 100$$

and expressed as a percent of cytotoxic activity (CI %).

Data analysis. Two independent experiments were conducted in quadruplicate for each concentration of tested compound. Mean values and standard deviations (SD) was calculated for each concentration. IC_{50} value defines the dose of compound that inhibits cell growth by 50%. The IC_{50} of each tested compound was determined by median effect analysis [19].

Acknowledgments

We would like to thank the Ministry of Education and Science of the Republic of Serbia for financial support (Grant No. 172021).

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.ejmech.2012.06.030>.

References

- [1] R.K. Dubey, S. Oparil, B. Imthurn, E.K. Jackson, Sex hormones and hypertension, *Cardiovasc. Res.* 53 (2002) 688–708.
- [2] (a) P.J. Sheridan, K. Blum, M. Trachtenberg, *Steroid Receptors and Disease*, Marcel Dekker, New York, 1988, pp. 289–564; (b) V.K. Moudgil, *Steroid Receptors in Health and Disease*, Plenum Press, New York/London, 1987.
- [3] G. Michels, U.C. Hoppe, Rapid actions of androgens, *Front. Neuroendocrinol* 29 (2008) 182–198.
- [4] T.V. Nguyen, M. Yao, C.J. Pike, Flutamide and Ciproterone acetate exert agonist effects: introduction of androgen receptor-dependent neuroprotection, *Endocrinology* 148 (2007) 2936–2943.
- [5] D. Gillatt, Antiandrogen treatments in locally advanced prostate cancer: are they all the same, *J. Cancer Res. Clin. Oncol.* 132 (2006) 17–26.
- [6] S. Aggarwal, S. Thareja, A. Verma, T.R. Bhardwaj, M. Kumar, An overview on 5 α -reductase inhibitors, *Steroids* 75 (2010) 109–153.
- [7] S.B. Karrothu, N. Kolla, A.R. Lekkala, C.M. Golla, R.V. Anand, V. Gandu, A. Bhattacharya, P.R. Padi, R. Bandichhor, Effect of solvent on Stereoselectivity in Pd/C (Type 39K)-Catalyzed Hydrogenation of methyl 3-oxo-4-aza-5-androstene-17-carboxylate, a key Intermediate for Finasteride and Dutasteride, *Org. Process Res. Dev.* 11 (2007) 889–891.
- [8] K.M. Penov-Gaši, M.Dj. Djurendić-Brenesel, E.A. Djurendić, M.N. Sakač, J.J. Čanadi, J.J. Daljev, T. Armbruster, S. Andrić, D.M. Sladić, T.T. Božić, I.T. Novaković, Z.D. Juranic, Synthesis and biological evaluation of some 17-picolyl and 17-picolinylidene androst-5-ene derivatives, *Steroids* 72 (2007) 31–40.
- [9] E. Djurendić, J. Daljev, M. Sakač, J. Čanadi, S. Jovanović-Šanta, S. Andrić, O. Klisurić, V. Kojić, G. Bogdanović, M. Djurendić-Brenesel, S. Novaković, K. Penov-Gaši, Synthesis of some epoxy and/or N-oxy 17-picolyl and

- 17-picolinylidene-androst-5-ene derivatives and evaluation of their biological activity, *Steroids* 73 (2008) 129–138.
- [10] E.A. Djurendić, J.J. Ajduković, M.N. Sakač, J.J. Čanadi, V.V. Kojić, G.M. Bogdanović, K.M. Penov Gaši, 17-Picolinylidene-substituted steroid derivatives and their antiaromatase and cytotoxic activity, *Arkivoc* 13 (2009) 311–323.
- [11] A.M.H. Brodie, V.C.O. Njar, Aromatase inhibitors and their application in breast cancer treatment, *Steroids* 65 (2000) 171–179.
- [12] D. Miljković, K. Gaši, M. Kindjer, S. Stanković, B. Ribár, G. Argay, Partial Syntheses of 21,27-Bisnordemissidine from Epiandrosterone and Dehydroepiandrosterone Acetates. Crystal and molecular structure of 21,27-Bisnordemissidine Hydrobromide, *Croat. Chem. Acta* 58 (1985) 721–736.
- [13] D. Miljković, K. Gaši, A novel synthetic approach to 3 β -acetoxy-17-picolinylidene-5-androstene-16-one, *Bull. Soc. Chim. Beograd* 47 (1982) 173–181.
- [14] A.C. Hunter, S.M. Priest, An efficient one-pot synthesis generating 4-ene-3,6-dione functionalized steroids from steroidal 5-ene-3 β -ols using a modified Jones oxidation methodology, *Steroids* 71 (2006) 30–33.
- [15] E. Ma, T. Choi, An efficient 4 β -hydroxylation of steroidal 5-en-3 β -ols and 1,4-conjugation of steroidal 4-en-3-ones using SeO₂ oxidation, *Bull. Korean Chem. Soc.* 30 (2009) 245–248.
- [16] L.C. King, M.J. Bigelow, Preparation, reactions and kinetics of reactions of epicholesteryl p-toluenesulfonate, *J. Am. Chem. Soc.* 74 (1952) 6238–6241.
- [17] M. Numazawa, A. Mutsumi, M. Tachibana, K. Hoshi, Synthesis of androst-5-en-7-ones and androsta-3,5-dien-7-ones and their related 7-deoxy analogs as conformational and catalytic probes for the active site of aromatase, *J. Med. Chem.* 37 (1994) 2198–2205.
- [18] P. Skehan, R. Storeng, D. Scudiero, A. Monks, J. McMahon, D. Visitica, T.J. Warren, H. Bokesch, S. Kenny, R.M. Boyd, New colorimetric anti-proliferative activity assay for anticancer-drug screening, *J. Natl. Cancer Inst.* 82 (1990) 1107–1112.
- [19] A. Sotto, V. Foulongne, D. Sirot, R. Labia, J. Jourdan, Experimental application of the median-effect principle for in vitro quantification of the combined inhibitory activities of clavulanic acid and imipenem against IRT-4 β -lactamase, *Int. J. Antimicrob. Agents* 19 (2002) 75–78.