

Review Article

A lipidologist perspective of global lipid guidelines and recommendations, part 2: Lipid treatment goals

Harold E. Bays, MD, FACC, FTOS, FACE, FNLA*

Louisville Metabolic and Atherosclerosis Research Center Inc, Louisville, KY, USA

KEYWORDS:

Low-density lipoprotein cholesterol;
Guidelines;
Lipid goals;
Recommendations;
Atherosclerotic cardiovascular disease risk

Abstract: Having knowledge of worldwide areas of harmonization and consensus regarding lipid guidelines and recommendations may provide clinicians a more global perspective on lipid management. This review examines 8 international scientific/medical organizations that have issued lipid guidelines, recommendations, and position papers: the National Lipid Association (2014), National Institute for Health and Care Excellence (2014), International Atherosclerosis Society (2013), American College of Cardiology/American Heart Association (2013), Canadian Cardiovascular Society (2013), Japan Atherosclerosis Society (2012), European Society of Cardiology/European Atherosclerosis Society (2012), and Adult Treatment Panel III (2001/2004). The focus of part 2 of this perspective was to review sentinel components of lipid guidelines and recommendations as applied to goals of lipid-altering therapy to reduce atherosclerotic cardiovascular disease (ASCVD) events. In general, lipid guidelines and recommendations have significant concordance regarding the need to reduce atherogenic lipoprotein cholesterol levels and are in general agreement on the primary lipid treatment targets. Some guidelines and recommendations differ with regard to ASCVD risk assessment and lipid treatment goals. Despite these differences, substantial agreement exists among lipid guidelines and recommendations in their emphasis on the need for aggressive treatment of hypercholesterolemia, for which the predominance of ASCVD outcomes studies suggests statins as currently the first-line treatment of choice.

© 2015 National Lipid Association. All rights reserved.

Introduction

A brief history of the emergence of lipid guidelines and recommendations was discussed in part 1 of this perspective, which examined the role of atherogenic lipoprotein cholesterol levels, primary lipid and lipoprotein targets of therapy, and other primary/secondary lipid treatment

targets, as well as assessment of atherosclerotic cardiovascular disease (ASCVD) risk. The purpose of Part 2 of this review was to summarize 8 worldwide lipid guidelines, recommendations, and position papers with respect to the goal of lipid-altering therapy in reducing ASCVD risk.

Do lipid-altering therapeutic agents reduce atherosclerotic cardiovascular disease risk?

According to most global lipid guidelines and recommendations, low-density lipoprotein cholesterol (LDL-C) level is the primary lipid treatment target. The rationale for

* Corresponding author. Louisville Metabolic and Atherosclerosis Research Center Inc, 3288 Illinois Avenue, Louisville, KY 40213, USA.

E-mail address: hbaysmd@aol.com

Submitted April 13, 2015; revised September 14, 2015. Accepted for publication October 16, 2015.

establishing LDL-C treatment goals is 2 fold: increased LDL-C levels are a strong and independent risk predictor of coronary heart disease (CHD) and ASCVD,¹ and interventions that lower LDL-C levels often reduce CHD and ASCVD risk. Meta-analyses of randomized clinical trials of statins suggest that every 1.0-mmol/L (~40 mg/dL) reduction in LDL-C level is associated with an approximately 20% reduction in cardiovascular disease (CVD) mortality, nonfatal myocardial infarction (MI), and overall cardiovascular events.^{2,3} In the Cochrane Database review of 56,934 individuals enrolled in 18 statin trials for primary prevention of CVD, statin therapy reduced the relative risk (RR) of all-cause mortality by 14% and of combined fatal and nonfatal CVD, CHD, and stroke by 25%, 27%, and 22%, respectively.⁴ In an analysis by the Cholesterol Treatment Trialists' (CTT) Collaboration of 27 randomized clinical trials of individuals grouped by predicted 5-year risk of a major vascular event and treated with statin therapy, the risk of a vascular event for patients without a history of vascular disease decreased by 15% (rate ratio per 1.0-mmol/L reduction in LDL-C) and all-cause mortality decreased by 9% (rate ratio per 1.0-mmol/L reduction in LDL-C).³ The benefits of LDL-C lowering are generally consistent in both primary and secondary prevention and in different patient subpopulations.^{2,3,5}

A question often arises about whether the ASCVD benefits noted in lipid-altering pharmacotherapy trials are due to cholesterol lowering or to some other "pleiotropic" properties of statins. However, non-statin, lipid-altering drug therapies that lower cholesterol are also associated with reduced ASCVD risk. When administered to patients without elevated triglyceride (TG) levels, fibrates can lower LDL-C levels. A number of outcomes clinical trials support fibrates as reducing ASCVD, although their benefit appears to be predominantly among patients with more elevated baseline TG levels.⁶⁻¹⁰ Niacin is a lipid-altering agent that can lower LDL-C at higher doses. Recent ASCVD outcomes trials (AIM-HIGH, HPS2-THRIVE) have not supported niacin as reducing ASCVD risk in statin-treated patients with very low baseline LDL-C levels.^{11,12} However, when administered as monotherapy to patients with higher baseline LDL-C levels, data (eg, Coronary Drug Project trial) support niacin as reducing ASCVD events.¹³ Bile acid sequestrants reduce cholesterol levels, and resin therapies such as cholestyramine and colestipol reduce ASCVD.¹⁴ Finally, in patients at very high ASCVD risk with baseline LDL-C level of ~95 mg/dL, patients who attained an LDL-C level below 70 mg/dL (mean value ~53 mg/dL) with the addition of ezetimibe to simvastatin experienced significantly reduced ASCVD risk compared with patients who attained an LDL-C level of ~70 mg/dL with simvastatin alone.¹⁵

Thus, clinical trial evidence based on ASCVD outcomes trials supports the benefit of non-statin in reducing ASCVD risk either as monotherapy or sometimes when combined with statins, depending on the patient population studied. In fact, a meta-analysis of non-statin therapies (ie,

diet, bile acid sequestrants, ileal bypass surgery) demonstrated that the degree of LDL-C lowering correlates one-to-one with reduction of CHD risk over 5 years.¹⁶

Author perspective

The ASCVD benefits of LDL-C lowering are generally consistent in both primary and secondary prevention trials and in different patient subpopulations. However, therapeutic interventions that lower LDL-C levels do not always reduce ASCVD risk. Examples noted previously include the lack of benefit of niacin when added to statins in patients having low LDL-C levels. Also previously noted is that the ASCVD benefits of fibrates appear to be mostly among patients with baseline elevations in TG levels (and lower high-density lipoprotein cholesterol [HDL-C] levels), which is how fibrates are most often used in clinical practice. Estrogens lower LDL-C levels and increase HDL-C levels, and they have a number of other cardiovascular effects that may reduce CVD risk.¹⁷ However, the clinical trial evidence suggests that when hormone therapy (including estrogen) is administered to postmenopausal women, CHD and thromboembolic complications may be increased, not decreased.¹⁸ Torcetrapib was an investigational cholesteryl ester transfer protein inhibitor that lowered LDL-C and substantially increased HDL-C levels,¹⁹ but increased (not decreased) ASCVD risk, which may or may not have been due to agent-specific, off-target effects.²⁰ What has emerged from these experiences is that not all interventions that lower LDL-C levels will reduce ASCVD risk. Therapeutic agents that lower LDL-C levels are most likely to reduce ASCVD if the agent has the following: (1) natural genetic mutation support,²¹ (2) a validated mechanism of action, (3) a lack of off-target harmful effects that might increase ASCVD risk, and (4) favorable signaling in pooled data during phase 2 and 3 clinical trial development.

What are the potential risks and benefits of lowering LDL-C levels below 70 mg/dL?

Although many guidelines recommend lipid treatment goals of LDL-C < 70 mg/dL for patients who have the highest risk of ASCVD, the long-term clinical risks and benefits of achieving even lower levels of LDL-C < 50 mg/dL (<1.3-1.8 mmol/L) are unknown. However, data from aboriginal populations and patients with gene mutations having LDL-C levels between 30 and 70 mg/dL suggest that ASCVD mortality is very low in these subpopulations.²²⁻²⁴ In Treating to New Targets (TNT), patients with ASCVD were administered atorvastatin 10 or 80 mg per day. The lowest on-treatment LDL-C levels were associated with the lowest rate of death from any cause and the lowest rate of death from ASCVD. Achievement of the lowest LDL-C levels did not result in clinically important differences in adverse experiences (eg, no increase in muscle

complaints, suicide, hemorrhagic stroke, or cancer deaths).²⁵ In the Pravastatin or Atorvastatin Evaluation and Infection Therapy (PROVE-IT) study, atorvastatin 80 mg was compared with pravastatin 40 mg per day in patients with acute coronary syndrome.²⁶ Atorvastatin-treated patients with on-treatment LDL-C levels < 40 and 40 to 60 mg/dL had fewer major cardiac events (ie, death, MI, stroke, recurrent ischemia, revascularization); however, they did not have significant differences in safety parameters (eg, muscle, liver, or retinal abnormalities; increased risk of intracranial hemorrhage; or death). A post hoc analysis of the JUPITER trial suggested that among adults with baseline LDL-C < 130 mg/dL and high-sensitivity C-reactive protein \geq 2 mg/L, rosuvastatin-allocated patients who attained LDL-C levels < 50 mg/dL had lower CHD risk compared with those without LDL-C < 50 mg/dL, without evidence of an increase in reported adverse experiences.^{27,28} This JUPITER analysis in conjunction with the rosuvastatin ASTEROID trial supports the safety of lower LDL-C (<50 mg/dL) in terms of statin side effects and overall safety and tolerability.^{27–29}

A newer class of lipid-altering agents are proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitors, which may substantially lower LDL-C levels, especially when added to statins. When alirocumab (a fully human monoclonal antibody that inhibits PCSK9) was added to a statin in a 78-week trial of 2341 high ASCVD-risk patients with LDL-C \geq 70 mg/dL, 37% of those who received alirocumab achieved a calculated LDL-C level < 25 mg/dL at 2 consecutive measurements. The percentage of patients with any adverse event was similar in the 2 study groups: 81.0% with alirocumab and 82.5% with placebo, with no clinically meaningful changes in fat-soluble vitamins or cortisol.³⁰ Although this was primarily an efficacy/safety trial (ie, not an ASCVD outcomes trial), a post hoc analysis suggested a reduction in the rate of cardiovascular events with alirocumab.^{30,31} Two open-label studies evaluated 4465 patients receiving either the fully human PCSK9 inhibitor evolocumab or standard therapy for a median of 11 months.³² Compared with those with higher LDL-C levels, rates of adverse experiences, serious adverse events, and elevations in aminotransferase or creatine kinase levels were similar among patients in the evolocumab group who had LDL-C < 40 mg/dL or <25 mg/dL. Again, although this was not an ASCVD outcome study, a prespecified exploratory analysis suggested evidence of reduction in ASCVD among patients receiving evolocumab. Finally, in a study of 354 patients receiving the humanized PCSK9 inhibitor bococizumab for 24 weeks, those who received bococizumab and who achieved LDL-C levels \leq 25 mg/dL, the incidence of adverse experiences was similar to those who had higher LDL-C levels.³³

Author perspective

Patients with very low LDL-C levels due to genetic hypobetalipoproteinemia or abetalipoproteinemia (inherent

lack of apolipoprotein b containing lipoproteins) have malabsorption of fat-soluble vitamins and often develop retinal disease, neurologic disorders, or other adverse complications.³⁴ Conversely, likely because the applicable mechanisms are entirely different, rare isolated case reports of individuals with PCSK9 loss of function may have very low LDL-C (<15 mg/dL) but can be generally healthy.^{35,36} Having said this, having a PCSK9 genetic abnormality leading to lower LDL-C levels may not have the same clinical implications as therapeutically induced reductions in PCSK9. Also, although the short-term data described previously regarding very low LDL-C levels found in clinical trials of statin and PCSK9 inhibitors are encouraging, these data are neither within the context of a prospective clinical trial to determine the efficacy and/or safety of very low LDL-C levels nor are they long-term data. More definitive information regarding the safety of very low levels of LDL-C will be available from results of ongoing, more long-term ASCVD outcomes trials of PCSK9 inhibitors.

Does clinical outcomes trial evidence support that lowering LDL-C to 70 mg/dL or less reduces ASCVD risk more than lowering LDL-C to 100 mg/dL or less?

Epidemiologically, a general principle is that the higher the cholesterol value, the higher the ASCVD risk. Conversely, the lower the cholesterol, the lower the ASCVD risk. These principles were key in the hypotheses and protocol design regarding sentinel ASCVD outcomes trials. With specific regard to lipid guidelines and recommendations, a central question is whether lowering LDL-C to 70 mg/dL reduces ASCVD risk more than lowering LDL-C to 100 mg/dL. Study of Effectiveness of Additional Reductions in Cholesterol and Homocysteine (SEARCH) was a clinical trial wherein patients with a baseline LDL-C of 97 mg/dL (2.5 mmol/L) were randomized to simvastatin 80 or 20 mg per day.³⁷ Simvastatin 80 mg reduced LDL-C 13 mg/dL (0.35 mmol/L) more than simvastatin 20 mg. The further reduction in LDL-C levels was associated with a 6% proportional risk reduction for major vascular events (RR, 0.94; 95% confidence interval [CI], 0.88–1.01; $P = .10$). A subsequent CTT meta-analysis² (which included the SEARCH trial and the PROVE-IT, TNT, IDEAL, and A to Z trials) showed that greater LDL-C lowering in patients with baseline LDL-C of 98 mg/dL (2.53 mmol/L) with higher-dose statins significantly reduced LDL-C by 19.7 mg/dL (0.51 mmol/L) compared with lower-dose statins. The more aggressive LDL-C reduction below \sim 100 mg/dL with the more-intensive statin regimen resulted in a 15% additional proportional risk reduction for first major vascular event (RR, 0.85; 95% CI, 0.82–0.89; $P < .0001$). Although SEARCH and the CTT meta-analysis supported ASCVD benefit by lowering LDL-C in patients with baseline LDL-C of \sim 100 mg/dL, they did not specifically address the benefit of achieving specific

LDL-C treatment goals because end-of-study LDL-C values were not reported. Therefore, it may be more illustrative to focus on the 4 sentinel CHD outcomes trials that better explored the LDL-C goal hypothesis: PROVE-IT, TNT, IDEAL, and A to Z.^{38–41} Table 1 summarizes randomized, controlled clinical outcomes trials supporting the hypothesis that achievement of an LDL-C of ~ 70 mg/dL further reduces ASCVD compared with achievement of an LDL-C of ~ 100 mg/dL.

Author perspective

Despite the perspective advocated by some that currently available CHD outcomes trials were not designed to evaluate the value of titration of lipid-altering therapy to specific LDL-C goals, evidence from randomized, controlled, clinical ASCVD outcomes trials supports the hypothesis that achievement of an LDL-C level of ~ 70 mg/dL offers greater protection against ASCVD events than achieving an LDL-C level of ~ 100 mg/dL with a good margin of safety.

Which ASCVD outcomes evidence supports a lipid-altering drug titration approach to reduce ASCVD risk?

Results from randomized, controlled clinical trials of lipid-altering drugs support the efficacy, and especially the safety, of various lipid-altering pharmacotherapy titration approaches to achieve lipid treatment goals.^{42–44}

Regarding ASCVD outcomes, perhaps the most important randomized, controlled, statin trial ever conducted was the Scandinavian Simvastatin Survival Study.⁴⁵ This secondary prevention study of 4444 patients is often considered the first study to provide convincing evidence that a statin (simvastatin) reduced not only CHD morbidity and mortality but also reduced overall all-cause mortality. In this randomized, controlled clinical trial, patients were initiated on placebo or simvastatin 20 mg/d. If subsequent total cholesterol exceeded 200 mg/dL (the lipid goal for this study), then, according to the “dosage titration” methods described in the article, the simvastatin 20 mg/d was increased (“titrated”) to 40 mg/d via a computer program at the central laboratory, which allowed for dosage adjustments without revealing lipid levels or treatment allocations. Although the lipid level metric at the time was total cholesterol, and not specifically LDL-C, one could make the case that the most important historic, randomized, controlled clinical statin trial was a titration study wherein the statin dose was increased based on attempts to achieve a lipid goal. Similarly, in the Management of Elevated Cholesterol in the Primary Prevention Group of Adult Japanese Trial, pravastatin was up-titrated from 10 mg to 20 mg to achieve a total cholesterol of <220 mg/dL.⁴⁶ In this titration study, CHD was significantly lower in the

diet plus pravastatin group than in the diet alone group (66 events vs 101 events; hazard ratio, 0.67; 95% CI, 0.49–0.91; $P = .01$).

In the AFCAPS-TexCAPS trial, individuals with average total cholesterol and LDL-C levels were randomized to lovastatin 20 mg or placebo.⁴⁷ The dose of lovastatin was titrated to 40 mg for individuals who did not achieve a goal LDL-C level of <110 mg/dL at 3 months. Lovastatin reduced LDL-C levels by 25% (to 115 mg/dL) from a baseline level of 150 mg/dL and decreased incidence of the first major coronary event (RR, 0.63; $P < .001$), MI (RR, 0.60; $P = .002$), unstable angina (RR, 0.68; $P = .02$), and revascularization procedures (RR, 0.67; $P = .001$).

The ALLIANCE trial evaluated patients with CHD from US managed care database records and implemented a real-world statin titration schedule wherein usual care was compared with titration of atorvastatin dose to achieve an LDL-C goal of <80 mg/dL or a maximum of atorvastatin 80 mg/d.⁴⁸ In this study, 72% of atorvastatin-treated individuals achieved goal compared with 40% of individuals receiving usual care. After an average of 51.5 months of follow-up, titration with atorvastatin significantly lowered LDL-C levels (147 mg/dL [3.8 mmol/L] to 95 mg/dL [2.5 mmol/L]) over usual care (146 mg/dL [3.8 mmol/L] to 111 mg/dL [2.9 mmol/L]). This greater LDL-C reduction was accompanied by improved ASCVD outcomes (-17% with atorvastatin vs usual care; $P = .02$).

Thus, whether it be randomized, controlled lipid-altering efficacy and safety trials or randomized, controlled ASCVD outcomes trials, the data support the efficacy—and especially the safety—of a titration strategy based on treatment to lipid goals. ASCVD outcomes trials support ASCVD risk reduction when lipid-altering drug therapy lowers LDL-C to ~ 70 mg/dL, compared to when the same lipid-altering drug therapy lowers LDL-C to ~ 100 mg/dL. Finally, as noted previously, in patients at very high ASCVD risk with baseline LDL-C level of ~ 95 mg/dL, patients who attained an LDL-C level below 70 mg/dL (mean value ~ 53 mg/dL) with the addition of ezetimibe to simvastatin experienced significantly reduced ASCVD risk compared with patients who attained an LDL-C level of ~ 70 mg/dL with simvastatin alone.¹⁵

Author perspective

Evidence-based data such as these from randomized, clinical outcomes trials help support a lipid treatment strategy of dose titration to a specified lipid goal. These data also support the LDL-C cutoff points established by many global scientific organizations—in this latter case, the reduction of LDL-C below 70 mg/dL in patients at very high ASCVD risk. Table 1 summarizes randomized, controlled clinical outcomes trials supporting the hypothesis that achievement of an LDL-C of ~ 70 mg/dL further reduces ASCVD compared with achievement of an LDL-C of ~ 100 mg/dL.

Table 1 Randomized, controlled ASCVD outcomes trials supporting an achievement of an end-of-study LDL-C of ~70 mg/dL as having improved ASCVD outcomes compared with end-of-study LDL-C levels ~100 mg/dL

Clinical trial	Intervention	LDL-C at baseline, mg/dL (mmol/L)	LDL-C at end of study, mg/dL (mmol/L)	LDL-C reduction (%)	ASCVD outcome
PROVE-IT ³⁸	Atorvastatin 80 mg/d	106 (2.74)	62* (1.60)	−41.5	Compared with achievement of an LDL-C of 95 mg/dL, achievement of an end-of-study LDL-C of 62 mg/dL reduced the risk of composite of death, MI, unstable angina, vascularization, and stroke by 16% ($P = .005$ atorvastatin vs pravastatin)
	Pravastatin 40 mg/d	106 (2.74)	95* (2.46)	−10.4	
TNT ⁴⁰	Atorvastatin 80 mg/d	97 (2.51)	77† (2.00)	−20.6	Compared with achievement of an LDL-C of 101 mg/dL, achievement of an end-of-study LDL-C of 77 mg/dL resulted in a 22% relative reduction in risk of major CV events (HR, 0.78; 95% CI, 0.69–0.89; $P < .001$)
	Atorvastatin 10 mg/d	98 (2.53)	101† (2.6)	3.1	
IDEAL ⁴¹	Atorvastatin 80 mg/d	121.6 (3.15)	80 (2.07)	−34.2‡	Compared with achievement of an LDL-C of 99.8 mg/dL, achievement of an LDL-C of 80 mg/dL reduced the risk for major coronary events by 11% (HR, 0.89; 95% CI, 0.78–1.01; $P = .07$)
	Simvastatin 20 mg/d	121.4 (3.14)	99.8 (2.58)	−17.8‡	
A to Z ³⁹	Simvastatin 40 mg/d then 80 mg/d	112 (2.90)	66 (1.71)	−41.1‡	Compared with an achievement of LDL-C of 81 mg/dL, an end-of-study achievement of an LDL-C of 66 mg/dL resulted in an 11% relative reduction in event rate (CV death, nonfatal MI, readmission for ACS, and stroke; HR, 0.89; 95% CI, 0.76–1.04; $P = .14$)
	Placebo then simvastatin 20 mg/d	111 (2.87)	81 (2.10)	−27.0‡	

ACS, acute coronary syndrome; ASCVD, atherosclerotic cardiovascular disease; CI, confidence interval; CV, cardiovascular; HR, hazard ratio; LDL-C, low-density lipoprotein cholesterol; MI, myocardial infarction.

To convert low-density lipoprotein to mmol/L, multiply by 0.0259.

*Median LDL-C achieved during treatment.

†Mean LDL-C levels during the study.

‡Calculated as percent change from aggregate baseline.

Table 2 Summary of guideline use of LDL-C treatment goals and associated clinical trial evidence*

Guideline† Country	Primary and secondary Goals	Evidence		Treatments/ regimens	Selected outcomes/recommendations
		Clinical trials	Design		
NLA (2014) ⁵² United States					<ul style="list-style-type: none"> Validates and formalizes non-HDL-C <100 mg/dL and LDL-C < 70 mg/dL goal in patients at very high ASCVD risk Therapy should continue in patients with LDL-C < 40 mg/dL based on evidence suggesting no harm from these lower levels Advocates either Framingham risk calculator, Pooled Cohort Equations (ACC/AHA), or Framingham long-term risk calculator for quantitative risk scoring Chronic kidney disease (stage ≥3) added as a high or very high long-term risk Preference for first-line statin therapy, with statins broken out by intensity (superimposable with 2013 ACC/AHA guidelines) Moderate- and high-intensity statin therapy is recommended as first line therapy for prevention of ASCVD but recommends in statin intolerant patients alternate regimens to achieve non-HDL-C and LDL-C goals. Advocates specific goals for patients with hypertriglyceridemia: For those with TG ≥ 500 mg/dL, first priority is lowering TG to <500 mg/dL; for those with TG 200–499, non-HDL-C and LDL-C goals should be achieved in accordance with the patient's risk status The HRs for major CV events per 1-SD ↑ in LDL-C (32 mg/dL), non-HDL-C (36 mg/dL), and apo B (27 mg/dL) were 1.13 (95% CI, 1.10–1.17), 1.16 (95% CI, 1.12–1.19), and 1.14 (95% CI, 1.11–1.18), respectively The association with future major CV events greater for non-HDL-C than for LDL-C and apo B On-treatment levels of non-HDL-C and apo B had a stronger association with CV outcome than levels of LDL-C Lovastatin ↓ LDL-C levels by 25% (to 115 mg/dL) from baseline of 150 mg/dL Lovastatin ↓ major coronary events (37%, $P < .001$) and revascularization (33%, $P = .001$) No significant effect on total mortality or coronary mortality Diet + pravastatin ↓ LDL-C by 18.0% vs 3.2% for diet alone
	Primary targets are non-HDL-C and LDL-C; apo B and HDL-C considered a secondary, optional target; TG not a target of therapy	Meta-analysis of 8 trials (38,153 individuals treated with statins) ⁵⁴	RCTs	Statin therapy in at least 1 study group	
		Meta-analysis of 2 trials ⁵⁵	RCTs (TNT-9319 individuals with stable CHD and LDL-C < 130 mg/dL; and IDEAL- 8699 individuals with a history of myocardial infarction)	High-dose and usual-dose statin (TNT-atorvastatin 10 or 80 mg; IDEAL-simvastatin 20 to 40 mg or atorvastatin 80 mg)	
	"Very high risk": Consider high- or moderate-intensity statin therapy, regardless of non-HDL-C or LDL-C; begin treatment at non-HDL-C ≥100 mg/dL and LDL-C ≥ 70 mg/dL. Treatment goals: non-HDL-C < 100 and LDL-C < 70 mg/dL. Optional apo B goal < 80 mg/dL.	AFCAPS/TexCAPS ⁴⁷	RCT	Lovastatin 20/40 mg or placebo	
	"High risk": begin treatment at non-HDL-C ≥ 130 and LDL-C ≥ 100 mg/dL. Treatment goals: non-HDL-C < 130 and LDL-C	MEGA ⁴⁶	RCT	Diet + pravastatin 10-20 mg or diet	

719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774

Table 2 (continued)

Guideline [†] Country	Primary and secondary Goals	Evidence			
		Clinical trials	Design	Treatments/ regimens	Selected outcomes/recommendations
IAS (2013) ^{†,62} Various countries outside the United States	Primary target is LDL-C (non-HDL-C expected to replace LDL-C in future guidelines) apo B considered an optional target Secondary prevention LDL-C < 70 mg/dL (1.8 mmol/L) Non-HDL-C < 100 mg/dL	SATURN ⁶⁰	Randomized, double-blind, active controlled (1039 individuals with coronary disease)	Atorvastatin 80 mg or rosuvastatin 40 mg	<ul style="list-style-type: none"> • Comparable overall mortality between groups • Percent atheroma volume ↓ by 0.99% (95% CI, −1.19 to −0.63) with atorvastatin and by 1.22% (95% CI, −1.52 to −0.90) with rosuvastatin ($P < .001$) • The frequency of cardiovascular events was similar in the two groups • Very low levels of LDL-C achieved in both treatment groups, with mean values ≤ 70 mg/dL; HDL-C in both groups ~ 50 mg/dL
		REVERSAL ⁶¹	Randomized, double-blind active controlled trial	Atorvastatin 80 mg or pravastatin 40 mg	<ul style="list-style-type: none"> • LDL-C levels reduced to 79 mg/dL in atorvastatin group vs 110 mg/dL in pravastatin group ($P < .001$) • Intensive lowering of LDL-C to <100 mg/dL ↓ progression of coronary atherosclerotic lesions compared with ↓ in LDL-C to 110 mg/dL
		IDEAL ⁴¹	Randomized, open-label, blinded	Atorvastatin 80 mg or simvastatin 20 mg	<ul style="list-style-type: none"> • LDL-C ↓ from 122 to 79 mg/dL in the 80-mg atorvastatin group, compared with 121 to 102 mg/dL in the simvastatin group (at 1 y) • Major coronary events occurred in 10.4% of simvastatin patients vs 9.3% of atorvastatin patients ($P = \text{NS}$)
		TNT ⁴⁰	Randomized, double-blind, active controlled	Atorvastatin 80 mg or atorvastatin 10 mg	<ul style="list-style-type: none"> • No between-group difference in mortality • Median on-treatment LDL-C levels were 77 mg/dL in the high-dose atorvastatin group and 101 mg/dL in the 10-mg group • 22% relative ↓ in risk of major CV events in 80-mg group
		IDEAL ⁴¹	Randomized, open-label, blinded	Atorvastatin 80 mg or simvastatin 20 mg	<ul style="list-style-type: none"> • Comparable overall mortality between groups • LDL-C ↓ from 122 to 79 mg/dL in the 80-mg atorvastatin group, compared with 121 to 102 mg/dL in the simvastatin group (at 1 y) • Major coronary events occurred in 10.4% of simvastatin patients vs 9.3% of atorvastatin patients ($P = \text{NS}$)
		PROVE-IT ³⁸	RCT (4162 individuals hospitalized for ACS)	Atorvastatin 80 mg or pravastatin 40 mg	<ul style="list-style-type: none"> • No between-group difference in mortality • Median on-treatment LDL-C with high-dose atorvastatin was 62 mg/dL vs 95 mg/dL for standard-dose pravastatin
		HPS ⁶³	RCT (20,536 individuals at high risk for CVD event)	Simvastatin 40 mg or placebo	<ul style="list-style-type: none"> • 16% ↓ in composite of death, MI, unstable angina resulting in hospitalization, revascularization, and stroke in intensive treatment group ($P = .005$) • Mean difference in LDL-C between groups was ~ 1 mmol/L favoring active treatment (from baseline of ~ 3.4 mmol/L) • Statin therapy ↓ all-cause mortality (12.9% vs 14.7%, $P = .0003$)

- ~24%–25% ↓ in first event rate for nonfatal MI or coronary death (8.7% vs 11.8%, $P < .0001$), nonfatal or fatal stroke (4.3% vs 5.7%, $P < .0001$), and coronary or non-coronary revascularization (9.1% vs 11.7%, $P < .0001$)
- Higher intensity statins ↓ coronary death of MI ($P < .00001$), and 16% odds reduction of coronary death or any CV event ($P < .00001$)
- Total cholesterol relationship with risk of CHD and death is curvilinear, with cholesterol levels above the 85th percentile ↑ RR by ~4 times
- Total cholesterol ↓ to 180 mg/dL or less over 5 y substantially ↓ risk of ASCVD (primary prevention)
- 2.6% of blacks and 3.2% of whites had sequence mutations in PCSK9
- Blacks had a 28% ↓ in LDL-C levels over 15 y and an 88% ↓ in risk of CHD ($P = .008$)
- Whites had a 15% ↓ in LDL-C levels over 15 y and a 47% ↓ in risk of CHD ($P = .003$)
- PCSK9 R46 L reduced risk of MI (OR, 0.40; $P = .00002$)
- PCSK9 loss-of-function allele ↓ LDL-C levels and provides protection against MI
- All 9 polymorphisms were associated with lower long-term LDL-C levels
- Long-term exposure to lower LDL-C was associated with a 54.5% ↓ in risk of CHD for each mmol/L lower LDL-C (3-fold greater CHD risk reduction per unit lower LDL-C than achieved with statin started later in life)
- Rosuvastatin ↓ LDL-C by 50% and CRP by 37%
- Mean 50% ↓ in LDL-C ↓ major CV events (HR, 0.56, $P < .00001$)
- Rosuvastatin ↓ risk of MI ($P = .0002$), stroke ($P = .002$), revascularization or unstable angina ($P < .00001$), and combined MI, stroke, or CV death ($P = .00001$)
- Atorvastatin ↓ LDL-C levels by 42 mg/dL from baseline of 131 mg/dL
- Nonsignificant trend toward ↓ in total mortality (13%, $P = .16$); ↓ in fatal and nonfatal stroke (27%, $P = .024$), total CV events (21%, $P = .0005$), and total coronary events (29%, $P = .0005$)
- Lovastatin ↓ LDL-C levels by 25% (to 115 mg/dL) from baseline of 150 mg/dL
- Lovastatin ↓ major coronary events (37%, $P < .001$) and revascularization (33%, $P = .001$)

(continued on next page)

Primary prevention LDL-C < 100 mg/dL (2.6 mmol/L) Non-HDL-C < 130 mg/dL	Meta-analysis of 4 trials (TNT, PROVE-IT, IDEAL, A to Z) ⁶⁴	RCTs	Intensive vs standard-dose statin therapy
	MRFIT ⁶⁵	Cohort study of 361,662 men	—
	Genetic (PCSK9) ARIC study ⁶⁶	Population-based samples of residents	Not taking lipid-lowering medication
	Genetic (PCSK9) ⁶⁷	Myocardial Infarction Genetics Consortium and matched free controls	Individuals with early-onset MI and age- and gender-matched controls
	Mendelian Genetic Analysis ⁶⁸	Meta-analysis of individuals with 9 polymorphisms in 6 different genes that ↓ LDL-C	—
	JUPITER ²⁸	RCT (17,802 healthy individuals w/LDL-C <130 mg/dL and CRP ≥ 2.0 mg/L)	Rosuvastatin 20 mg or placebo
	ASCOT-LLA ⁶⁹	RCT (10,305 hypertensive individuals randomized to lipid-lowering arm or placebo)	Atorvastatin 10 mg or placebo
	AFCAPS/TexCAPS ⁴⁷	RCT	Lovastatin 20/40 mg or placebo

Table 2 (continued)

Guideline [†] Country	Primary and secondary Goals	Evidence			
		Clinical trials	Design	Treatments/ regimens	Selected outcomes/recommendations
ACC/AHA (2013) ⁵¹ United States	"Secondary prevention" Primary target: LDL-C Non-HDL-C potential inclusion in future decision-making No recommendations are made for or against specific LDL-C targets Intensity of statin therapy considered the goal of treatment (applied to 1 of 4 major statin benefit groups): clinical ASCVD High-intensity statin therapy should be initiated or continued for adults aged ≤75 y with clinical ASCVD Moderate-intensity statin therapy should be initiated for adults aged >75 y with clinical ASCVD	WOSCOPS ⁷⁰	RCT (6595 men with hypercholesterolemia and no history of MI)	Pravastatin 40 mg or placebo	<ul style="list-style-type: none"> No significant effect on total mortality or coronary mortality Pravastatin ↓ LDL-C levels by 26% from baseline of 192 mg/dL Pravastatin ↓ RR of combined EP of nonfatal MI and death from CHD by 31% ($P < .001$); similar ↓ in nonfatal MI (31%, $P < .001$) and death from CHD ($P = .042$) No effect of pravastatin on deaths from non-CV causes Statin therapy ↓ recurrent CHD events by ~23% (rate ratio) and ↓ stroke events by 17% (CTT) 12% proportional ↓ in all-cause mortality per mmol/L ↓ in LDL-C (rate ratio, 0.88; $P < .0001$) 19% ↓ in coronary mortality (rate ratio, 0.81; $P < .0001$) Proportional ↓ in major vascular events differed significantly ($P < .0001$) according to absolute ↓ in LDL-C achieved Unknown how much additional benefit can be attained below LDL-C levels of 70 mg/dL (1.8 mmol/L)
		CTT collaboration meta-analysis ⁵	14 large, long-term RCT outcome studies	More or less intensive statin regimens or statin vs control	<ul style="list-style-type: none"> No benefit of atorvastatin on primary end point composite of death from cardiac causes, nonfatal MI, or stroke 14.2 event rate (CV death, nonfatal MI, readmission for ACS, and stroke) in intensive simvastatin group vs 16.5 in placebo-simvastatin control group ($P = \text{NS}$) Between-treatment first occurrences of nonfatal MI, nonfatal stroke or CV death were not different ($P = .32$) No significant differences between groups in any secondary outcomes Atorvastatin ↓ LDL-C levels by 34.3% vs 23.3% for usual care ($P < .0001$) 72.4% and 40.0% of atorvastatin-treated and usual-care patients achieved NCEP goal of LDL-C < 100 mg/dL ($P < .001$) Atorvastatin reduced time to first CV event compared with usual care (HR, 0.83; $P = .02$) Atorvastatin ↓ LDL-C levels by 29% vs placebo ($P < .0001$) over 4 y from baseline of 113 mg/dL
		4D ⁷¹	RCT (diabetics on hemodialysis)	Atorvastatin 20 mg or placebo	
		A to Z ⁷²	RCT	Simvastatin 40 mg →80 mg or placebo→ Simvastatin 20 mg	
		ACCORD ⁷³	RCT (diabetics at high risk of CVD)	Open-label simvastatin + fenofibrate or placebo	
		ALLIANCE ⁴⁸	RCT (patients with CHD)	Atorvastatin to achieve LDL-C < 80 mg/dL or up to 80 mg/dL or usual care	
		ASPEN ⁷⁴	RCT (patients with DM and LDL-C at target levels)	Atorvastatin 10 mg or placebo	

- No benefit of atorvastatin on primary EP of CV death, nonfatal MI and stroke, recanalization, CABG, resuscitated cardiac arrest, and angina (13.7% for atorvastatin vs 15% for placebo)
- Rosuvastatin ↓ LDL-C levels by 43% from baseline of 100 mg/dL at 3 mo
- No effect on primary EP of CV death, nonfatal MI, or stroke (9.2 and 9.5 events per 100 patient-y; $P = .59$)
- No effect of active treatment on all-cause mortality
- Pravastatin ↓ LDL-C levels by 32% (to 97–98 mg/dL) from baseline of 139 mg/dL
- Pravastatin ↓ major coronary events (24%, $P = .003$), revascularization (26%, $P = .005$), and stroke (31%, $P = .03$)
- Showed benefits of statin therapy in patients with average cholesterol levels
- Rosuvastatin ↓ LDL-C levels by 45% (difference between groups; $P < .001$) from baseline of ~ 3.5 mmol/L
- No effect on primary EP at 33 mo of CV death, nonfatal MI or stroke (HR, 0.92; $P = .12$)
- No effect of active treatment on all-cause mortality
- Atorvastatin ↓ LDL-C and non-HDL-C levels by 46% and 44%, respectively, over 3 y
- 95% and 97% of patients treated with atorvastatin achieved NCEP goals for LDL-C and non-HDL-C, compared with only 3% of usual-care patients (LDL-C)
- 24.5% of usual care patients vs 12% of atorvastatin patients had a CHD recurrent event or died (RR, 0.49; $P < .0001$)
- Simvastatin + niacin ↓ LDL-C by 42% and ↑ HDL-C by 26%, compared with no change in other groups
- Average stenosis regressed by 0.4% in the simvastatin + niacin group, compared with progression of 3.9% (placebos), 1.8% (antioxidants), and 0.7% (simvastatin + niacin + antioxidants; $P < .001$)
- Frequency of death, MI, stroke, or revascularization was lowest in simvastatin + niacin group (3% vs 14%–24% range in other groups)
- Median difference in LDL-C between groups was ~ 1 mmol/L favoring active treatment (from baseline of ~ 3.4 mmol/L)
- Statin therapy ↓ all-cause mortality (12.9% vs 14.7%, $P = .003$)

(continued on next page)

		Rosuvastatin 10 mg or placebo	
		Pravastatin 40 mg or placebo	
		Rosuvastatin 10 mg or placebo	
		Atorvastatin 10–80 mg to reach NCEP goal of LDL-C < 100 mg/dL or usual care	
		Simvastatin 10 mg (titrated to achieve LDL-C < 90 mg/dL during first year) + niacin 1000 mg b.i.d. or antioxidants or simvastatin + niacin + antioxidants or matching placebos	
		Simvastatin 40 mg or placebo	
AURORA ⁷⁵	RCT (patients undergoing maintenance hemodialysis)		
CARE ⁷⁶	RCT		
CORONA ⁷⁷	RCT (patients aged >60 y with systolic heart failure; NYHA II-IV)		
GREACE ⁷⁸	RCT (patients with established CHD)		
HATS ⁷⁹	RCT (patients with CHD and normal LDL-C levels)		
HPS ⁶³	RCT		

Table 2 (continued)

Guideline [†] Country	Primary and secondary Goals	Evidence			
		Clinical trials	Design	Treatments/ regimens	Selected outcomes/recommendations
		IDEAL ⁴¹	RCT	Atorvastatin 80 mg or simvastatin 20 mg	<ul style="list-style-type: none"> • ~24%–25% ↓ in first event rate for nonfatal MI or coronary death (8.7% vs 11.8%, $P < .0001$), nonfatal or fatal stroke (4.3% vs 5.7%, $P < .0001$), and coronary or non-coronary revascularization (9.1% vs 11.7%, $P < .0001$) • LDL-C ↓ from 122 to 79 mg/dL in the 80-mg atorvastatin group, compared with 121 to 102 mg/dL in the simvastatin group (at 1 y) • Major coronary events occurred in 10.4% of simvastatin patients vs 9.3% of atorvastatin patients ($P = \text{NS}$)
		LIPID ⁸⁰	RCT	Pravastatin 40 mg or placebo	<ul style="list-style-type: none"> • No between-group difference in mortality • Pravastatin ↓ LDL-C levels by 25% from baseline of 150 mg/dL • Pravastatin ↓ RR of death from CHD (24%, $P < .001$), overall mortality (22%, $P < .001$), MI (29%, $P < .001$), coronary revascularization (20%, $P < .001$), and stroke (19%, $P = .048$) • Pravastatin ↓ risk in a broad range of baseline LDL-C levels
		LIPS ⁸¹	RCT (patients with successful PCI and total cholesterol between 135 and 270 mg/dL)	Fluvastatin 80 mg or placebo	<ul style="list-style-type: none"> • Fluvastatin ↑ MACE-free survival time over ~4 y ($P = .01$) • 21.4% of fluvastatin group vs 26.7% of placebo group had at least 1 MACE (RR, 0.78; $P = .01$)
		MIRACL ⁸²	RCT	Atorvastatin 80 mg or placebo	<ul style="list-style-type: none"> • Mean LDL-C ↓ from 124 mg/dL to 72 mg/dL • Atorvastatin ↓ RR of death, nonfatal acute MI, cardiac arrest w/resuscitation, or recurrent symptomatic myocardial ischemia to 0.84 ($P = .048$) • Atorvastatin 80 mg/d ↓ recurrent ischemic events in first 16 wk
		MUSASHI-AMI ⁸³	RCT (Japanese patients with acute MI)	Various statins and doses vs no statin (early use after acute MI)	<ul style="list-style-type: none"> • Statins ↓ composite of CV death, nonfatal AMI, recurrent symptomatic myocardial ischemia, CHF, and stroke 6.1% vs 11.4% ($P = .043$) • Statins ↓ risk of CHF and symptomatic myocardial ischemia ($P = .0154$ and 0.0264, respectively)
		PROVE-IT ³⁸	RCT (4162 individuals hospitalized for ACS)	Atorvastatin 80 mg or pravastatin 40 mg	<ul style="list-style-type: none"> • Early lipid lowering is beneficial in reducing recurrent CV events • Median on-treatment LDL-C with high-dose atorvastatin was 62 mg/dL vs 95 mg/dL for standard-dose pravastatin

- 16% ↓ in composite of death, MI, unstable angina resulting in hospitalization, vascularization, and stroke in intensive treatment group ($P = .005$)
- Atorvastatin ↓ LDL-C by 45% to 73 mg/dL compared with a 4% change in placebo patients to 129 mg/dL
- Atorvastatin ↓ risk of second stroke (fatal or nonfatal) compared with placebo over 5 y (11.2% vs 13.1%; HR, 0.84; $P = .03$)
- ↓ in incidence of second stroke was specific to ischemic stroke vs hemorrhagic stroke
- Median on-treatment LDL-C levels were 77 mg/dL in the high-dose atorvastatin group and 101 mg/dL in the 10-mg group
- 22% relative ↓ in risk of major CV events in 80-mg group
- Comparable overall mortality between groups
- Lovastatin ↓ LDL-C levels by 25% (to 115 mg/dL) from baseline of 150 mg/dL
- Lovastatin ↓ major coronary events (37%, $P < .001$) and revascularization (33%, $P = .001$)
- No significant effect on total mortality or coronary mortality
- Atorvastatin ↓ LDL-C levels by 29% vs placebo ($P < .001$) from baseline of 113 mg/dL over 4 y
- No benefit of atorvastatin on primary EP of CV death, nonfatal MI and stroke, revascularization, CABG, resuscitated cardiac arrest, and angina (13.7% for atorvastatin vs 15% for placebo)
- Rosuvastatin ↓ LDL-C levels by 43% from baseline of 100 mg/dL at 3 mo
- No effect on primary EP of CV death, nonfatal MI or stroke (9.2 and 9.5 events per 100 patient-years; $P = .59$)
- No effect of active treatment on all-cause mortality
- 1.54 per 100 person-y at risk receiving atorvastatin experienced a major cardiovascular event vs 2.46 per 100 person-y at risk of patients allocated to placebo (rate reduction, 37%; $P = .001$)
- Atorvastatin ↓ acute CHD events by 36%, coronary revascularizations by 31%, stroke by 48%, and death rate by 27%
- Atorvastatin reduces risk of first CV disease event in patients with DM without high LDL-C
- Rosuvastatin ↓ LDL-C by 50% and CRP by 37%
- Mean 50% ↓ in LDL-C ↓ major CV events (HR, 0.56; $P < .0001$)
- Rosuvastatin ↓ risk of MI ($P = .0002$), stroke ($P = .002$), revascularization or

(continued on next page)

	SPARCL ⁵⁸	RCT (stroke patients without evidence of heart disease)	Atorvastatin 80 mg or placebo
	TNT ⁴⁰	RCT	Atorvastatin 80 mg or atorvastatin 10 mg
	AFCAPS/TexCAPS ⁴⁷	RCT	Lovastatin 20/40 mg or placebo
<p>"Primary prevention"</p> <p>Primary target is LDL-C</p> <p>No recommendations are made for or against specific LDL-C targets</p> <p>Intensity of statin therapy considered the goal of treatment (applied to 3 of 4 major statin-benefit groups):</p> <ol style="list-style-type: none"> 1. Primary LDL-C ≥ 190 mg/dL 2. Diabetes (40–75 y) with 3. LDL-C 70–189 mg/dL w/o ASCVD <p>10-y ASCVD risk $\geq 7.5\%$ and LDL-C 70–189 mg/dL w/o ASCVD</p> <p>For primary LDL-C ≥ 190 mg/dL, use high-intensity statin therapy to achieve $>50\%$ ↓</p> <p>For diabetes (40–75 y) with LDL-C 70–189 mg/dL w/o ASCVD, moderate-intensity statin therapy should be initiated or continued</p> <p>For patients with 10-y ASCVD risk $\geq 7.5\%$ and LDL-C 70–189 mg/dL w/o ASCVD, therapy should be initiated with moderate- or high-intensity statin</p>	ASPEN ⁷⁴	RCT (patients with DM and LDL-C at target)	Atorvastatin 10 mg or placebo
	AURORA ⁷⁵	RCT (patients undergoing maintenance hemodialysis)	Rosuvastatin 10 mg or placebo
	CARDS ⁸⁴	RCT (patients with type 2 DM, no history of CV disease, and normal LDL-C levels)	Atorvastatin 10 mg or placebo
	JUPITER ²⁸	RCT (17,802 healthy individuals w/LDL-C < 130 mg/dL and CRP ≥ 2.0 mg/L)	Rosuvastatin 20 mg or placebo

Table 2 (continued)

Guideline [†] Country	Primary and secondary Goals	Evidence		Treatments/ regimens	Selected outcomes/recommendations
		Clinical trials	Design		
CCS (2012/2013) ⁸⁵ Canada	<p>Primary target is LDL-C level Alternate lipid markers include non-HDL-C and apo B (newly added) High risk: LDL-C \leq 2 mmol/L or \geq50% \downarrow (LDL-C target of $<$1.8 mmol/L [70 mg/dL] is justified in some high-risk patients, in line with ATP III and ESC/EAS recommendations)^{86,87} Non-HDL-C \leq 2.6 mmol/L apo B \leq 0.8 mmol/L</p> <p>Intermediate risk: LDL-C \leq 2 mmol/L or \geq50% \downarrow Non-HDL-C \leq 2.6 mmol/L apo B \leq 0.8 g/L</p>	MEGA ⁴⁶	RCT	Diet + pravastatin 10-20 mg or diet	<p>unstable angina ($P < .00001$), and combined MI, stroke, or CV death ($P = .02$)</p> <ul style="list-style-type: none"> • Diet + pravastatin \downarrow LDL-C by 18.0% vs 3.2% for diet alone • Diet + pravastatin \downarrow TC by 11.5% vs 2.1% for diet alone • CHD was significantly \downarrow in the diet + pravastatin group (66 events vs 101 events for diet alone; HR, 0.67; $P = .01$) • Across all 26 trials, all-cause mortality was reduced by 10% per 1 mmol/L \downarrow in LDL-C • No increase in major side effect rates were observed at lower LDL-C levels • Simvastatin + ezetimibe produced an LDL-C difference of 0.85 mmol/L • The combination \downarrow major atherosclerotic events by 17% (11.3% vs 13.4%; rate ratio, 0.83; $P = .0021$) • Significant \downarrow in nonhemorrhagic stroke ($P = .01$) and arterial revascularization procedures ($P = .0036$) • Lovastatin \downarrow LDL-C levels by 25% (to 115 mg/dL) from baseline of 150 mg/dL • Lovastatin \downarrow major coronary events (37%, $P < .001$) and revascularization (33%, $P = .001$) • No significant effect on total mortality or coronary mortality • Pravastatin \downarrow LDL-C levels by 26% from baseline of 192 mg/dL • Pravastatin \downarrow RR of combined EP of nonfatal MI and death from CHD by 31% ($P < .001$); similar \downarrow in nonfatal MI (31%, $P < .001$) and deaths from CHD ($P = .042$) • No effect of pravastatin on deaths from non-CV causes • Atorvastatin \downarrow LDL-C levels by 42 mg/dL from baseline of 132 mg/dL • Nonsignificant trend toward \downarrow in total mortality (13%, $P = .16$); \downarrow in fatal and nonfatal stroke (27%, $P = .024$), total CV events (21%, $P = .0005$), and total coronary events (29%, $P = .0005$) • Rosuvastatin \downarrow LDL-C by 50% and CRP by 37% • Mean 50% \downarrow in LDL-C \downarrow major CV events (HR, 0.56; $P < .00001$) • Rosuvastatin \downarrow risk of MI ($P = .0002$), stroke ($P = .002$), revascularization or unstable angina ($P < .00001$), and combined MI, stroke, or CV death ($P = .02$)
		CTT collaboration meta-analysis ²	26 large, long-term outcome studies	More or less intensive statin regimens or statin or control	
		SHARP ⁸⁸ Additional clinical trials were reviewed but not cited in the guidelines	Randomized placebo-controlled trial in patients with CKD	Simvastatin 20 mg + ezetimibe 10 mg or placebo	
		AFCAPS/TexCAPS ⁴⁷	RCT	Lovastatin 20/40 mg or placebo	
		WOSCOPS ⁷⁰	RCT	Pravastatin 40 mg or placebo	
		ASCOT-LLA ⁶⁹	RCT (10,305 hypertensive individuals randomized to lipid-lowering arm or placebo)	Atorvastatin 10 mg or placebo	
		JUPITER ²⁸	RCT (17,802 healthy individuals w/LDL-C $<$ 130 mg/dL and CRP \geq 2.0 mg/L)	Rosuvastatin 20 mg or placebo	

1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670

Table 2 (continued)

Guideline [†] Country	Primary and secondary Goals	Evidence		Treatments/ regimens	Selected outcomes/recommendations
		Clinical trials	Design		
ESC/EAS (2011/2012) ⁸⁶ Europe, Mediterranean Region	"Primary prevention" ⁹⁴ Category I (<0.5% risk of CAD death) LDL-C < 160 mg/dL Category II (≥0.5%, <2.0% risk of CAD death) LDL-C < 140 mg/dL Category III (≥2% risk of CAD death) LDL-C < 120 mg/dL Primary target is LDL-C level Secondary targets include non-HDL-C and apo B Very high risk: LDL-C < 1.8 mmol/L (70 mg/dL) or ≥50% ↓ apo B < 80 mg/dL	MIRACL ⁸²	RCT (post-ACS)	Atorvastatin 80 mg or placebo	<ul style="list-style-type: none"> • Pravastatin ↓ RR of death from CHD (24%, $P < .001$), overall mortality (22%, $P < .001$), MI (29%, $P < .001$), coronary revascularization (20%, $P < .001$), and stroke (19%, $P = .048$) • Pravastatin ↓ risk in a broad range of baseline LDL-C levels • Mean LDL-C ↓ from 124 mg/dL to 72 mg/dL • Atorvastatin ↓ RR of death, nonfatal acute MI, cardiac arrest w/resuscitation, or recurrent symptomatic myocardial ischemia to 0.84 ($P = .048$) • Atorvastatin 80 mg ↓ recurrent ischemic events in first 16 wk • Pravastatin ↓ RR of death, recurrence of MI, or readmission to hospital for unstable angina (within 30 d) by 6.4% (11.6% vs 12.4%; $P=NS$) • 14.2 event rate (CV death, nonfatal MI, readmission for ACS, and stroke) in intensive simvastatin group vs 16.5 in placebo-simvastatin control group ($P=NS$) • At 1 y, adjusted mortality was 9.3% in the no-statin group and 4.0% in the statin group • Early statin treatment ↓ 1-y mortality (RR, 0.75; $P = .001$) • 6 mo after PCI, patients were treated to achieve LDL-C of <100 mg/dL • Early statin administration is a good predictor of MACCE (HR, 0.46; $P = .015$) • Cumulative event-free survival significantly higher in atorvastatin group ($P = .041$) • Early statin therapy for ACS ↓ rate of death and CV events over 2-y follow-up period (HR, 0.81; $P < .001$) • Survival benefits were evident within first 12 mo • Diet + pravastatin ↓ LDL-C by 18.0% vs 3.2% for diet alone • Diet + pravastatin ↓ TC by 11.5% vs 2.1% for diet alone • CHD was significantly ↓ in the diet + pravastatin group (66 events vs 101 events for diet alone; HR, 0.67; $P = .01$) • Every 1 mmol/L (~40 mg/dL) ↓ in LDL-C is associated with a 22% ↓ in CVD mortality and morbidity • Absolute ↓ to LDL-C level <1.8 mmol/L or at least a 50% relative ↓ provides the best reduction in CVD
		PACT ⁹⁰	RCT (post-acute MI or unstable angina)	Pravastatin 20-40 mg or placebo	
		A to Z ³⁹	Randomized, double-blind, controlled	Simvastatin 40 mg → 80 mg or placebo → Simvastatin 20 mg	
		RIKS-HIA ⁹¹	Prospective cohort study from Swedish Register of Cardiac Intensive Care	Various statin or no statin therapy	
		Ext-ESTABLISH ⁹²	RCT (post-acute coronary syndrome)	Atorvastatin 20 mg or standard care	
		Meta-analysis ⁹³	13 RCTs started within 14 d of hospitalization for acute coronary syndrome	Various statins at different dose intensities	
		MEGA ⁴⁶	RCT	Diet + pravastatin 10-20 mg or diet	
		CTT collaboration meta-analysis ²	26 large, long-term outcome studies	More or less intensive statin regimens or statin vs control	

(continued on next page)

Table 2 (continued)

Guideline† Country	Primary and secondary Goals	Evidence			
		Clinical trials	Design	Treatments/ regimens	Selected outcomes/recommendations
	High-risk CHD or CHD risk equivalents (10-y risk > 20%) LDL-C < 100 mg/dL	HPS ⁶³	RCT (20,536 individuals at high risk for CVD event)	Simvastatin 40 mg or placebo	<ul style="list-style-type: none"> Intensive lowering of LDL-C to <100 mg/dL ↓ progression of coronary atherosclerotic lesions compared with ↓ in LDL-C to 110 mg/dL Simvastatin ↓ all-cause mortality by 13% ($P = .0003$) Reductions in major vascular events (24%), coronary death rate (18%), nonfatal MI + coronary death (27%), nonfatal or fatal strokes (25%), and CV revascularization (24%) Benefits in older persons Composite EP of coronary death, nonfatal MI, and fatal or nonfatal stroke ↓ 15% ($P = .014$) in pravastatin group Reductions in major coronary events (19%, $P = .006$) and CHD mortality (24%, $P = .043$) in pravastatin group; no significant change in stroke occurrence Benefits in older persons Atorvastatin ↓ LDL-C levels by 42 mg/dL from baseline of 132 mg/dL Nonsignificant trend toward ↓ in total mortality (13%, $P = .16$); ↓ in fatal and nonfatal stroke (27%, $P = .024$), total CV events (21%, $P = .0005$), and total coronary events (29%, $P = .0005$) Atorvastatin LDL-C, 62 mg/dL vs 95 mg/dL with pravastatin 16% ↓ in composite of death, MI, unstable angina resulting in hospitalization, vascularization, and stroke in intensive treatment group ($P < .005$) Simvastatin ↓ LDL-C levels by 35% from baseline of 188 mg/dL Simvastatin ↓ RR of death by 30% ($P = .0003$) Significant ($P < .05$ or lower) ↓ in risk for major coronary events (35%), revascularization (37%), coronary mortality (42%), total mortality (30%), and stroke (27%) Pravastatin ↓ LDL-C levels by 32% (to 98–99 mg/dL) from baseline of 139 mg/dL Pravastatin ↓ major coronary events (24%, $P = .003$), revascularization (26%, $P = .005$), and stroke (31%, $P = .03$)
		PROSPER ⁹⁸	RCT (5804 individuals aged 70–82 y with history of vascular disease or CVD risk factors)	Pravastatin 40 mg or placebo	
		ASCOT-LLA ⁶⁹	RCT (10,305 hypertensive individuals randomized to lipid-lowering arm or placebo)	Atorvastatin 10 mg or placebo	
		PROVE-IT ³⁸	RCT (4162 individuals hospitalized for ACS)	Atorvastatin 80 mg or pravastatin 40 mg	
		4S ⁴⁵	RCT (4444 patients with CHD)	Simvastatin 10/40 mg or placebo	
		CARE ⁷⁶	RCT	Pravastatin 40 mg or placebo	

- Showed benefits of statin therapy in patients with average cholesterol levels
- Pravastatin ↓ LDL-C levels by 25% from baseline of 150 mg/dL
- Pravastatin ↓ RR of death from CHD (24%, $P < .001$), overall mortality (22%, $P < .001$), MI (29%, $P < .001$), coronary revascularization (20%, $P < .001$), and stroke (19%, $P = .048$)
- Pravastatin ↓ risk in a broad range of baseline LDL-C levels
- Mean LDL-C ↓ from 124 mg/dL to 72 mg/dL
- Atorvastatin ↓ RR of death, nonfatal acute MI, cardiac arrest w/resuscitation, or recurrent symptomatic myocardial ischemia to 0.84 ($P = .048$)
- Atorvastatin 80 mg ↓ recurrent ischemic events in first 16 wk
- 13% of patients in the atorvastatin group (↓ LDL-C by 46% to 77 mg/dL) had ischemic events, compared with 21% of patients who underwent angioplasty/usual care (36% ↓ over 18 mo; $P = .048$)
- Atorvastatin was associated with a significantly longer time to the first ischemic event ($P = .03$)
- No difference in percent diameter stenosis between groups ($P > .20$)
- For lesions $\geq 50\%$, average percent diameter ↓ 4.1% in lovastatin patients compared with an ↑ in stenosis diameter in the placebo group ($P = .005$)
- Lovastatin + diet slows the rate of progression and ↑ frequency of regression in coronary artery lesions
- Simvastatin ↓ LDL-C by 47% to 4.1 mmol/L compared with 63% to 3.0 mmol/L for simvastatin plus biweekly LDL-C apheresis
- There was no change in hyperemic mean transit time in the simvastatin group, compared with a ↓ hyperemic mean transit time in the simvastatin plus apheresis group
- Aggressive reduction in LDL-C has a favorable effect on regional myocardial perfusion and alleviates ischemia
- Aggressive lovastatin doses ↓ LDL-C levels to 93–97 mg/dL vs 132–136 mg/dL for moderate-dose lovastatin ($P < .001$)
- Percentage of grafts with progression to atherosclerosis was ↓ in the aggressive dosing group relative to moderate-dose group (27% vs 39%, respectively, $P < .001$)

(continued on next page)

LIPID ⁸⁰	RCT	Pravastatin 40 mg or placebo
MIRACL ⁸²	RCT	Atorvastatin 80 mg or placebo
AVERT ⁹⁹	RCT (341 patients with stable CAD referred for percutaneous revascularization)	Atorvastatin 80 mg or percutaneous revascularization followed by usual care
MARS ¹⁰⁰	RCT (270 patients with angiographically defined CAD)	Diet + lovastatin 80 mg or diet + placebo
LAARS ¹⁰¹	RCT (42 patients with severe hypercholesterolemia and extensive coronary artery disease)	Simvastatin 40 mg with or without LDL apheresis
Post-CABG ¹⁰²	RCT (1351 patients who had undergone bypass surgery and LDL-C between 130 and 175 mg/dL)	Lovastatin 40 mg (titrated to 80 mg to achieve LDL-C < 85 mg/dL) or lovastatin 5 mg (moderate-treatment group)

2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062

2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118

Table 2 (continued)

Guideline [†] Country	Primary and secondary Goals	Evidence		Treatments/ regimens	Selected outcomes/recommendations
		Clinical trials	Design		
		FATS extension ¹⁰³	RCT (29 patients with coronary artery disease and hyperlipidemia)	Triple therapy with lovastatin 20 mg b.i.d., niacin 500 mg q.i.d., and colestipol 10 g b.i.d.	<ul style="list-style-type: none"> Aggressive LDL-C lowering to <100 mg/dL reduced progression of atherosclerosis in grafts Target LDL-C level of \leq100 mg/dL at 8 mo achieved by 83% of patients (controlled-release niacin) vs 52% receiving regular niacin ($P < .01$) High proportion of patients achieved target goal of \leq100 mg/dL Controlled-release niacin preferred to standard niacin
		HATS ⁷⁹	RCT (patients with CHD and normal LDL-C levels)	Simvastatin 10 mg (titrated to achieve LDL-C <90 mg/dL during first year) + niacin 1000 mg b.i.d. or antioxidants or simvastatin + niacin + antioxidants or matching placebos	<ul style="list-style-type: none"> Simvastatin + niacin ↓ LDL-C by 42% and ↑ HDL-C by 26%, compared with no change in other groups Average stenosis regressed by 0.4% in the simvastatin + niacin group, compared with progression of 3.9% (placebos), 1.8% (antioxidants), and 0.7% (simvastatin + niacin + antioxidants; $P < .001$) Frequency of death, MI, stroke, or revascularization was lowest in simvastatin + niacin group (3% vs 14%–24% range in other groups)
	Moderately high risk 2 + risk factors (10-y risk 10%–20%) LDL-C < 130 mg/dL “Moderate risk” “2 + risk factors (10-y risk <10%)” LDL-C < 130 mg/dL “Low risk (0–1 risk factors)” LDL-C < 160 mg/dL	ASCOT-LLA ⁶⁹	RCT (10,305 hypertensive individuals randomized to lipid-lowering arm or placebo)	Atorvastatin 10 mg or placebo	<ul style="list-style-type: none"> Atorvastatin ↓ LDL-C levels by 42 mg/dL from baseline of 132 mg/dL Nonsignificant trend toward ↓ in total mortality (13%, $P = .16$); ↓ in fatal and nonfatal stroke (27%, $P = .024$), total CV events (21%, $P = .0005$), and total coronary events (29%, $P = .0005$) Clinical trial evidence suggests that an LDL-C goal < 100 mg/dL is an option LDL-C goal of <100 mg/dL extends to patients with baseline LDL-C of 100 to 129 mg/dL
		ALLHAT-LLT ¹⁰⁴	RCT (10,355 hypertensive individuals with moderate hypercholesterolemia)	Pravastatin or usual care	<ul style="list-style-type: none"> 6-y CHD event rates were similar between groups (9.3% for pravastatin and 10.4% for usual care) CHD events were significantly reduced in the African-American subgroup
		WOSCOPS ⁷⁰	RCT (6595 men with hypercholesterolemia and no history of MI)	Pravastatin 40 mg or placebo	<ul style="list-style-type: none"> Pravastatin ↓ LDL-C levels by 26% from baseline of 192 mg/dL Pravastatin ↓ RR of combined EP of nonfatal MI and death from CHD by 31% ($P < .001$); similar ↓ in nonfatal MI (31%, $P < .001$) and deaths from CHD ($P = .042$) No effect of pravastatin on deaths from non-CV causes
		Meta-analysis ¹⁰⁵	5 trials of 30,817 men and women	Various treated and control populations	<ul style="list-style-type: none"> Statin treatment ↓ LDL-C by 28% Statin treatment ↓ major coronary events (31%) and all-cause mortality (21%)
		AFCAPS/TexCAPS ⁴⁷	RCT	Lovastatin 20/40 mg or placebo	<ul style="list-style-type: none"> Lovastatin ↓ LDL-C levels by 25% (to 115 mg/dL) from baseline of 150 mg/dL

- Lovastatin ↓ major coronary events (37%,
P < .001) and revascularization (33%,
P = .001)
- No significant effect on total mortality or
coronary mortality

ACC, American College of Cardiology Foundation; AHA, American Heart Association; apo B, apolipoprotein B; ASCVD, atherosclerotic cardiovascular disease; b.i.d., twice daily; CABG, coronary artery bypass surgery; CAD, coronary artery disease; CHF, congestive heart failure; CI, confidence interval; CKD, chronic kidney disease; CRP, C-reactive protein; CTT, Cholesterol Treatment Trialists; CVD, cardiovascular disease; DM, diabetes mellitus; EAS, European Atherosclerosis Society; EP, end point; ESC, European Society of Cardiology; FRS, Framingham Risk Score; HDL-C, high-density lipoprotein cholesterol; HR, hazard ratio; JAS-G, Japan Atherosclerosis Society Guidelines; LDL-C, low-density lipoprotein cholesterol; MACCE, major adverse cardiac and cerebrovascular events; MACE, major adverse cardiac event; MI, myocardial infarction; NCEP, National Cholesterol Education Program; NICE, National Institute for Health and Care Excellence; NLA, National Lipid Association; NS, not significant; NYHA, New York Heart Association; OR, odds ratio; PCI, percutaneous coronary intervention; q.i.d., 4 times daily; RCT, randomized controlled trial; RR, relative risk; SCORE, Systemic Coronary Risk Estimation; SD, standard deviation; TC, total cholesterol; TG, triglyceride.

* Does not include an exhaustive review of various patient populations and demographic factors, including review of guidelines for diabetes, familial hypercholesterolemia, hemodialysis and CKD, peripheral artery disease, and considerations for gender, ethnicity, and age (eg, persons aged older than 75 years).

† Guidelines are generally listed in chronological order from most recent to least current.

‡ International Atherosclerosis Society is a position statement.

What are other potential advantages of lipid treatment goals, from a patient-centered perspective?

Beyond the scientific support for recommending lipid goals are practical and behavioral patient care considerations. In collaboration with the National Committee for Quality Assurance, American Society of Health-System Pharmacists, and American Medical Association, the American College of Cardiology (ACC), American Heart Association (AHA), American Association of Cardiovascular and Pulmonary Rehabilitation, American Academy of Family Physicians, and American Nurses Association have outlined concepts for clinician–patient shared accountability in performance measures.⁴⁹ The goal-related concepts included the following: (1) patients who are more actively engaged in self-care, defined as the ability to perform the activities necessary to achieve, maintain, or promote optimal health, are more likely to successfully achieve their treatment goals; (2) the general framework of shared accountability is predicated on partnerships between patients and clinicians, in which patients play an active role in setting goals, making treatment decisions, and assessing outcomes; (3) key conceptual issues for shared accountability are (a) shared goal setting, (b) shared decision-making, (c) shared care planning and monitoring, including patient feedback and self-care, and (d) assessment of patients' longitudinal outcomes; (4) in management of other metabolic diseases, such as hypertension, specific blood pressure goals are useful, as they allow for process measures, such as “treatment intensification or alteration by clinician(s) in response to elevated blood pressure levels,” if the first medication choice or dose titration does not achieve the desired goal, several iterations of this process may be required; (5) defining the appropriate period of evaluation is an important technical feature of performance measures and should be meaningful from both patient and provider perspectives; performance measures must define a discrete period of measurement consistent with the actual treatment goals for the measures; and (6) as a principle of shared accountability, performance on these measures should be reported back to both clinicians and patients in a timely fashion to facilitate shared care management and achievement of best outcomes. Thus, having lipid goals may enhance patient self-care and accountability, improve monitoring of the patient's response to treatment and clinical progress, and better allow for patient/clinician partnership in making treatment adjustments for the duration of lipid therapy.

Perhaps an unintended consequence of the removal of lipid treatment goals in the ACC/AHA guidelines was the proposed recommendation of “retirement” (or removal) of LDL-C measures for screening or assessment of lipid control (<http://www.ncqa.org/Portals/0/HomePage/CMC.pdf>, accessed December 15, 2014). The guidance that lipid levels should no longer be measured and the subsequent potential

reaction from payers that such lipid measures will not be reimbursed may severely impair the ability of clinicians to assess not only the lipid response to therapy but also the persistence of use of lipid-altering interventions within the context of clinical practice. It is based on challenges presented by the ACC/AHA guidelines, compared with other existing international guidelines that some European authors conclude that existing lipid guidelines and recommendations having lipid treatment goals (eg, European Society of Cardiology/European Atherosclerosis Society guidelines) “seem to be the most wide-ranging, pragmatic, and appropriate choice for European countries.”⁵⁰

Author perspective

With regard to the clinical utility of LDL-C and non-HDL-C levels in the management of dyslipidemic patients, the ACC/AHA Guideline authors stated: “This guideline does not recommend their use as performance measures.”⁵¹ However, ACC/AHA Guideline authors also recommended that a fasting lipid panel be performed 4 to 12 weeks after statin initiation or dose adjustment and every 3 to 12 months thereafter. Conversely, the National Lipid Association (NLA) recommendations recommend routine post-treatment lipid testing, because lipid “treatment goals facilitate effective communication between patients and clinicians, providing an easily interpretable means through which the clinician can communicate progress toward meeting treatment objectives, thus supporting efforts to maximize long-term adherence to the treatment plan.”⁵² Finally, with the continued development of novel lipid-altering drugs, some of those most involved in pharmaceutical administration have advocated for specific LDL-C goals, which serve to facilitate a stepwise approach and rational clinical decision-making, intended to enhance efficacy, safety, and evidence quality.⁵³

How do global lipid guidelines, recommendations, and position papers compare regarding primary and secondary treatment goals?

The previous discussion outlines the historical perspective of data supporting guidelines and recommendations. The clinical application of these data is reflected in the 8 international major lipid guidelines and recommendations summarized in Table 2.^{51,52,56,62,85,86,89,97}

Regardless of baseline lipid levels, patients at high or very high ASCVD risk may benefit from high-intensity statin therapy.¹⁰⁶ The ACC/AHA cholesterol guidelines (see in the following section) define specific statin benefit groups. Organizations such as the NLA recommend the consideration of moderate- to high-intensity statin for patients with ASCVD or diabetes mellitus, irrespective of baseline lipid levels.⁵²

In addition to the general recommendation of statin therapy for patients at high or very high ASCVD risk, 6 of 8 guidelines also provide explicit LDL-C goals aligned with appropriate risk categories.^{51,57,62,85,86,89,97} Consistent with the principle that achievement of lower cholesterol levels results in greater ASCVD reduction, an optional goal for LDL-C of <70 mg/dL (<1.8 mmol/L) for individuals at very high ASCVD risk is recommended in 3 guidelines (NLA, “European Society of Cardiology/European Atherosclerosis Society,” National Cholesterol Education Program Adult Treatment Panel III),^{52,62,86,97} and LDL-C <2 mmol/L (<77 mg/dL) in 1 lipid guideline (Canadian Cardiovascular Society).⁸⁵ The Japanese Atherosclerosis Society guidelines also have established explicit LDL-C goals; however, the lower level of LDL-C was set at <100 mg/dL (<2.6 mmol/L) for patients at high risk.⁸⁹

The NLA has dual lipid treatment goals, with respect to non-HDL-C and LDL-C, with non-HDL-C listed first to emphasize its primary importance (Table 2). The National Institute for Health and Care Excellence (NICE) recommends the use of non-HDL-C, rather than LDL-C, and sets a >40% reduction in non-HDL-C as the primary therapy goal.⁵⁶ Percentage reduction of LDL-C is used only to group statins into 3 different intensity levels: low intensity if the produced LDL-C reduction is 20% to 30%; medium intensity if the reduction is 31% to 40%; and high intensity if the reduction is >40%. According to NICE, individuals who have a 10-year risk of developing CVD ≥10% (estimated with the QRISK2 algorithm [a registered trademark of the University of Nottingham and EMIS] assessment tool) should be initiated with atorvastatin 20 mg for primary prevention. Statin treatment in people with CVD should be started with atorvastatin 80 mg for secondary prevention. Individuals with non-HDL-C concentration ≥7.5 mmol/L are advised to be assessed by a specialist.

In contrast to the other 7 sets of guidelines and recommendations, the ACC/AHA guidelines do not have lipid treatment goals.⁵¹ Rather, the ACC/AHA guidelines provide expectations regarding the percent LDL-C reduction with moderate- or high-intensity statins. A stated justification is that the use of LDL-C goals, and other risk assessment tools has the potential to undertreat some patients at high ASCVD risk,^{107–109} although some analyses suggest that most of the increased statin usage would be among older individuals without ASCVD.¹¹⁰ For individuals who require a 30% to <50% reduction in LDL-C levels, moderate-intensity statin therapy is used. High-intensity statin therapy is initiated in individuals to achieve LDL-C reductions of ≥50%. Instead of setting lipid treatment goals, ACC/AHA guidelines identify 4 major statin benefit groups that include individuals with: (1) clinical ASCVD, (2) primary elevations of LDL-C ≥190 mg/dL, (3) diabetes aged 40 to 75 years with LDL-C of 70 to 189 mg/dL and without clinical ASCVD, or (4) without clinical ASCVD or diabetes with LDL-C 70

to 189 mg/dL and estimated 10-year ASCVD risk $\geq 7.5\%$.⁵¹ In this case, the 10-year risk of ASCVD is determined from the Pooled Cohort Equations.

Author perspective

Recommendations for the 4 statin benefit groups described previously are among the most practical and clinically useful recommendations of the ACC/AHA guidelines. One of the listed statin benefit groups is individuals with untreated primary LDL-C levels ≥ 190 mg/dL. Within the ACC/AHA Guideline document, the National Heart and Lung Blood Institute Grade for this recommendation is "Expert Opinion." Thus, among the most sentinel and clinically useful recommendations made by the ACC/AHA Guidelines, allowance is given for Expert Opinion for clinical recommendations, presumably based on interpretation of available data and clinical judgment.

Conclusion and Summary

Part 2 of this review examined 8 illustrative, international lipid guidelines, and recommendations as they pertain to the goal of lipid-altering therapy in reducing ASCVD risk. These lipid guidelines and recommendations share a high degree of harmonization with respect to emphasizing the importance of treating LDL-C (primary target) and achieving lower LDL-C goals, whether through the approach of managing to an explicit goal, achieving a percentage reduction, or administering to treatment intensity based on the evidence of benefits. With the exception of the treatment-intensity approach recommended in the 2013 ACC/AHA⁵¹ and NICE⁵⁶ guidelines, all other assessed guidelines (including 2 US-based guidelines) suggest treatment to specific LDL-C goals (based on risk assessment). Despite some differences, the overall message is that cholesterol lowering, especially with statins, is beneficial in reducing ASCVD risk. Ultimately, the decision as to which guideline or recommendation the clinician chooses to follow should be based on which best fits the priorities of the patient.

Acknowledgment

The author would like to thank Dr. Carl Orringer for his review and helpful discussions in the development of this article.

Financial disclosure

Writing and editorial support for this article was provided by MicroMass Communications, Inc, with funding from Regeneron Pharmaceuticals, Inc, Tarrytown, NY, and Sanofi US, Bridgewater, NJ. Support for this educational article was provided by Regeneron Pharmaceuticals, Inc and Sanofi US.

The author discloses that he has received research grants from Arena Pharmaceuticals; Boehringer Ingelheim; Cargill; GlaxoSmithKline; Novo Nordisk; Orexigen Therapeutics; Shionogi; Takeda; Stratum Nutrition; California Raisin Board; Esperion; Essentialis; Forest; Gilead Sciences; Given; Hoffman-La Roche; Home Access; Novartis; Omthera; Pfizer; Trygg Pharmaceuticals; TWI Bio; Xoma; Ardea Inc; High Point Pharmaceuticals, LLC; Micropharma Limited; TransTech Pharma, Inc; TIMI; Pozen; Regeneron; and Elcelyx. He has received honoraria/research grants from Amarin; Amgen; AstraZeneca; Bristol-Myers Squibb; Catabasis; Daiichi-Sankyo; Eisai; Merck; Vivus; Zeomedex; and WPU.

References

1. Sharrett AR, Ballantyne CM, Coady SA, et al. Coronary heart disease prediction from lipoprotein cholesterol levels, triglycerides, lipoprotein(a), apolipoproteins A-I and B, and HDL density subfractions: the Atherosclerosis Risk in Communities (ARIC) Study. *Circulation*. 2001;104:1108–1113.
2. Cholesterol Treatment Trialists' (CTT) Collaboration. Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170,000 participants in 26 randomised trials. *Lancet*. 2010;376:1670–1681.
3. Cholesterol Treatment Trialists' (CTT) Collaborators. The effects of lowering LDL cholesterol with statin therapy in people at low risk of vascular disease: meta-analysis of individual data from 27 randomised trials. *Lancet*. 2012;380:581–590.
4. Taylor F, Huffman MD, Macedo AF, et al. Statins for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev*. 2013;1:CD004816.
5. Cholesterol Treatment Trialists' (CTT) Collaborators. Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90,056 participants in 14 randomised trials of statins. *Lancet*. 2005;366:1267–1278.
6. Frick MH, Elo O, Haapa K, et al. Helsinki Heart Study: primary-prevention trial with gemfibrozil in middle-aged men with dyslipidemia. Safety of treatment, changes in risk factors, and incidence of coronary heart disease. *N Engl J Med*. 1987;317:1237–1245.
7. Rubins HB, Robins SJ, Collins D, et al. Veterans Affairs High Density Lipoprotein Cholesterol Intervention Trial Study Group. Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol. *N Engl J Med*. 1999;341:410–418.
8. Bezafibrate Infarction Prevention (BIP) Study Group. Secondary prevention by raising HDL cholesterol and reducing triglycerides in patients with coronary artery disease. *Circulation*. 2000;102:21–27.
9. FIELD Study Investigators. Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): randomised controlled trial. *Lancet*. 2005;366:1849–1861.
10. Jun M, Foote C, Lv J, et al. Effects of fibrates on cardiovascular outcomes: a systematic review and meta-analysis. *Lancet*. 2010;375:1875–1884.
11. AIM-HIGH Investigators. Niacin in patients with low HDL cholesterol levels receiving intensive statin therapy. *N Engl J Med*. 2011;365:2255–2267.
12. HPS2-THRIVE Collaborative Group. Effects of extended-release niacin with laropiprant in high-risk patients. *N Engl J Med*. 2014;371:203–212.
13. Canner PL, Berge KG, Wenger NK, et al. Fifteen year mortality in Coronary Drug Project patients: long-term benefit with niacin. *J Am Coll Cardiol*. 1986;8:1245–1255.

14. Bays HE, Goldberg RB. The 'forgotten' bile acid sequestrants: is now a good time to remember? *Am J Ther*. 2007;14:567–580.
15. Cannon CP, Blazing MA, Giugliano RP, et al. IMPROVE-IT Investigators. Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med*. 2015;372:2387–2397.
16. Robinson JG, Smith B, Maheshwari N, Schrott H. Pleiotropic effects of statins: benefit beyond cholesterol reduction? A meta-regression analysis. *J Am Coll Cardiol*. 2005;46:1855–1862.
17. dos Santos RL, da Silva FB, Ribeiro RF Jr., Stefanon I. Sex hormones in the cardiovascular system. *Horm Mol Biol Clin Invest*. 2014;18:89–103.
18. Gurney EP, Nachtigall MJ, Nachtigall LE, Naftolin F. The Women's Health Initiative trial and related studies: 10 years later: a clinician's view. *J Steroid Biochem Mol Biol*. 2014;142:4–11.
19. Bays H, McKenney J, Davidson M. Torcetrapib/atorvastatin combination therapy. *Expert Rev Cardiovasc Ther*. 2005;3:789–820.
20. Johns DG, Duffy J, Fisher T, Hubbard BK, Forrest MJ. On- and off-target pharmacology of torcetrapib: current understanding and implications for the structure activity relationships (SAR), discovery and development of cholesteryl ester-transfer protein (CETP) inhibitors. *Drugs*. 2012;72:491–507.
21. Myocardial Infarction Genetics Consortium Investigators. Inactivating mutations in NPC1L1 and protection from coronary heart disease. *N Engl J Med*. 2014;371:2072–2082.
22. Abifadel M, Rabès JP, Devillers M, et al. Mutations and polymorphisms in the proprotein convertase subtilisin kexin 9 (PCSK9) gene in cholesterol metabolism and disease. *Hum Mutat*. 2009;30:520–529.
23. Martin SS, Blumenthal RS, Miller M. LDL cholesterol: the lower the better. *Med Clin North Am*. 2012;96:13–26.
24. Ramjee V, Eapen DJ, Sperling LS. Optimal lipid targets for the new era of cardiovascular prevention. *Ann N Y Acad Sci*. 2012;1254:106–114.
25. LaRosa JC, Grundy SM, Kastelein JJP, et al. Safety and efficacy of atorvastatin-induced very low-density lipoprotein cholesterol levels in patients with coronary heart disease (a post hoc analysis of the treating to new targets [TNT] study). *Am J Cardiol*. 2007;100:747–752.
26. Wiviott SD, Cannon CP, Morrow DA, et al. Can low-density lipoprotein be too low? The safety and efficacy of achieving very low low-density lipoprotein with intensive statin therapy: a PROVE IT-TIMI 22 substudy. *J Am Coll Cardiol*. 2005;46:1411–1416.
27. Hsia J, MacFadyen JG, Monyak J, Ridker PM. Cardiovascular event reduction and adverse events among subjects attaining low-density lipoprotein cholesterol <50 mg/dl with rosuvastatin. The JUPITER trial (Justification for the Use of Statins in Prevention: an Intervention Trial Evaluating Rosuvastatin). *J Am Coll Cardiol*. 2011;57:1666–1675.
28. Ridker PM, Danielson E, Fonseca FA, et al. JUPITER Study Group. Rosuvastatin to prevent vascular events in men and women with elevated C-reactive protein. *N Engl J Med*. 2008;359:2195–2207.
29. Wiviott SD, Mohanavelu S, Raichlen JS, Cain VA, Nissen SE, Libby P. Safety and efficacy of achieving very low low-density lipoprotein cholesterol levels with rosuvastatin 40 mg daily (from the ASTEROID Study). *Am J Cardiol*. 2009;104:29–35.
30. Robinson JG, Farnier M, Krempf M, et al. ODYSSEY LONG TERM Investigators. Efficacy and safety of alirocumab in reducing lipids and cardiovascular events. *N Engl J Med*. 2015;372:1489–1499.
31. US Food and Drug Administration, Center for Drug Evaluation and Research. The Endocrinologic and Metabolic Drugs Advisory Committee Meeting Briefing Document BLA 125559. Praluent (alirocumab) injection. 2015.
32. Sabatine MS, Giugliano RP, Wiviott SD, et al. Open-Label Study of Long-Term Evaluation Against LDL Cholesterol (OSLER) Investigators. Efficacy and safety of evolocumab in reducing lipids and cardiovascular events. *N Engl J Med*. 2015;372:1500–1509.
33. Ballantyne CM, Neutel J, Cropp A, et al. Results of bococizumab, a monoclonal antibody against proprotein convertase subtilisin/kexin type 9, from a randomized, placebo-controlled, dose-ranging study in statin-treated subjects with hypercholesterolemia. *Am J Cardiol*. 2015;115:1212–1221.
34. Welty FK. Hypobetalipoproteinemia and abetalipoproteinemia. *Curr Opin Lipidol*. 2014;25:161–168.
35. Hooper AJ, Marais AD, Tanyanyiwa DM, Burnett JR. The C679X mutation in PCSK9 is present and lowers blood cholesterol in a Southern African population. *Atherosclerosis*. 2007;193:445–448.
36. Zhao Z, Tuakli-Wosornu Y, Lagace TA, et al. Molecular characterization of loss-of-function mutations in PCSK9 and identification of a compound heterozygote. *Am J Hum Genet*. 2006;79:514–523.
37. Study of the Effectiveness of Additional Reductions in Cholesterol Homocysteine Collaborative Group. Intensive lowering of LDL cholesterol with 80 mg versus 20 mg simvastatin daily in 12,064 survivors of myocardial infarction: a double-blind randomised trial. *Lancet*. 2010;376:1658–1669.
38. Cannon CP, Braunwald E, McCabe CH, et al. Pravastatin or Atorvastatin Evaluation and Infection Therapy Thrombolysis in Myocardial Infarction 22 Investigators. Intensive versus moderate lipid lowering with statins after acute coronary syndromes. *N Engl J Med*. 2004;350:1495–1504.
39. de Lemos JA, Blazing MA, Wiviott SD, et al. A to Z Investigators. Early intensive vs a delayed conservative simvastatin strategy in patients with acute coronary syndromes: phase Z of the A to Z trial. *JAMA*. 2004;292:1307–1316.
40. LaRosa JC, Grundy SM, Waters DD, et al. Treating to New Targets (TNT) Investigators. Intensive lipid lowering with atorvastatin in patients with stable coronary disease. *N Engl J Med*. 2005;352:1425–1435.
41. Pedersen TR, Faergeman O, Kastelein JJ, et al. Incremental Decrease in End Points Through Aggressive Lipid Lowering (IDEAL) Study Group. High-dose atorvastatin vs usual-dose simvastatin for secondary prevention after myocardial infarction: the IDEAL study: a randomized controlled trial. *JAMA*. 2005;294:2437–2445.
42. Bays HE, Aversa M, Majul C, et al. Efficacy and safety of ezetimibe added to atorvastatin versus atorvastatin uptitration or switching to rosuvastatin in patients with primary hypercholesterolemia. *Am J Cardiol*. 2013;112:1885–1895.
43. Bays HE, Davidson MH, Massaad R, et al. Safety and efficacy of ezetimibe added on to rosuvastatin 5 or 10 mg versus up-titration of rosuvastatin in patients with hypercholesterolemia (the ACTE Study). *Am J Cardiol*. 2011;108:523–530.
44. Bays H, Farnier M, Gaudet D, et al. Efficacy and safety of combining alirocumab with atorvastatin or rosuvastatin versus statin intensification or adding ezetimibe in high cardiovascular risk patients: ODYSSEY OPTIONS I and II [abstract]. *Circulation*. 2014;130:2118–2119.
45. Scandinavian Simvastatin Survival Study Group. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet*. 1994;344:1383–1389.
46. Nakamura H, Arakawa K, Itakura H, et al. MEGA Study Group. Primary prevention of cardiovascular disease with pravastatin in Japan (MEGA Study): a prospective randomised controlled trial. *Lancet*. 2006;368:1155–1163.
47. Downs JR, Clearfield M, Weis S, et al. AFCAPS/TexCAPS Research Group. Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels: results of AFCAPS/TexCAPS. Air Force/Texas Coronary Atherosclerosis Prevention Study. *JAMA*. 1998;279:1615–1622.
48. Koren MJ, Hunninghake D, ALLIANCE Investigators. Clinical outcomes in managed-care patients with coronary heart disease treated aggressively in lipid-lowering disease management clinics: the alliance study. *J Am Coll Cardiol*. 2004;44:1772–1779.
49. Peterson ED, Ho PM, Barton M, et al. ACC/AHA/AACVPR/AAFP/ANA concepts for clinician-patient shared accountability in performance measures: a report of the American College of

- Cardiology/American Heart Association Task Force on Performance Measures. *J Am Coll Cardiol*. 2014;64:2133–2145.
50. Ray KK, Kastelein JJP, Boekholdt SM, et al. The ACC/AHA 2013 guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular disease risk in adults: the good the bad and the uncertain: a comparison with ESC/EAS guidelines for the management of dyslipidaemias 2011. *Eur Heart J*. 2014;35:960–968.
 51. Stone NJ, Robinson JG, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2014;63:2889–2934.
 52. Jacobson TA, Ito MK, Maki KC, et al. National Lipid Association recommendations for patient-centered management of dyslipidemia: part 1—full report. *J Clin Lipidols*. 2015;9:129–169.
 53. Shrank WH, Barlow JF, Brennan TA. New therapies in the treatment of high cholesterol: an argument to return to goal-based lipid guidelines. *JAMA*. 2015;314:1443–1444.
 54. Boekholdt SM, Arsenault BJ, Mora S, et al. Association of LDL cholesterol, non-HDL cholesterol, and apolipoprotein B levels with risk of cardiovascular events among patients treated with statins: a meta-analysis. *JAMA*. 2012;307:1302–1309.
 55. Kastelein JJP, van der Steeg WA, Holme I, et al, TNT and IDEAL Study Groups. Lipids, apolipoproteins, and their ratios in relation to cardiovascular events with statin treatment. *Circulation*. 2008;117:3002–3009.
 56. NICE clinical guideline 181 Lipid modification: cardiovascular risk assessment and the modification of blood lipids for the primary and secondary prevention of cardiovascular disease. National Institute for Health and Care Excellence; 2014.
 57. Lipid modification: cardiovascular risk assessment and the modification of blood lipids for the primary and secondary prevention of cardiovascular disease Clinical guideline: methods, evidence and recommendations; NICE clinical guideline CG181. National Clinical Guideline Centre; 2014.
 58. Schwertz DW, Badellino KO. High-dose statin therapy for secondary prevention of stroke: stroke prevention by aggressive reduction in cholesterol levels study review. *J Cardiovasc Nurs*. 2008;23:8–13.
 59. Deedwania P, Stone PH, Bairey Merz CN, et al. Effects of intensive versus moderate lipid-lowering therapy on myocardial ischemia in older patients with coronary heart disease: results of the Study Assessing Goals in the Elderly (SAGE). *Circulation*. 2007;115:700–707.
 60. Nicholls SJ, Ballantyne CM, Barter PJ, et al. Effect of two intensive statin regimens on progression of coronary disease. *N Engl J Med*. 2011;365:2078–2087.
 61. Nissen SE, Tuzcu EM, Schoenhagen P, et al, REVERSAL Investigators. Effect of intensive compared with moderate lipid-lowering therapy on progression of coronary atherosclerosis: a randomized controlled trial. *JAMA*. 2004;291:1071–1080.
 62. Expert Dyslipidemia Panel of the International Atherosclerosis Society Panel. An International Atherosclerosis Society Position Paper: global recommendations for the management of dyslipidemia—full report. *J Clin Lipidols*. 2014;8:29–60.
 63. Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of cholesterol lowering with simvastatin in 20,536 high-risk individuals: a randomised placebo-controlled trial. *Lancet*. 2002;360:7–22.
 64. Cannon CP, Steinberg BA, Murphy SA, Mega JL, Braunwald E. Meta-analysis of cardiovascular outcomes trials comparing intensive versus moderate statin therapy. *J Am Coll Cardiol*. 2006;48:438–445.
 65. Martin MJ, Hulley SB, Browner WS, Kuller LH, Wentworth D. Serum cholesterol, blood pressure, and mortality: implications from a cohort of 361,662 men. *Lancet*. 1986;2:933–936.
 66. Cohen JC, Boerwinkle E, Mosley TH Jr., Hobbs HH. Sequence variations in PCSK9, low LDL, and protection against coronary heart disease. *N Engl J Med*. 2006;354:1264–1272.
 67. Kathiresan S, Myocardial Infarction Genetics Consortium. A PCSK9 missense variant associated with a reduced risk of early-onset myocardial infarction [letter]. *N Engl J Med*. 2008;358:2299–2300.
 68. Ference BA, Yoo W, Alesh I, et al. Effect of long-term exposure to lower low-density lipoprotein cholesterol beginning early in life on the risk of coronary heart disease: a Mendelian randomization analysis. *J Am Coll Cardiol*. 2012;60:2631–2639.
 69. Sever PS, Dahlof B, Poulter NR, et al, ASCOT Investigators. Prevention of coronary and stroke events with atorvastatin in hypertensive patients who have average or lower-than-average cholesterol concentrations, in the Anglo-Scandinavian Cardiac Outcomes Trial—Lipid Lowering Arm (ASCOT-LLA): a multicentre randomised controlled trial. *Lancet*. 2003;361:1149–1158.
 70. Shepherd J, Cobbe SM, Ford I, et al, West of Scotland Coronary Prevention Study Group. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. West of Scotland Coronary Prevention Study Group. *N Engl J Med*. 1995;333:1301–1307.
 71. Wanner C, Krane V, Marz W, et al, German Diabetes and Dialysis Study Investigators. Atorvastatin in patients with type 2 diabetes mellitus undergoing hemodialysis. *N Engl J Med*. 2005;353:238–248.
 72. Wiviott SD, de Lemos JA, Cannon CP, et al. A tale of two trials: a comparison of the post-acute coronary syndrome lipid-lowering trials A to Z and PROVE IT-TIMI 22. *Circulation*. 2006;113:1406–1414.
 73. Accord Study Group. Effects of combination lipid therapy in type 2 diabetes mellitus. *N Engl J Med*. 2010;362:1563–1574.
 74. Knopp RH, d'Emden M, Smilde JG, Pocock SJ, ASPEN Study Group. Efficacy and safety of atorvastatin in the prevention of cardiovascular end points in subjects with type 2 diabetes: the Atorvastatin Study for Prevention of Coronary Heart Disease Endpoints in non-insulin-dependent diabetes mellitus (ASPEN). *Diabetes Care*. 2006;29:1478–1485.
 75. Fellström BC, Jardine AG, Schmieder RE, et al, AURORA Study Group. Rosuvastatin and cardiovascular events in patients undergoing hemodialysis. *N Engl J Med*. 2009;360:1395–1407.
 76. Sacks FM, Pfeffer MA, Moye LA, et al, Cholesterol and Recurrent Events Trial Investigators. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. *N Engl J Med*. 1996;335:1001–1009.
 77. Kjekshus J, Apetrei E, Barrios V, et al, CORONA Group. Rosuvastatin in older patients with systolic heart failure. *N Engl J Med*. 2007;357:2248–2261.
 78. Athyros VG, Papageorgiou AA, Mercouris BR, et al. Treatment with atorvastatin to the National Cholesterol Educational Program goal versus 'usual' care in secondary coronary heart disease prevention. The GREEK Atorvastatin and Coronary-heart-disease Evaluation (GREACE) study. *Curr Med Res Opin*. 2002;18:220–228.
 79. Brown BG, Zhao X-Q, Chait A, et al. Simvastatin and niacin, antioxidant vitamins, or the combination for the prevention of coronary disease. *N Engl J Med*. 2001;345:1583–1592.
 80. Long-Term Intervention with Pravastatin in Ischaemic Disease (LIPID) Study Group. Prevention of cardiovascular events and death with pravastatin in patients with coronary heart disease and a broad range of initial cholesterol levels. *N Engl J Med*. 1998;339:1349–1357.
 81. Serruys PW, de Feyter P, Macaya C, et al, Lescot Intervention Prevention Study (LIPS) Investigators. Fluvastatin for prevention of cardiac events following successful first percutaneous coronary intervention: a randomized controlled trial. *JAMA*. 2002;287:3215–3222.
 82. Schwartz GG, Olsson AG, Ezekowitz MD, et al, Myocardial Ischemia Reduction with Aggressive Cholesterol Lowering (MIRACL) Study Investigators. Effects of atorvastatin on early recurrent ischemic events in acute coronary syndromes: the MIRACL study: a randomized controlled trial. *JAMA*. 2001;285:1711–1718.
 83. Sakamoto T, Kojima S, Ogawa H, et al, Multicenter Study for Aggressive Lipid-Lowering Strategy by HMG-CoA Reductase Inhibitors in Patients With Myocardial Infarction Investigators. Effects of

- early statin treatment on symptomatic heart failure and ischemic events after acute myocardial infarction in Japanese. *Am J Cardiol*. 2006;97:1165–1171.
84. Colhoun HM, Betteridge DJ, Durrington PN, et al, CARDS Investigators. Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): multicentre randomised placebo-controlled trial. *Lancet*. 2004;364:685–696.
 85. Anderson TJ, Grégoire J, Hegele RA, et al. 2012 update of the Canadian Cardiovascular Society guidelines for the diagnosis and treatment of dyslipidemia for the prevention of cardiovascular disease in the adult. *Can J Cardiol*. 2013;29:151–167.
 86. European Association for Cardiovascular Prevention & Rehabilitation. ESC/EAS guidelines for the management of dyslipidaemias: the Task Force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS). *Eur Heart J*. 2011;32:1769–1818.
 87. Grundy SM, Cleeman JJ, Merz CN, et al. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. *Circulation*. 2004;110:227–239.
 88. Baigent C, Landray MJ, Reith C, et al, SHARP Investigators. The effects of lowering LDL cholesterol with simvastatin plus ezetimibe in patients with chronic kidney disease (Study of Heart and Renal Protection): a randomised placebo-controlled trial. *Lancet*. 2011;377:2181–2192.
 89. Teramoto T, Sasaki J, Ishibashi S, et al. Executive summary of the Japan Atherosclerosis Society (JAS) guidelines for the diagnosis and prevention of atherosclerotic cardiovascular diseases in Japan—2012 version. *J Atheroscler Thromb*. 2013;20:850–860.
 90. Thompson PL, Meredith I, Amerena J, et al. Effect of pravastatin compared with placebo initiated within 24 hours of onset of acute myocardial infarction or unstable angina: the Pravastatin in Acute Coronary Treatment (PACT) trial. *Am Heart J*. 2004;148:e2.
 91. Stenestrand U, Wallentin L, Swedish Register of Cardiac Intensive Care (RIKS-HIA). Early statin treatment following acute myocardial infarction and 1-year survival. *JAMA*. 2001;285:430–436.
 92. Dohi T, Miyauchi K, Okazaki S, et al. Early intensive statin treatment for six months improves long-term clinical outcomes in patients with acute coronary syndrome (Extended-ESTABLISH trial): a follow-up study. *Atherosclerosis*. 2010;210:497–502.
 93. Hultén E, Jackson JL, Douglas K, George S, Villines TC. The effect of early, intensive statin therapy on acute coronary syndrome: a meta-analysis of randomized controlled trials. *Arch Intern Med*. 2006;166:1814–1821.
 94. Hamazaki T, Okuyama H, Ogushi Y, Hama R. Cholesterol issues in Japan—why are the goals of cholesterol levels set so low? *Ann Nutr Metab*. 2013;62:32–36.
 95. Brugts JJ, Yetgin T, Hoeks SE, et al. The benefits of statins in people without established cardiovascular disease but with cardiovascular risk factors: meta-analysis of randomised controlled trials. *BMJ*. 2009;338:b2376.
 96. Mills EJ, Rachlis B, Wu P, Devereaux PJ, Arora P, Perri D. Primary prevention of cardiovascular mortality and events with statin treatments: a network meta-analysis involving more than 65,000 patients. *J Am Coll Cardiol*. 2008;52:1769–1781.
 97. Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III) final report. *Circulation*. 2002;106:3143–3421.
 98. Shepherd J, Blauw GJ, Murphy MB, et al, PROSPER Study Group. Pravastatin in elderly individuals at risk of vascular disease (PROSPER): a randomised controlled trial. *Lancet*. 2002;360:1623–1630.
 99. Pitt B, Waters D, Brown WV, et al, Atorvastatin Versus Revascularization Treatment Investigators. Aggressive lipid-lowering therapy compared with angioplasty in stable coronary artery disease. *N Engl J Med*. 1999;341:70–76.
 100. Blankenhorn DH, Azen SP, Krams DM, et al, MARS Research Group. Coronary angiographic changes with lovastatin therapy. The Monitored Atherosclerosis Regression Study (MARS). *Ann Intern Med*. 1993;119:969–976.
 101. Aengevaeren WR, Kroon AA, Stalenhoef AF, Uijen GJ, van der Werf T. Low density lipoprotein apheresis improves regional myocardial perfusion in patients with hypercholesterolemia and extensive coronary artery disease. LDL-Apheresis Atherosclerosis Regression Study (LAARS). *J Am Coll Cardiol*. 1996;28:1696–1704.
 102. Post Coronary Artery Bypass Graft Trial Investigators. The effect of aggressive lowering of low-density lipoprotein cholesterol levels and low-dose anticoagulation on obstructive changes in saphenous-vein coronary-artery bypass grafts. *N Engl J Med*. 1997;336:153–162.
 103. Brown BG, Bardsley J, Poulin D, et al. Moderate dose, three-drug therapy with niacin, lovastatin, and colestipol to reduce low-density lipoprotein cholesterol <100 mg/dl in patients with hyperlipidemia and coronary artery disease. *Am J Cardiol*. 1997;80:111–115.
 104. ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. Major outcomes in moderately hypercholesterolemic, hypertensive patients randomized to pravastatin vs usual care: The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT-LLT). *JAMA*. 2002;288:2998–3007.
 105. LaRosa JC, He J, Vupputuri S. Effect of statins on risk of coronary disease: a meta-analysis of randomized controlled trials. *JAMA*. 1999;282:2340–2346.
 106. Puri R, Nissen SE, Shao M, et al. Impact of baseline lipoprotein and C-reactive protein levels on coronary atheroma regression following high-intensity statin therapy. *Am J Cardiol*. 2014;114:1465–1472.
 107. Johnson KM, Dowe DA. Accuracy of statin assignment using the 2013 AHA/ACC Cholesterol Guideline versus the 2001 NCEP ATP III guideline: correlation with atherosclerotic plaque imaging. *J Am Coll Cardiol*. 2014;64:910–919.
 108. Karmali KN, Goff DC Jr., Ning H, Lloyd-Jones DM. A systematic examination of the 2013 ACC/AHA pooled cohort risk assessment tool for atherosclerotic cardiovascular disease. *J Am Coll Cardiol*. 2014;64:959–968.
 109. Paixao AR, Ayers CR, Berry JD, de Lemos JA, Khera A. Atherosclerotic cardiovascular disease prevention: a comparison between the third Adult Treatment Panel and the new 2013 treatment of blood cholesterol guidelines. *Circ Cardiovasc Qual Outcomes*. 2014;7:778–779.
 110. Pencina MJ, Navar-Boggan AM, D'Agostino RB Sr., et al. Application of new cholesterol guidelines to a population-based sample. *N Engl J Med*. 2014;370:1422–1431.