ELSEVIER

Contents lists available at ScienceDirect

# Forensic Science International: Genetics

journal homepage: www.elsevier.com/locate/fsig



## Letter to the Editor

## STR data for the AmpFISTR Identifiler from Dobruja region (SE Romania)

ARTICLE INFO

Keywords: DNA typing Short Tandem Repeats (STRs) AmpFISTR Identifiler Population data Dobruja

#### ABSTRACT

Allele frequencies for 15 STR loci included in AmpFISTR Identifiler kit (D8S1179, D21S11, D7S820, CSF1PO, D3S1358, TH01, D13S317, D16S539, D2S1338, D19S433, vWA, TPOX, D18S51, D5S818, and FGA) were determined in a sample of 569 unrelated individuals living in the region of Dobruja (SE Romania). No deviations from Hardy–Weinberg equilibrium were observed. Genetic parameters of forensic interest were calculated and comparison with geographically nearby populations was performed.

© 2008 Elsevier Ireland Ltd. All rights reserved.

Dear Editor,

We determined the allele frequencies for 15 STR loci included in AmpF/STR Identifiler kit (D8S1179, D21S11, D7S820, CSF1PO, D3S1358, TH01, D13S317, D16S539, D2S1338, D19S433, vWA, TPOX, D18S51, D5S818, and FGA) in a sample of 569 unrelated individuals living in the region of Dobruja (SE Romania – Tulcea and Constanta counties).

The territory of Dobruja (Dobrogea in Romanian) is an informal and also a historical region located between the lower Danube River and the Black Sea including the Danube Delta, Romanian coast and the northernmost part of the Bulgarian coast, and comprises Northern Dobruja, which is part of Romania, and Southern Dobruja, which belongs to Bulgaria. In the past this region had been conquered by Greeks, Romans, Tatars, Turks and Slavs for its economical and strategic value. The resulted population is an ethno – linguistic mixture in which Romanians are the majority [1].

DNA from buccal swabs was extracted, 35 samples by Chelex 100 method [2] and 532 samples by an adapted version of AGOWA mag DNA Isolation Kit Sputum for Freedom Evo 150 Liquid Handling Platform [3]. Approximately 1 ng target DNA was amplified using the commercial typing kit, AmpFISTR IdentifilerTM PCR Amplification Kit (Applied Biosystems, Foster City, CA, USA), according to the manufacturer's instructions. DNA Typing was done using an ABI Prism 3100 Genetic Analyzer and reference sequenced ladders (Applied Biosystems). Regarding analysis of data, Hardy-Weinberg equilibrium (P), expected heterozygosity (He), observed heterozygosity (Ho) and population differentiation tests were carried out with the Arlequin Software Version 3.1.1 [4]. Matching probability (MP), power of discrimination (PD), polymorphism information content (PIC), probability of exclusion (PE) and typical paternity index (TPI) were calculated with the Powerstats Version 1.2 (Promega Corp.) [5].

The observed allele frequencies and statistical parameters for forensic testing based on the 15 STR loci in Dobruja population are summarized in see supplementary Table 1. No deviations from Hardy–Weinberg equilibrium were observed. The combined power of discrimination (PD) and the combined power of exclusion (PE) for the 15 studied loci were 0.99999999999999997 and 0.999999. respectively.

Single locus comparisons with available published data on other geographically nearby populations (see supplementary Table 2), revealed significant differences between population from Dobruja and the Romania (Bucharest area) [6] at one locus – D19S433, between populations from Dobruja and Greece [7] at one locus – TPOX, between population from Dobruja and Turkey [8] at two loci – D7S820 and D18S51, and finally between population from Dobruja and Belarus [9] at six loci – TH01, D13S317, D16S539, TPOX, D18S51 and D5S818. Single locus comparisons did not show any statistically significant differences between populations from Dobruja and Italy [10] and also from Dobruja and Serbia [11].

For technical support in the DNA typing step (ABI3100) we would like to thank Daniela Cocioaba and Romica Potorac (Forensic Science Institute, Bucharest, Romania).

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.fsigen.2008.09.009.

## References

- [1] Dobruja, http://en.wikipedia.org/wiki/Dobruja.
- [2] P.S. Walsh, D.A. Metzger, R. Higuchi, Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material, Biotechniques 10 (1991) 506–513.

- [3] F. Stanciu, I.M. Stoian, O.R. Popescu, Evaluation of Freedom Evo 150 with AGOWA sep 9600 for Databasing Purposes, Forensica 2008 – Book of Abstracts, 2008, P17, http://www.dnahunt.ro/fronta/posters/one\_poster.cfm?posterID=4.
- [4] L. Excoffier, G. Laval, S. Schneider, Arlequin ver. 3.0: an integrated software package for population genetics data analysis, Evolutionary Bioinformatics Online 1 (2005) 47–50.
- [5] Powerstats version 1.2, Promega corporation website, http://www.promega.com/geneticidtools/powerstats/.
- [6] L.E. Barbarii, B. Rolf, C. Constantinescu, C. Hohoff, P. Calistru, D. Dermengiu, Allele frequencies of 13 short tandem repeat (STR) loci in the Romanian population, Forensic Science International 141 (2004) 171–174.
- [7] I. Skitsa, A. Salas, M.V. Lareu, A. Carracedó, STR-CODIS typing in Greece, Forensic Science International 137 (2003) 104–106.
- [8] U. Ulkuer, M. Kurtulus-Ulkuer, C. Elma, T. Kesici, S. Menevse, Short tandem repeat (STR) polymorphisms in Turkish population, Journal of Genetics 83 (2) (2004) 197–199
- [9] L.A. Zhivotovsky, V.M. Veremeichyk, A.I. Mikulich, I.G. Udina, L.A. Atramentova, S.A. Kotova, N.A. Kartel, I.S. Tsybovsky, A comprehensive population survey on the distribution of STR frequencies in Belarus, Forensic Science International 172 (2007) 156–160.
- [10] U. Ricci, I. Sani, L. Giunti, S. Guarducci, S. Coviello, M.L.G. Uzielli, Analysis of 13 tetrameric short tandem repeat loci in a population of Tuscany (Central Italy) performed by means of an automated infrared sequencer, Forensic Science International 125 (2002) 83–85.

[11] I. Veselinovic, M. Kubat, I. Furac, J. Skavic, I. Martinovic Klaric, M. Tasic, Allele frequencies of the 15 AmpFl STR Identifiler loci in the population of Vojvodina Province, Serbia and Montenegro, International Journal of Legal Medicine 118 (2004) 184–186.

Florin Stanciu\*
Oana Raluca Popescu
Ionel Marius Stoian
Forensic Science Institute, 13-15 Stefan cel Mare,
020123, Bucharest, Romania

\*Corresponding author at: Inspectoratul General al Politiei Romane, Institutul de Criminalistica, Sos. Stefan cel Mare, Nr.13-15, Sector 2, 020123, Bucharest, Romania. Tel.: +40 0726 211 280

E-mail address: staflorin@gmail.com (F. Stanciu)

29 August 2008