

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/266264454>

# Synthesis and metabolic studies of $1\alpha,2\alpha,25-$ , $1\alpha,4\alpha,25-$ and $1\alpha,4\beta,25-$ trihydroxyvitamin D<sub>3</sub>

ARTICLE in THE JOURNAL OF STEROID BIOCHEMISTRY AND MOLECULAR BIOLOGY · SEPTEMBER 2014

Impact Factor: 3.63 · DOI: 10.1016/j.jsbmb.2014.09.021 · Source: PubMed

CITATION

1

READS

24

## 11 AUTHORS, INCLUDING:



**Kaori Yasuda**

Toyama Prefectural University

32 PUBLICATIONS 534 CITATIONS

SEE PROFILE



**Kyohei Horie**

Teijin

9 PUBLICATIONS 58 CITATIONS

SEE PROFILE



**Tai C Chen**

Boston University

240 PUBLICATIONS 11,265 CITATIONS

SEE PROFILE




**Atsushi Kittaka**

Teikyo University

163 PUBLICATIONS 1,872 CITATIONS

SEE PROFILE



## Review

# Synthesis and metabolic studies of $1\alpha,2\alpha,25$ -, $1\alpha,4\alpha,25$ - and $1\alpha,4\beta,25$ -trihydroxyvitamin D<sub>3</sub>



Masashi Takano<sup>a</sup>, Daisuke Sawada<sup>a,1</sup>, Kaori Yasuda<sup>b</sup>, Miyu Nishikawa<sup>b</sup>, Akiko Takeuchi<sup>c</sup>, Ken-ichiro Takagi<sup>c</sup>, Kyohei Horie<sup>c</sup>, G. Satyanarayana Reddy<sup>d</sup>, Tai C. Chen<sup>e</sup>, Toshiyuki Sakaki<sup>b</sup>, Atsushi Kittaka<sup>a,\*</sup>

<sup>a</sup> Faculty of Pharmaceutical Sciences, Teikyo University, 2-11-1 Kaga, Itabashi, Tokyo 173-8605, Japan

<sup>b</sup> Faculty of Engineering, Toyama Prefectural University, Imizu, Toyama 939-0398, Japan

<sup>c</sup> Teijin Institute for Bio-medical Research, Teijin Pharma Ltd., Hino, Tokyo 191-8512, Japan

<sup>d</sup> Epimer LLC, North Smithfield, RI 02906, USA

<sup>e</sup> Section of Endocrinology, Diabetes & Nutrition, Boston University School of Medicine, Boston, MA 02118, USA

## ARTICLE INFO

## Article history:

Received 11 July 2014

Received in revised form 17 September 2014

Accepted 22 September 2014

Available online 26 September 2014

## Keywords:

Vitamin D<sub>3</sub>

Trihydroxyvitamin D<sub>3</sub>

Vitamin D receptor

CYP3A4

CYP24A1

UGT

## ABSTRACT

Three different A-ring perhydroxylated trihydroxyvitamin D<sub>3</sub> metabolites were synthesized from their appropriate A-ring precursors and CD-ring for their potential therapeutic applications. We first chemically synthesized  $1\alpha,2\alpha,25$ -trihydroxyvitamin D<sub>3</sub> [ $1\alpha,2\alpha,25(\text{OH})_3\text{D}_3$ ] to study its VDR binding affinity because this metabolite is a product of recombinant human CYP3A4 catalysis when  $2\alpha$ -(3-hydroxypropoxy)- $1\alpha,25$ -dihydroxyvitamin D<sub>3</sub> (O2C3), a more potent vitamin D receptor (VDR) binder than  $1\alpha,25$ -dihydroxyvitamin D<sub>3</sub> [ $1\alpha,25(\text{OH})_2\text{D}_3$ ], is used as the substrate. We found that this metabolite retained 27.3% of the VDR binding affinity compared to  $1\alpha,25(\text{OH})_2\text{D}_3$ . The  $k_{\text{cat}}/K_{\text{m}}$  value of CYP24A1 for  $1\alpha,2\alpha,25(\text{OH})_3\text{D}_3$  is 60% of that for  $1\alpha,25(\text{OH})_2\text{D}_3$ . Since the biological activity and the metabolic fate of a naturally occurring C4-hydroxylated vitamin D<sub>2</sub> metabolite found in the serum of rats treated with pharmacological doses of vitamin D<sub>2</sub> have never been described, we next synthesized  $1\alpha,4\alpha,25$ -trihydroxyvitamin D<sub>3</sub> and its diastereoisomer,  $1\alpha,4\beta,25$ -trihydroxyvitamin D<sub>3</sub>, to study their metabolism and biological activities. Both 4-hydroxylated isomers showed weaker VDR binding affinity than  $1\alpha,25(\text{OH})_2\text{D}_3$ . Although either 4-hydroxylated isomer can be metabolized by CYP24A1 almost at the same level as  $1\alpha,25(\text{OH})_2\text{D}_3$ , their metabolic patterns catalyzed by uridine 5'-diphosphoglucuronosyltransferase (UGT) are different; only the  $4\alpha$ -hydroxylated analog can be metabolized by UGT to produce a glucuronate conjugate. The results provide important information for the synthesis of new novel chemotherapeutic vitamin D analogs which would be less subjective to degradation and therefore more bioavailable than  $1\alpha,25(\text{OH})_2\text{D}_3$ .

This article is part of a Special Issue entitled '17th Vitamin D Workshop'.

© 2014 Elsevier Ltd. All rights reserved.

## Contents

1. Introduction	34
2. Results and discussion	36
3. Conclusion	37
Acknowledgments	37
References	37


## 1. Introduction

The physiologically active metabolite of vitamin D<sub>3</sub>,  $1\alpha,25$ -dihydroxyvitamin D<sub>3</sub> [ $1\alpha,25(\text{OH})_2\text{D}_3$ ], expresses its biological activity through binding to and modulation of the

\* Corresponding author. Tel.: +81 3 3964 8109; fax: +81 3 3964 8117.

E-mail address: [akittaka@pharm.teikyo-u.ac.jp](mailto:akittaka@pharm.teikyo-u.ac.jp) (A. Kittaka).


<sup>1</sup> Present address: Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University, Kitaku, Okayama 700-8530, Japan.




**Fig. 1.** Structures of 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> and its hydroxylated analogs O1C3, O2C3, 1 $\alpha$ ,2 $\alpha$ ,25(OH)<sub>3</sub>D<sub>3</sub>, 1 $\alpha$ ,4 $\alpha$ ,25(OH)<sub>3</sub>D<sub>3</sub>, and 1 $\alpha$ ,4 $\beta$ ,25(OH)<sub>3</sub>D<sub>3</sub>.

vitamin D receptor (VDR) [1]. Previously, we synthesized 2 $\alpha$ -(3-hydroxypropoxy)-1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> (O2C3) which possesses 1.8-times greater binding affinity for VDR than the natural hormone 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> [2–4]. The reason of being the better ligand can be explained by X-ray co-crystallographic analyses of the VDR-O2C3 complex, which shows additional hydrogen bonding between the terminal hydroxy group of the introduced 2 $\alpha$ -side chain of the O2C3 and Arg274 in the ligand binding domain (LBD) of the VDR [5]. Although 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> is inactivated by CYP24A1-dependent catabolism via C-24 and C-23 hydroxylation pathways to calcitroic acid and 26,23-lactone derivatives, respectively [6], it was found that some 2 $\alpha$ -substituted active vitamin D analogs like O2C3 were highly resistant to the CYP24A1 catabolism. The  $k_{\text{cat}}/K_m$  value of O2C3 was only 3% of

that for 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> [7]. Since CYP24A1 is a specific enzyme induced by the VDR-ligand [1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> or its analogs] complex in the target tissue and inactivates 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> and its analogs, CYP24A1-resistant ligands would have longer-term biological effects and bioavailability in the target tissues [8]. On the other hand, CYP3A4 is the most important drug-metabolizing cytochrome P450 enzyme with a broad catalytic spectrum [9]. Although 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> and its analog 2 $\alpha$ -(3-hydroxypropyl)-1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> (O1C3) are not the primary substrates for CYP3A4 [8], we demonstrated that O2C3 was metabolized by the human recombinant CYP3A4 to 1 $\alpha$ ,2 $\alpha$ ,25-trihydroxyvitamin D<sub>3</sub> [1 $\alpha$ ,2 $\alpha$ ,25(OH)<sub>3</sub>D<sub>3</sub>, Fig. 1] [8]. Its 2 $\beta$ -isomer, a known compound, shows potent 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub>-like activities [10], and is one of the eldecaltol (ED-71) metabolites [11]. To study



**Scheme 1.** Synthesis of 1 $\alpha$ ,2 $\alpha$ ,25(OH)<sub>3</sub>D<sub>3</sub>.



**Scheme 2.** Synthesis of  $1\alpha,4\alpha,25(\text{OH})_3\text{D}_3$  and  $1\alpha,4\beta,25(\text{OH})_3\text{D}_3$ .

biological activity of  $1\alpha,2\alpha,25(\text{OH})_3\text{D}_3$ , we chemically synthesized  $1\alpha,2\alpha,25(\text{OH})_3\text{D}_3$  using the Trost Pd-mediated coupling reaction between an A-ring precursor **1** from D-glucose and a known CD-ring bromoolefin **2** (Scheme 1) [12–14].

On the other hand, a number of hydroxylated vitamin D metabolites were found in the circulation of mammals. For example, 4,25-dihydroxyvitamin D<sub>2</sub> was isolated and identified from the serum of rats intoxicated with pharmacological amounts of vitamin D<sub>2</sub> [15]. However, the biological significance of the 4-hydroxy group remains unclear. We therefore synthesized  $1\alpha,4\alpha,25$ -trihydroxyvitamin D<sub>3</sub> [ $1\alpha,4\alpha,25(\text{OH})_3\text{D}_3$ ] and  $1\alpha,4\beta,25$ -trihydroxyvitamin D<sub>3</sub> [ $1\alpha,4\beta,25(\text{OH})_3\text{D}_3$ ] by the similar manner as  $1\alpha,2\alpha,25(\text{OH})_3\text{D}_3$ , and evaluated their hVDR binding affinity as well as their further metabolism by human CYP24A1 and human liver microsomal fraction to compare with those of the natural hormone  $1\alpha,25(\text{OH})_2\text{D}_3$ .

## 2. Results and discussion

Synthesis of  $1\alpha,2\alpha,25(\text{OH})_3\text{D}_3$  was accomplished as shown in Scheme 1. We prepared chiral epoxide from methyl α-D-glucoside [2], which was converted to the enyne A-ring precursor **1** in

11 steps [12]. The CD-ring bromoolefin **2** and enyne **1** were coupled using Pd-catalyst to give the seco-steroidal product [16,17]. Deprotection of O-silyl groups gave the target molecule  $1\alpha,2\alpha,25(\text{OH})_3\text{D}_3$ . The isolated product was re-purified by HPLC to test its biological activity.

The other A-ring hydroxylated vitamin D<sub>3</sub> analogs of  $1\alpha,4\alpha,25(\text{OH})_3\text{D}_3$  and  $1\alpha,4\beta,25(\text{OH})_3\text{D}_3$  were synthesized as shown in Scheme 2 [18]. The same chiral epoxide from D-glucose was converted to aldehyde **3** [19] followed by ethynylation afforded alcohols as a diastereomeric mixture in a 1:3 ratio. Stereochemistry of each alcohol was determined by the modified Mosher's method [19,20]. After manipulation of the protecting groups, enynes **4a** and **4b** were connected with CD-ring bromoolefin **2** using Pd-catalyst to give the coupling products, and TBAF-deprotection followed by HPLC-separation gave the target molecules  $1\alpha,4\alpha,25(\text{OH})_3\text{D}_3$  and  $1\alpha,4\beta,25(\text{OH})_3\text{D}_3$  [18,19].

Next, hVDR binding affinity was performed for the above three new A-ring perhydroxylated vitamin D<sub>3</sub> metabolites, and the results are shown in Table 1. Elcalcitol metabolite  $1\alpha,2\beta,25(\text{OH})_3\text{D}_3$  showed strong VDR binding affinity as compared with O2C3 metabolite  $1\alpha,2\alpha,25(\text{OH})_3\text{D}_3$  [10]. Although the 2-OH group is not a severe obstacle to VDR binding, the 4-OH group might be steric hindrance when this ligand binds to the ligand binding pocket of the VDR [12,18].

Regarding metabolism, O2C3 is resistant to CYP24A1, but its CYP3A4 metabolite  $1\alpha,2\alpha,25(\text{OH})_3\text{D}_3$  can be further metabolized by CYP24A1, similar to  $1\alpha,25(\text{OH})_2\text{D}_3$ . The  $k_{\text{cat}}/K_{\text{m}}$  value of hCYP24A1 for  $1\alpha,2\alpha,25(\text{OH})_3\text{D}_3$  was found to be 60% of that for  $1\alpha,25(\text{OH})_2\text{D}_3$  [8]. On the other hand,  $1\alpha,4\alpha,25(\text{OH})_3\text{D}_3$  and  $1\alpha,4\beta,25(\text{OH})_3\text{D}_3$  are metabolized to several metabolites by CYP24A1-dependent multi-step oxidation pathways similar to that for  $1\alpha,25(\text{OH})_2\text{D}_3$  or its A-ring diastereomers. Interestingly, HPLC profile of the metabolites of  $1\alpha,4\alpha,25(\text{OH})_3\text{D}_3$  is quite similar to that of  $1\alpha,25(\text{OH})_2\text{D}_3$ , whereas, the profile of  $1\alpha,4\beta,25(\text{OH})_3\text{D}_3$  is quite similar to that of 3-epimers of  $1\alpha,25(\text{OH})_2\text{D}_3$  (Fig. 2) [18,21]. After 60 min incubation, the % conversion of  $1\alpha,4\alpha,25(\text{OH})_3\text{D}_3$  and


**Table 1**  
Relative hVDR binding affinity of A-ring perhydroxylated analogs.

Compound	VDR Binding affinity <sup>a</sup>
$1\alpha,25(\text{OH})_3\text{D}_3$	100
O2C3	180 <sup>b</sup>
$1\alpha,2\alpha,25(\text{OH})_3\text{D}_3$	27.3
$1\alpha,2\beta,25(\text{OH})_3\text{D}_3$	110 <sup>c</sup>
$1\alpha,4\alpha,25(\text{OH})_3\text{D}_3$	0.9
$1\alpha,4\beta,25(\text{OH})_3\text{D}_3$	2.9

<sup>a</sup> The potency of  $1\alpha,25(\text{OH})_3\text{D}_3$  is normalised to 100.

<sup>b</sup> Ref. [2].

<sup>c</sup> Ref. [10].



**Fig. 2.** HPLC profiles of 1 $\alpha$ ,4 $\alpha$ ,25- (A) and 1 $\alpha$ ,4 $\beta$ ,25-trihydroxyvitamin D<sub>3</sub> (B) and their metabolites by human CYP24A1. The peaks with arrow heads show putative metabolites by human CYP24A1.

1 $\alpha$ ,4 $\beta$ ,25(OH)<sub>3</sub>D<sub>3</sub> are about 45% and 44%, respectively, which are nearly the same as those for 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub>. Both 4-hydroxyvitamin D<sub>3</sub> analogs were treated with human liver microsomal fraction containing drug-metabolizing enzymes, and no metabolite from either analog was detected. The results suggest that both analogs are poor substrates for human hepatic cytochrome P450s. However, addition of UDP-glucuronic acid produced a metabolite (glucuronide) of 1 $\alpha$ ,4 $\alpha$ ,25(OH)<sub>3</sub>D<sub>3</sub> whereas no metabolite of 1 $\alpha$ ,4 $\beta$ ,25(OH)<sub>3</sub>D<sub>3</sub> was observed [18]. Since  $\beta$ -glucuronidase treatment converted the metabolite of 1 $\alpha$ ,4 $\alpha$ ,25(OH)<sub>3</sub>D<sub>3</sub> back to 1 $\alpha$ ,4 $\alpha$ ,25(OH)<sub>3</sub>D<sub>3</sub> itself, this suggests that the metabolite is a glucuronide of 1 $\alpha$ ,4 $\alpha$ ,25(OH)<sub>3</sub>D<sub>3</sub>. Neither 1 $\alpha$ ,4 $\beta$ ,25(OH)<sub>3</sub>D<sub>3</sub> nor 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> was converted to its glucuronide. Therefore, the 4 $\alpha$ -OH group of 1 $\alpha$ ,4 $\alpha$ ,25(OH)<sub>3</sub>D<sub>3</sub> appeared to be glucuronidated by certain hepatic UGT(s) [18].

### 3. Conclusion

We synthesized three new A-ring perhydroxylated vitamin D<sub>3</sub> metabolites, 1 $\alpha$ ,2 $\alpha$ ,25(OH)<sub>3</sub>D<sub>3</sub>, the major O2C3 metabolite catalyzed by CYP3A4, and 1 $\alpha$ ,4 $\alpha$ ,25(OH)<sub>3</sub>D<sub>3</sub> and 1 $\alpha$ ,4 $\beta$ ,25(OH)<sub>3</sub>D<sub>3</sub>, the tentative 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> related metabolites. The position and stereochemistry of the synthetically introduced OH group to the A-ring influence hVDR binding affinity and their further metabolism by CYP enzymes. These findings would be useful for creating new novel potential chemotherapeutic ligands for VDR.

### Acknowledgments

This work was supported in part by a Grant-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology (No. 25860011 to M.T.) as well as Grants-in-Aid from the Japan Society for the Promotion of Science (No. 23590015 to D.S. and 24590021 to A.K.).

### References

- [1] D. Feldman, J.W. Pike, J.S. Adams, Vitamin D, third ed., Elsevier Academic Press, New York, 2011.
- [2] N. Saito, Y. Suhara, M. Kurihara, T. Fujishima, S. Honzawa, H. Takayanagi, T. Kozono, M. Matsumoto, M. Ohmori, N. Miyata, H. Takayama, A. Kittaka, Design and efficient synthesis of 2 $\alpha$ -( $\omega$ -hydroxyalkoxy)-1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> analogues, including 2-*epi*-ED-71 and their 20-epimers with HL-60 cell differentiation activity, *J. Org. Chem.* 69 (2004) 7463–7471.
- [3] For our review article, see: N. Saito, S. Honzawa, A. Kittaka, Recent results on A-ring modification of 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub>: design and synthesis of VDR-agonists and antagonists with high biological activity, *Curr. Topics Med. Chem.* 6 (2006) 1273–1288.
- [4] E. Takahashi, K. Nakagawa, Y. Suhara, A. Kittaka, K. Nihei, K. Konno, H. Takayama, K. Ozono, T. Okano, Biological activities of 2 $\alpha$ -substituted analogues of 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> in transcriptional regulation and human promyelocytic leukemia (HL-60) cell proliferation and differentiation, *Biol. Pharm. Bull.* 29 (2006) 2246–2250.
- [5] S. Hourai, T. Fujishima, A. Kittaka, Y. Suhara, H. Takayama, N. Rochel, D. Moras, Probing a water channel near the A-ring of receptor-bound 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> with selected 2 $\alpha$ -substituted analogues, *J. Med. Chem.* 49 (2006) 5199–5205.
- [6] T. Sakaki, K. Yasuda, A. Kittaka, K. Yamamoto, T.C. Chen, CYP24A1 as a potential target for cancer therapy, *Anti-Cancer Agents Med. Chem.* 14 (2014) 97–108.
- [7] D. Abe, T. Sakaki, T. Kusudo, A. Kittaka, N. Saito, Y. Suhara, T. Fujishima, H. Takayama, H. Hamamoto, M. Kamakura, M. Ohta, K. Inouye, Metabolism of 2 $\alpha$ -Propoxy-1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> and 2 $\alpha$ -(3-Hydroxypropoxy)-1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> by Human CYP27A1 and CYP24A1, *Drug Metab. Dispos.* 33 (2005) 778–784.
- [8] K. Yasuda, S. Ikushiro, M. Kamakura, M. Takano, N. Saito, A. Kittaka, T.C. Chen, M. Ohta, T. Sakaki, *J. Steroid Biochem. Mol. Biol.* 133 (2013) 84–92.
- [9] J. Zhu, H.F. DeLuca, Vitamin D 25-hydroxylase – four decades of searching are we there yet? *Arch. Biochem. Biophys.* 523 (2012) 30–36.
- [10] N. Tsugawa, K. Nakagawa, M. Kurobe, Y. Ono, N. Kubodera, K. Ozono, T. Okano, In vitro biological activities of a series of 2 $\beta$ -substituted analogues of 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub>, *Biol. Pharm. Bull.* 23 (2000) 66–71.
- [11] Y. Ono, Multifunctional and potent roles of the 3-hydroxypropoxy group provide eldelcalcitol's benefit in osteoporosis treatment, *J. Steroid Biochem. Mol. Biol.* 139 (2014) 88–97.
- [12] A part of this review has been published as in the following article, see: M. Takano, S. Ohya, K. Yasuda, M. Nishikawa, A. Takeuchi, D. Sawada, T. Sakaki, A. Kittaka, Synthesis and biological activity of 1 $\alpha$ ,2 $\alpha$ ,25-trihydroxyvitamin D<sub>3</sub>: active metabolite of 2 $\alpha$ -(3-hydroxypropoxy)-1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> by human CYP3A4, *Chem. Pharm. Bull.* 62 (2014) 182–184.
- [13] Synthesis of another A-ring precursor of 1 $\alpha$ ,2 $\alpha$ ,25(OH)<sub>3</sub>D<sub>3</sub> was reported, see: G.H. Posner, T.D. Nelson, Stereocontrolled synthesis of a trihydroxylated A-ring as an immediate precursor to 1 $\alpha$ ,2 $\alpha$ ,25-trihydroxyvitamin D<sub>3</sub>, *J. Org. Chem.* 56 (1991) 4339–4341.
- [14] (a) Synthesis of 1 $\alpha$ ,2 $\alpha$ ,25-trihydroxy-19-norvitamin D<sub>3</sub> was reported, see: R.R. Siczinski, K.L. Perlman, H.F. DeLuca, Synthesis and biological activity of 2-hydroxy and 2-alkoxy analogs of 1 $\alpha$ ,25-dihydroxy-19-norvitamin D<sub>3</sub>, *J. Med. Chem.* 37 (1994) 3730–3738; (b) M. Shimizu, Y. Iwasaki, Y. Shibamoto, M. Sato, H.F. DeLuca, S. Yamada, Novel synthesis of 2-substituted 19-norvitamin D A-ring phosphine oxide from *D*-glucose as a building block, *Bioorg. Med. Chem. Lett.* 13 (2003) 809–812; (c) K. Ono, A. Yoshida, N. Saito, T. Fujishima, S. Honzawa, Y. Suhara, S. Kishimoto, T. Sugiyama, K. Waku, H. Takayama, A. Kittaka, *J. Med. Chem.* 68 (2003) 7407–7415.
- [15] D.S. Rao, R. Dayal, M.-L. Siu-Caldera, R.L. Horst, M.R. Uskokovic, K.-Y. Tserng, G.S. Reddy, Isolation and identification of 4,25-dihydroxyvitamin D<sub>2</sub>: a novel A-ring hydroxylated metabolite of vitamin D<sub>2</sub>, *J. Steroid Biochem. Mol. Biol.* 71 (1999) 63–70.
- [16] M. Matsuo, A. Hasegawa, M. Takano, Y. Nakamura, H. Saito, S. Kakuda, K. Takagi, E. Ochiai, K. Horie, M. Takimoto-Kamimura, K. Takenouchi, D. Sawada, A. Kittaka, Synthesis of 2 $\alpha$ -heteroarylalkyl active vitamin D<sub>3</sub> with therapeutic effect on enhancing bone mineral density in vivo, *ACS Med. Chem. Lett.* 4 (2013) 671–674.
- [17] B.M. Trost, J. Dumas, M. Villa, New strategies for the synthesis of vitamin D metabolites via palladium-catalyzed reactions, *J. Am. Chem. Soc.* 114 (1992) 9836–9845.
- [18] A part of this review has been published as in the following article, see: D. Sawada, Y. Tsukuda, K. Yasuda, T. Sakaki, H. Saito, K. Takagi, K. Takenouchi, T.C. Chen, G.S. Reddy, A. Kittaka, Synthesis and biological activities of 1 $\alpha$ ,4 $\alpha$ ,25- and 1 $\alpha$ ,4 $\beta$ ,25-trihydroxyvitamin D<sub>3</sub> and their metabolism by human CYP24A1 and UDP-glucuronosyltransferase, *Chem. Pharm. Bull.* 60 (2012) 1343–1346.
- [19] D. Sawada, Y. Tsukuda, H. Saito, K. Takagi, K. Horie, E. Ochiai, K. Takenouchi, A. Kittaka, Synthesis and biological evaluation of 4-substituted vitamin D and 14-*epi*-previtamin D analogs, *Chem. Pharm. Bull.* 57 (2009) 1431–1433.
- [20] I. Ohtani, T. Kusumi, Y. Kashman, H. Kakisawa, High-field FT NMR application of Mosher's method. The absolute configurations of marine terpenoids, *J. Am. Chem. Soc.* 113 (1991) 4092–4096.
- [21] T. Kusudo, T. Sakaki, D. Abe, T. Fujishima, A. Kittaka, H. Takayama, S. Hatakeyama, M. Ohta, K. Inouye, Metabolism of A-ring diastereomers of 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> by CYP24A1, *Biochem. Biophys. Res. Commun.* 321 (2004) 774–782.