

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/261881979>

# Towards the development of 5-HT<sub>7</sub> ligands combining serotonin-like and arylpiperazine moieties

ARTICLE in EUROPEAN JOURNAL OF MEDICINAL CHEMISTRY · APRIL 2014

Impact Factor: 3.45 · DOI: 10.1016/j.ejmech.2014.04.034 · Source: PubMed

---

READS

105

## 13 AUTHORS, INCLUDING:



**Silvia Bartolucci**

Università degli Studi di Urbino "Carlo Bo"

8 PUBLICATIONS 35 CITATIONS

SEE PROFILE



**Marco Mor**

Università degli studi di Parma

183 PUBLICATIONS 6,004 CITATIONS

SEE PROFILE



**Walter Cabri**

Fresenius Kabi

127 PUBLICATIONS 2,577 CITATIONS

SEE PROFILE



**Diana Celona**

Sigma Tau Pharmaceuticals

7 PUBLICATIONS 27 CITATIONS

SEE PROFILE



## Original article

Towards the development of 5-HT<sub>7</sub> ligands combining serotonin-like and arylpiperazine moieties

Gilberto Spadoni<sup>a</sup>, Annalida Bedini<sup>a</sup>, Silvia Bartolucci<sup>a</sup>, Daniele Pala<sup>b</sup>, Marco Mor<sup>b</sup>,  
Teresa Riccioni<sup>c</sup>, Franco Borsini<sup>c</sup>, Walter Cabri<sup>c</sup>, Diana Celona<sup>c</sup>, Mauro Marzi<sup>c</sup>,  
Giorgio Tarzia<sup>a</sup>, Silvia Rivara<sup>b,\*</sup>, Patrizia Minetti<sup>c,\*</sup>

<sup>a</sup> Dipartimento di Scienze Biomolecolari, Università degli Studi di Urbino, Piazza Rinascimento 6, I-61029 Urbino, PU, Italy

<sup>b</sup> Dipartimento di Farmacia, Università degli Studi di Parma, Parco Area delle Scienze 27/A, I-43124 Parma, Italy

<sup>c</sup> Sigma-Tau Industrie Farmaceutiche Riunite S.p.A., Via Pontina Km 30,400, I-00040 Pomezia, Italy

## ARTICLE INFO

## Article history:

Received 18 October 2013

Received in revised form

18 March 2014

Accepted 10 April 2014

Available online 13 April 2014

## Keywords:

5-HT<sub>7</sub>

Medicinal chemistry

Chemical synthesis

Molecular modeling

Serotonin

Structure–activity relationships

## ABSTRACT

Many known 5-HT<sub>7</sub> ligands contain either a serotonin-like or an arylpiperazine structure that, in published SAR studies, are generally supposed to bind the same receptor pocket. Conversely, we explored the hypothesis that two such moieties can co-exist in the same ligand, binding to different pockets. We thus designed and synthesized a set of compounds including both a 5-hydroxyindol-3-ylethyl and a 1-aryl piperazine moieties connected by a short linker. The compounds were tested for their affinity for human 5-HT<sub>7</sub> serotonin receptor. We further prepared a novel series of 5-HT<sub>7</sub> ligands, where the 5-hydroxyindol-3-ylethyl moiety was bioisosterically replaced by a 3-hydroxyanilinoalkyl one. Among the newly synthesized compounds, potent ligands at the 5-HT<sub>7</sub> receptor, behaving as antagonists in functional tests, were identified, even if they showed limited subtype selectivity. Docking studies within a model of the 5-HT<sub>7</sub> receptor showed that the binding site can actually accommodate both moieties, with the serotonin-like one in the putative orthosteric site and the arylpiperazine one occupying an accessory pocket. The present results demonstrate that it is possible to devise and develop new 5-HT<sub>7</sub> ligands merging two privileged structures in the same molecule.

© 2014 Elsevier Masson SAS. All rights reserved.

## 1. Introduction

Serotonin (5-hydroxytryptamine, 5-HT, Fig. 1) is an important neurotransmitter that mediates a wide range of physiological functions by interacting with seven serotonergic receptor families (5-HT<sub>1–7</sub>) subdivided into at least 14 subpopulations [1]. Drugs which interact specifically with serotonin receptors are clinically approved for the treatment of migraine (sumatriptan: 5-HT<sub>1D/1F</sub> agonist), psychosis (risperidone: 5-HT<sub>2/D2</sub> antagonist), anxiety disorders (buspirone: 5-HT<sub>1A</sub> agonist) and emesis (ondansetron: 5-HT<sub>3</sub> antagonist) [2–4]. The most recently discovered and cloned member of the 5-HT receptor family is 5-HT<sub>7</sub>, which belongs to the G protein-coupled receptor (GPCR) superfamily and is positively coupled to adenylyl cyclase. The 5-HT<sub>7</sub> receptor was cloned independently by three laboratories in 1993

[5–7]. Since its discovery, the 5-HT<sub>7</sub> receptor has been cloned from different species, including rat, man, mouse, pig, guinea pig, honeybee, *Xenopus laevis*, *Aedes aegypti* and *Caenorhabditis elegans* [8]. It is expressed in the central and peripheral nervous system and also in the periphery. In the latter, the 5-HT<sub>7</sub> receptor has been detected predominantly in smooth muscle cells of the cardiovascular, gastrointestinal, and reproductive system, where it was consistently shown to induce smooth muscle relaxation [9]. In the central nervous system, the 5-HT<sub>7</sub> receptor is most abundant in hippocampus, thalamus, hypothalamus and cerebral cortex. The presence in the hypothalamus correlates with its involvement in circadian rhythm, thermoregulation, and endocrine regulation. Thalamic and cortical 5-HT<sub>7</sub> receptors may be of importance for sleep, mood regulation and epilepsy. The distribution of 5-HT<sub>7</sub> receptors in the hippocampus is of relevance for its role in learning and memory [10]. Finally, its localization in the spinal cord is in agreement with putative functions in nociception and locomotion. A consistent body of evidence, using both 5-HT<sub>7</sub> receptor knockout mice and selective ligands, supports the involvement of this receptor in various pathophysiological

\* Corresponding authors.

E-mail addresses: [silvia.rivara@unipr.it](mailto:silvia.rivara@unipr.it) (S. Rivara), [patrizia.minetti@virgilio.it](mailto:patrizia.minetti@virgilio.it) (P. Minetti).

processes. Therefore, this receptor has become an attractive target for drug discovery and in the past years intense efforts have led to the identification of different structural classes of 5-HT<sub>7</sub> receptor agents [for a recent review see Ref. [11]].

Ligands characterized by a 1-arylpiperazine moiety are well known for their affinity to several members of the family of serotonergic receptors, and recently different research groups have modified this template in order to identify selective 5-HT<sub>7</sub> receptor ligands [12]. On the other hand, it is known that appropriate molecular modifications of tryptamine can lead to derivatives that display an enhanced selectivity for different 5-HT receptor subtypes [13], even if the features conferring a subtype selective profile are not fully established.

A number of studies defining 5-HT<sub>7</sub> pharmacophore and homology (using rhodopsin crystal structures as templates) models have been published [12c,14], and they are characterized by at least four pharmacophoric features: a positive ionizable atom, a H-bonding acceptor group, and two hydrophobic regions. In these models, it is generally assumed that the 1-arylpiperazine and the indolyethylamine fragments are reciprocal bioisosteres, with their aromatic portions occupying the same hydrophobic region of the 5-HT<sub>7</sub> receptor.

In some structures of shorter 5-HT<sub>7</sub> ligands (**a–c** in Fig. 1), the basic amine group is bound to a serotonin-like fragment, i.e. a tryptamine or a 2-aminotetralin one, where the 5-hydroxyl and the 8-hydroxyl groups can be strictly superposed, and conserve a significant role for structure–activity relationships. On the other hand, in other ligands, such as aminochromanes (**d**, **e**), 5-aryl-2-aminotetralines (**f**) and mercaptopyridines (**g**) the similarity with the tryptamine scaffold of 5-HT is less evident, with the lack of a hydroxyl group or its bioisostere mimicking the 5-hydroxyl group of 5-HT. 1-Arylpiperazines (**h–j**), an important class of 5-HT<sub>7</sub> ligands, resemble more this second chemotype. In fact, the 7'-methoxy group of **h** can be replaced by a hydrogen atom with no loss of receptor affinity [14e]. Moreover, the 1-arylpiperazine moiety can be connected to a lipophilic tail with good 5-HT<sub>7</sub> affinity, as in compounds **i** and **j**, while no “longer” 5-HT<sub>7</sub> ligands with a tryptamine or 8-hydroxy-2-aminotetralin scaffold have been described.

These observations led us to hypothesize that a serotonin-like moiety could be combined with an arylpiperazine one, as they may occupy two distinct hydrophobic cavities. Although such distinct binding pockets have been proposed both in ligand-based and receptor-based models [14], compounds including both moieties have not been reported so far, nor has this hypothesis been investigated through structure–activity relationships.

We thus synthesized a first series of putative 5-HT<sub>7</sub> ligands, characterized by the presence of a serotonin-like scaffold, where the basic nitrogen is enclosed in a piperazine nucleus, linked to different lipophilic portions through its second nitrogen. As this lipophilic region showed SARs that resembled those of known 1-arylpiperazine ligands, we thus explored the possibility to bioisosterically replace the tryptamine moiety. In a previous work on melatonin receptor ligands, we had observed that a 2-(3-methoxyanilino)ethyl and a 3-(3-methoxyanilino)propyl fragment could replace the 5-methoxyindol-3-ylethyl portion of melatonin [15]. We therefore synthesized a novel series of putative 5-HT<sub>7</sub> ligands, composed by a 2-(3-hydroxyanilino)ethyl or a 3-(3-hydroxyanilino)propyl moiety, potentially mimicking the indole scaffold of 5-HT, and by an arylpiperazine moiety. Furthermore, we explored the SARs for this novel class of compounds and analyzed their possible binding modes at the 5-HT<sub>7</sub> receptor by docking simulations within a homology-based model of the receptor. Selected compounds, endowed with good receptor affinity, were further examined for their intrinsic activity and selectivity against different 5-HT receptor subtypes.

## 2. Chemistry

A previously reported method [16] was used to prepare the [(4-substituted-piperazin-1-yl)ethyl]-indole derivatives **5i–p** (Scheme 1); the procedure involves acylation of the suitable indole **1a–d** with oxalyl chloride, followed by reaction with the appropriate *N*-substituted-piperazine to give the (4-substituted-piperazin-1-yl) ethane-1,2-dione derivatives **3a–h**; these latter are then converted to the corresponding 5-hydroxyindole derivatives **5i–p** by reduction with LiAlH<sub>4</sub> and subsequent *O*-deprotection by hydrogenolysis or by treatment with boron tribromide.





Fig. 1. Representative 5-HT<sub>7</sub> ligands [12d,14b,55–58].



**Scheme 1.** Reagents and conditions: a) (COCl)<sub>2</sub>, THF, RT, 1 h; b) TEA, THF, RT, 1.5 h; c) LiAlH<sub>4</sub>, THF, reflux, 1 h; d) BBr<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, RT, 2 h (for **5i–k** and **5o–p**) or H<sub>2</sub>, 10% Pd/C, MeOH, RT, 6 h (for **5l–n**).

The (anilinoethyl)-piperazino derivatives **10h–l** were prepared as depicted in Scheme 2. Briefly, the (4-substituted-piperazine) acetamido derivatives **8d–g** were synthesized by *N*-acylation of anilines (**6a–c**) with bromoacetyl chloride in the presence of NaHCO<sub>3</sub> followed by substitution of the intermediate bromoacetamides **7a–c** with the suitable *N*-substituted-piperazine. The target compounds **10h–j** were then obtained by LiAlH<sub>4</sub> reduction of the amides **8d–f**, followed by *O*-deprotection of the intermediates **9d–f** by treatment with BBr<sub>3</sub> (for **10h–i**) or by hydrogenolysis catalyzed by palladium on carbon (for **10j**). The nitro-compound **9g**, obtained by BH<sub>3</sub> amide reduction of **8g**, was hydrogenated over 10% Pd/C to give the corresponding anilino derivative **10k** that could be converted into the *N*-methanesulfonamido target compound **10l**, by treatment with methanesulfonyl chloride.

Following a similar stepwise synthetic route (after changing the *N*-acylating reagent to 3-bromopropanoyl chloride), the (anilino-propyl)-piperazino derivatives **14b–o** were obtained from the corresponding anilines **6a**, **11b–i** as shown in Scheme 3. In a similar manner, the higher homologs **18a–d**, **18f–h** and the piperidine analog **18e** were synthesized starting from 3-benzyloxyaniline (**6b**) or 3-benzyloxy-*N*-methylaniline (**11i**) [17] and the suitable ω-alkanoylchloride (Scheme 4). The phenol-hexaneamido derivative **16i** was obtained by *O*-debenzylation with BCl<sub>3</sub> of the corresponding benzyloxy derivative **16f** (Scheme 4).


The *N*-acetamido- (**14p**) and the *N*-methanesulfonamido- (**14q**) target compounds were obtained by catalytic hydrogenation (10% Pd/C) of the nitro-compound **14f** (Scheme 3) and subsequent treatment of the crude aniline intermediate with acetic anhydride or methanesulfonyl chloride, respectively.

Reductive amination of the aniline derivative **17d** with acetaldehyde or benzaldehyde in the presence of NaBH<sub>3</sub>CN and subsequent *O*-debenzylation yielded, respectively, the *N*-ethyl- (**18j**) or *N*-benzyl- (**18k**) final product (Scheme 4).

Compounds **14r–u** were prepared from the ethyl ester derivative **14g** following the step sequence shown in Scheme 5. In particular **14g** was reduced by LiAlH<sub>4</sub> to the hydroxymethyl derivative **14t**, that was oxidized by MnO<sub>2</sub> to the carboxaldehyde derivative **14u**. By treatment of **14g** with magnesium nitride in methanol we obtained a mixture of the primary amide **14r** and of the methyl ester derivative **14s**, that was separated by chromatography.

*N*-substituted-[3-(4-phenylpiperazin-1-yl)propyl]anilines **24–27** were prepared from the aniline derivative **14h** (Scheme 6) that can be converted to the *N*-propyl-anilino derivative **24** by *N*-acylation with propionic anhydride, followed by BH<sub>3</sub> amide reduction and subsequent *O*-demethylation with boron tribromide. Alternatively, **14h** is converted to the *N*-benzyl- (**25**) or to the *N*-benzenesulfonyl- (**27**) target compound by reaction with benzyl bromide or benzenesulfonyl chloride, respectively, and subsequent *O*-demethylation with BBr<sub>3</sub>. The *N*-phenyl derivative **26** was achieved by palladium-catalyzed amination [Pd(OAc)<sub>2</sub>/BINAP/KOtBu] of **14h** with iodobenzene and subsequent *O*-demethylation with boron tribromide.

Preparation of the 4-substituted piperidine derivatives **30b–c** started with the reaction of **12a** with the suitable piperidine giving the corresponding piperidine-propanamido derivatives **28b–c**, that were then reduced with BH<sub>3</sub>, and finally *O*-deprotected by treatment with BBr<sub>3</sub> (Scheme 7).



**Scheme 2.** Reagents and conditions: a) 2-bromoacetyl chloride, NaHCO<sub>3</sub>, EtOAc, H<sub>2</sub>O, RT, 20 min; b) TEA, DMF, RT, 1–16 h; c) LiAlH<sub>4</sub>, THF, reflux, 2 h (for **9d–f**) or BH<sub>3</sub>, THF, RT, 16 h (for **9g**); d) BBr<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, RT, 2 h (for **10h–i**) or H<sub>2</sub>, 10% Pd/C, MeOH, 60 °C, 3 h (for **10j**); e) H<sub>2</sub>, 10% Pd/C, EtOH, EtOAc, RT, 5 h; f) methanesulfonyl chloride, TEA, THF, RT, 1 h.

Phenolalkyl-piperazino derivative **32a–c** were synthesized by condensation of the appropriate carboxylic acid with *N*-phenyl piperazine in the presence of 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC), followed by amide reduction with 1 M B<sub>2</sub>H<sub>6</sub> in THF and *O*-deprotection (for **32c**) (Scheme 8).


Compound **35** was prepared from 3-(benzyloxy)phenol following the three-step sequence (*O*-alkylation, amide reduction, *O*-debenzylation) illustrated in Scheme 9.

The tetralin derivative **37** was synthesized by reductive amination of 8-methoxy-2-tetralone with *N*-phenyl-piperazine and subsequent *O*-demethylation with BBr<sub>3</sub> (Scheme 10).

### 3. Results and discussion

Binding affinities at the human 5-HT<sub>7</sub> receptor for the newly synthesized compounds that combine a serotonin-like (5-hydroxyindol-3-ylethylamine) with an arylpiperazine moiety are reported in Table 1. Compared with 5-HT, the phenylpiperazinyll derivative **5n** showed moderate binding affinity (*K*<sub>i</sub> = 63 nM), suggesting that the presence of the phenyl-piperazine moiety did not prevent accommodation of the 5-hydroxyindole portion within the 5-HT<sub>7</sub> binding site. Bulkier aromatic substituents, in the *m*-

biphenyl and *p*-biphenyl derivatives **5l** and **5m**, abolished affinity for the receptor. On the other hand, an  $\alpha$ -naphthyl group (**5i**) had a favorable effect and led to a sixfold increase of binding affinity compared to the phenyl derivative **5n**. When small alkyl groups were inserted at position 2 of the indole ring in the  $\alpha$ -naphthyl derivative, a methyl group (**5k**) was tolerated and an ethyl one (**5j**) doubled 5-HT<sub>7</sub> binding affinity. This behavior differs from what observed for other tryptamine derivatives. In fact, insertion of a methyl or an ethyl group in position 2 of *N,N*-dimethyl-5-methoxytryptamine led to a decrease of binding affinity, with an increase from tenfold to fiftyfold in *K*<sub>i</sub> values [14b,18]. Replacement of the aromatic substituent on the piperazine nitrogen with an alkyl one, such as a *n*-butyl chain (**5o**) or a cyclopentyl ring (**5p**), led to derivatives devoid of any significant 5-HT<sub>7</sub> binding affinity. This observation is not in line with what reported for a series of 6-bromo-1-(homo-piperazinyl)ethylindoles, in which the presence of alkyl substituents on the homo-piperazine nitrogen afforded compounds with binding affinities in the nanomolar range [19]. Thus, both a phenyl- and an  $\alpha$ -naphthyl-piperazine moiety could be combined with a serotonin-like moiety with limited loss of receptor affinity. This preliminary result was confirmed preparing compound **37**, where the 5-hydroxyindol-3-ylethyl substructure




**Scheme 3.** Reagents and conditions: a) 3-bromopropionyl chloride, NaHCO<sub>3</sub>, EtOAc, H<sub>2</sub>O, RT, 20 min; b) TEA, DMF, RT, 1–16 h; c) BH<sub>3</sub>, THF, RT, 16 h; d) H<sub>2</sub>, 10% Pd/C, MeOH, 60 °C, 3 h; e) BBr<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, RT, 2 h; f) H<sub>2</sub>, 10% Pd/C, EtOH, EtOAc, RT, 5 h; g) Ac<sub>2</sub>O, TEA, THF, RT, 1 h; h) methanesulfonyl chloride, TEA, THF, RT, 1 h.

was replaced by a 8-hydroxytetralin one, which had been shown to be a bioisostere on several 5-HT receptors.

To investigate the binding mode of our indole derivatives, we built a homology model of the 5-HT<sub>7</sub> receptor based on the three-dimensional coordinates of the 5-HT<sub>1B</sub> serotonin receptor co-crystallized with the agonist dihydroergotamine. Crystal structures of different GPCRs have shown that agonist and antagonist compounds can be accommodated within binding sites having very similar location and size, with only subtle differences [20]. Therefore, given the level of accuracy of GPCR homology models, we considered the 5-HT<sub>1B</sub> crystal structure as a suitable template and we used the 5-HT<sub>7</sub> model for docking studies on our compounds, for which a functional characterization was not systematically carried out. As already observed by Kolaczowski et al. [14e] for a rhodopsin-based 5-HT<sub>7</sub> receptor model, the putative binding region is characterized by the presence of Asp162 (3.32 according to Ballesteros–Weinstein's nomenclature [21]) on transmembrane (TM) helix 3 which is supposed to be the counterpart for basic groups of ligands and which divides the binding site into two portions, one located within the helix bundle and delimited by TM3, 5 and 6 and one which lays towards the extracellular side of TM7. Accordingly, in the best induced-fit docking solution of

compound **5i** the protonated piperazine nitrogen interacts with Asp162<sup>3.32</sup> through a salt bridge and the ligand molecule spans the helix bundle from TM5 to TM7 (Fig. 2a). The indole ring of **5i** is located in a cavity delimited by TM3, TM5 and TM6 and is engaged in several hydrophobic contacts with Val163<sup>3.33</sup>, Ala247<sup>5.46</sup>, Phe343<sup>6.51</sup> and Phe344<sup>6.52</sup>; the α-naphthyl-piperazine moiety is accommodated towards the extracellular portion of the helix bundle, close to TM7, and undertakes a number of favorable interactions with Phe158<sup>3.28</sup>, Ile159<sup>3.29</sup>, Leu232 (extracellular loop 2) and Leu370<sup>7.39</sup>. Interestingly, the region occupied by the α-naphthyl substituent of compound **5i** is partially superposed to that occupied by the cyclic tripeptide of dihydroergotamine in the crystallized 5-HT<sub>1B</sub> receptor (Fig. 2a). A 100 ns long molecular dynamics (MD) simulation of the 5-HT<sub>7</sub>–**5i** complex was conducted in an explicit solvated lipid bilayer. The stability of the receptor structure was confirmed by the time evolution of the root mean squared deviation (RMSD) calculated for TM alpha carbons, which showed a plateau after 30 ns of simulation (Fig. S1). Compound **5i** retained its initial conformation during the whole MD simulation as well as the key ionic interaction with Asp162<sup>3.32</sup>, and subtle rearrangements involving the indole and the α-naphthyl rings were detected (Fig. 2b). In particular, an outward shift of the extracellular half of






	n	R <sup>1</sup>	X		n	R <sup>1</sup>	X
a	3	Me	N-Ph	f	5	Me	N- $\alpha$ -naphthyl
b	4	Me	N-Ph	g	5	Me	N- <i>m</i> -CF <sub>3</sub> -Ph
c	5	Me	N-Ph	h	5	Me	N-(7-Cl-quinolin-4-yl)
d	5	H	N-Ph	j	5	Et	N-Ph
e	5	Me	C-Ph	k	5	Bn	N-Ph

**Scheme 4.** Reagents and conditions: a)  $\omega$ -bromoalkanoate, NaHCO<sub>3</sub>, EtOAc, H<sub>2</sub>O, RT, 20 min; b) TEA, DMF, RT, 1–16 h; c) BCl<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, RT, 1 h; d) LiAlH<sub>4</sub>, THF, reflux, 2 h; e) BCl<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, RT, 2 h (for **18e–h**) or H<sub>2</sub>, 10% Pd/C, MeOH, 60 °C, 3 h (for **18a–c**); f) CH<sub>3</sub>CHO (for **17j**) or benzaldehyde (for **17k**), MeOH, AcOH, NaBH<sub>3</sub>CN, RT, 16 h.


TM5 was observed at the beginning of the MD simulation, allowing a slight rotation of the indole ring of **5i** and the consequent formation of a hydrogen bond between its 5-hydroxyl group and Ser243<sup>5,42</sup>, which was maintained throughout the MD simulation. Although no mutagenesis data are available for this amino acid, replacement of the corresponding serine of the 5-HT<sub>1A</sub> receptor

with an alanine decreased 5-HT binding affinity [22]. Moreover, the serine residue in position 5.42 of the  $\beta_1$  receptor is hydrogen bonded to the hydroxyl groups of isoprenaline [23], suggesting that the amino acid in this position may interact with polar groups of other aminergic neurotransmitters. The  $\alpha$ -naphthyl substituent of **5i** slightly rearranged during MD simulation. In particular, it moved slightly away from Phe158<sup>3,28</sup> on TM3 and was accommodated in a crevice located at the tips of TM6 and TM7, delimited by Phe343<sup>6,51</sup>, Leu346<sup>6,54</sup> and Leu370<sup>7,39</sup>. The shape and the location of this pocket could also explain the loss of binding affinity brought by the introduction of *meta*- or *para*-biphenyl substituents (compounds **5l** and **5m**). Indeed, these bulkier groups could not be accommodated, due to unfavorable steric clashes. This hydrophobic region, extending beyond the anionic site of Asp162<sup>3,32</sup>, can thus represent the binding pocket for the arylpiperazine moieties of our hybrid ligands.

The binding model depicted in Fig. 2b is also consistent with the possibility to replace the serotonin-like portion with bioisosteric substructures. In fact, compound **37**, having a 8-hydroxytetralin scaffold, showed binding affinity similar to that of the serotonin-like derivative **5n** ( $K_i$  = 81.4 nM and  $K_i$  = 63.1 nM, respectively). To further test this hypothesis, we replaced the indole scaffold with an aniline one and we prepared a small series of anilino-alkyl-piperazine derivatives, exemplified by compound **10h** (Table 2). In



**Scheme 5.** Reagents and conditions: a) Mg<sub>3</sub>N<sub>2</sub>, MeOH, 80 °C, 24 h; b) LiAlH<sub>4</sub>, THF, RT, 1 h; c) MnO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>, RT, 18 h.




**Scheme 6.** Reagents and conditions: a) (EtCO)<sub>2</sub>O, TEA, THF, RT, 24 h; b) BH<sub>3</sub>, THF, RT, 16 h; c) benzyl bromide, TEA, toluene, 80 °C, 16 h; d) Pd(OAc)<sub>2</sub>, (±)-BINAP, *tert*-BuO<sup>−</sup>K<sup>+</sup>, iodobenzene, toluene, reflux, 24 h; e) benzenesulfonyl chloride, TEA, THF, RT, 48 h; f) BBr<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, RT, 2 h.

fact, successful bioisosteric replacement of the 5-methoxyindol-3-ylethyl fragment with a *meta*-methoxyanilinoethyl or a *meta*-methoxyanilinopropyl one had already been described for melatonin (*N*-acetyl-5-methoxytryptamine) derivatives [15], and the anilino scaffold had resulted more versatile and synthetically feasible. Fig. 3 depicts *N,N*-dimethyl-5-hydroxytryptamine (**b**, Fig. 1) in the bioactive conformation proposed by Vermeulen et al. [14b], superposed to the tetralin derivative 8-OH-DPAT (**c**, Fig. 1) and to 3-[[2-(dimethylamino)ethyl](methyl)amino]phenol. The close superposition of this last compound highlights the possibility for a *meta*-hydroxyanilino-ethylamine fragment to bioisosterically replace the other two scaffolds.


Compound **10h** carries a hydroxyl group in *meta* position, topologically corresponding to position 5 of the indole ring of 5-HT, and a spacer with two methylene units connects the aniline nitrogen to the arylpiperazine portion. Compound **10h** showed a limited loss of binding affinity (Table 2), being about three times less potent than the corresponding indole derivative **5n**. An investigation on the importance of the hydroxyl group was carried out inserting in *meta* position substituents with different size and ability to form hydrogen bonds. While amino (**10k**) and nitro (**9g**) substituents led to a decrease of binding affinity, a methylsulfonylamino group had a positive effect, leading to a *K<sub>i</sub>* value of 57.0 nM for compound **10l**. Replacement of the phenyl ring on the piperazine nitrogen with a benzyl group markedly reduced binding affinity (**10i**), which stresses the role of the arylpiperazine moiety. Removal of the methyl substituent from the aniline nitrogen slightly reduced binding affinity (**10j**), while replacement of *N*-methyl with an oxygen atom (**35**) almost abolished binding affinity. On the other hand, replacement with a methylene, as in the phenylpropyl derivative **32a**, led to a recovery in potency (*K<sub>i</sub>* = 40.0 nM). SARs related to the lengthening of the alkyl spacer were also investigated, progressively increasing the number of methylene groups from two to six. Addition of one methylene group had a positive effect on binding affinity, since the

phenylpiperazine derivative **14o** had *K<sub>i</sub>* = 20.0 nM, compared to *K<sub>i</sub>* = 161.4 nM for the shorter analog **10h**. As already observed in the indole series, replacement of the phenyl substituent with an  $\alpha$ -naphthyl one significantly increased binding affinity (**14m**, *K<sub>i</sub>* = 2.9 nM), while a  $\beta$ -naphthyl group (**14n**) was hardly tolerated. When docked into the 5-HT<sub>7</sub> receptor binding site, compound **14m** showed an accommodation and a pattern of interactions similar to those described for the indole derivative **5i**. The *meta*-hydroxyl group is hydrogen-bonded to Ser243<sup>5,42</sup>, the protonated piperazine nitrogen interacts with Asp162<sup>3,32</sup> and the naphthyl substituent occupies the region defined by Phe343<sup>6,51</sup>, Leu346<sup>6,54</sup> and Leu370<sup>7,39</sup> (Fig. 2c). Within the anilinopropyl series, a slight increase in binding affinity was registered when the aniline nitrogen was demethylated (**14i**) or replaced by a methylene group, with *K<sub>i</sub>* = 3.1 nM for compound **32c**. The *meta*-hydroxyl group was fundamental to achieve high potency. In fact, its replacement with other substituents reduced binding affinity, as observed for a methyl group (**14b**), for halogens, such as fluoro (**14e**), chloro (**14d**) and bromo (**14c**), and for acetamino (**14p**) and methylsulfonylamino (**14q**) groups. A similar decrease of binding affinity was obtained, on the anilino-NH scaffold, with hydroxymethyl (**14t**), methoxycarbonyl (**14s**) and carboxamido (**14r**) groups, with only the formyl substituent (**14u**) maintaining the same binding affinity as the *meta*-hydroxyl derivative **14i**. Four different substituents were introduced on the aniline nitrogen to look for additional interactions with the binding site. Our receptor model suggested that the 5-HT<sub>7</sub> receptor binding site has room enough to accommodate substituents at nitrogen slightly bulkier than the methyl group. Indeed, a *n*-propyl chain (**24**) did not affect binding affinity, while the phenyl (**26**) and benzyl (**25**) groups reduced *K<sub>i</sub>* to about 90 nM. Interestingly, the phenylsulfonyl derivative **27** was the best one, with higher binding affinity than the reference **14o**. Docking of **27** into the receptor model gave two families of solutions. One of them was consistent with our hypothesis (Fig. 2d), while the other one showed inverted orientation, i.e. with the





**Scheme 7.** Reagents and conditions: a) TEA, DMF, RT, 16 h; b)  $\text{BH}_3$ , THF, RT, 16 h; c)  $\text{BBr}_3$ ,  $\text{CH}_2\text{Cl}_2$ , RT, 2 h.




**Scheme 8.** Reagents and conditions: a)  $\text{N}$ -phenylpiperazine, EDC, HOBT,  $\text{Et}_3\text{N}$ ,  $\text{CH}_2\text{Cl}_2$ , 2 h at 0 °C and 16 h at RT; b)  $\text{B}_2\text{H}_6$ , THF, RT, 4 h; c)  $\text{BBr}_3$ ,  $\text{CH}_2\text{Cl}_2$ , RT, 2 h.

phenyl-piperazine moiety in the serotonin-like binding cavity and vice versa. These alternative solutions can be the result of model errors, due to its homology-based nature, or simply a consequence of the choice to select a model adapted, by MD refinement, to a smaller ligand (**14m**), lacking the  $\text{N}$ -phenylsulfonyl group.


Attempts to probe the size of the binding site portion accommodating the arylpiperazine moiety with bulkier substituents were not successful, as derivatives carrying 4,4'-diphenylpiperidinyl (**30b**) and 4-( $\text{N,N}$ -diphenylamino)piperidinyl (**30c**) groups markedly reduced binding affinity. Lengthening of the alkyl spacer to four (**18a**), five (**18b**) and six (**18c**) methylene units did not

significantly affect binding affinity, ending up with the most potent compound **18c** having  $K_i = 12.0$  nM.

Selected anilino-alkyl-piperazines with two to six methylene spacers were tested in functional assays, to evaluate their agonist or antagonist behavior. Compound **18a** produced no increase of cAMP level when tested up to the concentration of  $10^{-5}$  M. On the other hand it decreased the accumulation of cAMP stimulated by 5-HT, with a  $K_B = 57$  nM. In the same test, compounds **10l**, **14o**, **18b** and **18c** at the concentration of 1  $\mu\text{M}$  produced a reduction of cAMP accumulation induced by 5-HT to 48%, 49%, 69% and 56%, respectively (Table 2), suggesting antagonist behavior.



**Scheme 9.** Reagents and conditions: a)  $\text{K}_2\text{CO}_3$ , DMF, 40 °C, 16 h; b)  $\text{B}_2\text{H}_6$ , THF, RT, 4 h; c)  $\text{BCl}_3$ ,  $\text{CH}_2\text{Cl}_2$ , RT, 1 h.



**Scheme 10.** Reagents and conditions: a) PTSA, toluene, reflux, 24 h; b) H<sub>2</sub>, EtOH/THF, PtO<sub>2</sub>, RT, 24 h; c) BBr<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, RT, 2 h.

The antagonists **18a** and **18c**, having a four-methylene and six-methylene spacers, respectively, were docked into the 5-HT<sub>7</sub> receptor binding site. Both were accommodated similarly to the shorter **14m**, with the same polar interactions formed by the piperazine nitrogen and hydroxyl substituent (Fig. 4). The hydroxylanilino moiety is accommodated within the serotonin-like binding pocket where agonists are supposed to bind, while the phenyl ring on the piperazine lies in the region of the binding site delimited by TM2, 3 and 7. This binding pose could be consistent with their antagonist behavior. Indeed, for the 5-HT<sub>7</sub> receptor [24] and, in general, for biogenic amine receptors [25] it has been proposed that antagonist and agonist compounds occupy the same binding region, with antagonists undertaking interactions with an additional portion of the receptor.

Other structural modifications were evaluated in the presence of the six-methylene spacer. Replacement of the phenyl-piperazine with a 4-phenyl-piperidine (**18e**) remarkably decreased binding

affinity, and insertion of an  $\alpha$ -naphthyl ring on the piperazine had a moderate negative effect (**18f**), opposite to what observed for shorter derivatives. Recovery of binding affinity could be attained by the anilide derivative **16i** having  $K_i = 8.1$  nM. The 5-HT<sub>7</sub> receptor was poorly tolerant to substituents on the piperidine nitrogen of compounds with a six-methylene linker. Indeed, insertion of a trifluoromethyl group in *meta* position on the phenyl ring (**18g**) reduced binding affinity, and a bulkier 7-chloro-quinolin-4-yl fragment replacing the phenyl ring led to compound **18h** devoid of any binding affinity. Substituents on the aniline nitrogen were tolerated also in this longer series, even if an ethyl chain (**18j**) and a benzyl group (**18k**) slightly decreased binding affinity compared to the methyl derivative **18c**.

Four compounds from both the indole (**5n**, **5i**) and the aniline series (**14o**, **16i**) were also tested on other 5-HT receptors (Table S1) [26]. These compounds interacted with different receptor subtypes with high affinity, in particular with the 5-HT<sub>1A</sub> receptor, which is known to bind different classes of 5-HT<sub>7</sub> ligands. This behavior highlights the necessity to optimize the structure of these compounds to gain receptor subtype selectivity while maintaining their good 5-HT<sub>7</sub> binding affinity.

## 4. Conclusions

Our data and models are consistent with the existence of two separate binding pockets in the 5-HT<sub>7</sub> receptor. The first one can interact with the serotonin-like 5-hydroxyindole substructure or its bioisosteres in the putative orthosteric site, and the second one can accommodate an arylpiperazine moiety. For this reason the two substructures can be combined to yield new ligands endowed with good receptor affinity and antagonist potency, while subtype selectivity still needs further investigations.

## 5. Experimental section


### 5.1. Chemistry

Melting points were determined on a Buchi B-540 capillary melting point apparatus and are uncorrected. <sup>1</sup>H NMR spectra and <sup>13</sup>C NMR were recorded on a Bruker AVANCE 200 spectrometer (<sup>1</sup>H: 200 MHz; <sup>13</sup>C: 50 MHz) using CDCl<sub>3</sub> as solvent unless stated otherwise. Chemical shifts ( $\delta$  scale) are reported in parts per million (ppm) relative to the central peak of the solvent. Coupling constants ( $J$  values) are given in Hz. EI-MS spectra (70 eV) were taken on a Fisons Trio 1000 instrument. Only molecular ions (M<sup>+</sup>) and base peaks are given. ESI-MS spectra were taken on a Waters Micromass Zq instrument. Only molecular ions (M + H)<sup>+</sup> are given. Infrared spectra were obtained on a Nicolet Avatar 360 FT-IR spectrometer; absorbances are reported in  $\nu$  (cm<sup>-1</sup>). Elemental analyses for C, H and N were performed on a Carlo Erba analyzer, and the results are within 0.4% of the calculated values. Column chromatography purifications were performed under “flash” conditions using Merck 230–400 mesh silica gel. Analytical thin-layer chromatography (TLC) was carried out on Merck silica gel 60 F<sub>254</sub> plates. The starting indoles **1a–d**, anilines **6a–c** and **11b–h**, 3-(3-hydroxyphenyl)propionic acid, 3-(benzyloxy)phenol, 8-methoxy-3,4-dihydronaphthalen-2(1H)-one, 4-phenylpiperidine and all the 4-substituted-piperazines were commercially available.

#### 5.1.1. General procedure for 1-(5-substituted-indol-3-yl)-2-(4-substituted-piperazin-1-yl)ethane-1,2-dione derivatives **3a–h**

A solution of oxalyl chloride (0.14 mL, 1.6 mmol) in dry THF (2.5 mL) was added dropwise to an ice-cooled stirred solution of the suitable indole **1a–c** (1.3 mmol) in dry THF (6.5 mL), under N<sub>2</sub>. Upon completion of the addition, the reaction mixture was allowed


**Table 1**  
Binding affinity of the newly synthesized indole and tetralin derivatives for human 5-HT<sub>7</sub> receptors.



Compd.	R <sup>1</sup>	R <sup>2</sup>	K <sub>i</sub> (nM) <sup>a</sup> (95% confidence interval) or % displacement at fixed conc.
<b>5-HT</b>			1.4 (0.8–2.2)
<b>5n</b>	Ph	H	63.1 (35.8–111.1)
<b>5l</b>	<i>m</i> -Biphenyl	H	88%@10 <sup>-5</sup> 42%@10 <sup>-6</sup>
<b>5m</b>	<i>p</i> -Biphenyl	H	32%@10 <sup>-5</sup> 10%@10 <sup>-7</sup>
<b>5i</b>	$\alpha$ -Naphthyl	H	11.6 (6.8–19.7)
<b>5k</b>	$\alpha$ -Naphthyl	Me	12.8 (8.7–18.8)
<b>5j</b>	$\alpha$ -Naphthyl	Et	6.6 (3.5–12.6)
<b>5o</b>	<i>n</i> -Butyl	H	–9%@10 <sup>-7</sup>
<b>5p</b>	<i>c</i> -Pentyl	H	1%@10 <sup>-7</sup>
<b>37<sup>b</sup></b>			81.4 (44.3–149.4)

<sup>a</sup> K<sub>i</sub> values were calculated from IC<sub>50</sub> values, obtained from competition curves by the method of Cheng and Prusoff [33], and are the mean of four determinations performed in duplicate.

<sup>b</sup> Tested as racemate.



**Fig. 2.** a) Induced-fit docking of compound **5i** (orange) within the 5-HT<sub>7</sub> receptor model (light gray). Dihydroergotamine is represented as transparent cyan carbons. b) MD simulation of the 5-HT<sub>7</sub>–**5i** complex: superposition of the starting (transparent sticks) and the final energy-minimized structures (opaque sticks and cartoons). c) Best docking pose of **14m** (green carbons) within the 5-HT<sub>7</sub> binding site. **5i** (transparent orange carbons) is depicted for comparison. d) Best docking pose of **27** (pink carbons) within the 5-HT<sub>7</sub> binding site. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

to warm to room temperature and stirred at this temperature for 1 h. The solvent and excess oxalyl chloride were removed under reduced pressure and the residue dissolved in dry THF. After cooling to 0 °C, a solution of TEA (0.42 mL) and of the suitable *N*-substituted-piperazine (1.95 mmol) in dry THF (6.5 mL) was added, and the resulting mixture was stirred at room temperature for 1.5 h. The solvent was removed by distillation *in vacuo*, the residue partitioned between EtOAc and water, and the aqueous phase extracted with EtOAc. The combined organic extracts were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure, to give a crude residue that was purified by flash-chromatography on silica gel: cyclohexane/EtOAc 1:1 as eluant (EtOAc/MeOH, 95:5 for **3g** and **3h**).

**5.1.1.1. 1-(5-Methoxy-1H-indol-3-yl)-2-[4-(naphthalen-1-yl)piperazin-1-yl]ethane-1,2-dione (3a).** White amorphous solid (0.33 g, 62%); <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>): δ = 3.15 (m, 4H, piperazine), 3.80 (m, 2H, piperazine), 3.87 (s, 3H, CH<sub>3</sub>), 4.01 (m, 2H, piperazine), 6.93 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 7.18 (m, 1H, Ar-H), 7.40–7.65 (m, 5H, Ar-H), 7.81 (d, 1H, *J* = 2.0, Ar-H), 7.89 (m, 1H, Ar-H), 8.16 (m, 1H, Ar-H), 8.29 (m, 1H, Ar-H), 11.23 (brs, 1H, NH); ESI-MS (*m/z*): 414 (M + H)<sup>+</sup>.

**5.1.1.2. 1-(2-Ethyl-5-methoxy-1H-indol-3-yl)-2-[4-(naphthalen-1-yl)piperazin-1-yl]ethane-1,2-dione (3b).** Yellowish amorphous solid (0.30 g, 52%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.41 (t, 3H, *J* = 7.5, CH<sub>2</sub>CH<sub>3</sub>), 3.18 (q, 2H, *J* = 7.5, CH<sub>2</sub>CH<sub>3</sub>), 3.22 (m, 4H, piperazine), 3.41 (m, 2H, piperazine), 3.75 (m, 2H, piperazine), 3.90 (s, 3H, OCH<sub>3</sub>), 6.90 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 7.08 (d, 1H, *J* = 8.0, Ar-H), 7.26 (m, 1H, Ar-H), 7.38–7.63 (m, 5H, Ar-H), 7.85 (m, 1H, Ar-H), 8.20 (m, 1H, Ar-H), 8.75 (brs, 1H, NH); ESI-MS (*m/z*): 442 (M + H)<sup>+</sup>.

**5.1.1.3. 1-(5-Methoxy-2-methyl-1H-indol-3-yl)-2-[4-(naphthalen-1-yl)piperazin-1-yl]ethane-1,2-dione (3c).** Yellowish amorphous solid (0.24 g, 43%); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ = 2.62 (s, 3H, CH<sub>3</sub>), 3.00 (m, 2H,


piperazine), 3.13 (m, 2H, piperazine), 3.61 (m, 2H, piperazine), 3.79 (s, 3H, OCH<sub>3</sub>), 3.94 (m, 2H, piperazine), 6.84 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 7.14 (m, 1H, Ar-H), 7.33 (d, 1H, *J* = 8.0, Ar-H), 7.39–7.65 (m, 5H, Ar-H), 7.90 (m, 1H, Ar-H), 8.19 (m, 1H, Ar-H), 12.01 (brs, 1H, NH); ESI-MS (*m/z*): 428 (M + H)<sup>+</sup>.

**5.1.1.4. 1-[5-(Benzyloxy)-1H-indol-3-yl]-2-[4-(biphenyl-3-yl)piperazin-1-yl]ethane-1,2-dione (3d).** White solid (0.45 g, 67%); mp: 221–222 °C (EtOAc); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 3.25 (m, 2H, piperazine), 3.36 (m, 2H, piperazine), 3.75 (m, 2H, piperazine), 3.96 (m, 2H, piperazine), 5.17 (s, 2H, CH<sub>2</sub>Ph), 6.95 (m, 1H, Ar-H), 7.06 (dd, 1H, *J* = 2.5 and 8.5, Ar-H), 7.13–7.17 (m, 2H, Ar-H), 7.31–7.60 (m, 12H, Ar-H), 7.98 (m, 2H, Ar-H), 8.80 (brs, 1H, NH); MS (EI, 70 eV): *m/z* 515 (M<sup>+</sup>), 175 (100).



**5.1.1.5. 1-[5-(Benzyloxy)-1H-indol-3-yl]-2-[4-(biphenyl-4-yl)piperazin-1-yl]ethane-1,2-dione (3e).** Yellowish solid (0.50 g, 75%); mp: 216–217 °C (EtOAc); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ = 3.16 (m, 2H, piperazine), 3.32 (m, 2H, piperazine), 3.49 (m, 2H, piperazine), 3.77 (m, 2H, piperazine), 5.14 (s, 2H, CH<sub>2</sub>Ph), 6.98 (dd, 1H, *J* = 2.5 and 8.5, Ar-H), 7.03 (d, 1H, *J* = 8.5, Ar-H), 7.23–7.70 (m, 15H, Ar-H), 7.74 (d, 1H, *J* = 2.5, Ar-H), 12.23 (brs, 1H, NH); ESI-MS (*m/z*): 516 (M + H)<sup>+</sup>.

**5.1.1.6. 1-[5-(Benzyloxy)-1H-indol-3-yl]-2-(4-phenylpiperazin-1-yl)ethane-1,2-dione (3f).** White solid (0.43 g, 75%); mp: 136–137 °C (EtOAc); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 3.16 (m, 2H, piperazine), 3.27 (m, 2H, piperazine), 3.68 (m, 2H, piperazine), 3.91 (m, 2H, piperazine), 5.14 (s, 2H, CH<sub>2</sub>Ph), 6.91–7.02 (m, 4H, Ar-H), 7.23–7.51 (m, 8H, Ar-H), 7.78 (d, 1H, *J* = 3.0, Ar-H), 7.95 (d, 1H, *J* = 2.0, Ar-H), 9.73 (brs, 1H, NH); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 185.5, 166.3, 156.1, 150.8, 137.1, 135.7, 131.5, 129.3, 128.6, 127.9, 127.7, 126.1, 120.8, 116.9, 115.4, 114.5, 112.8, 104.6, 70.6, 50.0, 49.6, 46.1, 41.5; MS (EI, 70 eV): *m/z* 439 (M<sup>+</sup>), 330 (100).

**Table 2**  
Binding affinity of the newly synthesized anilino-alkyl-piperazine derivatives for human 5-HT<sub>7</sub> receptors ( $K_i$  (nM) or % displacement at fixed concentration) and functional evaluation at the cAMP test.



Compd.	R <sup>1</sup>	X	n	R <sup>2</sup>	$K_i$ (nM) <sup>a</sup> (95% confidence interval) or % displacement at fixed conc.	% Serotonin response on cAMP accumulation at 1 $\mu$ M
<b>10h</b>	OH	N-Me	2	Ph	161.4 (78.3–332.9)	
<b>10k</b>	NH <sub>2</sub>	N-Me	2	Ph	85%@10 <sup>-5</sup> 40%@10 <sup>-6</sup>	
<b>9g</b>	NO <sub>2</sub>	N-Me	2	Ph	332.0 (96.3–1177.0)	
<b>10l</b>	NHSO <sub>2</sub> Me	N-Me	2	Ph	57.0 (38.9–83.2)	48%
<b>10i</b>	OH	N-Me	2	Bn	82%@10 <sup>-5</sup> 42%@10 <sup>-6</sup>	
<b>10j</b>	OH	NH	2	Ph	252.1 (144.4–555.5)	
<b>35</b>	OH	O	2	Ph	42%@10 <sup>-5</sup> 21%@10 <sup>-6</sup>	
<b>32a</b>	OH	CH <sub>2</sub>	2	Ph	40.0 (22.9–69.9)	
<b>14o</b>	OH	N-Me	3	Ph	20.0 (12.0–33.3)	49%
<b>14m</b>	OH	N-Me	3	$\alpha$ -Naphthyl	2.9 (1.0–8.3)	
<b>14n</b>	OH	N-Me	3	$\beta$ -Naphthyl	434.2 (117.6–1602.0)	
<b>32c</b>	OH	CH <sub>2</sub>	3	Ph	3.1 (1.3–7.6)	
<b>14l</b>	OH	NH	3	Ph	10.1 (5.2–19.6)	
<b>14b</b>	Me	N-Me	3	Ph	84.8 (26.1–275.2)	
<b>14e</b>	F	N-Me	3	Ph	111.7 (55.2–226.3)	
<b>14d</b>	Cl	N-Me	3	Ph	179.2 (72.9–440.5)	
<b>14c</b>	Br	N-Me	3	Ph	161.2 (49.5–525.3)	
<b>14p</b>	NHCOMe	N-Me	3	Ph	161.1 (73.8–351.5)	
<b>14q</b>	NHSO <sub>2</sub> Me	N-Me	3	Ph	39.6 (14.9–104.7)	
<b>14t</b>	CH <sub>2</sub> OH	NH	3	Ph	108.8 (8.9–1325.0)	
<b>14s</b>	COOMe	NH	3	Ph	77.9 (40.8–148.6)	
<b>14r</b>	CONH <sub>2</sub>	NH	3	Ph	63.3 (32.7–122.4)	
<b>14u</b>	CHO	NH	3	Ph	13.2 (7.2–24.1)	
<b>24</b>	OH	N- <i>n</i> Pr	3	Ph	20.8 (11.8–36.4)	
<b>26</b>	OH	N-Ph	3	Ph	90.1 (28.3–287.0)	
<b>25</b>	OH	N-Bn	3	Ph	92.5 (47.0–181.8)	
<b>27</b>	OH	N-SO <sub>2</sub> Ph	3	Ph	12.1 (6.2–23.8)	



<b>18a</b>	OH	N-Me	4	Ph	30.0 (14.0–59.3)	39% IC <sub>50</sub> = 540 nM K <sub>B</sub> = 57 nM
<b>18b</b>	OH	N-Me	5	Ph	16.7 (7.7–35.9)	69%
<b>18c</b>	OH	N-Me	6	Ph	12.0 (8.3–17.2)	56%
<b>18e</b>	OH	N-Me	6	$\alpha$ -Naphthyl	226.0 (141.9–360.0)	
<b>18f</b>	OH	N-Me	6	$\alpha$ -Naphthyl	39.4 (18.2–85.6)	
<b>16i</b>	OH	N-Me	6	$m$ -CF <sub>3</sub> -Ph	8.1 (4.6–14.2)	
<b>18g</b>	OH	N-Me	6	$m$ -CF <sub>3</sub> -Ph	95.6 (45.5–201.3)	

Table 2 (continued)

Compd.	R <sup>1</sup>	X	n	R <sup>2</sup>	K <sub>i</sub> (nM) <sup>a</sup> (95% confidence interval) or % displacement at fixed conc.	% Serotonin response on cAMP accumulation at 1 μM
<b>18h</b>	OH	N-Me	6	7-Cl-quinolin-4-yl	74%@10 <sup>-5</sup> 28%@10 <sup>-6</sup>	
<b>18j</b>	OH	N-Et	6	Ph	191.0 (51.8–703.7)	
<b>18k</b>	OH	N-Bn	6	Ph	137.5 (90.0–210.3)	

<sup>a</sup> K<sub>i</sub> values were calculated from IC<sub>50</sub> values, obtained from competition curves by the method of Cheng and Prusoff [33], and are the mean of four determinations performed in duplicate.

**5.1.1.7. 1-(4-Butylpiperazin-1-yl)-2-(5-methoxy-1H-indol-3-yl)ethane-1,2-dione (3g).** Yellowish solid (0.36 g, 82%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 0.92 (t, 3H, J = 7.0, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 1.39 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 2.43 (m, 6H, piperazine and CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 3.52 (m, 2H, piperazine), 3.76 (m, 2H, piperazine), 3.88 (s, 3H, OCH<sub>3</sub>), 6.89 (dd, 1H, J = 2.0 and 8.0, Ar-H), 7.24 (d, 1H, J = 8.0, Ar-H), 7.71 (brs, 1H, Ar-H), 7.79 (d, 1H, J = 2.0, Ar-H), 9.87 (brs, 1H, NH); ESI-MS (m/z): 344 (M + H)<sup>+</sup>.

**5.1.1.8. 1-(4-Cyclopentylpiperazin-1-yl)-2-(5-methoxy-1H-indol-3-yl)ethane-1,2-dione (3h).** Yellowish solid (0.28 g, 62%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.41–1.86 (m, 8H, c-pentyl), 2.55 (m, 5H, piperazine and CH), 3.55 (m, 2H, piperazine), 3.79 (m, 2H, piperazine), 3.90 (s, 3H, OCH<sub>3</sub>), 6.92 (dd, 1H, J = 2.0 and 10.0, Ar-H), 7.27 (d, 1H, J = 10, Ar-H), 7.77 (brs, 1H, Ar-H), 7.80 (d, 1H, J = 2.0, Ar-H), 9.65 (brs, 1H, NH); MS (EI, 70 eV) m/z: 355 (M<sup>+</sup>), 174 (100).

#### 5.1.2. General procedure for 3-[2-(4-substituted-piperazin-1-yl)ethyl]-1H-indoles **4a–h**


LiAlH<sub>4</sub> (0.2 g, 5.4 mmol) was added portionwise to an ice-cooled suspension of the suitable ethane-1,2-dione derivative **3a–h** (0.9 mmol) in dry THF (9 mL) under a N<sub>2</sub> atmosphere. Once the

addition was completed, the reaction mixture was refluxed for 1 h. After cooling to 0 °C, water was added dropwise to destroy the excess hydride and the resulting mixture was filtered on Celite. The filtrate was concentrated *in vacuo*, and partitioned between water and EtOAc. The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to afford a crude residue which was purified by flash chromatography on silica gel (cyclohexane/EtOAc 1:1 or EtOAc/MeOH 9:1 for **4g–h** as eluant).


**5.1.2.1. 5-Methoxy-3-[2-[4-(naphthalen-1-yl)piperazin-1-yl]ethyl]-1H-indole (4a).** Amorphous solid (0.23 g, 68%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 2.06 (m, 2H, CH<sub>2</sub>Ar), 2.82–3.09 (m, 6H, piperazine and CH<sub>2</sub>-piperazine), 3.24 (m, 4H, piperazine), 3.90 (s, 3H, OCH<sub>3</sub>), 6.88 (dd, 1H, J = 2.0 and 8.0, Ar-H), 7.07–7.30 (m, 4H, Ar-H), 7.39–7.60 (m, 4H, Ar-H), 7.84 (m, 1H, Ar-H), 7.92 (brs, 1H, NH), 8.23 (m, 1H, Ar-H); ESI-MS (m/z): 386 (M + H)<sup>+</sup>.

**5.1.2.2. 2-Ethyl-5-methoxy-3-[2-[4-(naphthalen-1-yl)piperazin-1-yl]ethyl]-1H-indole (4b).** Amorphous solid (0.32 g, 85%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.32 (t, 3H, J = 7.5, CH<sub>2</sub>CH<sub>3</sub>), 2.86 (m, 10H, CH<sub>2</sub>CH<sub>3</sub>, piperazine and ArCH<sub>2</sub>CH<sub>2</sub>N), 3.26 (m, 4H, piperazine), 3.90 (s, 3H, OCH<sub>3</sub>), 6.81 (dd, 1H, J = 2.0 and 8.0, Ar-H), 7.06 (d, 1H, J = 2.0, Ar-H), 7.19 (m, 2H, Ar-H), 7.50 (m, 4H, Ar-H), 7.73 (brs, 1H, NH), 7.85 (m, 1H, Ar-H), 8.24 (m, 1H, Ar-H); ESI-MS (m/z): 414 (M + H)<sup>+</sup>.

**5.1.2.3. 5-Methoxy-2-methyl-3-[2-[4-(naphthalen-1-yl)piperazin-1-yl]ethyl]-1H-indole (4c).** Amorphous solid (0.26 g, 74%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 2.40 (s, 3H, CH<sub>3</sub>), 2.70 (m, 2H, CH<sub>2</sub>Ar), 2.93 (m, 6H, piperazine and CH<sub>2</sub>-piperazine), 3.23 (m, 4H, piperazine), 3.88 (s,



**Fig. 3.** Superposition of *N,N*-dimethyl-5-hydroxytryptamine (**b** in Fig. 1, green carbons), 8-OH-DPAT (**c** in Fig. 1, orange carbons) and 3-[(2-(dimethylamino)ethyl)-(methylamino)phenol] (yellow carbons). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)



**Fig. 4.** Superposition of the best docking poses of **18a** (green carbons) and **18c** (yellow carbons) within the 5-HT<sub>7</sub> binding site. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)



3H, OCH<sub>3</sub>), 6.78 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 7.03 (d, 1H, *J* = 2.0, Ar-H), 7.15 (m, 2H, Ar-H), 7.38–7.59 (m, 4H, Ar-H), 7.69 (brs, 1H, NH), 7.83 (m, 1H, Ar-H), 8.23 (m, 1H, Ar-H); ESI-MS (*m/z*): 400 (*M* + *H*)<sup>+</sup>.

**5.1.2.4. 5-(Benzyloxy)-3-{2-[4-(biphenyl-3-yl)piperazin-1-yl]ethyl}-1H-indole (4d).** White solid (0.33 g, 75%); mp: 99–100 °C (EtOAc/n-hexane); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 2.78 (m, 6H, piperazine and CH<sub>2</sub>Ar), 3.00 (m, 2H, CH<sub>2</sub>-piperazine), 3.35 (m, 4H, piperazine), 5.14 (s, 2H, CH<sub>2</sub>Ph), 6.95–7.62 (m, 18H, Ar-H), 7.93 (brs, 1H, NH); MS (EI, 70 eV): *m/z* 487 (*M*<sup>+</sup>), 251 (100).

**5.1.2.5. 5-(Benzyloxy)-3-{2-[4-(biphenyl-4-yl)piperazin-1-yl]ethyl}-1H-indole (4e).** White solid (0.28 g, 64%); mp: 92–93 °C (EtOAc/n-hexane); <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>): δ = 2.70 (m, 6H, piperazine and CH<sub>2</sub>Ar), 2.97 (m, 2H, CH<sub>2</sub>-piperazine), 3.32 (m, 4H, piperazine), 5.19 (s, 2H, CH<sub>2</sub>Ph), 6.89 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 7.09 (m, 2H, Ar-H), 7.24 (m, 2H, Ar-H), 7.29–7.48 (m, 7H, Ar-H), 7.53–7.68 (m, 6H, Ar-H), 9.92 (brs, 1H, NH); ESI-MS (*m/z*): 488 (*M* + *H*)<sup>+</sup>.

**5.1.2.6. 5-(Benzyloxy)-3-{2-[4-(phenyl)piperazin-1-yl]ethyl}-1H-indole (4f).** White solid (0.26 g, 71%); mp: 133–134 °C (EtOAc/n-hexane); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 2.77 (m, 6H, piperazine and CH<sub>2</sub>Ar), 2.99 (m, 2H, CH<sub>2</sub>-piperazine), 3.29 (m, 4H, piperazine), 5.14 (s, 2H, CH<sub>2</sub>Ph), 6.85–7.00 (m, 4H, Ar-H), 7.05 (d, 1H, *J* = 2.0, Ar-H), 7.17 (d, 1H, *J* = 2.0, Ar-H), 7.25–7.53 (m, 8H, Ar-H), 7.92 (brs, 1H, NH); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 153.1, 151.3, 137.7, 131.6, 129.1, 128.5, 127.8, 127.6, 122.4, 119.7, 116.1, 114.1, 112.9, 111.8, 102.5, 71.0, 59.2, 53.3, 49.2, 23.0; MS (EI, 70 eV): *m/z* 411 (*M*<sup>+</sup>), 175 (100).

**5.1.2.7. 3-[2-(4-Butylpiperazin-1-yl)ethyl]-5-methoxy-1H-indole (4g).** Oil (0.18 g, 63%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 0.93 (t, 3H, *J* = 7.0, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 1.20–1.60 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 2.39–3.00 (m, 14H, piperazine, CH<sub>2</sub>Ar and 2 × CH<sub>2</sub>N), 3.87 (s, 3H, OCH<sub>3</sub>), 6.85 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 7.02 (d, 1H, *J* = 2.0, Ar-H), 7.03 (d, 1H, *J* = 2.0, Ar-H), 7.23 (d, 1H, *J* = 8.0, Ar-H), 8.04 (brs, 1H, NH); ESI-MS (*m/z*): 316 (*M* + *H*)<sup>+</sup>.

**5.1.2.8. 3-[2-(4-Cyclopentylpiperazin-1-yl)ethyl]-5-methoxy-1H-indole (4h).** Amorphous solid (0.21 g, 71%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.26–2.00 (m, 8H, *c*-pentyl), 2.45–3.00 (m, 13H, piperazine, CH<sub>2</sub>Ar, CH<sub>2</sub>N and CH), 3.86 (s, 3H, OCH<sub>3</sub>), 6.86 (dd, 1H, *J* = 2.5 and 8.0, Ar-H), 7.02 (d, 1H, *J* = 2.5, Ar-H), 7.05 (d, 1H, *J* = 2.5, Ar-H), 7.23 (d, 1H, *J* = 8.0, Ar-H), 7.93 (brs, 1H, NH); MS (EI, 70 eV): *m/z* 327 (*M*<sup>+</sup>), 167 (100).

### 5.1.3. O-Demethylation with BBr<sub>3</sub>. General procedure for 5-hydroxyindole derivatives **5i–k** and **5o–p**

A solution of BBr<sub>3</sub> (1 M in CH<sub>2</sub>Cl<sub>2</sub>, 3 mL, 3 mmol) diluted with dry CH<sub>2</sub>Cl<sub>2</sub> (6 mL) was added dropwise to a solution of the suitable methoxy derivative **4a–c** or **4g–h** (1 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (4 mL) at 0 °C, and the resulting mixture was stirred at room temperature for 2 h. The reaction mixture was neutralized with a 2 N aqueous solution of Na<sub>2</sub>CO<sub>3</sub> and extracted with EtOAc. The organic phases were combined, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure, to give a crude product that was purified by flash chromatography (silica gel; EtOAc for **5i–k** or EtOAc/MeOH 9:1 for **5o–p** as eluant) and crystallization.

**5.1.3.1. 3-[2-[4-(Naphthalen-1-yl)piperazin-1-yl]ethyl]-1H-indol-5-ol (5i).** Pink Solid (0.16 g, 44%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 2.96 (m, 8H, piperazine and CH<sub>2</sub>CH<sub>2</sub>Ar), 3.24 (m, 4H, piperazine), 6.80 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 6.99 (d, 1H, *J* = 2.0, Ar-H), 7.06 (d, 1H, *J* = 2.0, Ar-H), 7.10–7.27 (m, 2H, Ar-H), 7.48 (m, 4H, Ar-H and NH), 7.85 (m, 2H, Ar-H), 8.21 (m, 1H, Ar-H); <sup>13</sup>C NMR (acetone-*d*<sub>6</sub>): δ = 150.5, 149.9, 134.9, 128.9, 128.5, 128.3, 125.9, 125.7, 125.1, 123.6, 123.1,

122.9, 114.5, 112.7, 112.6, 111.5, 111.2, 102.5, 59.2, 53.6, 53.1, 22.9; MS (EI, 70 eV) *m/z*: 371 (*M*<sup>+</sup>), 146 (100); Anal. calcd for C<sub>24</sub>H<sub>25</sub>N<sub>3</sub>O: C 77.60, H 6.78, N 11.31, found C 77.81, H 6.90, N 11.03.

**5.1.3.2. 2-Ethyl-3-[2-[4-(naphthalen-1-yl)piperazin-1-yl]ethyl]-1H-indol-5-ol (5j).** White solid (0.28 g, 70%); mp: 200–201 °C (CH<sub>2</sub>Cl<sub>2</sub>/petroleum ether); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.29 (t, 3H, *J* = 7.5, CH<sub>3</sub>CH<sub>2</sub>), 2.76 (q, 2H, *J* = 7.5, CH<sub>3</sub>CH<sub>2</sub>), 2.77 (m, 2H, CH<sub>2</sub>Ar), 2.90 (m, 6H, piperazine and CH<sub>2</sub>CH<sub>2</sub>Ar), 3.25 (m, 4H, piperazine), 6.73 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 6.89 (d, 1H, *J* = 2.0, Ar-H), 7.14 (m, 2H, Ar-H), 7.37–7.59 (m, 4H, Ar-H), 7.69 (brs, 1H, NH), 7.85 (m, 1H, Ar-H), 8.21 (m, 1H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 149.7, 149.5, 138.3, 134.7, 130.3, 129.3, 128.9, 128.4, 125.9, 125.8, 125.3, 123.5, 123.5, 114.8, 111.1, 111.0, 107.9, 103.1, 59.4, 53.7, 52.6, 21.6, 19.5, 14.4; IR (nujol): ν = 3308 cm<sup>-1</sup>; ESI-MS (*m/z*): 400 (*M* + *H*)<sup>+</sup>; Anal. calcd for C<sub>26</sub>H<sub>29</sub>N<sub>3</sub>O: C 78.16, H 7.32, N 10.52, found C 78.01, H 7.21, N 10.29.

**5.1.3.3. 2-Methyl-3-[2-[4-(naphthalen-1-yl)piperazin-1-yl]ethyl]-1H-indol-5-ol (5k).** White solid (0.30 g, 79%); mp: 192–193 °C (CH<sub>2</sub>Cl<sub>2</sub>/petroleum ether); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 2.39 (s, 3H, CH<sub>3</sub>), 2.70 (m, 2H, CH<sub>2</sub>Ar), 2.90 (m, 6H, piperazine and CH<sub>2</sub>CH<sub>2</sub>Ar), 3.23 (m, 4H, piperazine), 6.68 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 6.91 (d, 1H, *J* = 2.0, Ar-H), 7.13 (m, 2H, Ar-H), 7.37–7.58 (m, 4H, Ar-H), 7.64 (brs, 1H, NH), 7.83 (m, 1H, Ar-H), 8.21 (m, 1H, Ar-H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>): δ = 150.6, 149.9, 134.8, 132.6, 130.1, 129.5, 128.7, 128.6, 126.5, 126.3, 125.8, 123.8, 123.4, 115.0, 111.1, 110.2, 107.9, 102.2, 59.4, 53.7, 53.2, 22.1, 11.8; IR (nujol): ν = 3238 cm<sup>-1</sup>; ESI-MS (*m/z*): 386 (*M* + *H*)<sup>+</sup>; Anal. calcd for C<sub>25</sub>H<sub>27</sub>N<sub>3</sub>O: C 77.89, H 7.06, N 10.90, found C 77.92, H 7.16, N 10.85.

**5.1.3.4. 3-[2-(4-Butylpiperazin-1-yl)ethyl]-1H-indol-5-ol (5o).** Amorphous solid (0.12 g, 42%); <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>): δ = 0.90 (t, 3H, *J* = 7.0, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 1.16–1.54 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 2.28–2.88 (m, 14H, piperazine, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub> and CH<sub>2</sub>CH<sub>2</sub>Ar), 6.68 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 6.96 (d, 1H, *J* = 2.0, Ar-H), 7.09 (d, 1H, *J* = 2.0, Ar-H), 7.18 (d, 1H, *J* = 8.0, Ar-H), 7.61 (brs, 1H, NH), 9.70 (brs, 1H, OH); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 150.0, 131.2, 128.0, 122.6, 112.8, 112.7, 111.9, 103.7, 58.9, 58.4, 52.6, 29.7, 28.6, 22.4, 20.7, 14.0; IR (nujol): ν = 3452, 3142 cm<sup>-1</sup>; ESI-MS (*m/z*): 302 (*M* + *H*)<sup>+</sup>; Anal. calcd for C<sub>18</sub>H<sub>27</sub>N<sub>3</sub>O: C 71.72, H 9.03, N 13.94, found C 71.63, H 9.21, N 14.02.

**5.1.3.5. 3-[2-(4-Cyclopentylpiperazin-1-yl)ethyl]-1H-indol-5-ol (5p).** Amorphous solid (0.15 g, 47%); <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>): δ = 1.32–1.89 (m, 8H, *c*-pentyl), 2.35–3.31 (m, 13H, piperazine, CH<sub>2</sub>CH<sub>2</sub>Ar and CH), 6.68 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 6.95 (d, 1H, *J* = 2.0, Ar-H), 7.09 (d, 1H, *J* = 2.0, Ar-H), 7.18 (d, 1H, *J* = 8.0, Ar-H), 7.63 (brs, 1H, NH), 9.70 (brs, 1H, OH); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 149.9, 131.2, 128.0, 122.6, 113.0, 112.8, 111.9, 103.7, 67.6, 59.0, 52.7, 51.8, 30.2, 24.1, 22.4; IR (nujol): ν = 3165, cm<sup>-1</sup>; ESI-MS (*m/z*): 314 (*M* + *H*)<sup>+</sup>; Anal. calcd for C<sub>19</sub>H<sub>27</sub>N<sub>3</sub>O: C 72.81, H 8.68, N 13.41, found C 72.85, H 8.74, N 13.58.

### 5.1.4. O-Debenzylation by hydrogenolysis. General procedure for 5-hydroxyindole derivatives **5l–n**

A solution of the benzyloxy derivative **4d–f** (0.73 mmol) was dissolved in anhydrous MeOH (12 mL) and then hydrogenated (4 atm) at room temperature in the presence of 10% Pd/C (0.053 g) for 6 h. The reaction mixture was filtered on Celite, the filtrate was evaporated under reduced pressure to yield a crude residue that was purified by crystallization.

**5.1.4.1. 3-[2-[4-(Biphenyl-3-yl)piperazin-1-yl]ethyl]-1H-indol-5-ol (5l).** White solid (148 mg, 51%); mp: 194–195 °C dec. (EtOAc/petroleum ether); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ = 2.64 (m, 6H, piperazine



and  $\text{CH}_2\text{Ar}$ ), 2.80 (m, 2H,  $\text{CH}_2\text{CH}_2\text{Ar}$ ), 3.25 (m, 4H, piperazine), 6.60 (dd, 1H,  $J = 2.0$  and 8.0, Ar-H), 6.85 (d, 1H,  $J = 2.0$ , Ar-H), 6.97–7.49 (m, 9H, Ar-H), 7.65 (m, 2H, Ar-H), 8.55 (brs, 1H, NH), 10.48 (brs, 1H, OH);  $^{13}\text{C}$  NMR (DMSO- $d_6$ ):  $\delta = 152.1, 150.6, 141.6, 141.4, 131.3, 129.9, 129.2, 128.4, 127.7, 127.3, 123.4, 117.8, 115.0, 114.1, 112.2, 112.1, 111.6, 102.7, 59.3, 53.3, 48.8, 23.1$ ; IR (nujol):  $\nu = 3198\text{ cm}^{-1}$ ; MS (EI, 70 eV)  $m/z$ : 397 ( $\text{M}^+$ ), 251 (100); Anal. calcd for  $\text{C}_{26}\text{H}_{27}\text{N}_3\text{O}$ : C 78.56, H 6.85, N 10.57, found C 78.77, H 6.97, N 10.75.

**5.1.4.2. 3-[2-[4-(Biphenyl-4-yl)piperazin-1-yl]ethyl]-1H-indol-5-ol (5m).** Gray solid (122 mg, 42%); mp: 216–217 °C dec. (MeOH);  $^1\text{H}$  NMR (acetone- $d_6$ ):  $\delta = 2.67$  (m, 6H, piperazine and  $\text{CH}_2\text{Ar}$ ), 2.88 (m, 2H,  $\text{CH}_2\text{CH}_2\text{Ar}$ ), 3.27 (m, 4H, piperazine), 6.68 (dd, 1H,  $J = 2.5$  and 8.0, Ar-H), 6.97 (d, 1H,  $J = 2.5$ , Ar-H), 6.97–7.61 (m, 12H, Ar-H and NH), 9.73 (brs, 1H, OH);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 150.8, 150.6, 140.5, 131.3, 130.8, 129.2, 128.4, 127.6, 126.7, 126.3, 123.5, 116.0, 112.1, 112.0, 111.7, 102.7, 59.0, 53.0, 48.3, 26.8$ ; IR (nujol):  $\nu = 3162\text{ cm}^{-1}$ ; ESI-MS ( $m/z$ ): 398 ( $\text{M} + \text{H}^+$ ); Anal. calcd for  $\text{C}_{26}\text{H}_{27}\text{N}_3\text{O}$ : C 78.56, H 6.85, N 10.57, found C 78.69, H 6.90, N 10.43.

**5.1.4.3. 3-[2-(4-Phenylpiperazin-1-yl)ethyl]-1H-indol-5-ol (5n).** White solid (145 mg, 62%); mp: 202–203 °C (MeOH/Et $_2\text{O}$ );  $^1\text{H}$  NMR (acetone- $d_6$ ):  $\delta = 2.69$  (m, 6H, piperazine and  $\text{CH}_2\text{Ar}$ ), 2.90 (m, 2H,  $\text{CH}_2\text{CH}_2\text{Ar}$ ), 3.22 (m, 4H, piperazine), 6.70 (ddd, 1H,  $J = 0.5, 2.0$  and 9.0, Ar-H), 6.78 (m, 1H, Ar-H), 6.94–6.99 (m, 3H, Ar-H), 7.13 (d, 1H,  $J = 2.0$ , Ar-H), 7.17–7.26 (m, 3H, Ar-H), 7.53 (brs, 1H, NH), 9.67 (brs, 1H, OH);  $^{13}\text{C}$  NMR (DMSO- $d_6$ ):  $\delta = 151.6, 150.6, 131.3, 129.3, 128.4, 123.4, 119.1, 115.7, 112.1, 112.1, 111.6, 102.7, 59.3, 53.2, 48.7, 23.1$ ; IR (nujol):  $\nu = 3205\text{ cm}^{-1}$ ; MS (EI, 70 eV)  $m/z$ : 321 ( $\text{M}^+$ ), 175 (100); Anal. calcd for  $\text{C}_{20}\text{H}_{23}\text{N}_3\text{O}$ : C 74.74, H 7.21, N 13.07, found C 74.96, H 7.19, N 13.32.

#### 5.1.5. N-Acylation of anilines with $\omega$ -bromoalkanoyl chlorides.

General procedure for  $\omega$ -bromoalkanamido derivatives **7a–c**, **12a–i** and **15a–d**

The suitable  $\omega$ -bromoalkanoyl chloride (0.7 mmol) was added dropwise to a stirred ice-cooled solution of the suitable aniline (0.33 mmol) and  $\text{NaHCO}_3$  (0.10 g, 1.22 mmol) in EtOAc (0.4 mL) and water (0.4 mL), and the resulting mixture was stirred at room temperature for 20 min. Water was added and the aqueous phase was extracted three times with EtOAc. The combined organic phases were washed with brine, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure to afford a crude residue that was used for the next step, without further purification.

**5.1.5.1. 2-Bromo-N-(3-methoxyphenyl)-N-methylacetamide (7a).** The product was prepared starting from 3-methoxy-N-methylaniline and 2-bromoethanoyl chloride (Scheme 2). Oil; MS (EI, 70 eV)  $m/z$ : 257–259 ( $\text{M}^+$ ), 136 (100) [27].

**5.1.5.2. N-[3-(Benzyloxy)phenyl]-2-bromo-N-methylacetamide (7b) [28].** The product was prepared starting from 3-benzyloxyaniline and 2-bromoethanoyl chloride (Scheme 2). Oil; ESI-MS ( $m/z$ ): 334–336 ( $\text{M} + \text{H}^+$ ).

**5.1.5.3. 2-Bromo-N-methyl-N-(3-nitrophenyl)acetamide (7c) [27].** The product was prepared starting from 3-nitro-N-methylaniline and 2-bromoethanoyl chloride (Scheme 2). Oil; ESI-MS ( $m/z$ ): 272–274 ( $\text{M} + \text{H}^+$ ).

**5.1.5.4. 3-Bromo-N-(3-methoxyphenyl)-N-methylpropanamide (12a).** The product was prepared starting from 3-methoxy-N-methylaniline and 3-bromopropanoyl chloride (Scheme 3). Oil; ESI-MS ( $m/z$ ): 272–274 ( $\text{M} + \text{H}^+$ ).

**5.1.5.5. 3-Bromo-N-methyl-N-m-tolylpropanamide (12b).** The product was prepared starting from 3-methyl-N-methylaniline and 3-bromopropanoyl chloride (Scheme 3). Oil; ESI-MS ( $m/z$ ): 256–258 ( $\text{M} + \text{H}^+$ ).

**5.1.5.6. 3-Bromo-N-(3-bromophenyl)-N-methylpropanamide (12c).** The product was prepared starting from 3-bromo-N-methylaniline and 3-bromopropanoyl chloride (Scheme 3). Oil; ESI-MS ( $m/z$ ): 320–322–324 ( $\text{M} + \text{H}^+$ ).

**5.1.5.7. 3-Bromo-N-(3-chlorophenyl)-N-methylpropanamide (12d).** The product was prepared starting from 3-chloro-N-methylaniline and 3-bromopropanoyl chloride (Scheme 3). Oil; ESI-MS ( $m/z$ ): 276–278–280 ( $\text{M} + \text{H}^+$ ).

**5.1.5.8. 3-Bromo-N-(3-fluorophenyl)-N-methylpropanamide (12e).** The product was prepared starting from 3-fluoro-N-methylaniline and 3-bromopropanoyl chloride (Scheme 3). Oil; ESI-MS ( $m/z$ ): 260–262 ( $\text{M} + \text{H}^+$ ).

**5.1.5.9. 3-Bromo-N-methyl-N-(3-nitrophenyl)propanamide (12f).** The product was prepared starting from 3-nitro-N-methylaniline and 3-bromopropanoyl chloride (Scheme 3). Oil; ESI-MS ( $m/z$ ): 300–302 ( $\text{M} + \text{H}^+$ ).

**5.1.5.10. Ethyl 3-(3-bromo-N-methylpropanamido)benzoate (12g).** The product was prepared starting from ethyl 3-aminobenzoate and 3-bromopropanoyl chloride (Scheme 3). Oil; ESI-MS ( $m/z$ ): 287–289 ( $\text{M} + \text{H}^+$ ).

**5.1.5.11. 3-Bromo-N-(3-methoxyphenyl)propanamide (12h).** The product was prepared starting from 3-methoxyaniline and 3-bromopropanoyl chloride (Scheme 3). Oil; ESI-MS ( $m/z$ ): 258–260 ( $\text{M} + \text{H}^+$ ).

**5.1.5.12. N-[3-(Benzyloxy)phenyl]-3-bromo-N-methylpropanamide (12i).** The product was prepared starting from 3-benzyloxy-N-methylaniline [17] and 3-bromopropanoyl chloride (Scheme 3). Oil; MS (EI, 70 eV)  $m/z$ : 347–349 ( $\text{M}^+$ ), 91 (100).

**5.1.5.13. N-[3-(Benzyloxy)phenyl]-4-bromo-N-methylbutanamide (15a).** The product was prepared starting from 3-benzyloxy-N-methylaniline [17] and 4-bromobutanoyl chloride (Scheme 4). Oil; MS (EI, 70 eV)  $m/z$ : 361–363 ( $\text{M}^+$ ), 91 (100).

**5.1.5.14. N-[3-(Benzyloxy)phenyl]-5-bromo-N-methylpentanamide (15b).** The product was prepared starting from 3-benzyloxy-N-methylaniline [17] and 5-bromopentanoyl chloride (Scheme 4). Oil;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 1.67$ – $1.87$  (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.09 (t, 2H,  $J = 7.0$ ,  $\text{CH}_2\text{CO}$ ), 3.25 (s, 3H,  $\text{CH}_3$ ), 3.31 (t, 2H,  $J = 6.5$ ,  $\text{CH}_2\text{Br}$ ), 5.09 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.76–6.79 (m, 2H, Ar-H), 7.00 (dd, 1H,  $J = 2.0$  and 8.0, Ar-H), 7.27–7.46 (m, 6H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 172.6, 159.6, 145.0, 136.3, 130.6, 128.7, 128.2, 127.5, 119.6, 114.3, 114.0, 70.2, 37.3, 33.3, 33.0, 32.3, 24.1$ ; MS (EI, 70 eV)  $m/z$ : 375–377 ( $\text{M}^+$ ), 91 (100).

**5.1.5.15. N-[3-(Benzyloxy)phenyl]-6-bromo-N-methylhexanamide (15c).** The product was prepared starting from 3-benzyloxy-N-methylaniline [17] and 3-bromohexanoyl chloride (Scheme 4). Oil; MS (EI, 70 eV)  $m/z$ : 389–391 ( $\text{M}^+$ ), 91 (100).

**5.1.5.16. N-[3-(Benzyloxy)phenyl]-6-bromohexanamide (15d).** The product was prepared starting from 3-benzyloxyaniline and 3-bromohexanoyl chloride (Scheme 4). Oil; ESI-MS ( $m/z$ ): 375–377 ( $\text{M} + \text{H}^+$ ).

#### 5.1.6. General procedure for $\omega$ -(4-substituted-piperazin-1-yl)alkanamides **8d–g**, **13b–k** and **16a–h**

The suitable *N*-substituted-piperazine (0.6 mmol) and TEA (0.83 mL) were added to a solution of the appropriate crude  $\omega$ -bromoalkanamide (from 0.5 mmol of substituted-aniline) in DMF (0.14 mL) and the resulting mixture was stirred at room temperature for 1–16 h, monitoring the progress of the reaction by TLC on silica gel. The reaction mixture was poured into water, and extracted with EtOAc. After drying over Na<sub>2</sub>SO<sub>4</sub>, the combined organic phases were concentrated *in vacuo* and the resulting crude product was purified by column flash chromatography on silica gel (EtOAc or cyclohexane/EtOAc, 1:1 for **8f** as eluant) and/or crystallization.

**5.1.6.1. *N*-(3-Methoxyphenyl)-*N*-methyl-2-(4-phenylpiperazin-1-yl)acetamide (**8d**).** The product was prepared starting from **7a** and *N*-phenylpiperazine (Scheme 2). Oil (154 mg, 91%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.67 (m, 4H, piperazine), 3.05 (s, 2H, CH<sub>2</sub>CO), 3.21 (m, 4H, piperazine), 3.29 (s, 3H, NCH<sub>3</sub>), 3.85 (s, 3H, OCH<sub>3</sub>), 6.86 (m, 6H, Ar-H), 7.30 (m, 3H, Ar-H); MS (EI, 70 eV): *m/z* 339 (M<sup>+</sup>), 175 (100).

**5.1.6.2. 2-(4-Benzylpiperazin-1-yl)-*N*-(3-methoxyphenyl)-*N*-methylacetamide (**8e**).** The product was prepared starting from **7a** and *N*-benzylpiperazine (Scheme 2). Oil (116 mg, 66%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.51 (m, 8H, piperazine), 2.96 (s, 2H, CH<sub>2</sub>CO), 3.26 (s, 3H, NCH<sub>3</sub>), 3.52 (s, 2H, CH<sub>2</sub>Ph), 3.83 (s, 3H, OCH<sub>3</sub>), 6.81 (m, 3H, Ar-H), 7.32 (m, 6H, Ar-H); MS (EI, 70 eV): *m/z* 353 (M<sup>+</sup>), 91 (100).

**5.1.6.3. *N*-[3-(Benzyloxy)phenyl]-2-(4-phenylpiperazin-1-yl)acetamide (**8f**).** The product was prepared starting from **7b** and *N*-phenylpiperazine (Scheme 2). White solid (160 mg, 80%); mp 119–120 °C (EtOAc/petroleum ether); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.82 (m, 4H, piperazine), 3.24 (s, 2H, CH<sub>2</sub>CO), 3.29 (m, 4H, piperazine), 5.09 (s, 2H, CH<sub>2</sub>Ph), 6.76 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 6.98 (m, 4H, Ar-H), 7.25–7.48 (m, 9H, Ar-H), 9.19 (brs, 1H, NH); ESI-MS (*m/z*): 402 (M + H)<sup>+</sup>.

**5.1.6.4. *N*-Methyl-*N*-(3-nitrophenyl)-2-(4-phenylpiperazin-1-yl)acetamide (**8g**).** The product was prepared starting from **7c** and *N*-phenylpiperazine (Scheme 2). Yellow oil (136 mg, 77%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.71 (m, 4H, piperazine), 3.14 (s, 2H, CH<sub>2</sub>CO), 3.20 (m, 4H, piperazine), 3.31 (s, 3H, NCH<sub>3</sub>), 6.87 (m, 3H, Ar-H), 7.28 (m, 2H, Ar-H), 7.62 (m, 2H, Ar-H), 8.14 (m, 1H, Ar-H), 8.24 (m, 1H, Ar-H); ESI-MS (*m/z*): 355 (M + H)<sup>+</sup>.

**5.1.6.5. *N*-Methyl-3-(4-phenylpiperazin-1-yl)-*N*-m-tolylpropanamide (**13b**).** The product was prepared starting from **12b** and *N*-phenylpiperazine (Scheme 3). Amorphous solid (106 mg, 63%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.33 (apt, 2H, *J* = 7.0, CH<sub>2</sub>CO), 2.39 (s, 3H, Ar-CH<sub>3</sub>), 2.51 (m, 4H, piperazine), 2.76 (apt, 2H, *J* = 7.0, COCH<sub>2</sub>CH<sub>2</sub>N), 3.15 (m, 4H, piperazine), 3.27 (s, 3H, NCH<sub>3</sub>), 6.82–7.35 (m, 9H, Ar-H); ESI-MS (*m/z*): 338 (M + H)<sup>+</sup>.

**5.1.6.6. *N*-(3-Bromophenyl)-*N*-methyl-3-(4-phenylpiperazin-1-yl)propanamide (**13c**).** The product was prepared starting from **12c** and *N*-phenylpiperazine (Scheme 3). Amorphous solid (140 mg, 70%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.34 (apt, 2H, *J* = 7.0, CH<sub>2</sub>CO), 2.52 (m, 4H, piperazine), 2.75 (apt, 2H, *J* = 7.0, COCH<sub>2</sub>CH<sub>2</sub>N), 3.16 (m, 4H, piperazine), 3.29 (s, 3H, NCH<sub>3</sub>), 6.93–7.38 (m, 9H, Ar-H); ESI-MS (*m/z*): 402–404 (M + H)<sup>+</sup>.

**5.1.6.7. *N*-(3-Chlorophenyl)-*N*-methyl-3-(4-phenylpiperazin-1-yl)propanamide (**13d**).** The product was prepared starting from **12d** and *N*-phenylpiperazine (Scheme 3). Amorphous solid (144 mg, 81%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.34 (apt, 2H, *J* = 7.0, CH<sub>2</sub>CO), 2.53 (m,

4H, piperazine), 2.76 (apt, 2H, *J* = 7.0, COCH<sub>2</sub>CH<sub>2</sub>N), 3.16 (m, 4H, piperazine), 3.28 (s, 3H, NCH<sub>3</sub>), 6.90 (m, 3H, Ar-H), 7.10–7.45 (m, 6H, Ar-H); ESI-MS (*m/z*): 358 (M + H)<sup>+</sup>.

**5.1.6.8. *N*-(3-Fluorophenyl)-*N*-methyl-3-(4-phenylpiperazin-1-yl)propanamide (**13e**).** The product was prepared starting from **12e** and *N*-phenylpiperazine (Scheme 3). Amorphous solid (128 mg, 75%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.36 (apt, 2H, *J* = 7.0, CH<sub>2</sub>CO), 2.53 (m, 4H, piperazine), 2.76 (apt, 2H, *J* = 7.0, COCH<sub>2</sub>CH<sub>2</sub>N), 3.16 (m, 4H, piperazine), 3.29 (s, 3H, NCH<sub>3</sub>), 6.82–7.47 (m, 9H, Ar-H); ESI-MS (*m/z*): 342 (M + H)<sup>+</sup>.

**5.1.6.9. *N*-Methyl-*N*-(3-nitrophenyl)-3-(4-phenylpiperazin-1-yl)propanamide (**13f**).** The product was prepared starting from **12f** and *N*-phenylpiperazine (Scheme 3). Yellow amorphous solid (147 mg, 80%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.38 (apt, 2H, *J* = 7.0, CH<sub>2</sub>CO), 2.58 (m, 4H, piperazine), 2.81 (apt, 2H, *J* = 7.0, COCH<sub>2</sub>CH<sub>2</sub>N), 3.18 (m, 4H, piperazine), 3.36 (s, 3H, NCH<sub>3</sub>), 6.88 (m, 3H, Ar-H), 7.25 (m, 2H, Ar-H), 7.64 (m, 2H, Ar-H), 8.13 (m, 1H, Ar-H), 8.23 (m, 1H, Ar-H); ESI-MS (*m/z*): 369 (M + H)<sup>+</sup>.

**5.1.6.10. Ethyl 3-[3-(4-phenylpiperazin-1-yl)propanamido]benzoate (**13g**).** The product was prepared starting from **12g** and *N*-phenylpiperazine (Scheme 3). White solid (86 mg, 45%); mp 127–128 °C (EtOAc) <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.30 (t, 3H, *J* = 7.0, CH<sub>2</sub>CH<sub>3</sub>), 2.60 (m, 2H, CH<sub>2</sub>CO), 2.79 (m, 6H, piperazine and COCH<sub>2</sub>CH<sub>2</sub>N), 3.31 (m, 4H, piperazine), 4.31 (q, 2H, *J* = 7.0, CH<sub>2</sub>CH<sub>3</sub>), 6.93 (m, 3H, Ar-H), 7.25–7.41 (m, 3H, Ar-H), 7.74 (m, 1H, Ar-H), 7.86 (m, 1H, Ar-H), 8.04 (m, 1H, Ar-H), 11.15 (brs, 1H, NH); ESI-MS (*m/z*): 382 (M + H)<sup>+</sup>.

**5.1.6.11. *N*-(3-Methoxyphenyl)-3-(4-phenylpiperazin-1-yl)propanamide (**13h**).** The product was prepared starting from **12h** and *N*-phenylpiperazine (Scheme 3). Amorphous solid (151 mg, 89%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.59 (m, 2H, CH<sub>2</sub>CO), 2.80 (m, 6H, piperazine and COCH<sub>2</sub>CH<sub>2</sub>N), 3.33 (m, 4H, piperazine), 3.80 (s, 3H, CH<sub>3</sub>), 6.64 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 6.95 (m, 4H, Ar-H), 7.16–7.43 (m, 4H, Ar-H), 10.95 (brs, 1H, NH); ESI-MS (*m/z*): 340 (M + H)<sup>+</sup>.

**5.1.6.12. *N*-[3-(Benzyloxy)phenyl]-*N*-methyl-3-(4-phenylpiperazin-1-yl)propanamide (**13i**).** The product was prepared starting from **12i** and *N*-phenylpiperazine (Scheme 3). Oil (131 mg, 61%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.35 (apt, 2H, *J* = 7.0, CH<sub>2</sub>CO), 2.52 (m, 4H, piperazine), 2.76 (apt, 2H, *J* = 7.0, COCH<sub>2</sub>CH<sub>2</sub>N), 3.16 (m, 4H, piperazine), 3.27 (s, 3H, NCH<sub>3</sub>), 5.09 (s, 2H, CH<sub>2</sub>Ph), 6.81–6.99 (m, 6H, Ar-H), 7.22–7.43 (m, 8H, Ar-H); MS (EI, 70 eV): *m/z* 429 (M<sup>+</sup>), 91 (100).

**5.1.6.13. *N*-(3-Methoxyphenyl)-*N*-methyl-3-[4-(naphthalen-1-yl)piperazin-1-yl]propanamide (**13j**).** The product was prepared starting from **12a** and 1-(naphthalen-1-yl)piperazine (Scheme 3). Amorphous solid (103 mg, 51%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.29 (apt, 2H, *J* = 7.0, CH<sub>2</sub>CO), 2.61 (m, 4H, piperazine), 2.70 (apt, 2H, *J* = 7.0, COCH<sub>2</sub>CH<sub>2</sub>N), 3.10 (m, 4H, piperazine), 3.24 (s, 3H, NCH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 6.82 (m, 3H, Ar-H), 7.09 (m, 1H, Ar-H), 7.30–7.61 (m, 5H, Ar-H), 7.85 (m, 1H, Ar-H), 8.21 (m, 1H, Ar-H); ESI-MS (*m/z*): 404 (M + H)<sup>+</sup>.

**5.1.6.14. *N*-(3-Methoxyphenyl)-*N*-methyl-3-[4-(naphthalen-2-yl)piperazin-1-yl]propanamide (**13k**).** The product was prepared starting from **12a** and 1-(naphthalen-2-yl)piperazine (Scheme 3). Oil (95 mg, 47%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.34 (apt, 2H, *J* = 7.0, CH<sub>2</sub>CO), 2.58 (m, 4H, piperazine), 2.79 (apt, 2H, *J* = 7.0, COCH<sub>2</sub>CH<sub>2</sub>N), 3.27 (m, 4H, piperazine), 3.28 (s, 3H, NCH<sub>3</sub>), 3.85 (s, 3H, OCH<sub>3</sub>), 6.75–6.93 (m, 3H, Ar-H), 7.08 (m, 1H, Ar-H), 7.22–7.44 (m, 4H, Ar-H), 7.70 (m, 3H, Ar-H); ESI-MS (*m/z*): 404 (M + H)<sup>+</sup>.

5.1.6.15. *N*-[3-(Benzyloxy)phenyl]-*N*-methyl-4-(4-phenylpiperazin-1-yl)butanamide (**16a**). The product was prepared starting from **15a** and *N*-phenylpiperazine (Scheme 4). Oil (51 mg, 23%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.81 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.14 (apt, 2H,  $J$  = 7.0,  $\text{CH}_2\text{CO}$ ), 2.32 (apt, 2H,  $J$  = 7.0,  $\text{COCH}_2\text{CH}_2\text{CH}_2\text{N}$ ), 2.55 (m, 4H, piperazine), 3.16 (m, 4H, piperazine), 3.26 (s, 3H,  $\text{NCH}_3$ ), 5.08 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.79–6.99 (m, 6H, Ar-H), 7.26–7.43 (m, 8H, Ar-H); MS (EI, 70 eV):  $m/z$  443 ( $\text{M}^+$ ), 91 (100).

5.1.6.16. *N*-[3-(Benzyloxy)phenyl]-*N*-methyl-5-(4-phenylpiperazin-1-yl)pentanamide (**16b**). The product was prepared starting from **15b** and *N*-phenylpiperazine (Scheme 4). Oil (94 mg, 41%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.54 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.13 (apt, 2H,  $J$  = 7.0,  $\text{CH}_2\text{CO}$ ), 2.34 (t, 2H,  $J$  = 7.5,  $\text{CH}_2$ -piperazine), 2.57 (m, 4H, piperazine), 3.19 (m, 4H, piperazine), 3.25 (s, 3H,  $\text{CH}_3$ ), 5.08 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.78–6.99 (m, 6H, Ar-H), 7.23–7.46 (m, 8H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 172.9, 159.6, 151.3, 145.3, 136.4, 130.4, 129.1, 128.7, 128.2, 127.5, 119.8, 119.6, 116.0, 114.2, 114.0, 70.2, 58.3, 53.2, 49.0, 37.2, 33.8, 26.4, 23.5; MS (EI, 70 eV):  $m/z$  457 ( $\text{M}^+$ ), 91 (100).

5.1.6.17. *N*-[3-(Benzyloxy)phenyl]-*N*-methyl-6-(4-phenylpiperazin-1-yl)hexanamide (**16c**). The product was prepared starting from **15c** and *N*-phenylpiperazine (Scheme 4). Oil (94 mg, 40%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.27 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.57 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.09 (apt, 2H,  $J$  = 7.5,  $\text{CH}_2\text{CO}$ ), 2.42 (apt, 2H,  $J$  = 7.5,  $\text{CH}_2\text{N}$ ), 2.66 (m, 4H, piperazine), 3.24 (m, 4H, piperazine), 3.25 (s, 3H,  $\text{CH}_3$ ), 5.09 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.77–6.99 (m, 6H, Ar-H), 7.24–7.45 (m, 8H, Ar-H); MS (EI, 70 eV):  $m/z$  471 ( $\text{M}^+$ ), 91 (100).

5.1.6.18. *N*-[3-(Benzyloxy)phenyl]-6-(4-phenylpiperazin-1-yl)hexanamide (**16d**). The product was prepared starting from **15d** and *N*-phenylpiperazine (Scheme 4). Oil (103 mg, 45%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.28 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.61 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.13 (apt, 2H,  $J$  = 7.5,  $\text{CH}_2\text{CO}$ ), 2.43 (apt, 2H,  $J$  = 7.5,  $\text{CH}_2$ -piperazine), 2.71 (m, 4H, piperazine), 3.21 (m, 4H, piperazine), 5.08 (s, 2H,  $\text{CH}_2\text{-Ph}$ ), 6.73–6.98 (m, 5H, Ar-H), 7.21–7.55 (m, 9H, Ar-H), 9.03 (brs, 1H, NH); ESI-MS:  $m/z$  458 ( $\text{M} + \text{H}^+$ ).

5.1.6.19. *N*-[3-(Benzyloxy)phenyl]-*N*-methyl-6-(4-phenylpiperidin-1-yl)hexanamide (**16e**). The product was prepared starting from **15e** and 4-phenylpiperidine (Scheme 4). Oil (94 mg, 40%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.19 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.58 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.81 (m, 2H, piperidine), 2.01–2.25 (m, 6H,  $\text{CH}_2\text{N}$ ,  $\text{CH}_2\text{CO}$  and piperidine), 2.45 (m, 1H, CH), 2.60 (m, 4H, piperidine), 3.20 (s, 3H,  $\text{CH}_3$ ), 5.01 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.71 (m, 2H, Ar-H), 6.92 (m, 1H, Ar-H), 7.16–7.42 (m, 11H, Ar-H); ESI-MS:  $m/z$  471 ( $\text{M} + \text{H}^+$ ).

5.1.6.20. *N*-[3-(Benzyloxy)phenyl]-*N*-methyl-6-[4-(naphthalen-1-yl)piperazin-1-yl]hexanamide (**16f**). The product was prepared starting from **15c** and 1-(naphthalen-1-yl)piperazine (Scheme 4). Oil (99 mg, 38%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.32 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.61 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.78 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.13 (apt, 2H,  $J$  = 7.0,  $\text{CH}_2\text{CO}$ ), 2.79 (m, 2H,  $\text{CH}_2$ -piperazine), 2.91–3.51 (m, 8H, piperazine), 3.22 (s, 3H,  $\text{CH}_3$ ), 5.09 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.80 (m, 2H, Ar-H), 6.99 (m, 1H, Ar-H), 6.38 (m, 1H, Ar-H), 7.28–7.55 (m, 9H, Ar-H), 7.61 (m, 1H, Ar-H), 7.84 (m, 1H, Ar-H), 8.13 (m, 1H, Ar-H); ESI-MS:  $m/z$  522 ( $\text{M} + \text{H}^+$ ).

5.1.6.21. *N*-[3-(Benzyloxy)phenyl]-*N*-methyl-6-[4-[3-(trifluoromethyl)phenyl]piperazin-1-yl]hexanamide (**16g**). The product was prepared starting from **15c** and 1-[3-(trifluoromethyl)phenyl]piperazine (Scheme 4). Oil (124 mg, 46%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.19 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.56 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.02 (apt, 2H,  $J$  = 7.0,  $\text{CH}_2\text{CO}$ ), 2.55 (m, 2H,  $\text{CH}_2$ -piperazine), 2.68 (m, 4H,

piperazine), 3.19 (s, 3H,  $\text{CH}_3$ ), 3.31 (m, 4H, piperazine), 5.01 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.76 (m, 2H, Ar-H), 6.86–7.08 (m, 4H, Ar-H), 7.19–7.40 (m, 7H, Ar-H). ESI-MS:  $m/z$  540 ( $\text{M} + \text{H}^+$ ).

5.1.6.22. *N*-[3-(Benzyloxy)phenyl]-6-[4-(7-chloroquinolin-4-yl)piperazin-1-yl]-*N*-methylhexanamide (**16h**). The product was prepared starting from **15c** and 7-chloro-4-(piperazin-1-yl)quinoline (Scheme 4). Oil (81 mg, 29%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.27 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.58 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.97 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.14 (m, 2H,  $\text{CH}_2\text{CO}$ ), 3.01–3.68 (m, 6H, 4.18 (m, 2H), 4.42 (m, 2H), 3.18 (s, 3H,  $\text{CH}_3$ ), 5.01 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.75 (m, 2H, Ar-H), 6.92 (m, 1H, Ar-H), 7.16 (m, 1H, Ar-H), 7.18–7.41 (m, 6H, Ar-H), 7.48 (m, 1H, Ar-H), 7.88 (m, 1H, Ar-H), 8.22 (m, 1H, Ar-H), 8.32 (m, 1H, Ar-H). ESI-MS:  $m/z$  557 ( $\text{M} + \text{H}^+$ ).

#### 5.1.7. Reduction of the alkanamido derivatives with $\text{LiAlH}_4$ . General procedure for compounds **9d–f** and **17a–h**

$\text{LiAlH}_4$  (0.59 g, 15.5 mmol) was added portionwise to an ice-cooled solution of the suitable alkanamido derivative **8d–f**, **16a–h** (2.5 mmol) in dry THF (30 mL) under a  $\text{N}_2$  atmosphere. Once the addition was completed, the reaction mixture was refluxed for 2 h. After cooling to 0 °C, water was added dropwise to destroy the excess hydride and the resulting mixture was filtered on Celite. The filtrate was concentrated *in vacuo* and partitioned between water and EtOAc. The combined organic layers were washed with brine, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated under reduced pressure to afford a crude residue which was purified by flash chromatography on silica gel.

5.1.7.1. 3-Methoxy-*N*-methyl-*N*-[2-(4-phenylpiperazin-1-yl)ethyl]aniline (**9d**). The product was prepared starting from **8d** (Scheme 2). Purification by flash chromatography on silica gel (EtOAc as eluant). Oil (292 mg, 36%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 2.63 (apt, 2H,  $J$  = 7.5,  $\text{CH}_2\text{CH}_2\text{NCH}_3$ ), 2.70 (m, 4H, piperazine), 2.98 (s, 3H,  $\text{NCH}_3$ ), 3.23 (m, 4H, piperazine), 3.54 (apt, 2H,  $J$  = 7.5,  $\text{CH}_2\text{NCH}_3$ ), 3.82 (s, 3H,  $\text{OCH}_3$ ), 6.35 (m, 3H, Ar-H), 6.91 (m, 3H, Ar-H), 7.23 (m, 3H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 160.9, 151.3, 150.6, 130.0, 129.1, 119.7, 116.0, 105.3, 101.0, 98.7, 55.1, 54.9, 53.7, 50.5, 49.1, 38.7; MS (EI, 70 eV):  $m/z$  325 ( $\text{M}^+$ ), 150 (100).

5.1.7.2. *N*-[2-(4-Benzyloxy)piperazin-1-yl]ethyl-3-methoxy-*N*-methyl-aniline (**9e**). The product was prepared starting from **8e** (Scheme 2). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 8:2 as eluant). Oil (246 mg, 29%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 2.53 (m, 10H, piperazine and  $\text{CH}_2\text{CH}_2\text{NCH}_3$ ), 2.93 (s, 3H,  $\text{NCH}_3$ ), 3.46 (m, 2H,  $\text{CH}_2\text{NCH}_3$ ), 3.51 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 3.78 (s, 3H,  $\text{OCH}_3$ ), 6.29 (m, 3H, Ar-H), 7.08–7.33 (m, 6H, Ar-H); MS (EI, 70 eV):  $m/z$  339 ( $\text{M}^+$ ), 91 (100).

5.1.7.3. 3-(Benzyloxy)-*N*-[2-(4-phenylpiperazin-1-yl)ethyl]aniline (**9f**). The product was prepared starting from **8f** (Scheme 2). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 4:6 as eluant). Oil (822 mg, 85%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 2.69 (m, 6H, piperazine and  $\text{CH}_2\text{CH}_2\text{NH}$ ), 3.22 (m, 6H, piperazine and  $\text{CH}_2\text{NH}$ ), 4.38 (brs, 1H, NH), 5.06 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.35 (m, 3H, Ar-H), 6.93 (m, 3H, Ar-H), 7.07–7.45 (m, 8H, Ar-H); MS (EI, 70 eV):  $m/z$  387 ( $\text{M}^+$ ), 91 (100).

5.1.7.4. 3-(Benzyloxy)-*N*-methyl-*N*-[4-(4-phenylpiperazin-1-yl)butyl]aniline (**17a**). The product was prepared starting from **16a** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 7:3 as eluant). Amorphous solid (257 mg, 24%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.63 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.46 (m, 2H,  $\text{CH}_2$ -piperazine), 2.65 (m, 4H, piperazine), 2.93 (s, 3H,  $\text{CH}_3$ ), 3.26 (m, 4H, piperazine), 3.34 (m, 2H,  $\text{CH}_2\text{NCH}_3$ ), 5.06 (s, 2H,  $\text{CH}_2\text{Ar}$ ), 6.35



(m, 3H, Ar-H), 6.92 (m, 3H, Ar-H), 7.10–7.48 (m, 8H, Ar-H); MS (EI, 70 eV):  $m/z$  429 ( $M^+$ ), 91 (100).

**5.1.7.5. 3-(Benzyloxy)-N-methyl-N-[5-(4-phenylpiperazin-1-yl)pentyl]aniline (17b).** The product was prepared starting from **16b** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 7:3 as eluant). Oil (764 mg, 69%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.36 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.61 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.44 (apt, 2H,  $J$  = 8.0,  $\text{CH}_2$ -piperazine), 2.65 (m, 4H, piperazine), 2.92 (s, 3H,  $\text{CH}_3$ ), 3.25 (m, 4H, piperazine), 3.31 (t, 2H,  $J$  = 7.5,  $\text{CH}_2\text{NCH}_3$ ), 5.07 (s, 2H,  $\text{CH}_2\text{Ar}$ ), 6.35 (m, 3H, Ar-H), 6.91 (m, 3H, Ar-H), 7.11–7.49 (m, 8H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 160.1, 151.2, 150.7, 137.4, 129.8, 129.1, 128.6, 127.9, 127.6, 119.8, 116.1, 105.5, 101.4, 99.6, 69.9, 58.6, 53.2, 52.7, 49.0, 38.4, 26.7, 26.6, 25.1; MS (EI, 70 eV):  $m/z$  443 ( $M^+$ ), 91 (100).

**5.1.7.6. 3-(Benzyloxy)-N-methyl-N-[6-(4-phenylpiperazin-1-yl)hexyl]aniline (17c).** The product was prepared starting from **16c** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 7:3 as eluant). Oil (320 mg, 28%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.35 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.58 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.45 (m, 2H,  $\text{CH}_2$ -piperazine), 2.66 (m, 4H, piperazine), 2.92 (s, 3H,  $\text{CH}_3$ ), 3.29 (m, 6H, piperazine and  $\text{CH}_2\text{NCH}_3$ ), 5.06 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.34 (m, 3H, Ar-H), 6.86 (m, 3H, Ar-H), 7.10–7.48 (m, 8H, Ar-H); MS (EI, 70 eV):  $m/z$  457 ( $M^+$ ), 91 (100).

**5.1.7.7. 3-(Benzyloxy)-N-methyl-N-[6-(4-phenylpiperazin-1-yl)hexyl]aniline (17d).** The product was prepared starting from **16d** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc as eluant). Oil (443 mg, 40%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.38 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.59 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.50 (m, 2H,  $\text{CH}_2$ -piperazine), 2.71 (m, 4H, piperazine), 3.11 (m, 2H,  $\text{CH}_2\text{NH}$ ), 3.30 (m, 4H, piperazine), 3.65 (brs, 1H, NH), 5.05 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.21 (m, 3H, Ar-H), 6.83–7.61 (m, 11H, Ar-H); ESI-MS:  $m/z$  444 ( $M + \text{H}$ ) $^+$ .

**5.1.7.8. 3-(Benzyloxy)-N-methyl-N-[6-(4-phenylpiperidin-1-yl)hexyl]aniline (17e).** The product was prepared starting from **16e** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 8:2 as eluant). Oil (513 mg, 45%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.22 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.57 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.83 (m, 2H, piperidine), 1.88–2.57 (m, 9H, piperidine, and  $\text{CH}_2\text{N}$ ), 2.95 (s, 3H,  $\text{CH}_3$ ), 3.24 (m, 2H,  $\text{CH}_2\text{NCH}_3$ ), 5.02 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.70 (m, 2H, Ar-H), 6.95 (m, 1H, Ar-H), 7.13–7.45 (m, 11H, Ar-H); ESI-MS:  $m/z$  457 ( $M + \text{H}$ ) $^+$ .

**5.1.7.9. 3-(Benzyloxy)-N-methyl-N-[6-(4-(naphthalen-1-yl)piperazin-1-yl)hexyl]aniline (17f).** The product was prepared starting from **16f** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 7:3 as eluant). Oil (532 mg, 42%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.36 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.60 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.65 (m, 2H,  $\text{CH}_2$ -piperazine), 2.71 (m, 4H, piperazine), 2.90 (s, 3H,  $\text{CH}_3$ ), 3.29 (m, 6H, piperazine and  $\text{CH}_2\text{NCH}_3$ ), 5.08 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.78 (m, 2H, Ar-H), 6.99 (m, 1H, Ar-H), 7.13 (m, 1H, Ar-H), 7.25–7.59 (m, 9H, Ar-H), 7.63 (m, 1H, Ar-H), 7.82 (m, 1H, Ar-H), 8.15 (m, 1H, Ar-H); ESI-MS:  $m/z$  508 ( $M + \text{H}$ ) $^+$ .

**5.1.7.10. 3-(Benzyloxy)-N-methyl-N-(6-{4-[3-(trifluoromethyl)phenyl]piperazin-1-yl}hexyl)aniline (17g).** The product was prepared starting from **16g** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 7:3 as eluant). Oil (670 mg, 51%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.37 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.59 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.43 (m, 2H,  $\text{CH}_2$ -piperazine), 2.69 (m, 4H, piperazine), 2.90 (s, 3H,  $\text{CH}_3$ ), 3.31 (m, 6H, piperazine and  $\text{CH}_2\text{NCH}_3$ ), 5.02 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.76 (m,

2H, Ar-H), 6.83–7.09 (m, 4H, Ar-H), 7.15–7.42 (m, 7H, Ar-H). ESI-MS:  $m/z$  526 ( $M + \text{H}$ ) $^+$ .

**5.1.7.11. 3-(Benzyloxy)-N-{6-[4-(7-chloroquinolin-4-yl)piperazin-1-yl]hexyl}-N-methylaniline (17h).** The product was prepared starting from **16h** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc as eluant). Oil (528 mg, 39%);  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  = 1.42 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.58 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.81 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 3.20 (s, 3H,  $\text{CH}_3$ ), 3.23 (m, 2H,  $\text{CH}_2\text{NCH}_3$ ), 3.49 (m, 4H, piperazine), 3.79 (m, 2H), 3.92 (m, 2H), 4.38 (m, 2H), 5.18 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 7.10 (m, 3H, Ar-H), 7.30–7.51 (m, 7H, Ar-H), 7.78 (m, 1H, Ar-H), 8.05 (m, 1H, Ar-H), 8.24 (m, 1H, Ar-H), 8.78 (m, 1H, Ar-H). ESI-MS:  $m/z$  543 ( $M + \text{H}$ ) $^+$ .

#### 5.1.8. Reduction of the alkanamido derivatives with $\text{BH}_3$ . General procedure for compounds **9g**, **14b–k**

A 1 M solution of  $\text{BH}_3$  in THF (3 equiv) was added dropwise to an ice-cooled solution of the suitable alkanamido derivative **8g**, **13b–k** (1 mmol) in dry THF (7 mL) under a  $\text{N}_2$  atmosphere. Once the addition was completed, the reaction mixture was stirred at room temperature for 16 h; 6 N HCl (0.7 mL) [a 1.25 N solution of HCl in EtOH for **14g**] was added and the resulting mixture was heated to reflux for 1 h. After cooling to room temperature the reaction mixture was made alkaline with 1 N NaOH (2.7 mL) and then extracted with EtOAc. The combined organic layers were washed with brine, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated to afford a crude residue which was purified by flash chromatography on silica gel and/or crystallization.

**5.1.8.1. N-Methyl-3-nitro-N-[2-(4-phenylpiperazin-1-yl)ethyl]aniline·HCl (9g).** The product was prepared starting from **8g** (Scheme 2). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 1:1 as eluant); the product was then dissolved in a solution of HCl in MeOH and the solvent was removed by distillation under vacuum obtaining the final product as white solid (207 mg, 55%);  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  = 3.01 (s, 3H,  $\text{CH}_3$ ), 3.20 (m, 4H, piperazine), 3.31 (m, 2H,  $\text{CH}_2$ -piperazine), 3.61 (m, 2H, piperazine), 3.82 (m, 2H, piperazine), 3.96 (m, 2H,  $\text{CH}_2\text{NCH}_3$ ), 6.86 (m, 1H, Ar-H), 7.02 (m, 2H, Ar-H), 7.25–7.34 (m, 3H, Ar-H), 7.43–7.52 (m, 3H, Ar-H), 11.50 (brs, 1H,  $\text{NH}^+$ ); ESI-MS ( $m/z$ ): 341 ( $M + \text{H}$ ) $^+$ . Anal. calcd for  $\text{C}_{19}\text{H}_{25}\text{ClN}_4\text{O}_2 \cdot 0.15 \text{H}_2\text{O}$ : C 60.12, H 6.72, N 14.76, found C 60.28, H 6.93, N 14.55.

**5.1.8.2. N,3-Dimethyl-N-[3-(4-phenylpiperazin-1-yl)propyl]aniline (14b).** The product was prepared starting from **13b** (Scheme 3). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 7:3 as eluant). Oil (145 mg, 45%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.82 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.34 (s, 3H, Ar- $\text{CH}_3$ ), 2.46 (t, 2H,  $J$  = 7.5,  $\text{CH}_2$ -piperazine), 2.63 (m, 4H, piperazine), 2.95 (s, 3H,  $\text{NCH}_3$ ), 3.25 (m, 4H, piperazine), 3.41 (t, 2H,  $J$  = 7.5,  $\text{CH}_2\text{NCH}_3$ ), 6.57 (m, 3H, Ar-H), 6.90 (m, 3H, Ar-H), 7.22 (m, 1H, Ar-H), 7.23–7.34 (m, 2H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 151.3, 149.5, 138.8, 129.1, 129.0, 119.7, 118.6, 117.0, 116.0, 113.0, 109.5, 55.8, 53.3, 50.7, 49.2, 38.2, 24.3, 22.0; ESI-MS ( $m/z$ ): 324 ( $M + \text{H}$ ) $^+$ . Anal. calcd for  $\text{C}_{21}\text{H}_{29}\text{N}_3$ : C 77.97, H 9.04, N 12.99, found C 77.90, H 9.18, N 13.07.

**5.1.8.3. 3-Bromo-N-methyl-N-[3-(4-phenylpiperazin-1-yl)propyl]aniline (14c).** The product was prepared starting from **13c** (Scheme 3). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 3:7 as eluant). Oil (314 mg, 81%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.80 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.43 (t, 2H,  $J$  = 7.0,  $\text{CH}_2$ -piperazine), 2.62 (m, 4H, piperazine), 2.94 (s, 3H,  $\text{CH}_3$ ), 3.25 (m, 4H, piperazine), 3.41 (t, 2H,  $J$  = 7.0,  $\text{CH}_2\text{NCH}_3$ ), 6.65 (dd, 1H,  $J$  = 2.0 and 8.0, Ar-H), 6.78–7.00 (m, 5H, Ar-H), 7.07 (dd, 1H,  $J_1 = J_2 = 8.0$ , Ar-H), 7.27 (d, 1H,  $J$  = 7.0, Ar-H), 7.31 (d, 1H,  $J$  = 8.0, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 151.3, 150.6,

130.3, 129.1, 123.5, 119.7, 118.6, 116.1, 114.8, 110.5, 55.4, 53.3, 50.3, 49.2, 38.1, 24.1. ESI-MS ( $m/z$ ): 388–390 ( $M + H$ )<sup>+</sup>. Anal. calcd for C<sub>20</sub>H<sub>26</sub>BrN<sub>3</sub>: C 61.86, H 6.75, N 10.82, found C 61.69, H 6.69, N 12.63.

**5.1.8.4. 3-Chloro-N-methyl-N-[3-(4-phenylpiperazin-1-yl)propyl]aniline (14d).** The product was prepared starting from **13d** (Scheme 3). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 3:7 as eluant). Oil (250 mg, 73%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.82 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.45 (t, 2H,  $J$  = 7.0, CH<sub>2</sub>-piperazine), 2.64 (m, 4H, piperazine), 2.96 (s, 3H, CH<sub>3</sub>), 3.28 (m, 4H, piperazine), 3.44 (t, 2H,  $J$  = 7.0, CH<sub>2</sub>NCH<sub>3</sub>), 6.70 (m, 3H, Ar-H), 6.94 (m, 3H, Ar-H), 7.16 (t, 1H, Ar-H), 7.28–7.37 (m, 2H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 151.4, 150.5, 135.1, 130.1, 129.2, 119.7, 116.1, 115.7, 111.9, 110.1, 55.5, 53.3, 50.3, 49.2, 38.2, 24.2. ESI-MS ( $m/z$ ): 344 ( $M + H$ )<sup>+</sup>. Anal. calcd for C<sub>20</sub>H<sub>26</sub>ClN<sub>3</sub>: C 69.85, H 7.62, N 12.22, found C 70.01, H 7.78, N 12.35.

**5.1.8.5. 3-Fluoro-N-methyl-N-[3-(4-phenylpiperazin-1-yl)propyl]aniline (14e).** The product was prepared starting from **13e** (Scheme 3). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 3:7 as eluant). White solid (245 mg, 75%); mp: 64–65 °C (Et<sub>2</sub>O/petroleum ether); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.79 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.42 (t, 2H,  $J$  = 7.0, CH<sub>2</sub>-piperazine), 2.61 (m, 4H, piperazine), 2.94 (s, 3H, CH<sub>3</sub>), 3.23 (m, 4H, piperazine), 3.40 (t, 2H,  $J$  = 7.0, CH<sub>2</sub>NCH<sub>3</sub>), 6.31–6.53 (m, 3H, Ar-H), 6.84–7.32 (m, 6H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 164.2 (d,  $J$  = 239.5), 151.3, 151.1 (d,  $J$  = 10.5), 130.1 (d,  $J$  = 10.5), 129.1, 119.7, 116.0, 107.5 (d,  $J$  = 2), 102.3 (d,  $J$  = 21.5), 99.0 (d,  $J$  = 26), 55.5, 53.3, 50.4, 49.2, 38.3, 24.2; ESI-MS ( $m/z$ ): 328 ( $M + H$ )<sup>+</sup>. Anal. calcd for C<sub>20</sub>H<sub>26</sub>FN<sub>3</sub>: C 73.36, H 8.00, N 12.83, found C 73.21, H 7.91, N 12.85.

**5.1.8.6. N-Methyl-3-nitro-N-[3-(4-phenylpiperazin-1-yl)propyl]aniline (14f).** The product was prepared starting from **13f** (Scheme 3). Purification by flash chromatography on silica gel (EtOAc as eluant). Yellow oil (241 mg, 68%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.86 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.47 (t, 2H,  $J$  = 7.5, CH<sub>2</sub>-piperazine), 2.65 (m, 4H, piperazine), 3.03 (s, 3H, CH<sub>3</sub>), 3.27 (m, 4H, piperazine), 3.51 (t, 2H,  $J$  = 7.5, CH<sub>2</sub>NCH<sub>3</sub>), 6.95 (m, 4H, Ar-H), 7.31 (m, 3H, Ar-H), 7.51 (m, 2H, Ar-H); ESI-MS ( $m/z$ ): 355 ( $M + H$ )<sup>+</sup>.

**5.1.8.7. Ethyl 3-[3-(4-phenylpiperazin-1-yl)propylamino]benzoate (14g).** The product was prepared starting from **13g** (Scheme 3). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 1:1 as eluant). Amorphous solid (253 mg, 69%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.37 (t, 3H,  $J$  = 7.0, CH<sub>2</sub>CH<sub>3</sub>), 1.87 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.58 (t, 2H,  $J$  = 7.0, CH<sub>2</sub>-piperazine), 2.66 (m, 4H, piperazine), 3.26 (m, 6H, piperazine and CH<sub>2</sub>NH), 4.35 (q, 2H,  $J$  = 7.0, CH<sub>2</sub>CH<sub>3</sub>), 4.89 (brs, 1H, NH), 6.75–6.99 (m, 4H, Ar-H), 7.17–7.43 (m, 5H, Ar-H); ESI-MS ( $m/z$ ): 368 ( $M + H$ )<sup>+</sup>.

**5.1.8.8. 3-Methoxy-N-[3-(4-phenylpiperazin-1-yl)propyl]aniline (14h).** The product was prepared starting from **13h** (Scheme 3). Purification by flash chromatography on silica gel (EtOAc as eluant). Amorphous solid (292 mg, 90%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.86 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.57 (t, 2H,  $J$  = 7.0, CH<sub>2</sub>-piperazine), 2.66 (m, 4H, piperazine), 3.24 (m, 6H, piperazine and CH<sub>2</sub>NH), 3.79 (s, 3H, CH<sub>3</sub>), 4.78 (brs, 1H, NH), 6.16–6.30 (m, 3H, Ar-H), 6.92 (m, 3H, Ar-H), 7.09 (dd, 1H,  $J_1 = J_2 = 8.0$ , Ar-H), 7.30 (m, 2H, Ar-H); ESI-MS ( $m/z$ ): 326 ( $M + H$ )<sup>+</sup>.

**5.1.8.9. 3-(Benzyloxy)-N-methyl-N-[3-(4-phenylpiperazin-1-yl)propyl]aniline (14i).** The product was prepared starting from **13i** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc/cyclohexane, 7:3 as eluant). Oil (286 mg, 69%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.81 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.43 (apt, 2H,  $J$  = 7.0, CH<sub>2</sub>-

piperazine), 2.62 (m, 4H, piperazine), 2.93 (s, 3H, CH<sub>3</sub>), 3.23 (m, 4H, piperazine), 3.39 (apt, 2H,  $J$  = 7.0, CH<sub>2</sub>NCH<sub>3</sub>), 5.06 (s, 2H, CH<sub>2</sub>Ph), 6.37 (m, 3H, Ar-H), 6.90 (m, 3H, Ar-H), 7.11–7.49 (m, 8H, Ar-H); MS (EI, 70 eV):  $m/z$  415 ( $M^+$ ), 91 (100).

**5.1.8.10. 3-Methoxy-N-methyl-N-[3-[4-(naphthalen-1-yl)piperazin-1-yl]propyl]aniline (14j).** The product was prepared starting from **13j** (Scheme 3). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 1:1 as eluant). Oil (323 mg, 83%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.86 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.52 (apt, 2H,  $J$  = 7.0, CH<sub>2</sub>-piperazine), 2.77 (m, 4H, piperazine), 2.97 (s, 3H, NCH<sub>3</sub>), 3.19 (m, 4H, piperazine), 3.44 (apt, 2H,  $J$  = 7.0, CH<sub>2</sub>NCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 6.28 (m, 2H, Ar-H), 6.41 (m, 1H, Ar-H), 7.15 (m, 2H, Ar-H), 7.38–7.59 (m, 4H, Ar-H), 7.84 (m, 1H, Ar-H), 8.22 (m, 1H, Ar-H); ESI-MS ( $m/z$ ): 390 ( $M + H$ )<sup>+</sup>.

**5.1.8.11. 3-Methoxy-N-methyl-N-[3-[4-(naphthalen-2-yl)piperazin-1-yl]propyl]aniline (14k).** The product was prepared starting from **13k** (Scheme 3). Purification by flash chromatography on silica gel (EtOAc/cyclohexane 1:1 as eluant). Oil (222 mg, 57%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.84 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.46 (apt, 2H,  $J$  = 7.0, CH<sub>2</sub>-piperazine), 2.67 (m, 4H, piperazine), 2.96 (s, 3H, NCH<sub>3</sub>), 3.34 (m, 4H, piperazine), 3.42 (apt, 2H,  $J$  = 7.0, CH<sub>2</sub>NCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 6.29 (m, 2H, Ar-H), 6.40 (m, 1H, Ar-H), 7.16 (m, 2H, Ar-H), 7.25–7.45 (m, 3H, Ar-H), 7.73 (m, 3H, Ar-H); ESI-MS ( $m/z$ ): 390 ( $M + H$ )<sup>+</sup>.

#### 5.1.9. O-Demethylation of methoxyanilino-derivatives with BBr<sub>3</sub>. General procedure for compounds **10h–i**, **14l–n**

A solution of BBr<sub>3</sub> (1 M in CH<sub>2</sub>Cl<sub>2</sub>, 3 mL, 3 mmol) diluted with dry CH<sub>2</sub>Cl<sub>2</sub> (6 mL) was added dropwise to a solution of the suitable methoxy derivative **9d–e**, **14h**, **14j–k** (1 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (4 mL) at 0 °C, and the resulting mixture was stirred at room temperature for 2 h. The reaction mixture was neutralized with a 2 N aqueous solution of Na<sub>2</sub>CO<sub>3</sub> and extracted with EtOAc. The organic phases were combined, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure, to give a crude product that was purified by flash chromatography on silica gel and/or crystallization.

**5.1.9.1. 3-[Methyl[2-(4-phenylpiperazin-1-yl)ethyl]amino]phenol (10h).** The product was prepared starting from **9d**. Purification by flash chromatography on silica gel (cyclohexane/EtOAc 1:1) and subsequent crystallization. White solid (106 mg, 34%); mp: 155–156 °C (Et<sub>2</sub>O); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.63 (t, 2H,  $J$  = 8.0, CH<sub>2</sub>CH<sub>2</sub>NCH<sub>3</sub>), 2.66 (m, 4H, piperazine), 2.95 (s, 3H, NCH<sub>3</sub>), 3.24 (m, 4H, piperazine), 3.53 (t, 2H,  $J$  = 8.0, CH<sub>2</sub>NCH<sub>3</sub>), 5.51 (brs, 1H, OH), 6.24 (m, 3H, Ar-H), 6.90 (m, 3H, Ar-H), 7.08 (t, 1H, Ar-H), 7.28 (m, 2H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 157.0, 151.1, 150.6, 130.2, 129.1, 119.9, 116.1, 104.7, 103.4, 99.2, 54.7, 53.6, 50.2, 49.0, 38.7; MS (EI, 70 eV):  $m/z$  311 ( $M^+$ ), 136 (100); Anal. calcd for C<sub>19</sub>H<sub>25</sub>N<sub>3</sub>O: C 73.28, H 8.09, N 13.49, found C 73.52, H 8.33, N 13.20.

**5.1.9.2. 3-[2-(4-Benzylpiperazin-1-yl)ethyl]methylamino]phenol (10i).** The product was prepared starting from **9e**. Purification by flash chromatography on silica gel (cyclohexane/EtOAc 2:8) and subsequent crystallization. White solid (159 mg, 49%); mp: 99–100 °C (Et<sub>2</sub>O/petroleum ether); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.56 (m, 10H, piperazine and CH<sub>2</sub>CH<sub>2</sub>NCH<sub>3</sub>), 2.92 (s, 3H, NCH<sub>3</sub>), 3.47 (apt, 2H,  $J$  = 7.5, CH<sub>2</sub>NCH<sub>3</sub>), 3.54 (s, 2H, CH<sub>2</sub>-Ar), 6.15 (m, 2H, Ar-H), 6.27 (m, 1H, Ar-H), 7.06 (m, 1H, Ar-H), 7.22–7.34 (m, 5H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 157.3, 150.5, 137.5, 130.1, 129.4, 128.3, 127.2, 104.3, 103.7, 99.5, 63.0, 54.5, 53.3, 52.7, 50.0, 38.6; ESI-MS ( $m/z$ ): 326 ( $M + H$ )<sup>+</sup>; Anal. calcd for C<sub>20</sub>H<sub>27</sub>N<sub>3</sub>O: C 73.81, H 8.36, N 12.91, found C 73.98, H 8.44, N 12.95.

**5.1.9.3. 3-[3-(4-Phenylpiperazin-1-yl)propylamino]phenol (14i).** The product was prepared starting from **14h**. Purification by flash chromatography on silica gel (cyclohexane/EtOAc 3:7) and subsequent crystallization. White solid (208 mg, 67%); mp: 140–141 °C (EtOAc);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.87 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.56 (t, 2H,  $J$  = 7.0,  $\text{CH}_2$ -piperazine), 2.66 (m, 4H, piperazine), 3.17 (t, 2H,  $J$  = 6.5,  $\text{CH}_2\text{NH}$ ), 3.23 (m, 4H, piperazine), 6.05 (m, 1H, Ar-H), 6.12–6.21 (m, 2H, Ar-H), 6.85–7.05 (m, 4H, Ar-H), 7.25–7.33 (m, 2H, Ar-H); IR (nujol):  $\nu$  = 3390, 3340  $\text{cm}^{-1}$ ;  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 157.1, 151.1, 150.2, 130.2, 129.1, 119.9, 116.1, 105.7, 104.5, 99.7, 57.1, 53.2, 49.1, 43.4, 25.7. ESI-MS ( $m/z$ ): 312 ( $\text{M} + \text{H}$ ) $^+$ ; Anal. calcd for  $\text{C}_{19}\text{H}_{25}\text{N}_3\text{O}$ : C 73.28, H 8.09, N 13.49, found C 73.09, H 7.78, N 13.55.

**5.1.9.4. 3-(Methyl[3-[4-(naphthalen-1-yl)piperazin-1-yl]propyl]amino)phenol (14m).** The product was prepared starting from **14j**. Purification by flash chromatography on silica gel (EtOAc as eluant) and subsequent crystallization. Beige solid (255 mg, 68%); mp: 124–125 °C ( $\text{Et}_2\text{O}$ /petroleum ether);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.85 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.56 (apt, 2H,  $J$  = 7.0,  $\text{CH}_2$ -piperazine), 2.83 (m, 4H, piperazine), 2.89 (s, 3H,  $\text{CH}_3$ ), 3.21 (m, 4H, piperazine), 3.38 (apt, 2H,  $J$  = 7.0,  $\text{CH}_2\text{NCH}_3$ ), 6.18 (m, 2H, Ar-H), 6.31 (m, 1H, Ar-H), 7.09 (m, 2H, Ar-H), 7.37–7.59 (m, 4H, Ar-H), 7.85 (m, 1H, Ar-H), 8.21 (m, 1H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 157.2, 150.9, 149.4, 134.7, 130.1, 128.8, 128.4, 125.9, 125.8, 125.3, 123.6, 123.5, 114.8, 104.7, 103.4, 99.5, 56.0, 53.8, 52.6, 50.6, 38.3, 24.2. ESI-MS ( $m/z$ ): 376 ( $\text{M} + \text{H}$ ) $^+$ ; Anal. calcd for  $\text{C}_{24}\text{H}_{29}\text{N}_3\text{O}$ : C 76.76, H 7.78, N 11.19, found C 76.99, H 7.83, N 11.32.

**5.1.9.5. 3-(Methyl[3-[4-(naphthalen-2-yl)piperazin-1-yl]propyl]amino)phenol (14n).** The product was prepared starting from **14k**. Purification by flash chromatography on silica gel (gradient cyclohexane/EtOAc 1:1 to EtOAc as eluant) and subsequent crystallization. White solid (198 mg, 53%); mp: 161–162 °C (acetone);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.84 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.47 (apt, 2H,  $J$  = 7.5,  $\text{CH}_2$ -piperazine), 2.69 (m, 4H, piperazine), 2.92 (s, 3H,  $\text{CH}_3$ ), 3.37 (m, 6H, piperazine and  $\text{CH}_2\text{NCH}_3$ ), 6.18 (m, 2H, Ar-H), 6.33 (m, 1H, Ar-H), 7.04–7.14 (m, 2H, Ar-H), 7.25–7.45 (m, 3H, Ar-H), 7.72 (m, 3H, Ar-H); MS (EI, 70 eV):  $m/z$  325 ( $\text{M}^+$ ), 91 (100);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 156.8, 150.9, 149.0, 134.5, 130.0, 128.7, 128.5, 126.7, 126.2, 125.2, 123.4, 119.4, 110.3, 104.9, 103.1, 99.2, 55.8, 53.2, 50.5, 49.4, 38.3, 24.2. ESI-MS ( $m/z$ ): 376 ( $\text{M} + \text{H}$ ) $^+$ ; Anal. calcd for  $\text{C}_{24}\text{H}_{29}\text{N}_3\text{O}$ : C 76.76, H 7.78, N 11.19, found C 76.59, H 7.71, N 11.27.

**5.1.10.  $N^1$ -Methyl- $N^1$ -[2-(4-phenylpiperazin-1-yl)ethyl]benzene-1,3-diamine-HCl (10k)**

A solution of the nitro compound **9g** (1.0 mmol) in EtOH (10 mL) and EtOAc (10 mL) was hydrogenated over 10% Pd/C (0.075 g) at 1 atm of  $\text{H}_2$  for 5 h at room temperature. The catalyst was filtered on Celite, the filtrate was concentrated under reduced pressure to give a crude oily amine that was dissolved in a solution of HCl in EtOH and stirred for 1 h. After removing the solvent by distillation under vacuum, the resulting solid was purified by crystallization in EtOH/ $\text{Et}_2\text{O}$ . White solid (280 mg, 81%).  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  = 2.98 (s, 3H,  $\text{CH}_3$ ), 3.20 (m, 4H, piperazine), 3.30 (apt, 2H,  $J$  = 7.0,  $\text{CH}_2$ -piperazine), 3.63 (m, 2H, piperazine), 3.82–3.95 (m, 4H, piperazine and  $\text{CH}_2\text{NCH}_3$ ), 6.66 (m, 1H, Ar-H), 6.82 (m, 1H, Ar-H), 6.87 (m, 1H, Ar-H), 6.93 (m, 1H, Ar-H), 7.10 (m, 2H, Ar-H), 7.27 (m, 3H, Ar-H), 10.25 (brs, 2H,  $\text{NH}_2$ ), 11.50 (brs, 1H,  $\text{NH}^+$ ); ESI-MS ( $m/z$ ): 311 ( $\text{M} + \text{H}$ ) $^+$ . Anal. calcd for  $\text{C}_{19}\text{H}_{27}\text{ClN}_4 \cdot 0.3 \text{ EtOH}$ : C 65.26, H 8.05, N 15.53, found C 65.12, H 8.24, N 15.31.

**5.1.11.  $N$ -(3-{Methyl[2-(4-phenylpiperazin-1-yl)ethyl]amino}phenyl)methanesulfonamide-HCl (10l)**

Methanesulfonyl chloride (0.05 mL, 0.65 mmol) was added to a solution of **10k** (0.25 g, 0.63 mmol) and TEA (0.18 mL) in dry THF

(5 mL), and the resulting mixture was stirred at room temperature for 1 h. After removing the solvent by distillation at reduced pressure, the residue was taken up in EtOAc, and washed with 1 N NaOH and with brine. After drying over  $\text{Na}_2\text{SO}_4$ , the organic layer was concentrated under reduced pressure and the resulting crude product was purified by column flash chromatography on silica gel (EtOAc/MeOH, 95:5 as eluant) to give a crude oily amine that was dissolved in a solution of HCl in EtOH and stirred for 1 h. After removing the solvent by distillation under vacuum, the resulting solid was purified by crystallization in EtOH/ $\text{Et}_2\text{O}$ . White solid (216 mg, 81%).  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  = 2.91 (s, 3H,  $\text{CH}_2\text{NCH}_3$ ), 2.97 (s, 3H,  $\text{SO}_2\text{CH}_3$ ), 3.10 (m, 4H, piperazine), 3.25 (apt, 2H,  $J$  = 7.0,  $\text{CH}_2$ -piperazine), 3.58 (m, 2H, piperazine), 3.80 (m, 2H, piperazine), 3.84 (apt, 2H,  $J$  = 7.0,  $\text{CH}_2\text{NCH}_3$ ), 6.57–6.69 (m, 3H, Ar-H), 6.87 (t, 1H, Ar-H), 7.01 (m, 2H, Ar-H), 7.14 (t, 1H, Ar-H), 7.24–7.29 (m, 2H, Ar-H), 9.51 (s, 1H, NH), 11.60 (brs, 1H,  $\text{NH}^+$ ). ESI-MS ( $m/z$ ): 389 ( $\text{M} + \text{H}$ ) $^+$ . Anal. calcd for  $\text{C}_{20}\text{H}_{29}\text{ClN}_4\text{O}_2\text{S} \cdot 0.5 \text{ H}_2\text{O}$ : C 55.35, H 6.97, N 12.91, found C 55.49, H 7.13, N 12.79.

**5.1.12.  $O$ -Debenzylation of benzyloxyanilino-derivatives by hydrogenolysis. General procedure for compounds 10j, 14o, 18a–c**

A solution of the suitable benzyloxy-anilino derivative **9f**, **14i**, **17a–c** (1 mmol) in dry MeOH (18 mL) was hydrogenated over 10% Pd/C (0.075 g) at 4 atm of  $\text{H}_2$  for 3 h at 60 °C. The catalyst was filtered on Celite, the filtrate was concentrated under reduced pressure to give a crude residue that was purified by flash chromatography on silica gel and/or crystallization.

**5.1.12.1. 3-[2-(4-Phenylpiperazin-1-yl)ethylamino]phenol (10j).** The product was prepared starting from **9f**. Purification by flash chromatography on silica gel (EtOAc/cyclohexane, 6:4 as eluant) and subsequent crystallization. Beige solid (119 mg, 40%); mp: 159–160 °C (EtOAc);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 2.69 (m, 6H, piperazine and  $\text{CH}_2\text{CH}_2\text{NH}$ ), 3.21 (m, 6H, piperazine and  $\text{CH}_2\text{NH}$ ), 4.35 (brs, 1H, NH), 6.10 (dd, 1H,  $J_1 = J_2 = 2.0$ , Ar-H), 6.21 (m, 2H, Ar-H), 6.84–6.98 (m, 3H, Ar-H), 7.03 (dd, 1H,  $J_1 = J_2 = 8.0$ , Ar-H), 7.29 (m, 2H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 156.9, 151.2, 149.9, 147.9, 130.2, 129.1, 119.9, 116.2, 106.0, 99.8, 56.6, 52.9, 49.1, 40.2; IR (nujol):  $\nu$  = 3423  $\text{cm}^{-1}$ ; MS (EI, 70 eV):  $m/z$  297 ( $\text{M}^+$ ), 122 (100). Anal. calcd for  $\text{C}_{18}\text{H}_{23}\text{N}_3\text{O}$ : C 72.70, H 7.80, N 14.13, found C 72.39, H 8.03, N 14.00.

**5.1.12.2. 3-{Methyl[3-(1-phenylpiperazin-4-yl)propyl]amino}phenol (14o).** The product was prepared starting from **14i** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc as eluant) and subsequent crystallization. Gray solid (172 mg, 53%); mp: 177–178 °C (EtOAc);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.82 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.44 (apt, 2H,  $J$  = 7.0,  $\text{CH}_2$ -piperazine), 2.63 (m, 4H, piperazine), 2.90 (s, 3H,  $\text{CH}_3$ ), 3.24 (m, 4H, piperazine), 3.37 (t, 2H,  $J$  = 7.0,  $\text{CH}_2\text{NCH}_3$ ), 6.16 (m, 2H, Ar-H), 6.31 (m, 1H, Ar-H), 6.83–7.32 (m, 6H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 156.7, 151.3, 150.9, 130.0, 129.1, 119.7, 116.0, 105.0, 103.0, 99.1, 55.7, 53.3, 50.5, 49.1, 38.3, 24.2; ESI-MS ( $m/z$ ): 326 ( $\text{M} + \text{H}$ ) $^+$ ; Anal. calcd for  $\text{C}_{20}\text{H}_{27}\text{N}_3\text{O}$ : C 73.81, H 8.36, N 12.91, found C 74.10, H 8.65, N 12.89.

**5.1.12.3. 3-{Methyl[4-(1-phenylpiperazin-4-yl)butyl]amino}phenol (18a).** The product was prepared starting from **17a** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc, as eluant) and subsequent crystallization. White solid (217 mg, 64%); mp: 169–170 °C (EtOAc);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.59 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.45 (t, 2H,  $J$  = 7.0,  $\text{CH}_2$ -piperazine), 2.65 (m, 4H, piperazine), 2.90 (s, 3H,  $\text{CH}_3$ ), 3.26 (m, 6H, piperazine and  $\text{CH}_2\text{NCH}_3$ ), 6.13 (m, 2H, Ar-H), 6.32 (m, 1H, Ar-H), 6.83–7.32 (m, 6H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 156.9, 151.2, 150.8, 130.0, 129.1, 119.8, 116.1, 104.8, 103.1, 99.3, 58.4, 53.2, 52.6, 49.0, 38.4, 24.7, 24.2; ESI-



MS (*m/z*): 340 (*M* + *H*)<sup>+</sup>; Anal. calcd for C<sub>21</sub>H<sub>29</sub>N<sub>3</sub>O: C 74.30, H 8.61, N 12.38, found C 74.15, H 8.44, N 12.27.

**5.1.12.4. 3-{Methyl[5-(1-phenylpiperazin-4-yl)pentyl]amino}phenol (18b).** The product was prepared starting from **17b** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc, as eluant) and subsequent crystallization. White solid (141 mg, 40%); mp: 144–145 °C (EtOAc); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.33 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.61 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.45 (apt, 2H, *J* = 8.0, CH<sub>2</sub>-piperazine), 2.69 (m, 4H, piperazine), 2.88 (s, 3H, CH<sub>3</sub>), 3.25 (m, 6H, CH<sub>2</sub>NCH<sub>3</sub> and piperazine), 6.14 (m, 2H, Ar-H), 6.25 (m, 1H, Ar-H), 6.84–7.10 (m, 4H, Ar-H), 7.24–7.32 (m, 2H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 157.2, 151.1, 150.8, 130.0, 129.1, 119.9, 116.2, 104.5, 103.3, 99.4, 58.7, 53.3, 52.5, 48.9, 38.4, 26.4, 26.3, 25.0; ESI-MS (*m/z*): 354 (*M* + *H*)<sup>+</sup>; Anal. calcd for C<sub>22</sub>H<sub>31</sub>N<sub>3</sub>O: C 74.75, H 8.84, N 11.89, found C 74.55, H 8.71, N 11.96.

**5.1.12.5. 3-{Methyl[6-(4-phenylpiperazin-1-yl)hexyl]amino}phenol (18c).** The product was prepared starting from **17c** (Scheme 4). Purification by flash chromatography on silica gel (EtOAc as eluant) and subsequent crystallization. White solid (183 mg, 50%); mp: 125–126 °C (EtOAc/petroleum ether); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.35 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.57 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.41 (apt, 2H, *J* = 7.5, CH<sub>2</sub>-piperazine), 2.62 (m, 4H, piperazine), 2.89 (s, 3H, CH<sub>3</sub>), 3.25 (m, 4H, piperazine), 3.28 (apt, 2H *J* = 7.0, CH<sub>2</sub>NCH<sub>3</sub>), 6.13 (m, 2H, Ar-H), 6.26 (m, 1H, Ar-H), 6.82–7.10 (m, 4H, Ar-H), 7.22–7.30 (m, 2H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 157.1, 151.2, 150.8, 130.0, 129.1, 119.8, 116.1, 104.6, 103.2, 99.4, 58.7, 53.2, 52.6, 48.8, 38.4, 27.3, 26.8, 26.4, 26.3; ESI-MS (*m/z*): 368 (*M* + *H*)<sup>+</sup>; Anal. calcd for C<sub>23</sub>H<sub>33</sub>N<sub>3</sub>O: C 75.16, H 9.05, N 11.43, found C 74.91, H 8.86, N 11.56.

#### 5.1.13. O-Debenzylation of benzyloxyanilino-derivatives with BCl<sub>3</sub>.

##### General procedure for compounds **16i**, **18e–h**, **18j–k**

A solution of BCl<sub>3</sub> (1 M in CH<sub>2</sub>Cl<sub>2</sub>, 3 mL, 3 mmol) was added dropwise to a solution of the suitable benzyloxy-anilino derivative **16f**, **17e–h**, or **17j–k** (1 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (6 mL) at 0 °C, and the resulting mixture was stirred at room temperature for 1 h. The reaction mixture was neutralized with 2 N NaOH, the organic layer was washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure, to give a crude product that was purified by flash chromatography on silica gel (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 8: 2 as eluant). The product was dissolved in a solution of HCl in MeOH and the solvent was removed by distillation under vacuum to give the desired final product as hydrochloride.

**5.1.13.1. N-(3-Hydroxyphenyl)-N-methyl-6-[4-(naphthalen-1-yl)piperazin-1-yl]hexanamide·HCl (16i).** The product was prepared starting from **16f** (Scheme 4). White solid (398 mg, 85%); <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 1.35 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.70 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.20 (apt, 2H, CH<sub>2</sub>CO), 3.20 (m, 4H, piperazine), 3.22 (s, 3H, CH<sub>3</sub>), 3.50 (m, 4H, piperazine), 3.70 (m, 2H, CH<sub>2</sub>-piperazine), 6.67 (m, 2H, Ar-H), 6.82 (m, 1H, Ar-H), 7.21–7.58 (m, 5H, Ar-H), 7.62 (m, 1H, Ar-H), 7.90 (m, 1H, Ar-H), 8.22 (m, 1H, Ar-H). ESI-MS (*m/z*): 432 (*M* + *H*)<sup>+</sup>. Anal. calcd for C<sub>27</sub>H<sub>34</sub>ClN<sub>3</sub>O<sub>2</sub>·0.15 H<sub>2</sub>O: C 69.29, H 7.32, N 8.98, found C 69.49, H 7.51, N 8.69.

**5.1.13.2. 3-{Methyl[6-(4-phenylpiperidin-1-yl)hexyl]amino}phenol·HCl (18e).** The product was prepared starting from **17e** (Scheme 4). White solid (322 mg, 80%); <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 1.58 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.70 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.90 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.10–2.30 (m, 4H, piperidine), 3.10 (m, 1H, CH), 3.26 (m, 4H, CH<sub>2</sub>N), 3.38 (s, 3H, CH<sub>3</sub>), 3.83 (m, 4H, CH<sub>2</sub>N), 7.19 (m, 3H, Ar-H), 7.44–7.58 (m, 6H, Ar-H). ESI-MS (*m/z*): 367 (*M* + *H*)<sup>+</sup>. Anal. calcd for

C<sub>24</sub>H<sub>35</sub>ClN<sub>2</sub>O·0.5 H<sub>2</sub>O: C 71.05, H 8.77, N 6.90, found C 70.88, H 8.61, N 6.69.

**5.1.13.3. 3-(Methyl[6-[4-(naphthalen-1-yl)piperazin-1-yl]hexyl]amino)phenol·HCl (18f).** The product was prepared starting from **17f** (Scheme 4). White solid (385 mg, 85%); <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 1.60 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.72 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.00 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 3.36 (m, 4H, piperazine), 3.40 (s, 3H, CH<sub>3</sub>), 3.62–3.95 (m, 8H, piperazine, CH<sub>2</sub>N and CH<sub>2</sub>NCH<sub>3</sub>), 7.18 (dd, 1H, Ar-H), 7.23 (m, 2H, Ar-H), 7.38 (m, 1H, Ar-H), 7.52–7.75 (m, 4H, Ar-H), 7.81 (m, 1H, Ar-H), 8.06 (m, 1H, Ar-H), 8.40 (m, 1H, Ar-H); ESI-MS (*m/z*): 418 (*M* + *H*)<sup>+</sup>. Anal. calcd for C<sub>27</sub>H<sub>36</sub>ClN<sub>3</sub>O·0.2 MeOH: C 70.95, H 8.06, N 9.13, found C 71.21, H 8.28, N 9.33.

**5.1.13.4. 3-[Methyl(6-[4-[3-(trifluoromethyl)phenyl]piperazin-1-yl]hexyl)amino]phenol·HCl (18g).** The product was prepared starting from **17g** (Scheme 4). White solid (368 mg, 78%); <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 1.42 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.57 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.81 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 3.19 (m, 6H, piperazine and CH<sub>2</sub>N), 3.22 (s, 3H, CH<sub>3</sub>), 3.62 (m, 4H, piperazine), 3.91 (m, 2H, CH<sub>2</sub>NCH<sub>3</sub>), 6.94 (m, 1H, Ar-H), 7.08 (m, 2H, Ar-H), 7.21 (m, 1H, Ar-H), 7.32 (m, 2H, Ar-H), 7.42 (m, 2H, Ar-H). ESI-MS (*m/z*): 436 (*M* + *H*)<sup>+</sup>. Anal. calcd for C<sub>24</sub>H<sub>33</sub>ClF<sub>3</sub>N<sub>3</sub>O·0.3 MeOH: C 60.60, H 7.16, N 8.72, found C 60.65, H 7.01, N 8.63.

**5.1.13.5. 3-([6-[4-(7-Chloroquinolin-4-yl)piperazin-1-yl]hexyl]methylanilino)phenol (18h)·HCl.** The product was prepared starting from **17h** (Scheme 4). White solid (381 mg, 78%); <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 1.41 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.58 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.82 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 3.22 (s, 3H, CH<sub>3</sub>), 3.25 (m, 2H), 3.51 (m, 4H, piperazine), 3.79 (m, 2H), 3.92 (m, 2H), 4.38 (m, 2H), 6.98 (m, 1H, Ar-H), 7.08 (m, 2H, Ar-H), 7.42 (m, 2H, Ar-H), 7.80 (m, 1H, Ar-H), 8.08 (m, 1H, Ar-H), 8.22 (d, 1H, Ar-H), 8.78 (d, 1H, Ar-H). ESI-MS (*m/z*): 453.5 (*M* + *H*)<sup>+</sup>. Anal. calcd for C<sub>26</sub>H<sub>34</sub>Cl<sub>2</sub>N<sub>4</sub>O·0.1 H<sub>2</sub>O·0.15 MeOH: C 63.31, H 7.07, N 11.29, found C 63.60, H 7.24, N 11.19.

**5.1.13.6. 3-{Ethyl[6-(4-phenylpiperazin-1-yl)hexyl]amino}phenol (18j)·HCl.** The product was prepared starting from **17j** (Scheme 4). White solid (342 mg, 82%); <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 1.19 (t, 3H, CH<sub>2</sub>CH<sub>3</sub>), 1.40 (m, 4H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.50 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.76 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 3.19 (m, 6H), 3.60 (m, 6H), 3.85 (m, 2H), 6.90–7.10 (m, 6H, Ar-H), 7.30 (m, 2H, Ar-H), 7.43 (m, 1H, Ar-H); ESI-MS (*m/z*): 382 (*M* + *H*)<sup>+</sup>. Anal. calcd for C<sub>24</sub>H<sub>36</sub>ClN<sub>3</sub>O·0.3 H<sub>2</sub>O: C 68.08, H 8.71, N 9.92, found C 67.91, H 8.52, N 9.78.

**5.1.13.7. 3-{Benzyl[6-(4-phenylpiperazin-1-yl)hexyl]amino}phenol (18k)·HCl.** The product was prepared starting from **17k** (Scheme 4). White solid (379 mg, 79%); <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 1.42 (m, 6H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.75 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 3.20 (m, 6H, piperazine and CH<sub>2</sub>N), 3.78 (m, 6H, piperazine and CH<sub>2</sub>N), 4.77 (s, 2H, CH<sub>2</sub>Ph), 6.81–7.08 (m, 6H, Ar-H), 7.10–7.23 (m, 8H, Ar-H); ESI-MS (*m/z*): 444 (*M* + *H*)<sup>+</sup>. Anal. calcd for C<sub>29</sub>H<sub>38</sub>ClN<sub>3</sub>O: C 72.55, H 7.98, N 8.75, found C 72.81, H 8.15, N 8.90.

#### 5.1.14. 3-(Benzyloxy)-N-ethyl-N-[6-(4-phenylpiperazin-1-yl)hexyl]aniline (17j)

A solution of acetaldehyde (132 mg, 3.0 mmol) in MeOH (16 mL) was added dropwise to a stirred cooled (0 °C) solution of **17d** (1.33 g, 3.0 mmol), AcOH (0.47 mL) and sodium cyanoborohydride (0.35 g, 5.55 mmol) and the resulting mixture was stirred at room temperature for 16 h. MeOH was removed by distillation *in vacuo*, then a 1N NaOH aqueous solution (5 mL) was added, and the

aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$ ; after drying over  $\text{Na}_2\text{SO}_4$ , the combined organic layers were concentrated under reduced pressure to give a crude residue which was purified by flash chromatography on silica gel (EtOAc/cyclohexane 8:2 as eluant). Oil (848 mg, 60%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.15 (t, 3H,  $J$  = 7.0,  $\text{CH}_2\text{CH}_3$ ), 1.38 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.56 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.83 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.62–3.24 (m, 8H, piperazine and  $2 \times \text{CH}_2\text{N}$ ), 3.30 (q, 2H,  $J$  = 7.0,  $\text{CH}_2\text{CH}_3$ ), 3.56 (m, 4H, piperazine), 5.02 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.22 (m, 3H, Ar-H), 6.82–7.52 (m, 11H, Ar-H). ESI-MS:  $m/z$  472 ( $\text{M} + \text{H}$ ) $^+$ .

#### 5.1.15. *N*-Benzyl-3-(benzyloxy)-*N*-[6-(4-phenylpiperazin-1-yl)hexyl]aniline (**17k**)

The product was prepared according to the above described procedure for **17j**, using benzaldehyde instead of acetaldehyde. Purification by flash chromatography on silica gel (EtOAc/cyclohexane 7:3 as eluant). Oil (1.04 g, 65%)  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.31 (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.58 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 1.92 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.91 (m, 4H, piperazine), 3.38 (m, 2H,  $\text{CH}_2$ -piperazine), 3.60 (m, 6H, piperazine and  $\text{CH}_2\text{NAr}$ ), 4.78 (d, 2H,  $J$  = 9.0,  $\text{NCH}_2\text{Ph}$ ), 4.98 (s, 2H,  $\text{OCH}_2\text{Ph}$ ), 6.30 (m, 2H, Ar-H), 6.90–7.41 (m, 17H, Ar-H). ESI-MS:  $m/z$  534 ( $\text{M} + \text{H}$ ) $^+$ .

#### 5.1.16. *N*<sup>1</sup>-Methyl-*N*<sup>1</sup>-[3-(4-phenylpiperazin-1-yl)propyl]benzene-1,3-diamine

A solution of the nitro compound **14f** (0.38 g, 1.07 mmol) in EtOH (10 mL) and EtOAc (10 mL) was hydrogenated over 10% Pd/C (0.075 g) at 1 atm of  $\text{H}_2$  for 5 h at room temperature. The catalyst was filtered on celite, the filtrate was concentrated under reduced pressure to give a crude oily amine that was used for the next step, without further purification (342 mg, near quantitative yield). ESI-MS ( $m/z$ ): 325 ( $\text{M} + \text{H}$ ) $^+$ .

#### 5.1.17. *N*-[3-(Methyl[3-(4-phenylpiperazin-1-yl)propyl]amino)phenyl]acetamide (**14p**)

$\text{Ac}_2\text{O}$  (0.045 mL, 0.48 mmol) was added to a solution of *N*<sup>1</sup>-methyl-*N*<sup>1</sup>-[3-(4-phenylpiperazin-1-yl)propyl]benzene-1,3-diamine (0.15 g, 0.46 mmol) and TEA (0.07 mL) in dry THF (4 mL), and the resulting mixture was stirred at room temperature for 1 h. After removing the solvent by distillation at reduced pressure, the residue was taken up in EtOAc, and washed with 1 N NaOH and with brine. After drying over  $\text{Na}_2\text{SO}_4$ , the organic layer was concentrated under reduced pressure and the resulting crude product was purified by column flash chromatography on silica gel (EtOAc/MeOH, 95:5 as eluant). Amorphous solid (145 mg, 86%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.80 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.15 (s, 3H,  $\text{COCH}_3$ ), 2.42 (apt, 2H,  $J$  = 7.0,  $\text{CH}_2$ -piperazine), 2.59 (m, 4H, piperazine), 2.93 (s, 3H,  $\text{NCH}_3$ ), 3.21 (m, 4H, piperazine), 3.39 (t, 2H,  $J$  = 7.0,  $\text{CH}_2\text{NCH}_3$ ), 6.49 (dd, 1H,  $J$  = 2.0 and 8.0, Ar-H), 6.66 (dd, 1H,  $J$  = 1.5 and 8.0, Ar-H), 6.81–6.97 (m, 3H, Ar-H), 7.04–7.08 (m, 2H, Ar-H), 7.13 (dd, 1H,  $J_1 = J_2 = 8.0$ , Ar-H), 7.21–7.32 (m, 1H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 168.2, 151.3, 150.0, 138.9, 129.5, 129.1, 119.6, 116.0, 108.3, 107.5, 103.7, 55.7, 53.3, 50.6, 49.1, 38.3, 24.8, 24.2; IR (nujol):  $\nu$  = 3311, 1657  $\text{cm}^{-1}$ ; ESI-MS ( $m/z$ ): 367 ( $\text{M} + \text{H}$ ) $^+$ ; Anal. calcd for  $\text{C}_{22}\text{H}_{30}\text{N}_4\text{O}$ : C 72.10, H 8.25, N 15.29, found C 72.28, H 8.19, N 15.41.

#### 5.1.18. *N*-[3-(Methyl[3-(4-phenylpiperazin-1-yl)propyl]amino)phenyl]methanesulfonamide (**14q**)

The product was prepared according the above described procedure for **14p**, using methanesulfonyl chloride (0.05 mL, 0.65 mmol) instead of  $\text{Ac}_2\text{O}$ . Purification by column flash chromatography on silica gel (EtOAc as eluant). Oil (157 mg, 85%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.79 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.42 (apt, 2H,  $J$  = 7.0,  $\text{CH}_2$ -piperazine), 2.60 (m, 4H, piperazine), 2.94 (s, 3H,  $\text{NCH}_3$ ), 2.99 (s, 3H,  $\text{NH}_2\text{SO}_2\text{CH}_3$ ), 3.21 (m, 4H, piperazine), 3.39 (apt, 2H,  $J$  = 7.0,

$\text{CH}_2\text{NCH}_3$ ), 6.47–6.58 (m, 3H, Ar-H), 6.83–6.97 (m, 3H, Ar-H), 7.13–7.32 (m, 3H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 151.2, 150.3, 138.0, 130.2, 129.1, 119.8, 116.1, 109.1, 108.0, 104.4, 55.7, 53.2, 50.5, 49.1, 38.9, 38.4, 24.1; IR (nujol):  $\nu$  = 3248, 1148  $\text{cm}^{-1}$ ; ESI-MS ( $m/z$ ): 403 ( $\text{M} + \text{H}$ ) $^+$ ; Anal. calcd for  $\text{C}_{21}\text{H}_{30}\text{N}_4\text{O}_2\text{S}$ : C 62.66, H 7.51, N 13.92, found C 62.54, H 7.38, N 13.78.

#### 5.1.19. 3-[3-(4-Phenylpiperazin-1-yl)propylamino]benzamide (**14r**)

To a stirred ice-cooled solution of the ester **14g** (0.12 g, 0.33 mmol) in MeOH (3 mL) in a 5 mL microwave vial, was added magnesium nitride (0.165 g, 1.64 mmol). The vial was sealed immediately and allowed to warm to room temperature in a water bath. After approximately 1 h, the reaction was heated at 80 °C for 24 h. After cooling to room temperature the reaction mixture was diluted with dichloromethane (12 mL) and water (10 mL), the aqueous phase was neutralized with 1 N HCl and then extracted ( $2 \times 10$  mL) with dichloromethane. After drying over  $\text{Na}_2\text{SO}_4$ , the combined organic layers were concentrated under reduced pressure and the resulting crude product was purified by column flash chromatography on silica gel (EtOAc/MeOH, 9:1 as eluant) and subsequent crystallization. White solid (14 mg, 13%); mp: 147–148 °C (EtOAc);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.87 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.57 (t, 2H,  $J$  = 6.5,  $\text{CH}_2$ -piperazine), 2.65 (m, 4H, piperazine), 3.23–3.28 (m, 6H, piperazine and  $\text{CH}_2\text{NH}$ ), 4.95 (brs, 1H, NH), 5.71 (brs, 1H, NH), 6.07 (brs, 1H, NH), 6.75 (dd, 1H,  $J$  = 2.0 and 8.0, Ar-H), 6.85–7.34 (m, 8H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 170.0, 151.2, 149.0, 134.5, 129.3, 129.1, 119.8, 116.4, 116.0, 115.1, 111.1, 57.3, 53.3, 49.3, 43.4, 25.5; IR (nujol):  $\nu$  = 3321, 3151, 1661  $\text{cm}^{-1}$ ; ESI-MS ( $m/z$ ): 339 ( $\text{M} + \text{H}$ ) $^+$ ; Anal. calcd for  $\text{C}_{20}\text{H}_{26}\text{N}_4\text{O}$ : C 70.98, H 7.74, N 16.55, found C 71.11, H 7.86, N 16.37.

From chromatographic purification we also isolated the methyl ester derivative **14s** (87 mg, 75%) as a result of **14g** transesterification.

#### 5.1.20. Methyl 3-[3-(4-phenylpiperazin-1-yl)propylamino]benzoate (**14s**)

White solid; mp: 102–103 °C (EtOAc/petroleum ether);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.88 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.59 (t, 2H,  $J$  = 6.5,  $\text{CH}_2$ -piperazine), 2.67 (m, 4H, piperazine), 3.24–3.29 (m, 6H, piperazine and  $\text{CH}_2\text{NH}$ ), 3.90 (s, 3H,  $\text{CH}_3$ ), 4.93 (brs, 1H, NH), 6.79 (ddd, 1H,  $J$  = 1.0, 2.5, 8.0, Ar-H), 6.89 (m, 1H, Ar-H), 6.95 (m, 1H, Ar-H), 6.99 (m, 1H, Ar-H), 7.18–7.38 (m, 5H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 167.6, 151.2, 148.7, 131.0, 129.2, 129.1, 119.8, 118.2, 117.5, 116.1, 112.8, 57.3, 53.3, 52.0, 49.2, 43.4, 25.5; IR (nujol):  $\nu$  = 3384, 1707  $\text{cm}^{-1}$ ; ESI-MS ( $m/z$ ): 354 ( $\text{M} + \text{H}$ ) $^+$ ; Anal. calcd for  $\text{C}_{21}\text{H}_{27}\text{N}_3\text{O}_2$ : C 71.36, H 7.70, N 11.89, found C 71.02, H 7.58, N 12.05.

#### 5.1.21. 3-[3-(4-Phenylpiperazin-1-yl)propylamino]phenyl methanol (**14t**)

$\text{LiAlH}_4$  (0.085 g, 2.24 mmol) was added portionwise to an ice-cooled stirred solution of **14g** (1.13 mmol) in dry THF (10 mL) under nitrogen atmosphere, and the resulting mixture was stirred for 1 h at room temperature. Water was added dropwise to destroy the excess of hydride, the resulting mixture was filtered on Celite, and the filtrate was concentrated *in vacuo* and partitioned between EtOAc and water. The combined organic phases were washed with brine, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated to afford a crude residue which was purified by crystallization. White solid (334 mg, 91%); mp: 142–143 °C (EtOAc);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 1.87 (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.57 (t, 2H,  $J$  = 6.5,  $\text{CH}_2$ -piperazine), 2.65 (m, 4H, piperazine), 3.23–3.28 (m, 6H, piperazine and  $\text{CH}_2\text{NH}$ ), 4.62 (s, 2H,  $\text{CH}_2\text{OH}$ ), 4.74 (brs, 1H, NH), 6.55 (dd, 1H,  $J$  = 2.0 and 8.0, Ar-H), 6.61 (m, 2H, Ar-H), 6.92 (m, 3H, Ar-H), 7.17 (dd, 1H,  $J_1 = J_2 = 8.0$ , Ar-H), 7.31 (m, 2H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  = 151.2, 149.0, 142.3, 129.4, 129.1, 119.8, 116.0, 115.6, 112.1, 111.0, 65.5, 57.2, 53.3, 49.2, 43.4, 25.8;

IR (nujol):  $\nu = 3332, 3121 \text{ cm}^{-1}$ ; ESI-MS ( $m/z$ ): 326 ( $M + H$ )<sup>+</sup>; Anal. calcd for  $C_{20}H_{27}N_3O$ : C 73.81, H 8.36, N 12.91, found C 73.59, H 8.18, N 12.77.

#### 5.1.22. 3-[3-(4-Phenylpiperazin-1-yl)propylamino]benzaldehyde (**14u**)

Activated manganese dioxide (1.28 g) was added to a solution of the alcohol derivative **14t** (0.35 g, 1.08 mmol) in dry dichloromethane (15 mL), and the resulting mixture was stirred for 18 h at room temperature. The mixture was filtered, the filter cake was washed with hot acetone and the combined filtrates were concentrated under reduced pressure to yield a crude residue that was purified by flash chromatography (EtOAc as eluant) and crystallization. Yellow solid (261 mg, 75%); mp: 87–88 °C (EtOAc); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.88$  (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.58 (t, 2H,  $J = 6.5$ , CH<sub>2</sub>-piperazine), 2.66 (m, 4H, piperazine), 3.23–3.32 (m, 6H, piperazine and CH<sub>2</sub>NH), 5.13 (brs, 1H, NH), 6.84–7.35 (m, 9H, Ar-H), 9.93 (s, 1H, CHO); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 193.1, 151.2, 149.2, 137.5, 129.7, 129.2, 120.0, 119.9, 119.5, 116.1, 110.8, 57.3, 53.3, 49.3, 43.4, 25.4$ ; IR (nujol):  $\nu = 3346, 1677 \text{ cm}^{-1}$ ; ESI-MS ( $m/z$ ): 324 ( $M + H$ )<sup>+</sup>; Anal. calcd for  $C_{20}H_{25}N_3O$ : C 74.27, H 7.79, N 12.99, found C 74.45, H 7.92, N 12.63.

#### 5.1.23. N-(3-Methoxyphenyl)-N-[3-(4-phenylpiperazin-1-yl)propyl]propionamide (**19**)

TEA (0.186 mL, 1.34 mmol) and propionic anhydride (0.171 mL, 1.34 mmol) were added to a solution of **14h** (0.29 g, 0.89 mmol) in dry THF (12 mL) and the resulting mixture was stirred at room temperature for 24 h. The solvent was removed by distillation *in vacuo*, water was added and the aqueous phase was extracted (3×) with EtOAc. The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to give a crude residue that was purified by flash chromatography on silica gel (EtOA as eluant). Oil (308 mg, 91%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.06$  (t, 3H,  $J = 7.0$ , CH<sub>2</sub>CH<sub>3</sub>), 1.82 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.09 (m, 2H, CH<sub>2</sub>-piperazine), 2.41 (q, 2H,  $J = 7.0$ , CH<sub>2</sub>CH<sub>3</sub>), 2.57 (m, 4H, piperazine), 3.17 (m, 4H, piperazine), 3.77 (t, 2H,  $J = 7.5$ , CH<sub>2</sub>NAr), 3.82 (s, 3H, OCH<sub>3</sub>), 6.71–6.93 (m, 6H, Ar-H), 7.21–7.32 (m, 3H, Ar-H); ESI-MS ( $m/z$ ): 382 ( $M + H$ )<sup>+</sup>.

#### 5.1.24. 3-Methoxy-N-[3-(4-phenylpiperazin-1-yl)propyl]-N-propylaniline (**20**)

A 1 M solution of BH<sub>3</sub> in THF (2.43 mL) was added dropwise to an ice-cooled solution of **19** (0.31 g, 0.81 mmol) in dry THF (6 mL) under a N<sub>2</sub> atmosphere. Once the addition was completed, the reaction mixture was stirred at room temperature for 16 h; 6 N HCl (0.7 mL) was cautiously added and the resulting mixture was heated to reflux for 1 h. After cooling to room temperature the reaction mixture was made alkaline with 1 N NaOH (2.7 mL) and then extracted with EtOAc. The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to afford a crude residue which was purified by flash chromatography on silica gel (cyclohexane/EtOAc, 7:3 as eluant). Oil (193 mg, 65%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 0.92$  (t, 3H,  $J = 7.0$ , CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 1.63 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 1.80 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.43 (t, 2H,  $J = 7.0$ , CH<sub>2</sub>-piperazine), 2.62 (m, 4H, piperazine), 3.23 (m, 6H, piperazine and CH<sub>2</sub>N), 3.35 (t, 2H,  $J = 7.0$ , CH<sub>2</sub>NAr), 3.80 (s, 3H, OCH<sub>3</sub>), 6.22 (m, 2H, Ar-H), 6.33 (dd, 1H,  $J = 2.0$  and 8.5, Ar-H), 6.83–6.98 (m, 3H, Ar-H), 7.12 (dd, 1H,  $J_1 = J_2 = 8.5$ , Ar-H), 7.23–7.32 (m, 2H, Ar-H); ESI-MS ( $m/z$ ): 368 ( $M + H$ )<sup>+</sup>.

#### 5.1.25. N-Benzyl-3-methoxy-N-[3-(4-phenylpiperazin-1-yl)propyl]aniline (**21**)

TEA (0.5 mL, 3.6 mmol) and benzyl bromide (0.37 mL, 3.11 mmol) were added to a solution of **14h** (0.20 g, 0.61 mmol) in toluene (4 mL), and the resulting mixture was heated at 80 °C for

16 h. After cooling to room temperature, water was added and the aqueous phase was extracted (3×) with EtOAc. The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to give a crude residue that was purified by flash chromatography on silica gel (EtOAc/cyclohexane, 3:7 as eluant). Oil (207 mg, 82%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.90$  (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.45 (t, 2H,  $J = 7.0$ , CH<sub>2</sub>-piperazine), 2.61 (m, 4H, piperazine), 3.23 (m, 4H, piperazine), 3.49 (t, 2H,  $J = 7.0$ , CH<sub>2</sub>NCH<sub>2</sub>Ph), 3.75 (s, 3H, OCH<sub>3</sub>), 4.57 (s, 2H, CH<sub>2</sub>Ph), 6.27 (m, 2H, Ar-H), 6.38 (dd, 1H,  $J = 2.0$  and 8.0, Ar-H), 6.92 (m, 3H Ar-H), 7.11 (dd, 1H,  $J_1 = J_2 = 8.0$ , Ar-H), 7.22–7.35 (m, 7H, Ar-H); ESI-MS ( $m/z$ ): 416 ( $M + H$ )<sup>+</sup>.

#### 5.1.26. 3-Methoxy-N-phenyl-N-[3-(4-phenylpiperazin-1-yl)propyl]aniline (**22**)

A Schlenk flask was charged with palladium acetate (0.01 g, 0.044 mmol), (±)-BINAP (0.031 g, 0.05 mmol), potassium *tert*-butoxide (0.147 g, 1.3 mmol) and toluene (2 mL) under a nitrogen atmosphere. To this mixture a solution of **14h** (0.3 g, 0.92 mmol) and iodobenzene (0.1 mL, 0.92 mmol) in dry toluene (2 mL) was added dropwise via syringe. After the addition was completed, the mixture was heated at 110 °C for 24 h. After cooling to room temperature EtOAc was added, the mixture was filtered and the filtrate was washed with brine. After drying over Na<sub>2</sub>SO<sub>4</sub>, the organic layer was concentrated under reduced pressure and the resulting crude product was purified by column flash chromatography on silica gel (cyclohexane/EtOAc 7:3 as eluant). Oil (162 mg, 44%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.91$  (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.47 (t, 2H,  $J = 7.0$ , CH<sub>2</sub>-piperazine), 2.61 (m, 4H, piperazine), 3.23 (m, 4H, piperazine), 3.76 (s, 3H, OCH<sub>3</sub>), 3.80 (t, 2H,  $J = 7.0$ , CH<sub>2</sub>NPh), 6.54 (m, 3H, Ar-H), 6.84–7.32 (m, 11H, Ar-H); MS (EI, 70 eV):  $m/z$  401 ( $M^+$ ), 226 (100).

#### 5.1.27. N-(3-Methoxyphenyl)-N-[3-(4-phenylpiperazin-1-yl)propyl]benzenesulfonamide (**23**)

TEA (0.23 mL, 1.8 mmol) and benzenesulfonyl chloride (0.18 mL, 1.41 mmol) were added to a solution of **14h** (0.25 g, 0.77 mmol) in dry THF (7.5 mL), and the resulting mixture was stirred at room temperature under nitrogen for 48 h. After removing the solvent by distillation *in vacuo*, the residue was partitioned between water and EtOAc. The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure to afford a crude residue which was purified by flash chromatography on silica gel (EtOAc/cyclohexane, 7:3 as eluant). White solid (250 mg, 70%); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.60$  (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.48 (t, 2H,  $J = 7.0$ , CH<sub>2</sub>-piperazine), 2.56 (m, 4H, piperazine), 3.18 (m, 4H, piperazine), 3.62 (t, 2H,  $J = 7.0$ , CH<sub>2</sub>NSO<sub>2</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 6.62 (m, 2H, Ar-H), 6.88 (m, 4H, Ar-H), 7.24 (m, 3H, Ar-H), 7.54 (m, 5H, Ar-H); ESI-MS ( $m/z$ ): 466 ( $M + H$ )<sup>+</sup>.

#### 5.1.28. 3-[[3-(4-Phenylpiperazin-1-yl)propyl]propylamino]phenol (**24**)

The product was prepared starting from **20** using the above general *O*-demethylation procedure with boron tribromide. Purification by flash chromatography on silica gel (cyclohexane/EtOAc 1:1 as eluant) and subsequent crystallization. White solid (339 mg, 96%); mp: 132–133 °C (Et<sub>2</sub>O/petroleum ether); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 0.91$  (t, 3H,  $J = 7.5$ , CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 1.67 (q, 2H,  $J = 7.5$ , CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 1.84 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.45 (t, 2H,  $J = 7.0$ , CH<sub>2</sub>-piperazine), 2.66 (m, 4H, piperazine), 3.16–3.36 (m, 8H, piperazine, 2 × CH<sub>2</sub>NAr), 5.86 (brs, 1H, OH), 6.10 (m, 2H, Ar-H), 6.25 (dd, 1H,  $J = 2.0$  and 8.0, Ar-H), 6.84–7.07 (m, 4H, Ar-H), 7.26–7.33 (m, 2H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 157.1, 151.2, 149.7, 130.0, 129.1, 119.9, 116.1, 104.5, 102.6, 99.0, 55.9, 53.3, 52.8, 49.0, 48.9, 24.6, 20.3, 11.5$ ; ESI-MS ( $m/z$ ): 354 ( $M + H$ )<sup>+</sup>; Anal. calcd for  $C_{22}H_{31}N_3O$ : C 74.75, H 8.84, N 11.89, found C 74.98, H 8.59, N 11.66.

### 5.1.29. 3-{Benzyl[3-(4-phenylpiperazin-1-yl)propyl]amino}phenol (**25**)

The product was prepared starting from **21** using the above general *O*-demethylation procedure with boron tribromide. Purification by flash chromatography on silica gel (cyclohexane/EtOAc 3:7 as eluant) and subsequent crystallization. White solid (228 mg, 57%); mp: 166–167 °C (EtOAc); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.89 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.44 (t, 2H, *J* = 7.0, CH<sub>2</sub>-piperazine), 2.54 (m, 4H, piperazine), 3.16 (m, 4H, piperazine), 3.45 (t, 2H, *J* = 7.0, CH<sub>2</sub>NCH<sub>2</sub>Ph), 4.44 (s, 2H, CH<sub>2</sub>Ph), 6.08 (m, 1H, Ar-H), 6.16 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 6.31 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 6.83–6.94 (m, 3H, Ar-H), 7.06 (dd, 1H, *J*<sub>1</sub> = *J*<sub>2</sub> = 8.0, Ar-H), 7.13–7.35 (m, 7H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 157.3, 151.1, 150.3, 138.5, 130.2, 129.1, 128.6, 126.7, 126.4, 119.9, 116.1, 104.4, 103.6, 99.5, 55.6, 53.5, 53.1, 48.8, 48.5, 24.7; ESI-MS (*m/z*): 402 (M + H)<sup>+</sup>; Anal. calcd for C<sub>26</sub>H<sub>31</sub>N<sub>3</sub>O: C 77.77, H 7.78, N 10.46, found C 77.39, H 7.52, N 10.30.

### 5.1.30. 3-{Phenyl[3-(4-phenylpiperazin-1-yl)propyl]amino}phenol (**26**)

The product was prepared starting from **22** using the above general *O*-demethylation procedure with boron tribromide. Purification by flash chromatography on silica gel (cyclohexane/EtOAc 1:1 as eluant) and subsequent crystallization. White solid (248 mg, 64%); mp: 172–173 °C (Et<sub>2</sub>O); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.87 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.44 (t, 2H, *J* = 7.0, CH<sub>2</sub>-piperazine), 2.57 (m, 4H, piperazine), 3.19 (m, 4H, piperazine), 3.74 (t, 2H, *J* = 7.0, CH<sub>2</sub>NPh), 6.33 (m, 2H, Ar-H), 6.51 (m, 1H, Ar-H), 6.83–7.13 (m, 7H, Ar-H), 7.21–7.32 (m, 4H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 156.8, 151.1, 149.5, 147.6, 130.1, 129.4, 129.1, 123.4, 122.7, 119.9, 116.1, 110.8, 107.1, 105.9, 55.6, 53.1, 50.0, 48.9, 24.8; ESI-MS (*m/z*): 387 (M + H)<sup>+</sup>; Anal. calcd for C<sub>25</sub>H<sub>29</sub>N<sub>3</sub>O: C 77.48, H 7.54, N 10.84, found C 77.13, H 7.30, N 10.96.

### 5.1.31. *N*-(3-Hydroxyphenyl)-*N*-[3-(4-phenylpiperazin-1-yl)propyl]benzenesulfonamide (**27**)

The product was prepared starting from **23** using the above general *O*-demethylation procedure with boron tribromide. Purification by flash chromatography on silica gel (cyclohexane/EtOAc 3:7 as eluant). Amorphous solid (333 mg, 74%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.69 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.55 (t, 2H, *J* = 7.0, CH<sub>2</sub>-piperazine), 2.68 (m, 4H, piperazine), 3.19 (m, 4H, piperazine), 3.55 (t, 2H, *J* = 7.0, CH<sub>2</sub>NSO<sub>2</sub>), 6.42 (m, 2H, Ar-H), 6.55 (dd, 1H, *J* = 2.0 and 8.0, Ar-H), 6.83–6.93 (m, 3H, Ar-H), 7.06 (dd, 1H, *J*<sub>1</sub> = *J*<sub>2</sub> = 8.0, Ar-H), 7.22–7.30 (m, 2H, Ar-H), 7.36–7.58 (m, 5H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 156.9, 151.0, 139.3, 137.7, 132.7, 129.8, 129.1, 128.8, 127.6, 120.0, 119.1, 116.1, 115.5, 55.5, 53.1, 48.6, 48.4, 24.9; IR (nujol): ν = 1164 cm<sup>-1</sup>; ESI-MS (*m/z*): 452 (M + H)<sup>+</sup>; Anal. calcd for C<sub>25</sub>H<sub>29</sub>N<sub>3</sub>O<sub>3</sub>S: C 66.49, H 6.47, N 9.31, found C 66.20, H 6.55, N 9.66.

### 5.1.32. 3-(4,4-Diphenylpiperidin-1-yl)-*N*-(3-methoxyphenyl)-*N*-methylpropanamide (**28b**)

4,4-diphenylpiperidine [**29**] (0.6 mmol) and TEA (0.83 mL) were added to a solution of 3-bromo-*N*-(3-methoxyphenyl)-*N*-methylpropanamide (**12a**) (0.5 mmol) in DMF (0.14 mL) and the resulting mixture was stirred at room temperature for 16 h, monitoring the progress of the reaction by TLC on silica gel. The reaction mixture was poured into water, and extracted with EtOAc. After drying over Na<sub>2</sub>SO<sub>4</sub>, the combined organic phases were concentrated *in vacuo* and the resulting crude product was purified by column flash chromatography on silica gel (EtOAc as eluant). Amorphous solid (103 mg, 48%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 2.28 (t, 2H, *J* = 7.5, CH<sub>2</sub>CO), 2.41 (m, 8H, piperidine), 2.60 (t, 2H, *J* = 7.5, COCH<sub>2</sub>CH<sub>2</sub>N), 3.24 (s, 3H, NCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 6.68–6.76 (m, 2H, Ar-H), 6.87 (ddd, 1H, *J* = 1.0, 3.0 and 8.0, Ar-H), 7.09–7.34 (m, 11H, Ar-H); ESI-MS (*m/z*): 429 (M + H)<sup>+</sup>.

### 5.1.33. 3-[4-(Diphenylamino)piperidin-1-yl]-*N*-(3-methoxyphenyl)-*N*-methylpropanamide (**28c**)

This compound was prepared using the above described procedure for **28b**, using diphenylpiperidin-4-ylamine [**30**] instead of 4,4-diphenylpiperidine. Purification by flash chromatography on silica gel (EtOAc as eluant). Oil (113 mg, 51%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.39 (m, 2H, piperidine), 1.99 (m, 4H, piperidine), 2.27 (t, 2H, *J* = 7.5, CH<sub>2</sub>CO), 2.66 (t, 2H, *J* = 7.5, COCH<sub>2</sub>CH<sub>2</sub>N), 2.82 (m, 2H, piperidine), 3.23 (s, 3H, NCH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 3.81 (m, 1H, CH), 6.67–7.01 (m, 9H, Ar-H), 7.19–7.33 (m, 5H, Ar-H); ESI-MS (*m/z*): 444 (M + H)<sup>+</sup>.

### 5.1.34. *N*-[3-(4,4-Diphenylpiperidin-1-yl)propyl]-3-methoxy-*N*-methylaniline (**29b**)

A 1 M solution of BH<sub>3</sub> in THF (3 mL) was added dropwise to an ice-cooled solution of **28b** (0.43 g, 1 mmol) in dry THF (7 mL) under a N<sub>2</sub> atmosphere. Once the addition was completed, the reaction mixture was stirred at room temperature for 16 h; 6 N HCl (0.9 mL) was cautiously added and the resulting mixture was heated to reflux for 1 h. After cooling to room temperature the reaction mixture was made alkaline with 1 N NaOH (3 mL) and then extracted with EtOAc. The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to afford a crude residue which was purified by flash chromatography on silica gel (EtOAc as eluant). Oil (270 mg, 65%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.78 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.34 (t, 2H, *J* = 7.5, CH<sub>2</sub>-piperidine), 2.54 (m, 8H, piperidine), 2.79 (s, 3H, NCH<sub>3</sub>), 3.28 (t, 2H, *J* = 7.5, CH<sub>2</sub>NCH<sub>3</sub>), 3.83 (s, 3H, OCH<sub>3</sub>), 6.16 (m, 3H, Ar-H), 7.00–7.32 (m, 11H, Ar-H); ESI-MS (*m/z*): 415 (M + H)<sup>+</sup>.

### 5.1.35. 1-{3-[(3-Methoxyphenyl)methylamino]propyl}-*N,N*-diphenylpiperidin-4-amine (**29c**)

The product was prepared using the above described procedure starting from **28c**. Purification by flash chromatography on silica gel (EtOAc/cyclohexane 1:1 as eluant). Oil (202 mg, 47%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.50 (m, 2H, piperidine), 1.72 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.03 (m, 4H, piperidine), 2.34 (t, 2H, *J* = 7.0, CH<sub>2</sub>-piperidine), 2.88 (s, 3H, NCH<sub>3</sub>), 2.97 (m, 2H, piperidine), 3.30 (t, 2H, *J* = 7.0, CH<sub>2</sub>NCH<sub>3</sub>), 3.77 (s, 3H, OCH<sub>3</sub>), 3.85 (m, 1H, CH), 6.29 (m, 3H, Ar-H), 6.84 (m, 4H, Ar-H), 7.05 (m, 3H, Ar-H), 7.22–7.32 (m, 4H, Ar-H); ESI-MS (*m/z*): 430 (M + H)<sup>+</sup>.

### 5.1.36. 3-{3-[(4,4-Diphenylpiperidin-1-yl)propyl]methylamino}phenol (**30b**)

The product was prepared starting from **29b** using the above general *O*-demethylation procedure with boron tribromide. Purification by flash chromatography on silica gel (EtOAc as eluant). Amorphous solid (116 mg, 29%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.76 (m, 2H, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.31 (t, 2H, *J* = 7.5, CH<sub>2</sub>-piperidine), 2.52 (m, 8H, piperidine), 2.82 (s, 3H, NCH<sub>3</sub>), 3.29 (t, 2H, *J* = 7.0, CH<sub>2</sub>NCH<sub>3</sub>), 6.13 (m, 2H, Ar-H), 6.25 (ddd, 1H, *J* = 1.5, 3.0 and 8.0, Ar-H), 7.04 (dd, 1H, *J*<sub>1</sub> = *J*<sub>2</sub> = 8.0, Ar-H), 7.11–7.18 (m, 2H, Ar-H), 7.21–7.30 (m, 8H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 157.6, 150.8, 130.0, 128.4, 127.1, 125.8, 104.3, 103.7, 99.5, 56.1, 50.6, 50.5, 44.6, 38.1, 35.6, 29.7, 24.2; ESI-MS (*m/z*): 401 (M + H)<sup>+</sup>; Anal. calcd for C<sub>27</sub>H<sub>32</sub>N<sub>2</sub>O: C 80.96, H 8.05, N 6.99, found C 81.24, H 8.22, N 6.68.

### 5.1.37. 3-{3-[4-(Diphenylamino)piperidin-1-yl]propyl}methylamino}phenol (**30c**)

The product was prepared starting from **29c** using the above general *O*-demethylation procedure with boron tribromide. Purification by flash chromatography on silica gel (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 95:5). Amorphous solid (166 mg, 40%); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.45–1.81 (m, 4H, piperidine and CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.94–2.16 (m, 4H, piperidine), 2.36 (t, 2H, *J* = 7.0, CH<sub>2</sub>-piperidine), 2.86 (s, 3H, NCH<sub>3</sub>), 3.00 (m, 2H,



piperidine), 3.29 (t, 2H,  $J = 7.0$ ,  $\text{CH}_2\text{NCH}_3$ ), 3.87 (m, 1H,  $\text{CH}$ ), 6.12 (m, 2H, Ar-H), 6.25 (ddd, 1H,  $J = 1.5$ , 3.0 and 8.0, Ar-H), 6.80–6.86 (m, 4H, Ar-H), 6.95–7.07 (m, 3H, Ar-H), 7.23–7.31 (m, 4H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 157.1$ , 150.8, 146.0, 130.0, 129.3, 122.7, 121.8, 104.7, 103.5, 99.5, 55.7, 55.0, 53.4, 50.7, 38.2, 30.2, 24.2; ESI-MS ( $m/z$ ): 416 ( $\text{M} + \text{H}$ )<sup>+</sup>; Anal. calcd for  $\text{C}_{27}\text{H}_{33}\text{N}_3\text{O}$ : C 78.03, H 8.00, N 10.11, found C 78.36, H 8.35, N 10.01.

#### 5.1.38. 3-(3-Hydroxyphenyl)-1-(4-phenylpiperazin-1-yl)propan-1-one (**31a**)

Phenyl piperazine (500 mg, 3.08 mmol), 3-(3-hydroxyphenyl) propionic acid (570 mg, 3.4 mmol) and  $\text{Et}_3\text{N}$  (1.3 mL, 9.24 mmol) were added to an ice-cooled solution of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide [EDC] (885 mg, 4.62 mmol) and *N*-hydroxybenzotriazole [HOBT] (582 mg, 4.3 mmol) in  $\text{CH}_2\text{Cl}_2$  (16.6 mL). The resulting mixture was stirred for 2 h at 0 °C and at room temperature overnight. The organic solution was washed with water, dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated under vacuum, obtaining a crude residue that was used for the next step without further purification. Oil (827 mg, 86%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 2.41$  (t, 2H,  $J = 7.5$ ,  $\text{CH}_2\text{CO}$ ), 2.67 (t, 2H,  $J = 7.5$ ,  $\text{CH}_2\text{Ar}$ ), 3.20 (m, 4H, piperazine), 3.61 (m, 2H, piperazine), 3.85 (m, 2H, piperazine), 6.68–6.99 (m, 6H, Ar-H), 7.17–7.35 (m, 3H, Ar-H); ESI-MS ( $m/z$ ): 333 ( $\text{M} + \text{Na}$ )<sup>+</sup>.

#### 5.1.39. 4-(3-Methoxyphenyl)-1-(4-phenylpiperazin-1-yl)butan-1-one (**31b**)

This product was obtained according to the above described procedure for **31a** starting from 4-(3-methoxyphenyl)butanoic acid [31] (0.7 g, 3.6 mmol) instead of 3-(3-hydroxyphenyl)propionic acid. Purification by flash chromatography on silica gel (cyclohexane/EtOAc, 6:4 as eluant). Oil (770 mg, 74%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 2.03$  (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.37 (apt, 2H,  $J = 7.5$ ,  $\text{CH}_2\text{CO}$ ), 2.69 (apt, 2H,  $J = 7.5$ ,  $\text{CH}_2\text{Ar}$ ), 3.15 (m, 4H, piperazine), 3.57 (m, 2H, piperazine), 3.79 (m, 2H, piperazine), 3.80 (s, 3H,  $\text{OCH}_3$ ), 6.72–6.85 (m, 3H, Ar-H), 6.86–6.95 (m, 3H, Ar-H), 7.18–7.34 (m, 3H, Ar-H); ESI-MS ( $m/z$ ): 339 ( $\text{M} + \text{H}$ )<sup>+</sup>.

#### 5.1.40. 3-[3-(4-Phenylpiperazin-1-yl)propyl]phenol·HCl (**32a**)

A 1 M solution of  $\text{B}_2\text{H}_6$  in THF (20 mL, 16 mmol) was added dropwise to an ice-cooled solution of **31a** (827 mg, 2.5 mmol) in THF (18 mL) and the reaction mixture underwent stirring for 4 h at room temperature. 6 M HCl (7 mL) was added to the solution at 0 °C and the resulting mixture was evaporated under vacuum. The solid was dissolved in EtOAc and washed with an 7% aqueous solution of  $\text{NaHCO}_3$ . The organic layer was washed with water, dried ( $\text{Na}_2\text{SO}_4$ ), and evaporated under vacuum obtaining a crude product (1.02 g, 3.6 mmol). The oily product was dissolved in a solution of EtOH (10 mL) and conc. HCl (300 mL, 3.6 mmol) and the mixture was stirred for 1 h. After removing the solvent by distillation under vacuum, the resulting solid was purified by crystallization in EtOH. White solid (196 mg, 59%);  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta = 2.01$  (m, 2H,  $\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.57 (m, 2H,  $\text{CH}_2$ -piperazine), 3.01 (m, 6H, piperazine and  $\text{CH}_2\text{Ar}$ ), 3.58 (m, 2H, piperazine), 3.78 (m, 2H, piperazine), 6.62 (m, 2H, Ar-H), 6.82 (ddd, 1H,  $J = 1.5$ , 3.0 and 8.0, Ar-H), 7.00 (m, 3H, Ar-H), 7.15 (dd, 1H,  $J_1 = J_2 = 8.0$ , Ar-H), 7.30 (m, 2H, Ar-H); 9.38 (s, 1H, OH), 11.00 (brs, 1H,  $\text{NH}^+$ ); ESI-MS ( $m/z$ ): 297 ( $\text{M} + \text{H}$ )<sup>+</sup>. Anal. calcd for  $\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}$ : C 76.99, H 8.16, N 9.45, found C 76.79, H 8.01, N 9.39.

#### 5.1.41. 1-[4-(3-Methoxyphenyl)butyl]-4-phenylpiperazine (**32b**)

This compound was obtained by reduction of 4-(3-methoxyphenyl)-1-(4-phenylpiperazin-1-yl)butan-1-one (0.2 g, 0.6 mmol) with 1 M  $\text{B}_2\text{H}_6$  in THF according to the above described procedure for compound **32a**. Purification by flash chromatography

on silica gel (cyclohexane/EtOAc, 7:3 as eluant). Oil (130 mg, 67%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 1.68$  (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.42 (t, 2H,  $J = 7.0$ ,  $\text{CH}_2\text{Ar}$ ), 2.63 (m, 6H, piperazine and  $\text{CH}_2\text{N}$ ), 3.22 (m, 4H, piperazine), 3.81 (s, 3H,  $\text{OCH}_3$ ), 6.73–6.98 (m, 6H, Ar-H), 7.17–7.32 (m, 3H, Ar-H); ESI-MS ( $m/z$ ): 325 ( $\text{M} + \text{H}$ )<sup>+</sup>.

#### 5.1.42. 3-[4-(4-Phenylpiperazin-1-yl)butyl]phenol (**32c**)

The product was prepared starting from **32b** using the above described general *O*-demethylation procedure with boron tribromide. Purification by flash chromatography on silica gel (cyclohexane/EtOAc 1:1 as eluant) and subsequent crystallization. White solid (226 mg, 73%); mp: 143–144 °C ( $\text{Et}_2\text{O}$ /petroleum ether);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 1.63$  (m, 4H,  $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2$ ), 2.48 (t, 2H,  $J = 7.0$ ,  $\text{CH}_2\text{Ar}$ ), 2.58 (t, 2H,  $J = 7.0$ ,  $\text{CH}_2\text{N}$ ), 2.69 (m, 4H, piperazine), 3.26 (m, 4H, piperazine), 6.59 (m, 2H, Ar-H), 6.70 (d, 1H,  $J = 7.5$ , Ar-H), 6.82–6.98 (m, 3H, Ar-H), 7.12 (dd, 1H,  $J_1 = J_2 = 7.5$ , Ar-H), 7.24–7.32 (m, 2H, Ar-H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 156.2$ , 151.1, 144.0, 129.4, 129.1, 120.2, 119.9, 116.2, 115.5, 113.0, 58.6, 53.2, 48.8, 35.6, 29.2, 26.0; ESI-MS ( $m/z$ ): 311 ( $\text{M} + \text{H}$ )<sup>+</sup>; Anal. calcd for  $\text{C}_{20}\text{H}_{26}\text{N}_2\text{O}$ : C 77.38, H 8.44, N 9.02, found C 77.02, H 8.12, N 9.33.

#### 5.1.43. 2-[3-(Benzyloxy)phenoxy]-1-(4-phenylpiperazin-1-yl)ethanone (**33**)

2-Bromo-1-(4-phenylpiperazin-1-yl)ethanone [32] (1.5 g, 6.28 mmol) was added to a solution of 3-benzyloxyphenol (1.9 g, 9.42 mmol) and  $\text{K}_2\text{CO}_3$  (1.35 g, 9.42 mmol) in DMF (5 mL), and the resulting mixture was stirred at 40 °C overnight. After removing the solvent by distillation under vacuum, EtOAc was added and the organic solution was washed with 1 M NaOH and then with brine. After drying over  $\text{Na}_2\text{SO}_4$ , the organic layer was concentrated under reduced pressure and the resulting crude product was purified by column flash chromatography on silica gel (EtOAc/hexane, 1:1 as eluant). White solid (0.88 g, 35%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 3.10$  (m, 4H, piperazine), 3.71 (m, 4H, piperazine), 4.62 (s, 2H,  $\text{OCH}_2\text{CO}$ ), 4.98 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.52 (m, 3H, Ar-H), 6.85 (m, 3H, Ar-H), 7.05–7.40 (m, 8H, Ar-H). ESI-MS ( $m/z$ ): 425 ( $\text{M} + \text{Na}$ )<sup>+</sup>.

#### 5.1.44. 1-[2-[3-(Benzyloxy)phenoxy]ethyl]-4-phenylpiperazine·HCl (**34**)

This compound was obtained by reduction of **33** (0.63 g, 1.56 mmol) with 1 M  $\text{B}_2\text{H}_6$  in THF according to the above described procedure for compound **32a**. The crude product was dissolved in a saturated solution of HCl in MeOH and then crystallized by MeOH/ $\text{Et}_2\text{O}$ . White solid (0.58 g, 87%);  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta = 3.02$ –3.37 (m, 4H, piperazine), 3.57 (m, 4H, piperazine), 3.83 (m, 2H,  $\text{CH}_2$ -piperazine), 4.40 (m, 2H,  $\text{OCH}_2$ ), 5.10 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 6.64 (m, 3H, Ar-H), 6.85 (m, 1H, Ar-H), 7.01 (m, 2H, Ar-H), 7.01 (m, 3H, Ar-H), 7.05–7.50 (m, 8H, Ar-H), 10.81 (brs, 1H). ESI-MS ( $m/z$ ): 389 ( $\text{M} + \text{H}$ )<sup>+</sup>.

#### 5.1.45. 3-[2-(4-Phenylpiperazin-1-yl)ethoxy]phenol·HCl (**35**)

This compound was obtained starting from compound **34** and using the above described general *O*-debenzylation procedure with  $\text{BCl}_3$ . White solid (237 mg, 71%);  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta = 3.15$  (m, 4H, piperazine), 3.60 (m, 4H, piperazine), 3.81 (m, 2H,  $\text{CH}_2$ -piperazine), 4.37 (m, 2H,  $\text{OCH}_2$ ), 6.42 (m, 3H, Ar-H), 6.85 (m, 1H, Ar-H), 6.98 (m, 2H, Ar-H), 7.08 (m, 1H, Ar-H), 7.25 (m, 2H, Ar-H); ESI-MS ( $m/z$ ): 299 ( $\text{M} + \text{H}$ )<sup>+</sup>. Anal. calcd for  $\text{C}_{18}\text{H}_{23}\text{ClN}_2\text{O}_2 \cdot 0.5 \text{ MeOH}$ : C 63.33, H 7.18, N 7.98, found C 63.51, H 7.40, N 7.83.

#### 5.1.46. 1-(8-Methoxy-1,2,3,4-tetrahydronaphthalen-2-yl)-4-phenylpiperazine (**36**)

*N*-Phenylpiperazine (1.4 g, 8.7 mmol) and *p*-toluenesulfonic acid (0.053 g, 0.29 mmol) were added to a solution of 8-methoxy-3,4-dihydronaphthalen-2(1H)-one (0.5 g, 2.85 mmol) in dry toluene (25 mL). This solution was heated to reflux in a Dean Stark

apparatus for 24 h, removing the water of reaction. The solvent was removed by distillation under vacuum, the resulting crude residue was dissolved in EtOH/THF (20 mL + 20 mL) and then hydrogenated (4 atm) at room temperature in the presence of PtO<sub>2</sub> (0.05 g) for 24 h. The reaction mixture was filtered on celite, the filtrate was evaporated under reduced pressure to yield a crude residue that was purified by flash-chromatography on silica gel (cyclohexane/EtOAc 8:2 as eluant). White solid (688 mg, 75%); mp: 119–120 °C (Et<sub>2</sub>O/petroleum ether); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.65 (dddd, 1H, *J* = 5.0, 6.0, 11.0, 16.5, H<sub>3a</sub>), 2.17 (m, 1H, H<sub>2</sub>), 2.53 (dd, 1H, *J* = 11.0, 16.5, H<sub>4a</sub>), 2.73 (m, 1H, H<sub>3b</sub>), 2.90 (m, 6H, NCH<sub>2</sub>CH<sub>2</sub>N, H<sub>1a</sub> and H<sub>1b</sub>), 3.10 (ddd, 1H, *J* = 1.5, 5.0, 16.5, H<sub>4b</sub>), 3.28 (m, 4H, NCH<sub>2</sub>CH<sub>2</sub>N), 3.84 (s, 3H, CH<sub>3</sub>), 6.69 (d, 2H, *J* = 8.0, Ar-H), 6.74 (d, 1H, *J* = 8.0, Ar-H), 6.88 (m, 1H, Ar-H), 6.97 (m, 2H, Ar-H), 7.11 (dd, 1H, *J*<sub>1</sub> = *J*<sub>2</sub> = 8.0, Ar-H), 7.25–7.33 (m, 2H, Ar-H); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 157.6, 151.4, 137.7, 129.1, 126.1, 124.6, 120.7, 119.6, 116.1, 106.7, 60.4, 55.2, 49.6, 49.2, 29.7, 25.9, 25.8; MS (EI, 70 eV): *m/z* 322 (M<sup>+</sup>, 100).

#### 5.1.47. 7-(4-Phenylpiperazin-1-yl)-5,6,7,8-tetrahydronaphthalen-1-ol (**37**)

The product was prepared according to the above described general *O*-demethylation procedure with BBr<sub>3</sub> starting from **36**. Purification by flash-chromatography on silica gel (cyclohexane/EtOAc 1:1 as eluant) and crystallization. White solid (194 mg, 63%); mp: 211 °C dec. (EtOAc); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>): δ = 1.50 (dddd, 1H, *J* = 4.5, 6.0, 11.0, 16.5, H<sub>6a</sub>), 2.02 (m, 1H, H<sub>7</sub>), 2.39 (dd, 1H, *J* = 11.0, 16.0, H<sub>5a</sub>), 2.60–2.85 (m, 7H, NCH<sub>2</sub>CH<sub>2</sub>N + H<sub>8a</sub> + H<sub>8b</sub> + H<sub>6b</sub>), 2.86 (dd, 1H, *J* = 4.5, 16.0, H<sub>5b</sub>), 3.15 (m, 4H, NCH<sub>2</sub>CH<sub>2</sub>N), 6.52 (d, 1H, *J* = 7.5, Ar-H), 6.58 (d, 1H, *J* = 8.0), 6.77 (dd, 1H, *J*<sub>1</sub> = *J*<sub>2</sub> = 7.0, Ar-H), 6.87 (dd, 1H, *J* = 7.5, 8.0, Ar-H), 6.93 (m, 2H, Ar-H), 7.21 (m, 2H, Ar-H), 9.17 (brs, 1H, NH); <sup>13</sup>C NMR (CD<sub>3</sub>OD): δ = 154.9, 151.3, 137.2, 128.6, 125.8, 121.9, 119.7, 119.1, 116.0, 111.1, 60.8, 49.1, 47.8, 29.0, 25.6, 25.4; MS (EI, 70 eV): *m/z* 308 (M<sup>+</sup>, 120 (100)); Anal. calcd for C<sub>20</sub>H<sub>24</sub>N<sub>2</sub>O: C 77.89, H 7.84; N 9.08 found C 77.95, H 7.99, N 8.97.

## 5.2. Pharmacology

### 5.2.1. HEK293 cell culture and membrane preparation

HEK293 cells, stably expressing the human serotonin 5-HT<sub>7</sub> receptor gene, were grown in Falcon flasks in DMEM (Cambrex, Verviers, Belgium) supplemented with 10% FBS (Cambrex), and 2 µg/mL puromycin (Sigma–Aldrich, St Louis, MO) at 37 °C in a 5% CO<sub>2</sub> atmosphere. For radioligand binding experiments, the cells, at 80% of confluence, were collected in PBS/EDTA (Lonza, Basel, Switzerland, cat.# BE02-017F) and centrifuged at 1000 × *g* for 10 min. The pellet obtained was resuspended in the same buffer used in binding experiments and homogenized with Ultra Turrax. The homogenate was centrifuged at 48,000 × *g* for 90 min. The final pellet was stored at –80 °C until the day of experiment. The protein concentration of membrane suspension was determined using the Bradford method (Pierce, Rockford, IL, USA) with bovine albumin as standard.

### 5.2.2. Competition binding assay

Competition binding experiments were performed incubating membranes (5–10 µg of protein/sample) with a single concentration of [<sup>3</sup>H]5-HT (5 nmol/L) (Biotrend, Cologne, Germany), in the presence of test compounds, in 96-well filter plates (MultiScreen system, cat # MAFBN0B10, Millipore, Billerica, MA, USA) for 30 min at 25 °C in a total volume of 200 µL/well of Tris–HCl 50 mM, pH 7.4, containing 10 µmol/L pargyline, 4 mmol/L CaCl<sub>2</sub> and 0.05% ascorbic acid. Non-specific binding was determined in the presence of 100 µM cold 5-HT (Sigma–Aldrich). At the end of incubation, bound and free radioligand were separated by filtering the 96-well filter plates using a Millipore filtration apparatus (Multiscreen<sub>HTS</sub>

vacuum manifold). Filter plates were then washed several times with ice-cold buffer (50 mM Tris–HCl, pH 7.4) and filter-bound radioactivity measured using a MicroBeta counter (PerkinElmer) after addition of 30 µL/well of OptiPhase SuperMix scintillation cocktail (PerkinElmer). Compounds were first tested at fixed (10–1 and/or 0.1 µmol/L) concentrations and those active at these concentrations (50% inhibition vs control) were repeated at scalar concentrations (ranging from 10<sup>–5</sup> to 10<sup>–10</sup> mol/L).

### 5.2.3. Data evaluation

Data were analyzed using nonlinear regression with GraphPad PRISM commercial software. Test compound concentrations causing a half maximal inhibition of control values (IC<sub>50</sub> calculated by GraphPad Prism software) were calculated. Inhibitory binding constant (*K*<sub>i</sub>) values were calculated from IC<sub>50</sub> values, according to the Cheng and Prusoff equation [33], *K*<sub>i</sub> = IC<sub>50</sub>/(1 + [C]/*K*<sub>d</sub>), where [C] is the concentration of the radioligand and *K*<sub>d</sub> its dissociation constant.

### 5.2.4. Human 5-HT<sub>7</sub> functional assays

Agonist and antagonist behavior were evaluated by Cerep [26] in cAMP functional assays performed in CHO cells, according to the method of Adham et al. [34]. Agonist behavior was determined by evaluating the concentration-dependent elevation of intracellular cAMP levels produced by test compounds. Serotonin showed EC<sub>50</sub> = 29 nM. Antagonist behavior of test compounds was determined evaluating the effect on cAMP accumulation stimulated by 300 nM 5-HT.

IC<sub>50</sub> values (concentration causing a half-maximal inhibition of the control specific agonist response) were determined by nonlinear regression analysis of the concentration–response curves generated with mean replicate values using Hill equation curve fitting ( $Y = D + [(A - D)/(1 + (C/IC_{50})^{nH})]$ , where *Y* = specific response, *D* = minimum specific response, *A* = maximum specific response, *C* = compound concentration, and *nH* = slope factor). This analysis was performed using a software developed at Cerep (Hill software). The apparent dissociation constants (*K*<sub>B</sub>) were calculated using the modified Cheng and Prusoff equation ( $K_B = IC_{50}/(1 + (A/EC_{50}A))$ ), where *A* = concentration of reference agonist in the assay, and EC<sub>50</sub>*A* = EC<sub>50</sub> value of the reference agonist). The reference compound mesulergine had *K*<sub>B</sub> = 16 nM.

### 5.2.5. Binding assays on other serotonin receptors

Affinity of compounds **5n**, **5i**, **14a** and **16i** for 5-HT<sub>1A</sub>, 5-HT<sub>1B</sub>, 5-HT<sub>1D</sub>, 5-HT<sub>2A</sub>, 5-HT<sub>2B</sub>, 5-HT<sub>2C</sub>, 5-HT<sub>3</sub>, 5-HT<sub>4e</sub>, 5-HT<sub>5a</sub> and 5-HT<sub>6</sub> receptors was evaluated by Cerep [26]. % Displacement of reference ligands at the concentration of 1 µM of **5n**, **5i**, **14a** and **16i** are reported in Supplementary Material Table S1.

## 5.3. Molecular modeling

The X-ray structure of the 5-HT<sub>1B</sub> serotonin receptor in complex with the agonist dihydroergotamine (PDB code: 4IAQ) [35] was used as the template for 5-HT<sub>7</sub> receptor modeling. The thermostabilized apocytochrome b<sub>562</sub> RIL segment and all non-protein atoms were removed and hydrogen atoms were added using the Protein Preparation Wizard [36] implemented in Maestro 9.2 [37]. The amino acid sequence of the human 5-HT<sub>7</sub> receptor was retrieved from the Universal Protein Resource (UniProt ID: P34969) [38,39]. Initial alignment was carried out with Prime 3.0 [40] and subsequently optimized considering conserved sequence motifs of class A GPCRs [21,41] (Fig. S2). Since the intracellular loop (ICL) 3 of the 5-HT<sub>1B</sub> receptor was replaced with the thermostabilized apocytochrome b<sub>562</sub> RIL, the ICL3 sequence of the 5-HT<sub>7</sub> receptor was excluded from homology modeling and Ala270, at the N-



terminus of ICL3, was directly linked to Ser317 belonging to the C-terminus of ICL3. This should not influence docking results, since the putative binding pocket of the 5-HT<sub>7</sub> receptor is located in the extracellular portion of the helix bundle. The all-hydrogen 5-HT<sub>7</sub> model structure obtained with Prime 3.0 was subjected to a restrained minimization using Impact 5.7 [42] and the OPLS2005 force field [43] to an RMSD value of 0.3 Å calculated on the protein heavy atoms. The stereochemical quality of the minimized structure was evaluated using Procheck [44]. The Ramachandran plot of the final refined structure showed only Val230 in extracellular loop (ECL) 3 and Ala270 at the N terminus of IL3 in disallowed regions (Fig. S3).

Compound **5i** was built with Maestro 9.2 and subjected to an energy minimization procedure using the OPLS2005 force field to a convergence threshold of 0.05 kJ mol<sup>-1</sup> Å<sup>-1</sup>. Induced-Fit Docking (IFD) [45,46] was then applied to account for receptor flexibility during docking calculations. During the initial softened-potential docking, a Van der Waals (VdW) radii scaling of 0.7 and 0.5 were applied on protein and ligand non-polar atoms, respectively. Moreover, Asp142<sup>2,65</sup>, Arg367<sup>7,36</sup> and Leu370<sup>7,39</sup> were temporarily mutated to alanines, to favor initial ligand placement. Asp162<sup>3,32</sup> was used to define the location of the binding site and the dimensions of bounding and enclosing boxes were set to 10 Å and 30 Å, respectively. Initial docking runs were performed in standard precision (SP) mode, applying a hydrogen-bond constraint between the protonated piperazine nitrogen of **5i** and Asp162<sup>3,32</sup>. A core constraint was also applied on compound **5i**, to induce the heavy atoms of the indole ring to superpose to the corresponding atoms of dihydroergotamine (co-crystallized with the 5-HT<sub>1B</sub> receptor), with a tolerance of 1 Å. Twenty ligand poses were retained for subsequent refinement. Each ligand–receptor complex obtained at the end of the softened-potential docking run was then submitted to side chain and backbone refinements with Prime 3.0. During this phase, the previously removed side chains were re-introduced and all residues located within 5.0 Å from the ligand molecule were refined by a side chain conformational search, followed by energy minimization of the residues and the ligand molecule. To prevent Glide from searching for alternative accommodations of the ligand during the final docking stage of the IFD workflow, each ligand molecule obtained at the end of the Prime stage was not re-docked into its induced-fit receptor structure, but it was only optimized in the field of the receptor and subsequently scored using default Glide settings (VdW radii scaling of 1.0 and 0.8 for receptor and ligand non-polar atoms, respectively). The resulting ligand–protein complexes were ranked according to their IFD score and the 5-HT<sub>7</sub> receptor–**5i** complex showing the best IFD score was selected. To prevent ECL3 from interpenetrating into the lipid bilayer, a loop optimization procedure was conducted on the segment Pro351<sup>6,59</sup>–Pro362<sup>7,31</sup> using Prime 3.0. To this end, the selected 5-HT<sub>7</sub> receptor–**5i** complex was aligned to the 4IAQ three-dimensional structure deposited in the Orientations of Protein in Membranes (OPM) database [47], which includes the coordinates of the putative membrane boundary planes. The orientation and position of the membrane boundaries were used to generate a low-dielectric slab-shaped region in Prime 3.0, which simulates the lipid bilayer environment during loop refinement. The resulting 5-HT<sub>7</sub> receptor–**5i** complex bearing the optimized ECL3 portion was energy-minimized to a convergence threshold of 0.5 kJ mol<sup>-1</sup> Å<sup>-1</sup>, applying the OPLS2005 force field and the GB/SA water solvation treatment. The resulting complex is represented in Fig. 2a.

The energy-minimized 5-HT<sub>7</sub> receptor–**5i** complex was embedded in a pre-equilibrated POPC lipid bilayer using Desmond version 2.2 [48,49] and subsequently solvated with TIP3P water molecules. Chlorine atoms were added to neutralize the net charge of the system. The simulation box measured 64 × 76 × 93 Å<sup>3</sup> and

contained 87 lipid molecules, 12 chlorine atoms and 8284 water molecules. The system was initially relaxed using a customized version of a membrane relaxation protocol implemented in the Desmond package and subsequently simulated for 100 ns in the NPT ensemble. Temperature and pressure were set to 310 K and 1 atm, respectively, applying the Langevin coupling scheme [50], and the M-SHAKE algorithm [51] was used to constrain all bond lengths to hydrogen atoms. VdW and short-range electrostatic interactions were cut off at 9 Å and long-range electrostatic interactions were computed using the smooth Particle Mesh Ewald method [52]. A multistep RESPA scheme [53] was used to integrate the equations of motion, applying a timestep of 2 fs for the bonded and the short-range nonbonded interactions and of 6 fs for the long-range nonbonded interactions. The OPLS2005 force field was used for both the relaxation and the production phase of the MD simulation.

The last snapshot of the MD simulation was submitted to energy minimization with Desmond 2.2, applying the OPLS2005 force field to a convergence threshold of 0.5 kcal mol<sup>-1</sup> Å<sup>-1</sup>. The energy-minimized 5-HT<sub>7</sub> receptor–**5i** complex was then extracted from the solvated lipid bilayer and used for docking studies of compounds **14m**, **27**, **18a** and **18c**. Docking grids were generated with Glide 5.7 [54] and were centered on the **5i** molecule, setting bounding and enclosing boxes to 10 Å and 30 Å on each side, respectively. To retain the key ionic interaction with Asp162<sup>3,32</sup>, a hydrogen-bond constraint was applied between the protonated piperazine nitrogen of ligands and Asp162<sup>3,32</sup> during docking runs. Twenty docking poses were collected for each ligand and the top-ranked pose according to the Emodel value was merged into the 5-HT<sub>7</sub> model structure. The resulting complexes were subjected to an energy minimization process using the OPLS2005 force field and the GB/SA water solvation treatment to a convergence threshold of 0.5 kJ mol<sup>-1</sup> Å<sup>-1</sup>. The energy-minimized complexes are depicted in Fig. 2c, d and 4.

## Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.ejmech.2014.04.034>.

## References

- [1] M. Pytliak, V. Vargová, V. Mechírová, M. Felšöci, Serotonin receptors – from molecular biology to clinical applications, *Physiological Research* (Prague, Czech Republic) 60 (2011) 15–25.
- [2] W.H. Visser, R.H. de Vriend, M.W. Jaspers, M.D. Ferrari, Sumatriptan in clinical practice: a 2-year review of 453 migraine patients, *Neurology* 47 (1996) 46–51.
- [3] R.B. Lydiard, An overview of generalized anxiety disorder: disease state–appropriate therapy, *Clinical Therapeutics* 22 (Suppl. A) (2000) A3–A19 discussion A20–24.
- [4] R.J. Milne, R.C. Heel, Ondansetron. Therapeutic use as an antiemetic, *Drugs* 41 (1991) 574–595.
- [5] T.W. Lovenberg, B.M. Baron, L. de Lecea, J.D. Miller, R.A. Prosser, M.A. Rea, P.E. Foye, M. Racke, A.L. Slone, B.W. Siegel, A novel adenylyl cyclase-activating serotonin receptor (5-HT<sub>7</sub>) implicated in the regulation of mammalian circadian rhythms, *Neuron* 11 (1993) 449–458.
- [6] M. Ruat, E. Traiffort, R. Leurs, J. Tardivel-Lacombe, J. Diaz, J.M. Arrang, J.C. Schwartz, Molecular cloning, characterization, and localization of a high-affinity serotonin receptor (5-HT<sub>7</sub>) activating cAMP formation, *Proceedings of the National Academy of Sciences of the United States of America* 90 (1993) 8547–8551.
- [7] J.A. Bard, J. Zgombick, N. Adham, P. Vaysse, T.A. Branchek, R.L. Weinshank, Cloning of a novel human serotonin receptor (5-HT<sub>7</sub>) positively linked to adenylyl cyclase, *The Journal of Biological Chemistry* 268 (1993) 23422–23426.
- [8] P. Vanhoenacker, G. Haegeman, J.E. Leysen, 5-HT<sub>7</sub> receptors: current knowledge and future prospects, *Trends in Pharmacological Sciences* 21 (2000) 70–77.
- [9] D.R. Thomas, J.J. Hagan, 5-HT<sub>7</sub> receptors, *CNS & Neurological Disorders – Drug Targets* 3 (2004) 81–90.

- [10] A. Matthys, G. Haegeman, K. Van Craenenbroeck, P. Vanhoenacker, Role of the 5-HT<sub>7</sub> receptor in the central nervous system: from current status to future perspectives, *Molecular Neurobiology* 43 (2011) 228–253.
- [11] M. Leopoldo, E. Lacivita, F. Berardi, R. Perrone, P.B. Hedlund, Serotonin 5-HT<sub>7</sub> receptor agents: structure–activity relationships and potential therapeutic applications in central nervous system disorders, *Pharmacology & Therapeutics* 129 (2011) 120–148.
- [12] (a) C. Kikuchi, H. Nagaso, T. Hiranuma, M. Koyama, Tetrahydrobenzindoles: selective antagonists of the 5-HT<sub>7</sub> receptor, *Journal of Medicinal Chemistry* 42 (1999) 533–535;  
(b) C. Kikuchi, T. Ando, T. Watanabe, H. Nagaso, M. Okuno, T. Hiranuma, M. Koyama, 2a-[4-(Tetrahydropyridoindol-2-yl)butyl]tetrahydrobenzindole derivatives: new selective antagonists of the 5-hydroxytryptamine 7 receptor, *Journal of Medicinal Chemistry* 45 (2002) 2197–2206;  
(c) M.L. López-Rodríguez, E. Porras, B. Benhamú, J.A. Ramos, M.J. Morcillo, J.L. Lavandera, First pharmacophoric hypothesis for 5-HT<sub>7</sub> antagonism, *Bioorganic & Medicinal Chemistry Letters* 10 (2000) 1097–1100;  
(d) R. Perrone, F. Berardi, N.A. Colabufo, E. Lacivita, M. Leopoldo, V. Tortorella, Synthesis and structure–affinity relationships of 1-[omega-(4-aryl-1-piperazinyl)alkyl]-1-aryl ketones as 5-HT<sub>7</sub> receptor ligands, *Journal of Medicinal Chemistry* 46 (2003) 646–649;  
(e) M. Leopoldo, E. Lacivita, M. Contino, N.A. Colabufo, F. Berardi, R. Perrone, Structure–activity relationship study on N-(1,2,3,4-tetrahydronaphthalen-1-yl)-4-aryl-1-piperazinehexanamides, a class of 5-HT<sub>7</sub> receptor agents. 2, *Journal of Medicinal Chemistry* 50 (2007) 4214–4221;  
(f) M. Leopoldo, E. Lacivita, P. De Giorgio, C. Fracasso, S. Guzzetti, S. Caccia, M. Contino, N.A. Colabufo, F. Berardi, R. Perrone, Structural modifications of N-(1,2,3,4-tetrahydronaphthalen-1-yl)-4-aryl-1-piperazinehexanamides: influence on lipophilicity and 5-HT<sub>7</sub> receptor activity. Part III, *Journal of Medicinal Chemistry* 51 (2008) 5813–5822;  
(g) B. Volk, J. Barkóczy, E. Hegedus, S. Udvari, I. Gacsályi, T. Mezei, K. Pallagi, H. Kompagne, G. Lévy, A. Egyed, L.G. Hársing Jr., M. Spedding, G. Simig, (Phenylpiperazinyl-butyl)oxindoles as selective 5-HT<sub>7</sub> receptor antagonists, *Journal of Medicinal Chemistry* 51 (2008) 2522–2532.
- [13] R.A. Glennon, Higher-end serotonin receptors: 5-HT<sub>5</sub>, 5-HT<sub>6</sub>, and 5-HT<sub>7</sub>, *Journal of Medicinal Chemistry* 46 (2003) 2795–2812.
- [14] (a) R.E. Wilcox, J.E. Ragan, R.S. Pearlman, M.Y. Brusniak, R.M. Eglén, D.W. Bonhaus, T.E. Tenner Jr., J.D. Miller, High-affinity interactions of ligands at recombinant guinea pig 5HT<sub>7</sub> receptors, *Journal of Computer-aided Molecular Design* 15 (2001) 883–909;  
(b) E.S. Vermeulen, A.W. Schmidt, J.S. Sprouse, H.V. Wikström, C.J. Grol, Characterization of the 5-HT<sub>7</sub> receptor. Determination of the pharmacophore for 5-HT<sub>7</sub> receptor agonism and CoMFA-based modeling of the agonist binding site, *Journal of Medicinal Chemistry* 46 (2003) 5365–5374;  
(c) E.S. Vermeulen, M. van Smeden, A.W. Schmidt, J.S. Sprouse, H.V. Wikström, C.J. Grol, Novel 5-HT<sub>7</sub> receptor inverse agonists. Synthesis and molecular modeling of arylpiperazine- and 1,2,3,4-tetrahydroisoquinoline-based arylsulfonamides, *Journal of Medicinal Chemistry* 47 (2004) 5451–5466;  
(d) A. Lepailleur, R. Bureau, S. Lemaître, F. Dauphin, J.-C. Lancelot, V. Contesse, S. Lenglet, C. Delarue, H. Vaudry, S. Rault, Molecular design based on 3D pharmacophores. Applications to 5-HT<sub>7</sub> receptors, *Journal of Chemical Information and Computer Science* 44 (2004) 1148–1152;  
(e) M. Kołaczowski, M. Nowak, M. Pawłowski, A.J. Bojarski, Receptor-based pharmacophores for serotonin 5-HT<sub>7</sub>R antagonists-implications to selectivity, *Journal of Medicinal Chemistry* 49 (2006) 6732–6741;  
(f) M.L. López-Rodríguez, E. Porras, M.J. Morcillo, B. Benhamú, L.J. Soto, J.L. Lavandera, J.A. Ramos, M. Olivella, M. Campillo, L. Pardo, Optimization of the pharmacophore model for 5-HT<sub>7</sub>R antagonism. Design and synthesis of new naphtholactam and naphthosultam derivatives, *Journal of Medicinal Chemistry* 46 (2003) 5638–5650;  
(g) E. Badarau, R. Bugno, F. Suzenet, A.J. Bojarski, A.-L. Finaru, G. Guillaumet, SAR studies on new bis-aryls 5-HT<sub>7</sub> ligands: synthesis and molecular modeling, *Bioorganic & Medicinal Chemistry* 18 (2010) 1958–1967;  
(h) T. Varin, H. Gutiérrez-de-Terán, M. Castro, J. Brea, F. Fabis, F. Dauphin, J. Aqvist, A. Lepailleur, P. Perez, J. Burgueno, J.M. Vela, M.I. Loza, J. Rodrigo, Phe369(7.38) at human 5-HT<sub>7</sub> receptors confers interspecies selectivity to antagonists and partial agonists, *British Journal of Pharmacology* 159 (2010) 1069–1081;  
(i) E. Lacivita, D. Patarnello, N. Stroth, A. Caroli, M. Niso, M. Contino, P. De Giorgio, P. Di Pilato, N.A. Colabufo, F. Berardi, R. Perrone, P. Svenningsson, P.B. Hedlund, M. Leopoldo, Investigations on the 1-(2-biphenyl)piperazine motif: identification of new potent and selective ligands for the serotonin(7) (5-HT<sub>7</sub>) receptor with agonist or antagonist action in vitro or ex vivo, *Journal of Medicinal Chemistry* 55 (2012) 6375–6380.
- [15] S. Rivara, A. Lodola, M. Mor, A. Bedini, G. Spadoni, V. Lucini, M. Pannacci, A. Caronno, F. Scaglione, G. Gobbi, G. Spadoni, A. Bedini, P. Orlando, S. Lucarini, G. Tarzia, N-(substituted-anilinoethyl)amides: design, synthesis, and pharmacological characterization of a new class of melatonin receptor ligands, *Journal of Medicinal Chemistry* 50 (2007) 6618–6626.
- [16] M.E. Speeter, W.C. Anthony, The action of oxalyl chloride on indoles: a new approach to tryptamines, *Journal of the American Chemical Society* 76 (1954) 6208–6210.
- [17] X. Piao, X. Zhang, Y. Mori, M. Koishi, A. Nakaya, S. Inoue, I. Aoki, A. Otomo, S. Yokoyama, Nonlinear optical side-chain polymers post-functionalized with high- $\beta$  chromophores exhibiting large electro-optic property, *Journal of Polymer Science Part A: Polymer Chemistry* 49 (2011) 47–54.
- [18] R.A. Glennon, M. Lee, J.B. Rangisetty, M. Dukat, B.L. Roth, J.E. Savage, A. McBride, L. Rauser, S. Hufeisen, D.K. Lee, 2-Substituted tryptamines: agents with selectivity for 5-HT<sub>6</sub> serotonin receptors, *Journal of Medicinal Chemistry* 43 (2000) 1011–1018.
- [19] M.B. Isaac, T. Xin, A. O'Brien, D. St-Martin, A. Naismith, N. MacLean, J. Wilson, L. Demchysyn, A. Tehim, A. Slassi, 1-(Bicyclopiperazinyl)ethylindoles and 1-(homopiperazinyl)ethyl-indoles as highly selective and potent 5-HT<sub>7</sub> receptor ligands, *Bioorganic & Medicinal Chemistry Letters* 12 (2002) 2451–2454.
- [20] (a) G. Lebon, T. Warne, P.C. Edwards, K. Bennett, C.J. Langmead, A.G.W. Leslie, C.G. Tate, Agonist-bound adenosine A2A receptor structures reveal common features of GPCR activation, *Nature* 474 (2011) 521–525;  
(b) S.G.F. Rasmussen, H.-J. Choi, J.J. Fung, E. Pardon, P. Casarosa, P.S. Chae, B.T. Devree, D.M. Rosenbaum, F.S. Thian, T.S. Koblika, A. Schnapp, I. Konetzki, R.K. Sunahara, S.H. Gellman, A. Pautsch, W.I. Weis, B.K. Koblika, Structure of a nanobody-stabilized active state of the  $\beta_2$  adrenoceptor, *Nature* 469 (2011) 175–180.
- [21] J.A. Ballesteros, H. Weinstein, Integrated methods for the construction of three-dimensional models and computational probing of structure–function relations in G protein-coupled receptors, in: S.C. Sealfon, P.M. Conn (Eds.), *Methods in Neurosciences*, vol. 25 Academic Press, San Diego (CA), 1995, pp. 366–428.
- [22] B.Y. Ho, A. Karschin, T. Branchek, N. Davidson, H.A. Lester, The role of conserved aspartate and serine residues in ligand binding and in function of the 5-HT<sub>1A</sub> receptor: a site-directed mutation study, *FEBS Letters* 312 (1992) 259–262.
- [23] T. Warne, R. Moukhametzianov, J.G. Baker, R. Nehmé, P.C. Edwards, A.G.W. Leslie, G.F.X. Schertler, C.G. Tate, The structural basis for agonist and partial agonist action on a  $\beta(1)$ -adrenergic receptor, *Nature* 469 (2011) 241–244.
- [24] J. Kim, Y. Kim, J. Tae, M. Yeom, B. Moon, X.-P. Huang, B.L. Roth, K. Lee, H. Rhim, I.H. Choo, Y. Chong, G. Keum, G. Nam, H. Choo, Aryl biphenyl-3-ylmethylpiperazines as 5-HT<sub>7</sub> receptor antagonists, *ChemMedChem* 8 (2013) 1855–1864.
- [25] J.-S. Surgand, J. Rodrigo, E. Kellenberger, D. Rognan, A chemogenomic analysis of the transmembrane binding cavity of human G-protein-coupled receptors, *Proteins* 62 (2006) 509–538.
- [26] Tests Performed by Cerep, <http://www.cerep.fr>.
- [27] G. Himbert, D. Fink, K. Diehl, P. Rademacher, A.J. Bittner, Cycloadditionen, 15. Einfluß von Substituenten in p-, m- und o-Position am Aromaten auf die intramolekulare Diels-Alder-Reaktion von Allencarbonsäure-aniliden und -phenylestern, *Chemische Berichte* 122 (1989) 1161–1173.
- [28] H. Xie, D. Ng, S.N. Savinov, B. Dey, P.D. Kwong, R. Wyatt, A.B. Smith 3rd, W.A. Hendrickson, Structure–activity relationships in the binding of chemically derivatized CD4 to gp120 from human immunodeficiency virus, *Journal of Medicinal Chemistry* 50 (2007) 4898–4908.
- [29] D.A. Klump, M. Garza, A. Jones, S. Mendoza, Synthesis of aryl-substituted piperidines by superacid activation of piperidones, *The Journal of Organic Chemistry* 64 (1999) 6702–6705.
- [30] G. Thoma, F. Nuninger, M. Schaefer, K.G. Akyel, R. Albert, C. Beerli, C. Bruns, E. Francotte, M. Luyten, D. MacKenzie, L. Oberer, M.B. Streiff, T. Wagner, H. Walter, G. Weckbecker, H.-G. Zervos, Orally bioavailable competitive CCR5 antagonists, *Journal of Medicinal Chemistry* 47 (2004) 1939–1955.
- [31] P. Kumar, Synthesis of substituted 1-tetralones, *Organic Preparations and Procedures International* 29 (1997) 477–480.
- [32] D. Lauffer, R. Tomlinson, E. Ottow, M. Botfield, Acyclic Piperazine and Piperidine Derivatives Which are Useful for Treating Neuronal Damage (2002-12-27) WO2002102381.
- [33] Y. Cheng, W.H. Prusoff, Relationship between the inhibition constant ( $K_i$ ) and the concentration of inhibitor which causes 50 per cent inhibition ( $I_{50}$ ) of an enzymatic reaction, *Biochemical Pharmacology* 22 (1973) 3099–3108.
- [34] N. Adham, J.M. Zgombick, J. Bard, T.A. Branchek, Functional characterization of the recombinant human 5-hydroxytryptamine<sub>7</sub>(a) receptor isoform coupled to adenylate cyclase stimulation, *The Journal of Pharmacology and Experimental Therapeutics* 287 (1998) 508–514.
- [35] C. Wang, Y. Jiang, J. Ma, H. Wu, D. Wacker, V. Katritch, G.W. Han, W. Liu, H.-P. Huang, E. Vardy, J.D. McCorvy, X. Gao, X.E. Zhou, K. Melcher, C. Zhang, F. Bai, H. Yang, L. Yang, H. Jiang, B.L. Roth, V. Cherezov, R.C. Stevens, H.E. Xu, Structural basis for molecular recognition at serotonin receptors, *Science* 340 (2013) 610–614.
- [36] Schrödinger Suite, Protein Preparation Wizard; Epik Version 2.2, Schrödinger, LLC, New York, NY, 2011. Impact Version 5.7, Schrödinger, LLC, New York, NY, 2011; Prime version 3.0, Schrödinger, LLC, New York, NY, 2011.
- [37] Maestro, version 9.2, Schrödinger, LLC, New York, NY, 2011.
- [38] UniProt Consortium, Reorganizing the protein space at the Universal Protein Resource (UniProt), *Nucleic Acids Research* 40 (2012) D71–D75.
- [39] E. Jain, A. Bairoch, S. Duvaud, I. Phan, N. Redaschi, B.E. Suzek, M.J. Martin, P. McGarvey, E. Gasteiger, Infrastructure for the life sciences: design and implementation of the UniProt website, *BMC Bioinformatics* 10 (2009) 136.
- [40] Prime, Version 3.0, Schrödinger, LLC, New York, NY, 2011.

- [41] T. Mirzadegan, G. Benkö, S. Filipek, K. Palczewski, Sequence analyses of G-protein-coupled receptors: similarities to rhodopsin, *Biochemistry* 42 (2003) 2759–2767.
- [42] Impact Version 5.7, Schrödinger, LLC, New York, NY, 2011.
- [43] W.L. Jorgensen, D.S. Maxwell, J. Tirado-Rives, Development and testing of the OPLS all-atom force field on conformational energetics and properties of organic liquids, *Journal of the American Chemical Society* 118 (1996) 11225–11236.
- [44] R.A. Laskowski, M.W. MacArthur, D.S. Moss, J.M. Thornton, PROCHECK: a program to check the stereochemical quality of protein structures, *Journal of Applied Crystallography* 26 (1993) 283–291.
- [45] W. Sherman, T. Day, M.P. Jacobson, R.A. Friesner, R. Farid, Novel procedure for modeling ligand/receptor induced fit effects, *Journal of Medicinal Chemistry* 49 (2006) 534–553.
- [46] Schrödinger Suite, Induced Fit Docking Protocol; Glide Version 5.7, Schrödinger, LLC, New York, NY, 2011. Prime Version 3.0, Schrödinger, LLC, New York, NY, 2011.
- [47] M.A. Lomize, A.L. Lomize, I.D. Pogozheva, H.I. Mosberg, OPM: orientations of proteins in membranes database, *Bioinformatics* 22 (2006) 623–625.
- [48] Desmond Molecular Dynamics System, Version 3.0, D. E. Shaw Research, New York, NY, 2011. Maestro-Desmond Interoperability Tools, version 3.0, Schrödinger, New York, NY, 2011.
- [49] K.J. Bowers, F.D. Sacerdoti, J.K. Salmon, Y. Shan, D.E. Shaw, E. Chow, H. Xu, R.O. Dror, M.P. Eastwood, B.A. Gregersen, J.L. Klepeis, I. Kolossvary, M.A. Moraes, Scalable algorithms for molecular dynamics simulations on commodity clusters, in: SC '06 Proceedings of the 2006 ACM/IEEE Conference on Supercomputing, ACM New York, NY, 2006, p. 84.
- [50] S.E. Feller, Y. Zhang, R.W. Pastor, B.R. Brooks, Constant pressure molecular dynamics simulation: the Langevin piston method, *The Journal of Chemical Physics* 103 (1995) 4613–4621.
- [51] R.A. Lippert, K.J. Bowers, R.O. Dror, M.P. Eastwood, B.A. Gregersen, J.L. Klepeis, I. Kolossvary, D.E. Shaw, A common, avoidable source of error in molecular dynamics integrators, *The Journal of Chemical Physics* 126 (2007) 046101.
- [52] U. Essmann, L. Perera, M.L. Berkowitz, T. Darden, H. Lee, L.G. Pedersen, A smooth particle mesh Ewald method, *The Journal of Chemical Physics* 103 (1995) 8577–8593.
- [53] M. Tuckerman, B.J. Berne, G.J. Martyna, Reversible multiple time scale molecular dynamics, *The Journal of Chemical Physics* 97 (1992) 1990–2001.
- [54] Glide, Version 5.7, Schrödinger, LLC, New York, NY, 2011.
- [55] P. Holmberg, D. Sohn, R. Leideborg, P. Caldirola, P. Zlatoidsky, S. Hanson, N. Mohell, S. Rosqvist, G. Nordvall, A.M. Johansson, R. Johansson, Novel 2-aminotetralin and 3-aminochroman derivatives as selective serotonin 5-HT<sub>7</sub> receptor agonists and antagonists, *Journal of Medicinal Chemistry* 47 (2004) 3927–3930.
- [56] P. Holmberg, L. Tedenborg, S. Rosqvist, A.M. Johansson, Novel 3-aminochromans as potential pharmacological tools for the serotonin 5-HT<sub>7</sub> receptor, *Bioorganic & Medicinal Chemistry Letters* 15 (2005) 747–750.
- [57] C.G. Thomson, M.S. Beer, N.R. Curtis, H.J. Diggle, E. Handford, J.J. Kulagowski, Thiazoles and thiopyridines: novel series of high affinity h5HT<sub>7</sub> ligands, *Bioorganic & Medicinal Chemistry Letters* 14 (2004) 677–680.
- [58] G. Chłoń-Rzepa, P. Zmudzki, G. Satała, B. Duszyńska, A. Partyka, D. Wróbel, M. Jastrzębska-Więsek, A. Wesołowska, A.J. Bojarski, M. Pawłowski, P. Zajdel, New 8-aminoalkyl derivatives of purine-2,6-dione with arylalkyl, allyl or propynyl substituents in position 7, their 5-HT<sub>1A</sub>, 5-HT<sub>2A</sub>, and 5-HT<sub>7</sub> receptor affinity and pharmacological evaluation, *Pharmacological Reports: PR* 65 (2013) 15–29.