

REVIEW

B. Nürnberg · T. Gudermann · G. Schultz

Receptors and G proteins as primary components of transmembrane signal transduction

Part 2. G proteins: structure and function

Received: 20 December 1994/Accepted: 18 January 1995

Abstract Seven-transmembrane receptors signal through nucleotide-binding proteins (G proteins) into the cell. G proteins are membrane-associated proteins composed of three subunits termed α , β and γ , of which the $G\alpha$ subunit classifies the heterotrimer. So far, 23 different mammalian $G\alpha$ subunits are known, which are grouped in four subfamilies (G_s , G_i , G_q , G_{12}) on the basis of their amino acid similarity. They carry an endogenous GTPase activity allowing reversible functional coupling between ligand-bound receptors and effectors such as enzymes and ion channels. In addition, five $G\beta$ and seven $G\gamma$ subunits have been identified which form tightly associated $\beta\gamma$ heterodimers. Upon activation by a ligand-bound receptor the G protein dissociates into $G\alpha$ and $G\beta\gamma$, which both transmit signal by interacting with effectors. On the G protein level, specificity and selectivity of the incoming signal is accomplished by G protein trimers composed of distinct subunits. On the other hand, many receptors have been shown to activate different G proteins, thereby regulating diverse signal transduction pathways.

Key words Cellular effectors · Cellular signaling · G protein · Heptahelical receptor · Signal transduction

Abbreviations CT Cholera toxin · PT Pertussis toxin

Introduction

In mammals, intercellular transmembrane signaling *via* hormones, neurotransmitters, autacoids and growth factors is accomplished by various signal-transducing mechanisms. In many cases primary cellular substrates are membrane-spanning receptors. Some of these (e.g., ligand-gated ion channels, protein tyrosine kinases, phosphoprotein tyrosine phosphatases, or guanylyl cyclases) transmit signals by intramolecular coupling be-




Fig. 1 Transmembrane receptor-effector systems. Extracellular signals activate transmembrane receptors. Receptors can be coupled intramolecularly to an effector domain generating intracellular signals (*left*). Examples are ligand-gated ion channels and enzyme-linked receptors. In contrast, the G protein coupled signal transduction cascade consists of a heptahelical transmembrane receptor, a heterotrimeric G protein, and effectors such as enzymes or ion channels (*right*)

tween an extracellular ligand-binding domain and an intracellular (enzyme) or transmembrane effector domain (ion channel pore) [1–7]. Principally, their common feature is a directly coupled receptor-effector system generating intracellular signals upon agonist binding. In contrast, G protein-coupled transmembrane signal requires the sequential and reversible interaction of a machinery composed of three elements (Fig. 1): (a) a receptor (see part I [8]), (b) a transducer (i.e., a heterotrimeric G protein), and (c) an effector (i.e., an enzyme, ion channel or transporter) [9]. This review focuses on the structures and functions of heterotrimeric G proteins. G protein-coupled effectors are reviewed only briefly since a large number of recent excellent reviews are available (e.g., [10–14]).

Heterotrimeric G proteins

Heterotrimeric regulatory guanine nucleotide-binding proteins (**G proteins**) belong to the superfamily of GTP-binding proteins or GTPases and are extremely conserved throughout the animal kingdom, underlining their pivotal role in cellular regulations [15, 16]. Although more than 20 years have passed since Rodbell, Birnbaumer, and associates discovered that GTP regulates the specific binding of glucagon to liver membranes, and the fact that GTP is required for receptor-dependent activation of adenylyl cyclase, at present the structures of G proteins are much better understood than their functions (Fig. 2) [17–19]. G proteins are composed of three different subunits termed α , β and γ , with molecular masses of approximately 39–45, 35–39, and 6–8 kDa, respectively. Currently, the heterotrimer is classified by the nature of its $G\alpha$ subunit (Table 1). G proteins are membrane-associated proteins attached to the cytoplasmic surface from which they can be solubilized by detergents (see below). Exceptions are the retinal transducins (G_t) which are more hydrophilic and consequently extractable from rod outer segment membranes without detergents. Convincing evidence suggests that G proteins are also located on intracellular membranes, involved in membrane trafficking and vesicular transport mechanisms of the cell [20–23]. Two newly identified members of G proteins belonging to the G_s and G_i subfamilies have recently been reported to be specifically involved in the regulation of intracellular transport processes (see below). Whereas functionally active monomeric $G\alpha$ subunits can be isolated, dissociation of the $G\beta\gamma$ dimer does not occur under physiological conditions without resulting in a loss of function of the monomers. G proteins are enzymes bearing a GTPase activity on the $G\alpha$ subunit (EC 3.6.1.-). On the basis of their amino acid similarity G protein α subunits are at present classified in four families: G_s , G_i , G_q , and G_{12} (see Table 1) [19, 24, 25]. Nucleotide sequence analysis shows that the genes of two subfamilies, G_i and G_q , are closely related, suggesting that the members of these subfamilies segregated as pairs of closely linked genes whereas members of the G_s and G_{12} subfamily segregated as unlinked genes during evolution [26].

All G proteins follow the same principal **activation/inactivation cycle** allowing reversible and specific transmission of hormonal signals (Fig. 2) [27]. Activation of the G protein is initiated by its interaction with distinct cytoplasmic segments of a ligand-activated heptahelical receptor [8], resulting in the release of bound GDP. Most likely the former event is the rate-limiting step of the G protein activation reaction followed by high-affinity binding (nM affinity) of cytosolic GTP or nonhydrolyzable analogs [e.g., guanosine 5'-(3-*O*-thio)-triphosphate or guanosine 5'-($\beta\gamma$ -imido)-triphosphate under experimental conditions]. During this reaction Mg^{2+} is very closely associated with GTP and $G\alpha$ in this complex and is required for activation of the $G\alpha$ subunit and for subsequent hydrolytic activity [28]. Up-




Fig. 2 Activation/Inactivation cycle of G proteins. α , β , and γ represent the $G\alpha$, $G\beta$, and $G\gamma$ subunits of the G protein. Pertussis toxin functionally uncouples the G protein from the receptor whereas cholera toxin abolishes the GTPase activity of the G protein

on G protein activation, dissociation of the $G\alpha$ subunit from the $G\beta\gamma$ complex occurs, which does not necessarily result in a physical separation. Both the activated $G\alpha$ subunit and the $G\beta\gamma$ complex modulate effector proteins until hydrolysis of the GTP bound to the $G\alpha$ subunit terminates signaling (intrinsic GTPase activity). The GTPase activity of the $G\alpha$ subunit is modulated by some effectors (e.g., cGMP-dependent phosphodiesterase, phospholipase C- β , calcium channel) which thereby function as GAPs (such as the GTPase-activating proteins of the ras family). Following GTP hydrolysis the inactive GDP-bound $G\alpha$ subunit dissociates from the effector and reassociates with the $G\beta\gamma$ complex to form a $G\alpha\beta\gamma$ heterotrimer. This inactive heterotrimer becomes available for subsequent activation cycles. Interestingly, the receptor-stimulated GDP/GTP exchange reaction varies among different G proteins: G_i/G_o proteins are fast nucleotide exchangers, G_t , G_s , G_z and G proteins of the G_q subfamily show smaller rate constants, and for G_{12} and G_{13} very small guanine nucleotide exchange rates are reported [29, 30]. Supplementary to this well established mechanism for activation of G proteins, an alternative route has recently been postulated [31]. In addition to the above reaction, it involves $G\beta$ subunits as high-energy phosphate acceptors. It is thought that following the GTPase reaction the phosphate is transferred to a histine residue of a $G\beta$ subunit, resulting in a high-energy phosphate bond which allows rapid phosphate transfer to a GDP bound to a second $G\alpha$ subunit. Furthermore, it has been speculated that the phosphorylated $G\beta$ subunit represents an activated state of the $G\beta\gamma$ dimer similar to the GTP-bound $G\alpha$ subunit [32]. Significant differences between various cells and tissues apparently exist with respect to $G\beta$ phosphorylation efficiency, pointing to the need of further studies.

Table 1 Heterotrimeric regulatory guanine nucleotide binding proteins (G proteins)

G proteins are named according to their α -subunits and are listed with their effector systems so far known (effector given in parentheses indicates that the indicated effect may be without physiological relevance). CT and PT indicate whether the α subunits are substrates of cholera or pertussis toxins, respectively. \uparrow , \downarrow , Increased and decreased effector activity, respectively

^a Receptor-mediated activation is required for ADP ribosylation by CT

	Occurrence	Effector	PT/CT
G_s subfamily			
G_s (S, L)	Ubiquitous	Adenylyl cyclases \uparrow	CT
G_{olf}	Olfactory epithelium	Adenylyl cyclases \uparrow	CT
G_i subfamily			
Transducin _{r,c}	Retinal rods and cones	cGMP PDE \uparrow	CT, PT
Gustducin	Taste cells	?	CT, PT
G_{i1}	Mainly neuronal cells	?	PT (CT) ^a
G_{i2}	Ubiquitous	Adenylyl cyclases \downarrow	PT (CT) ^a
G_{i3}	Mainly nonneuronal cells	?	PT (CT) ^a
$G_{o1,2}$	Neuronal, endocrine cells	Calcium channels \downarrow	PT (CT) ^a
G_z	Neuronal cells, platelets	?	—
G_q subfamily			
G_q	Ubiquitous	Phospholipases C- β \uparrow	—
G_{q1}	Nonhematopoietic cells	Phospholipases C- β \uparrow	—
G_{q2}	Many cells	Phospholipases C- β \uparrow	—
G_{q3}	Hematopoietic cells	(Phospholipases C- β \uparrow)	—
G_{12} subfamily			
G_{12}	Ubiquitous	?	—
G_{13}	Ubiquitous	?	—

Table 2 Covalent modifications of G protein α subunits

^a Functional uncoupling of receptor-G protein interaction

^b Also modification of G_{α_i} by lauric acid and unsaturated fatty acids ($C_{14:2}$, $C_{14:1}$)

^c In platelets additionally acylation of G_{α_i} , G_{α_q} , G_{α_z} , $G_{\alpha_{13}}$ by arachidonic acid

^d Substoichiometric phosphorylation; unknown physiological role

^e Unknown physiological role

^f Inhibition of pertussis toxin sensitive PI response

ADP ribosylation	
Cholera toxin: constitutively activated G-protein	G_{α_s} (ARG-186/201), G_{α_i}
Pertussis toxin: inactivated $\alpha\beta\gamma$ heterotrimer ^a	$G_{\alpha_{i1-3}}$, $G_{\alpha_{o1-2}}$
sequence motif: CGA Φ (C-terminus)	$G_{\alpha_{i1-3}}$, $G_{\alpha_{o1-2}}$, G_{α_i}
Acylation	
Myristoylation ^b (cotranslational, irreversible) sequence motif: MGXXXS/T? (N-terminus)	$G_{\alpha_{i1-3}}$, $G_{\alpha_{o1-2}}$, G_{α_i} , G_{α_z}
enzyme: <i>N</i> -myristyl transferase	
Palmitoylation ^c (posttranslational, reversible) sequence motif: MGC? (N-terminus)	$G_{\alpha_{i1-3}}$, G_{α_z} , $G_{\alpha_{o1-2}}$, G_{α_s} , $G_{\alpha_{i1-13}}$
Phosphorylation	
cAMP-dependent protein kinase ^d	G_{α_i} ?, G_{α_s} ?
Protein kinase C ^e	G_{α_z} (Ser-16/27), $G_{\alpha_{i2}}$
cGMP-dependent protein kinase ^f	$G_{\alpha_{i1-3}}$

In contrast to transmembrane signal transduction by intramolecularly linked receptor effector domains, intermolecular functional coupling of heptahelical receptors to G proteins allows **amplification of the hormone signal** since a ligand-bound receptor activates many G proteins within a short time [17, 33]. In fact, one light-activated rhodopsin light receptor couples to 3000 G proteins per second [34]. Pharmacological modulation of this G protein-conveyed signal transduction cascade is currently regulated almost exclusively at the receptor level, i.e., receptor agonists or antagonists. On the G protein level direct modulation by drugs is rather limited [35]. However, some bacterial toxins are frequently used for studying the function of G proteins. Of particular experimental importance are cholera toxin (CT) and pertussis toxin (PT), which catalyze ADP-ribosylation of various G_{α} subunits [36]. CT modifies easily an arginine residue located in one of the nucleotide-binding regions of the G_{α} subunit of G_s and G_i . A functional consequence is the abolishment of the endogenous GTPase activity resulting in continuous activation of the G_{α} sub-

unit (see Table 2). In contrast, PT catalyzes the transfer of an ADP moiety from NAD⁺ to a cysteine residue of the α subunit of G_i/G_o and G_t . The amino acid motif required for PT-mediated ADP ribosylation is similar to the consensus site for isoprenylation of $G\gamma$ subunits (see below) except for the C-terminal aromatic amino acid (see Tables 2, 4). Since the modified cysteine is located four residues upstream the carboxy-terminus, PT-catalyzed ADP-ribosylation prevents interaction of the receptor with the G protein representing a functional receptor-G protein uncoupling. Nevertheless, PT-modified G proteins still possess the capability to interact with effectors.

G_{α} subunits

Currently 23 different α subunits including several splice variants of G_s , G_i and G_o are known, which are the products of 17 different genes (see Table 1) [37]. The properties and functions of the CT- and PT-sensitive members

of the G_s and G_i subfamilies are well understood as a result of studying purified proteins and employing sophisticated tools and techniques, for example, the photoreactive GTP-analog 4'-azidoanilido- $[\alpha\text{-}^{32}\text{P}]\text{GTP}$, bacterial toxins, specific antibodies, and antisense technology [29, 38–40].

The CT-sensitive G_s belongs to the **G_s subfamily**, and four splice variants are ubiquitously expressed. All G_s isoforms directly activate adenylyl cyclases and calcium channels [12, 13]. G_{olf} , which is closely related to G_s on the basis of its amino acid similarity (88% identity), appears to be expressed only in the sensory neurons within the olfactory epithelium. Similarly to G_s , it has the capacity to stimulate adenylyl cyclases. Most recently an "extra-large" 92-kDa G_s protein [$G\alpha_{s(XL)}$] has been reported which consists of a new 51-kDa XL portion linked to an N-terminally truncated $G\alpha_s$ [41]. It is suggested to be specifically associated with the trans-golgi-network, hence mediating CT-sensitive effects on secretory vesicle formation.

Members of the PT-sensitive **G_i subfamily** include at least six gene products, of which G_{i2} is expressed ubiquitously [42]. It is involved in the hormonal inhibition of adenylyl cyclases. However, knock-out of the $G\alpha_{i2}$ gene does not produce fatal disturbances in mice. This is consistent with previous observations showing that murine embryonic $G\alpha_{i2}$ -null cell lines regulate adenylyl cyclase, grew, and differentiated in vitro similar to wild-type embryonic stem cells [43, 44]. Nevertheless, these observations do not rule out the possibility that vital functions of G_{i2} are substituted by other closely related G proteins, for example, G_{i1} and G_{i3} . Recently a splice variant of G_{i2} was discovered [$G_{i2(L)}$] which is resistant to PT-mediated ADP ribosylation and appears to be important for intracellular traffic processes [45]. It lacks the PT-sensitive carboxy-terminal peptide motif $CGA\Phi$ [36]. The physiological functions of G_{i1} and G_{i3} are less defined although they show a slightly different expression pattern. Both proteins are detectable throughout the organism: G_{i1} is particularly abundant in neural tissues whereas G_{i3} is predominantly found in nonneuronal systems. They are very homologous to G_{i2} (85–94% identity based on amino acid similarity) despite being transcripts of different genes. Further analysis revealed that some of the $G\alpha_i$ subunit genes were generated by tandem gene duplication which occurred prior to the divergence of rodents and primates [26]. Interestingly, the four subfamilies of mammalian $G\alpha$ genes evolved prior to the divergence of vertebrate and invertebrate lineages. All three G_i proteins stimulate potassium channels most likely *via* their $\beta\gamma$ subunits [19]. In contrast, results obtained in the neuroendocrine cell line GH₃ show that thyrotropin-releasing hormone receptor induced stimulation of calcium channels is apparently transduced *via* activation of G_{i2} [46]. G_{o1} and G_{o2} are alternatively spliced-products of the same gene, exhibiting an expression pattern restricted to endocrine and neuronal systems. The amino acid sequences differ in their carboxy-terminus which is proposed to represent an effector-interacting domain. Either protein mediates receptor-induced inhibitory effects on

calcium channels [10]. The combination of intranuclear microinjection of specific antisense oligonucleotides and the patch clamp technique allowed identification of specific $G_o \alpha\beta\gamma$ heterotrimers involved in signal transduction pathways regulating voltage-activated calcium channels [47–50]. In the rat pituitary cell line, GH₃, somatostatin, and carbachol (through M_4 muscarinic receptors) cause inhibition of prolactin secretion and of L-type calcium currents. Here, the two splice variants of the $G\alpha_o$ subunit, $G\alpha_{o1}$ and $G\alpha_{o2}$, are differentially involved in the coupling of M_4 muscarinic and somatostatin receptors to L type calcium channels, respectively. Moreover, there is a high specificity of $G\beta$ and $G\gamma$ subunits. In this system, $G\alpha_{o1}\beta_3\gamma_4$ and $G\alpha_{o2}\beta_1\gamma_3$ appear to represent the active G_o subforms transferring inhibitory signals from muscarinic and somatostatin receptors to the calcium channel, respectively. Infusion of purified $G\alpha_o$ subunits into PT-pretreated GH₃ cells identify the same $G\alpha_o$ subunits as the antisense oligonucleotide technique [51]. *Transducins* (the rod and cone photoreceptor G proteins) also belonging to the G_i subfamily are found solely in the retina; they couple rhodopsin functionally to a cGMP-phosphodiesterase [52].

$G\alpha_i$ was the first G protein to be crystalized in both its GDP-bound and guanosine 5'-(3-*O*-thio)-triphosphate bound conformations and subjected to X-ray structural crystallographic analysis [53, 54]. Conformation of 86% of the amino acids were found identical for either condition. The remaining 14% of amino acids, which showed flexibility depending on the nucleotide binding state, were identified or confirmed as important domains for interaction with $\beta\gamma$ dimers (amino-terminus) or receptors and effectors (carboxy-terminus). Very recently Gilman and associates [55] presented the three-dimensional structure of native $G\alpha_{i1}$ and two mutants. Based on the crystal structures it was found that glutamine²⁰⁴ and arginine¹⁷⁸ (Glu²⁰⁰ and Arg¹⁷⁴ in $G\alpha_i$) are apparently mobile residues which do not bind nucleotides in the ground state but interact closely with substrates (GTP and bound water) in the transition state, i.e. during hydrolysis of GTP *via* a second-order nucleophilic substitution reaction (S_N2). Both residues are present on comparable loci found in all heterotrimeric $G\alpha$ subunits. They were previously predicted as a prerequisite for GTPase activity of various G proteins based on site-directed mutagenesis experiments [56]. Furthermore, this arginine residue represents the substrate which is sensitive to CT (see above). Interestingly, the monomeric GTPase p21^{ras} does not have an arginine in a position analogous to Arg-178 (in $G\alpha_{i1}$), thus explaining the enzymes' weak hydrolytic activity and its inability to bind the G protein activating AlF_4^- complex. In addition to previous assumptions that $\text{GDP}\cdot\text{AlF}_4^-$ is a GTP analog, it now became clear that $\text{GDP}\cdot\text{AlF}_4^-$ rather mimics the transition state of the hydrolytic second-order nucleophilic substitution reaction, thereby fixing the $G\alpha$ subunit in an active conformation which allows effector interaction [57]. These data are confirmed by analysis of the crystal structure of AlF_4^- -bound transducin [58].

Gustducin, as G_{olf} and transducins, belongs to the

group of G proteins solely expressed in distinct sensory neurons [59]. It is detectable in taste buds of all taste papillae of the tongue. Its function is not yet known, but it is speculated on the basis of analogy and high structural conservation to G_i that gustducin transduces taste receptor activation into activation of a taste cell phosphodiesterase.

Over the past 5 years cDNAs of a number of G protein α subunits have been cloned and sequenced which do not serve as substrates for PT because they lack the carboxy-terminal cysteine. The first entity discovered, called G_z (or G_x), belongs to the G_i subfamily. It exhibits an extremely slow basal GTPase activity similar to monomeric GTPases, for example, ras. G_z was purified partially from bovine brain membranes [60], G_z -activating receptors or G_z -regulated effectors are currently unknown.

The ubiquitously expressed G proteins G_q and G_{11} , G_{14} , G_{15} , and G_{16} form the **G_q subfamily** [37]. Native G_q and G_{11} were initially purified and characterized by the laboratories of John Exton and Paul Sternweis [11]. G_{14} has not been purified so far, but mammalian G_{16} has been isolated from cell membranes gained from the insect baculovirus/Sf9 overexpression system [61]. G_q and G_{11} exhibit low basal GTPase activity and guanosine 5'-(3-*O*-thio)-triphosphate binding. Based on a large body of evidence it is now accepted that G_q and G_{11} stimulate phospholipases C- β which are responsible for triggering the PT-insensitive PI response [11, 62]. Utilizing photoaffinity labeling of receptor-activated G proteins it has been shown that activation of these G proteins is induced by vasopressin receptors in liver cell membranes and by thromboxane A_2 and thrombin receptors in platelets [63, 64]. Additional evidence came from reconstitution and transfection experiments showing the functional coupling of various receptors inducing a PT-insensitive PI response and proteins of the G_q subfamily. G_{16} appears to be the human homolog of murine G_{15} , both expressed in hematopoietic cells, whereas G_{11} is only detectable in nonhematopoietic cells [65–67]. The physiological role of the $G_{15/16}$ proteins remains to be elucidated. Notably, expression of the proteins is reduced with differentiation of leukocytes [65, 66]. In cotransfection experiments Simon and associates [68, 69] demonstrated the functional coupling between interleukin-8 and C5a receptors and G_{16} but not G_q and G_{11} .

G_{12} and G_{13} , belonging to the **G_{12} subfamily**, are expressed ubiquitously at very low abundance [70, 71]. Native $G\alpha_{13}$ and $G\alpha_{12}$ have recently been purified from rat and bovine brain membranes [72–75]. Their $G\alpha$ subunits share less than 50% amino acid identity than other G protein subfamilies. However, the amino acid identity between $G\alpha_{12}$ and $G\alpha_{13}$ amounts to only 67%, which is less than that observed among members of the other G protein subfamilies. Both G proteins appear to be relatively slowly activated by thromboxane A_2 and thrombin receptors in human platelets [64]. In addition, there are reports that G_{13} mediates slow inhibition of calcium currents induced by bradykinin in a neuroblastoma/glioma cell line, and that G_{13} stimulates a Na^+/H^+ exchanger

[76–78]. Interestingly, members of the G_{12} subfamily have been proposed as oncogenes exhibiting the most remarkable transforming potential of all heterotrimeric G proteins. Expression of mutants of G_{12} and G_{13} lacking GTPase activity resulted in a malignant transformation of cells [79–81]. Mice injected with transformed cells overexpressing these $G\alpha$ subunits rapidly developed large tumors. Even the wild-type gene products of G_{12} and G_{13} harbor significant oncogenic activity [81, 82].

Co- and posttranslational modifications of $G\alpha$ subunits

Most G protein α subunits are modified by co- and posttranslational events (see Table 2) [83, 84]. PT-sensitive proteins of the G_i subfamily contain an N-terminal *MGCXXS/T* amino acid motif. During elongation of the nascent protein, the initial methionine is cleaved, followed by **myristoyl amidation** of the glycine residue. In addition to myristate (C_{14} carbon acid), various unsaturated C_{14} fatty acids and the C_{12} lauryl acid are linked to the glycine residue of the retinal G protein transducin [85]. Based on competition experiments utilizing acylated peptides from the N-terminus of $G\alpha_i$, it was found that lauryl-modified peptides are much less potent competitors of $G\alpha$ - $G\beta\gamma$ interactions than myristoyl peptides, suggesting that laurylated $G\alpha_i$ interacts with $G\beta\gamma$ less efficiently than its myristoylated counterpart. Myristoylation appears to be irreversible. This protein modification, on the one hand, enhances lipophilicity of the $G\alpha$ subunit and, on the other, contributes to the specificity of protein-protein interactions. For instance, the affinity of a myristoylated $G\alpha$ subunit to $G\beta\gamma$ complexes is significantly higher than that of the nonmyristoylated counterpart. Simultaneously, myristoylation enhances the potency of the $G\alpha$ subunit to activate effectors.

Palmytoylation of all $G\alpha$ subunits except transducin was recently reported as a posttranslational, reversible modification occurring at a cysteine residue near the N-terminus via a labile thioester bond and possibly at other sites [86–89]. After cell fractionation nonmyristoylated $G\alpha$ subunits carrying a palmitate (C_{16} carbon acid) accumulated in the particulate fraction whereas the protein devoid of fatty acid modification was found in the cytosol. However, it remains questionable whether the only function of palmitoyl acylation is to anchor the $G\alpha$ subunit in the membrane. For instance, the degree of palmytoylation of $G\alpha$ subunits is receptor-dependently regulated [90]. Furthermore, nonpalmytoylated chimeras of $G\alpha_s/G\alpha_i$ and $G\alpha_q/G\alpha_i$ are unable to stimulate their respective effectors, i.e., adenylyl cyclases and phospholipases C [91]. Recently, arachidonate-acylated $G\alpha$ subunits ($G\alpha_i$, $G\alpha_q$, $G\alpha_z$, $G\alpha_{13}$) were detected in platelets [92].

Reports on **phosphorylation of G proteins** are scarce. For the PT-insensitive G protein G_z , selective phosphorylation of a serine residue near the N-terminus by protein kinase C is well established [93, 94]. However, function-

al consequences remain cryptic. Other $G\alpha$ subunits also possess consensus sites sensitive to protein kinases which allow in vitro phosphorylation under certain experimental conditions [83]. In neuroblastoma/glioma cells the activation of protein kinase C stimulated phosphorylation of G_{i2} with kinetics similar to its effect on adenylyl cyclase [95]. Stable transfection of CHO cells with cGMP protein kinase I α (cGMP-pk) suppressed the thrombin-induced increase in inositol 1,4,5-trisphosphate and cytosolic calcium. This regulatory step was found to be PT-sensitive. Further studies showed that cGMP-pk phosphorylates a PT-sensitive G protein in these cells, and that purified cGMP-pk is able to phosphorylate reconstituted G_i proteins but not G_o [96] (see Table 2).

$G\beta$ and γ subunits

Much attention has been focused in recent years on $G\beta\gamma$ dimers, which were previously thought to function solely as membrane anchors. Meanwhile, the number of purified, cloned, and sequenced $G\beta$ and $G\gamma$ subunits has increased substantially (Table 3), comprising five G protein β and seven $G\gamma$ subunits [25, 97]. Among the $G\beta$ subunits $G\beta_5$ exhibits the largest difference regarding amino acid sequence homology, with 53% identity to other $G\beta$ subunits [98]. In contrast, comparison of the deduced amino acid sequences of $G\beta_1$ – $G\beta_4$ displays a high degree of homology of at least 79% identity. In contrast, $G\gamma$ subunits demonstrate a higher structural diversity, for example, the bovine retinal $G\gamma_1$ subunit and the $G\gamma_2$ subunit exhibit only 36% amino acid identity [99]. These observations raise the question of whether $G\gamma$ subunits (beside $G\alpha$ subunits) have a larger impact on the specificity of receptor–G protein interactions than $G\beta$ subunits [19, 97]. Support for this view is coming from a recent study employing isoprenylated (see below) C-terminal peptides derived from the $G\gamma_1$ amino acid sequence which are sufficient to stabilize the active form of the rhodopsin photoreceptor and uncouple rhodopsin– G_t interaction in a concentration-dependent manner [100]. On the other hand, $G\beta$ subunits contain essential structural attributes necessary for G protein–subunit and G protein–protein interactions. Evidence arises from computer-based analysis of the structural features of $G\beta$ subunits. They belong to the family of **WD repeat proteins** [101]. Their common feature is a conserved core of the repeating unit ($n = 4$ –8) of 36–46 amino acids usually ending with an Trp-Asp (WD). All 27 members of this ancient regulatory protein family are found in eukaryotes but not in prokaryotes. They regulate different cellular functions, such as cell division, cell fate determination, gene transcription, transmembrane signaling, mRNA modification, and vesicle fusion. Furthermore, $G\beta$ subunits contain **coiled-coil** forming structures at their N-terminal domain (Met-1 to Ser-31 of $G\beta_1$) [102–104]. Coiled coils are stabilized protein structures resulting from the interaction of two or more right-handed α -helices that wind around each other in a left-handed super-

Table 3 G protein β and γ subunits

β_1	Ubiquitous
β_2	Many cells
β_3	Many cells
β_4	Many cells
β_5	Neuronal cells
γ_1	Retinal rods
γ_2	Many cells
γ_3	Many cells
γ_4	Many cells
γ_5	Many cells
γ_7	Many cells
γ_8	Retinal cones

$\beta_1\gamma_1$ and $\beta_3\gamma_8$ have been demonstrated as $\beta\gamma$ complexes occurring in retinal rods and cones, respectively. Whether all conceivable $\beta\gamma$ combinations occur in other cells is not known. According to co-expression studies, $\beta_2\gamma_1$ is an impossible combination

coil [105]. $G\beta$ subunits are predicted to form a three-stranded coiled-coil of the G protein heterotrimer *via* N-terminal domains of each the $G\alpha$, $G\beta$, and the $G\gamma$ subunit. Experimental evidence supports this view in that the N-termini of the $G\beta$ subunits are elementary for dimerization with $G\gamma$ subunits [104, 106]. Furthermore, while certain domains of $G\beta$ subunits are qualitatively essential for protein interactions, they presumably do not contribute to the specificity of $G\beta\gamma$ complexing since certain $G\beta\gamma$ combinations are known not to dimerize, for example, $G\beta_2\gamma_1$ [107]. However, the specificity of the interaction between G protein subunits, for example, $G\beta_1$, $G\beta_2$ and $G\gamma_1$, $G\gamma_2$, appears to be accomplished by a 14 amino acid region of $G\gamma_1$ (Cys-36 to Gly-49) and $G\gamma_2$ (Ala-33 to Lys-46) [99].

Posttranslational modifications of $G\gamma$ subunits

G protein $\beta\gamma$ dimers exhibit a higher lipophilicity than $G\alpha$ subunits. This is correlated with **isoprenylation of $G\gamma$ subunits** [83, 84]. The signal sequence causing this modification is a CAAX box motif ($X \neq \Phi$, see above) at the C-terminus common to all $G\gamma$ subunits and some monomeric GTPases (Table 4). This motif triggers a number of posttranslational events initiated by prenylation of the cysteine through a thioether bond followed by endoproteolytic cleavage of the three carboxy-terminal amino acids (AAX) and methylation at the new C-terminus, which is the prenylated cysteine residue. Efforts are currently being made to elucidate the mechanism for $G\beta\gamma$ assembly and functional consequences of the various posttranslational modifications. Initial results suggest that prenylation is not required for $G\beta\gamma$ assembly which occurs prior to the proteolytic processing of $G\gamma$ [108]. Interestingly, both isoprenylation and carboxymethylation of $G\gamma_1$ appear to contribute to the efficiency of membrane association, subunit interaction between $G\alpha$ and $G\beta\gamma$, and the functional coupling of the G protein with receptors or effectors [84, 109, 110]. Only the retina-spe-

Table 4 Posttranslational modifications of G protein γ subunits

Sequence motif: CAAX ^a (C-terminus)		
Cysteine polyisoprenylation		
Farnesyl transferase (C ₁₅)		
CAAX: X=Ala, Cys, Glu, Met, Ser		G γ_1 , ras-family ^b
Geranylgeranyl transferase I (C ₂₀)		
CAAX: X=Leu		G γ_2 , G γ_3 , G γ_5 ?, G γ_7 ?
Geranylgeranyl transferase II (C ₂₀)		
CCXX; CXC		rab-family ^b
Endoproteolytic cleavage of three C-terminal amino acids		
Cysteine carboxymethylation		

^a A Aliphatic amino acid, X=Φ^b Members of monomeric GTPases**Table 5** Effectors regulated by G protein α subunits and $\beta\gamma$ complexes

	α subunit	$\beta\gamma$ complex
cGMP phosphodiesterase	↑	
βARK		↑↑
ras-Regulating proteins		↑↑
PI-3-kinase		↑↑
Phospholipases C-β	↑	↑
Adenylyl cyclases	↑, ↓	↑, ↓
Ca ²⁺ channels	↑, ↓	
Cl ⁻ channels	↑↑	
Na ⁺ channels	↑, ↓	
K ⁺ ch. (inw. rect., ATP regul.)	↑	↑

cific G γ_1 subunit is modified by a C₁₅ sesquiterpene (*farnesyl moiety*) whereas the nonretinal G γ s are modified by a C₂₀ diterpene (*geranylgeranyl group*). This difference in isoprenylation of G γ affects lipophilicity of the G $\beta\gamma$ dimer, i.e., retinal G $\beta_1\gamma_1$ dimers are soluble without detergents, in contrast to all other known G $\beta\gamma$ dimers including those of G $\beta_1\gamma_x$. Furthermore, the type of isoprenylation has an impact on all functions of G $\beta\gamma$ dimers studied so far. For instance, PT-catalyzed ADP-ribosylation of G α subunits is less efficient in the presence of G $\beta_1\gamma_1$ dimers than in the presence of geranylgeranyl-modified G $\beta_1\gamma_x$ complexes. In general, C₂₀ isoprenylated G $\beta\gamma$ complexes appear to exhibit a higher potency in modulating effectors than C₁₅-isoprenylated G $\beta_1\gamma_1$.

G protein regulated effectors

Effectors regulated by G $\beta\gamma$ dimers include *adenylyl cyclases*, *phospholipases C-β*, *potassium channels* and *phosphoinositide 3 kinase* (Table 5) [19, 24, 111, 112]. Additionally, G $\beta\gamma$ dimers also appear to participate in the regulation of receptors. There is reasonable evidence to propose that G protein $\beta\gamma$ subunits contribute to desensitization of receptors by a negative feedback mechanism *via* receptor kinases [113]. Evidence suggests that G $\beta\gamma$ dimers also activate the monomeric GTPase *ras* through binding to ras-regulatory proteins hence signaling into the growth factor signal transduction pathway [114, 115]. Very recently a region of homology termed the

pleckstrin homology domain (PH) was discovered which is detectable in a plethora of proteins including signal transduction components, for example, receptor and protein kinases, phospholipases C, GTPases, GAPs, and nucleotide exchange factors [116–119]. PH domains are apparently required for effective binding of G $\beta\gamma$ subunits. If this is true generally, it would facilitate searching for yet unidentified signal transduction components involved in G-protein-mediated cellular signaling.

The number of cellular effector systems affected by G protein α and/or $\beta\gamma$ subunits are increasing rapidly. Well-established effector enzymes and ion channels are already being discussed (see above and see Table 5) [10–13, 52]. Recent experiments suggest that also amiloride-sensitive renal epithelial *sodium channels* are regulated differently by PT-sensitive G proteins depending on the state of the channel and/or G protein [120, 121]. Further potential candidates belonging to the class of G protein-regulated cellular effector enzymes are phospholipases D and A₂ [122–124]. In addition, nonselective cation channels expressed in a wide range of tissues may be modulated by PT-insensitive G proteins [125].

Acknowledgements The authors' own research reported herein was supported by Deutsche Forschungsgemeinschaft, Sonderforschungsbereich 366, and Fonds der Chemischen Industrie. Many important references could not be included because of space limitations but are indicated in the reviews cited.

References

- Koesling D, Böhme E, Schultz G (1993) Guanylyl cyclases as effectors of hormone and neurotransmitter receptors. In: Huch F (ed) Neurotransmitter receptors. Elsevier, pp 325–338
- Jan LY, Jan YN (1992) Tracing the roots of ion channels. Cell 69:715–718
- Unwin N (1993) Neurotransmitter action: opening of ligand-gated ion channels. Neuron [Suppl] 10:31–41
- Schlessinger J, Ullrich A (1992) Growth factor signaling by receptor tyrosine kinase. Neuron 9: 383–391
- Egan SE, Weinberg A (1993) The pathway to signal achievement. Nature (London) 365:781–783
- Charbonneau H, Tonks NK (1992) 1002 protein phosphatases? Annu Rev Cell Biol 8:463–493
- Garbers DL (1992) Guanylyl cyclase receptors and their endocrine, paracrine, and autocrine ligands. Cell 71:1–4
- Gudermann T, Nürnberg B, Schultz G (1995) Receptors and G proteins as primary components of transmembrane signal transduction. I. G protein-coupled receptors: structure and function. J Mol Med 73:51–63
- Watson S, Arkinstall S (1994) The G protein linked receptor facts book. Academic, New York
- Schultz G (1994) G proteins involved in hormonal regulations of the cytoplasmatic calcium concentrations. Ernst Schering Research Foundation, vol 19, Berlin
- Sternweis PC, Smrcka AV (1992) Regulation of phospholipase C by G proteins. Trends Biochem Sci 17:502–506
- Iyengar R (1993) Molecular and functional diversity of mammalian G α -stimulated adenylyl cyclases. FASEB J 7:768–775
- Tang WJ, Gilman AG (1992) Adenylyl cyclases. Cell 70:869–972
- Dickey BF, Birnbaumer L (eds) (1993) Handbook of experimental pharmacology, vol 108. Springer, Berlin Heidelberg New York
- Bourne HR, Sanders DA, McCormick F (1990) The GTPase superfamily: a conserved switch for diverse cell functions. Nature (London) 348:125–132

16. Bourne HR, Sanders DA, McCormick F (1991) The GTPase superfamily: a conserved switch for diverse cell functions. *Nature* (London) 349:117–127
17. Rodbell M, Krans HMJ, Pohl SL, Birnbaumer L (1971) The glucagon-sensitive adenylyl cyclase system in plasma membranes of rat liver. IV. Binding of glucagon: effect of guanyl nucleotides. *J Biol Chem* 246:1872–1876
18. Bourne HR, Nicoll R (1993) Molecular machines integrate coincident synaptic signals. *Neuron* [Suppl] 10:65–75
19. Clapham DE, Neer EJ (1993) New roles for G protein $\beta\gamma$ -dimers in transmembrane signalling. *Nature* (London) 365:403–406
20. Leyte A, Barr FA, Kehlenbach RH, Huttner WB (1992) Multiple trimeric G proteins on the trans-golgi network exert stimulatory and inhibitory effects on secretory vesicle formation. *EMBO J* 11:4795–4804
21. Pimplikar SW, Simons K (1993) Regulation of apical transport in epithelial cells by a G_i class of heterotrimeric G protein. *Nature* (London) 362:456–458
22. Bomsel M, Mostov K (1992) Role of heterotrimeric G proteins in membrane traffic. *Mol Biol Cell* 3:1317–1328
23. Barr FA, Leyte A, Huttner WB (1992) Trimeric G proteins and vesicle formation. *Trends Cell Biol* 2:91–94
24. Birnbaumer L (1992) Receptor-to-effector signaling through G proteins: roles for $\beta\gamma$ dimers as well as α subunits. *Cell* 71:1069–1072
25. Hepler JR, Gilman AG (1992) G proteins. *Trends Biochem Sci* 17:383–387
26. Wilkie TM, Gilbert DJ, Olsen AS, Chen XN, Amatruda TT, Korenberg JR, Trask BJ, de Jong P, Reed RR, Simon MI, Jenkins NA, Copeland NG (1992) Evolution of the mammalian G protein α subunit multigene family. *Nature Genetics* 1:85–91
27. Gilman AG (1987) G proteins: transducers of receptor-generated signals. *Annu Rev Biochem* 56:615–649
28. Higashijima T, Ferguson KM, Sternweis PC, Smigel MD, Gilman AG (1987) Effects of Mg^{2+} and the $\beta\gamma$ -subunit complex on the interactions of guanine nucleotides with G proteins. *J Biol Chem* 262:762–766
29. Fields TA, Linder ME, Casey PJ (1994) Subtype-specific binding of azidoanilido-GTP by purified G protein α subunits. *Biochemistry* 33:6877–6883
30. Laugwitz KL, Spicher K, Schultz G, Offermanns S (1994) Identification of receptor-activated G proteins: selective immunoprecipitation of photolabeled G protein α -subunits. *Methods Enzymol* 237:283–294
31. Kaldenberg-Stasch S, Baden M, Fessler B, Jakobs KH, Wieland T (1994) Receptor-stimulated guanine nucleoside triphosphate binding to G proteins: nucleotide exchange and β -subunit-mediated phosphotransfer reactions. *Eur J Biochem* 221:25–33
32. Sternweis PC (1994) The active role of $\beta\gamma$ in signal transduction. *Curr Opin Cell Biol* 6:198–203
33. Hille B (1992) G protein-coupled mechanisms and nervous signaling. *Neuron* 9:187–195 (1992)
34. Hargrave PA, Hamm HE, Hofmann KP (1993) Interaction of rhodopsin with the G protein, transducin. *Bioessays* 15:43–50
35. Nürnberg B (1994) Signaltransduktion durch heterotrimere G Proteine. *Pharmazie* 49:795–800
36. Gierschik P, Jakobs KH (1992) ADP-ribosylation of signal-transducing guanine nucleotide binding proteins by cholera and pertussis toxin. In: Herken H, Hucho F (eds) *Selective neurotoxicity. Handbook of experimental pharmacology*, vol 102. Springer, Berlin Heidelberg New York, pp 807–839
37. Simon MI, Strathmann MP, Gautam N (1991) Diversity of G proteins in signal transduction. *Science* 252:802–808
38. Milligan G (1988) Techniques used in the identification and analysis of function of pertussis toxin-sensitive guanine nucleotide binding proteins. *Biochem J* 255:1–13
39. Johnson RA, Corbin JD (eds) (1991) Adenylyl cyclase, G proteins and guanylyl cyclase. *Methods Enzymol* 195
40. Iyengar R (ed) (1994) Heterotrimeric G proteins. *Methods Enzymol* 237
41. Kehlenbach RH, Matthey J, Huttner WB (1995) $XL\alpha$ s is a new type of G protein. *Nature* (London) 372:804–809
42. Nürnberg B, Spicher K, Harhammer R, Bosserhoff A, Frank R, Hiltz H, Schultz G (1994) Purification of a novel G protein α_o -subtype from mammalian brain. *Biochem J* 300:387–394
43. Rudolph U, Brabet P, Kaplan J, Hasty P, Bradley A, Birnbaumer L (1993) targeting of the $G_{i2}\alpha$ gene in IS cells with replacements and insertion vectors. *J Recept Res* 13:619–629
44. Mortensen RM, Zubiaur M, Neer EJ, Seidman JG (1991) Embryonic stem cells lacking a functional inhibitory G protein subunit (α_i) produced by gene targeting of both alleles. *Proc Natl Acad Sci USA* 88:7036–7040
45. Montmayeur JP, Borelli E (1994) Targeting of $G\alpha_{i2}$ to the golgi by alternative spliced carboxyl-terminal region. *Science* 263:95–98
46. Gollasch M, Kleuss C, Hescheler J, Wittig B, Schultz G (1993) G_{i2} and protein kinase C are required for thyrotropin-releasing hormone-induced stimulation of voltage-dependent Ca^{2+} channels in rat pituitary GH₃ cells. *Proc Natl Acad Sci USA* 90:5587–5591
47. Kleuss C, Hescheler J, Ewel C, Rosenthal W, Schultz G, Wittig B (1991) Assignment of G protein subtypes to specific receptors inducing inhibition of calcium currents. *Nature* (London) 353:43–48
48. Kleuss C, Scherübl H, Hescheler J, Schultz G, Wittig B (1992) Different β subunits determine G protein interaction with transmembrane receptors. *Nature* (London) 358:424–426
49. Kleuss C, Scherübl H, Hescheler J, Schultz G, Wittig B (1993) Selectivity in signal transduction determined by γ subunits of heterotrimeric G proteins. *Science* 259:832–834
50. Kalkbrenner F, Degtyar V, Schenker M, Hescheler J, Brendel S, Wittig B, Schultz G (1994) Subunit composition of the G protein coupling galanin receptors to L-type calcium channels. *Naunyn Schmiedeberg's Arch Pharmacol* 349:R13
51. Nürnberg B, Degtiar VE, Harhammer R, Uhde M, Hescheler J, Schultz G (1994) Hormone-induced G_o -subtype-specific inhibition of calcium currents. *Naunyn Schmiedeberg's Arch Pharmacol* 349:R13
52. Kaupp UB, Koch KW (1992) Role of cGMP and Ca^{2+} in vertebrate photoreceptor excitation and adaptation. *Annu Rev Physiol* 54:153–175
53. Noel JP, Hamm HE, Sigler PB (1993) The 2.2 Å crystal structure of transducin- α complexed with GTP γ S. *Nature* (London) 366:654–663
54. Lambbright DG, Noel JP, Hamm HE, Sigler PB (1994) Structural determinants for activation of a heterotrimeric G protein. *Nature* (London) 369:621–628
55. Coleman DE, Berghuis AM, Lee E, Linder ME, Gilman AG, Sprang SR (1994) Structures of active conformations of $G_{i\alpha 1}$ and the mechanism of GTP hydrolysis. *Science* 265:1405–1412
56. Goody RS (1994) How G proteins turn off. *Nature* (London) 372:220–221
57. Kleuss C, Raw AS, Lee E, Sprang SR, Gilman AG (1994) Mechanism of GTP hydrolysis by G protein α subunits. *Proc Natl Acad Sci USA* 91:9828–9831
58. Sondek J, Lambright DG, Noel JP, Hamm HE, Sigler PB (1994) GTPase mechanism of G protein from the 1.7 Å crystal structure of transducin α -GDP-AlF₄⁻. *Nature* (London) 372:276–279
59. McLaughlin SK, McKinnon PJ, Margolskee RF (1992) Gustducin as a taste-cell specific G protein closely related to the transducins. *Nature* (London) 357:563–569
60. Casey PJ, Fong HKW, Simon MI, Gilman AG (1990) G_{α_s} , a guanine nucleotide-binding protein with unique biochemical properties. *J Biol Chem* 265:2383–2390
61. Kozasa T, Hepler JR, Smrcka AV, Simon MI, Rhee SG, Sternweis PC, Gilman AG (1993) Purification and characterization of recombinant $G_{16\alpha}$ from Sf9 cells: activation of purified phospholipase C isoenzymes by G protein α subunits. *Proc Natl Acad Sci USA* 90:9176–9180
62. Dennis EA, Rhee SG, Billah MM, Hannun YA (1991) Role of phospholipases in generating lipid second messengers in signal transduction. *FASEB J* 5:2068–2077
63. Wange RL, Smrcka AV, Sternweis PC, Exton JH (1991) Photoaffinity labeling of two rat liver plasma membrane proteins

- with [32 P] γ -azidoanilido GTP in response to vasopressin. *J Biol Chem* 266:11409–11412
64. Offermanns S, Laugwitz KL, Spicher K, Schultz G (1994) G proteins of the G_{12} family are activated via thromboxane A_2 and thrombin receptors in human platelets. *Proc Natl Acad Sci USA* 91:504–508
 65. Wilkie TM, Scherle PA, Strathmann MP, Slepak VZ, Simon MI (1991) Characterization of G protein α subunits in the G_q class: expression in murine tissues and in stromal and hematopoietic cell lines. *Proc Natl Acad Sci USA* 88:10049–10053
 66. Amatruda TT, Steele DA, Slepak VZ, Simon MI (1991) $G\alpha_{16}$, a G protein α subunit specifically expressed in hematopoietic cells. *Proc Natl Acad Sci USA* 88:5587–5591
 67. Milligan G, Mullaney I, McCallum JF (1993) Distribution and relative levels of expression of the phosphoinositidase-C-linked G proteins $G_q\alpha$ and $G_{11}\alpha$: absence of $G_{11}\alpha$ in human platelets and hematopoietically derived cell lines. *Biochim Biophys Acta* 1179:208–212
 68. Amatruda TT, Gerard NP, Gerard C, Simon MI (1993) Specific interactions of chemoattractant factor receptors with G proteins. *J Biol Chem* 268:10139–10144
 69. Wu D, LaRosa GJ, Simon MI (1993) G protein-coupled signal transduction pathways for interleukin-8. *Science* 261:101–103
 70. Strathmann MP, Simon MI (1991) $G\alpha_{12}$ and $G\alpha_{13}$ subunits define a fourth class of G protein α subunits. *Proc Natl Acad Sci USA* 88:5582–5586
 71. Spicher K, Kalkbrenner F, Zobel A, Harhammer R, Nürnberg B, Söling A, Schultz G (1994) G_{12} and G_{13} α -subunits are immunochemically detectable in membranes of most tissues of various mammalian species. *Biochem Biophys Res Comm* 198:906–914
 72. Harhammer R, Nürnberg B, Spicher K, Schultz G (1994): Purification G protein $G\alpha_{13}$ from rat brain membranes. *Biochem J* 303:135–140
 73. Singer WD, Miller RT, Sternweis PC (1994) Purification and characterization of the α subunit of G_{13} . *J Biol Chem* 269:19769–19802
 74. Harhammer R, Nürnberg B, Spicher K, Schultz G (1994) Rapid purification of $G\alpha_{13}$ from bovine brain membranes: supportive effect of ethylene glycol. *Biochem Biophys Res Comm* 204:835–840
 75. Harhammer R, Nürnberg B, Spicher K, Schultz G (1995) Differential properties of the purified members of the G protein G_{12} -subfamily. *Naunyn Schmiedeberg Arch Pharmacol* (in press)
 76. Voyno-Yasenetskaya T, Conklin BR, Gilbert RL, Hooley R, Bourne HR, Barber DL (1994) $G\alpha_{13}$ stimulates Na-H exchange. *J Biol Chem* 269:4721–4724
 77. Dhanasekaran N, Vara Prasad MVVS, Wadsworth SJ, Dermott JM, van Rossum G (1994) protein kinase C-dependent and -independent activation of Na^+/H^+ exchanger by $G\alpha_{12}$ class of G proteins. *J Biol Chem* 269:11802–11806
 78. Wilk-Blaszczak MA, Singer WD, Gutowski S, Sternweis PC, Berladetti F (1994) The G protein G_{13} mediates bradykinin inhibition of voltage-dependent calcium current. *Neuron* 13:1215–1224
 79. Xu N, Bradley L, Ambdukar I, Gutkind JS (1993) A mutant α subunit of G_{12} potentiates the eicosanoid pathway and is highly oncogenic in NIH 3T3 cells. *Proc Natl Acad Sci USA* 90:6741–6745
 80. Jiang H, Wu D, Simon MI (1993) The transforming activity of activated $G\alpha_{12}$. *FEBS Lett* 330:319–322
 81. Xu N, Voyno-Yasenetskaya T, Gutkind JS (1994) Potent transforming activity of the G_{13} α subunit defines a novel family of oncogenes. *Biochem Biophys Res Comm* 201:603–609
 82. Chan AML, Fleming TP, McGovern ES, Chedid M, Miki T, Aaronson SA (1993) Expression cDNA cloning of a transforming gene encoding the wild-type $G\alpha_{12}$ gene product. *Mol Cell Biol* 13:762–768
 83. Yamane HK, Fung BKK (1993) Covalent modifications of G proteins. *Annu Rev Pharmacol Toxicol* 32:201–241
 84. Casey PJ (1994) Lipid modifications of G proteins. *Curr Opin Cell Biol* 6:219–225
 85. Casey P (1992) Visual differences. *Nature (London)* 359:671–672
 86. Degtyarev MY, Spiegel AM, Jones TL (1993) Increased palmitoylation of the G_s protein α subunit after activation by the β -adrenergic receptor or cholera toxin. *J Biol Chem* 268:23769–23772
 87. Linder ME, Middleton P, Hepler JR, Taussig R, Gilman AG, Mumby SM (1993) Lipid modifications of G proteins: α subunits are palmitoylated. *Proc Natl Acad Sci USA* 90:3675–3679
 88. Parenti M, Vigano MA, Newman CMH, Milligan G, Magee AI (1993) A novel N-terminal motif for palmitoylation of G protein α subunits. *Biochem J* 291:349–353
 89. Veit M, Nürnberg B, Spicher K, Harteneck C, Ponimaskin E, Schultz G, Schmidt MFG (1994) The α -subunits of G_{12} and G_{13} are palmitoylated, but not amidically myristoylated. *FEBS Lett* 339:160–164
 90. Mumby SM, Kleuss C, Gilman AG (1994) Receptor regulation of G protein palmitoylation. *Proc Natl Acad Sci USA* 91:2800–2804
 91. Wedegaertner PB, Chu DH, Wilson PT, Levis MJ, Bourne HR (1993) Palmitoylation is required for signaling functions and membrane attachment of $G_q\alpha$ and $G_{12}\alpha$. *J Biol Chem* 268:25001–25008
 92. Hallak H, Muszbek L, Laposata M, Belmonte E, Brass LF, Manning DR (1994) Covalent binding of arachidonate to G protein α subunits of human platelets. *J Biol Chem* 269:4713–4716
 93. Lounsbury KM, Casey PJ, Brass LF, Manning DR (1991) Phosphorylation of G_i in human platelets. selectivity and site of modification. *J Biol Chem* 266:22051–22056
 94. Lounsbury KM, Schlegel B, Poncz M, Brass LF, Manning DR (1993) Analysis of $G_{12}\alpha$ by site-directed mutagenesis. *J Biol Chem* 268:3494–3498
 95. Strassheim D, Malbon CC (1994) Phosphorylation of $G\alpha_{12}$ attenuates inhibitory adenylyl cyclase in neuroblastoma/glioma hybrid (NG-108-15) cells. *J Biol Chem* 269:14307–14313
 96. Pfeifer A, Nürnberg B, Kamm S, Uhde U, Schultz G, Ruth P, Hofmann F (1995): Cyclic GMP-dependent protein kinase blocks pertussis toxin-sensitive hormone receptor signaling pathways in CHO-cells. *J Biol Chem* (in press)
 97. Iñiguez-Lluhi J, Kleuss C, Gilman AG (1993) The importance of G protein $\beta\gamma$ subunits. *Trends Cell Biol* 3:230–236
 98. Watson AJ, Katz A, Simon MI (1994) A fifth member of the mammalian G protein β -subunit family. *J Biol Chem* 269:22150–22156
 99. Spring DJ, Neer EJ (1994) A 14-amino acid region of the G protein γ subunit is sufficient to confer selectivity of γ binding to the β subunit. *J Biol Chem* 269:22882–22886
 100. Kisselev OG, Ermolaeva MV, Gautam N (1994) A farnesylated domain in the G protein γ subunit is a specific determinant of receptor coupling. *J Biol Chem* 269:21399–21402
 101. Neer EJ, Schmidt CJ, Nambudripad R, Smith TF (1994) The ancient regulatory-protein family of WD-repeat proteins. *Nature (London)* 371:297–300
 102. Lupas A, van Dyke M, Stock J (1991) Predicting coiled coils from protein sequences. *Science* 252:1162–1164
 103. Lupas AN, Lupas JM, Stock JB (1992) Do G protein subunits associate via a three-stranded coiled coil? *FEBS Lett* 314:105–108
 104. Garritsen A, van Galen PJM, Simonds WF (1993) The N-terminal coiled-coil domain of β is essential for γ association: a model for G protein $\beta\gamma$ subunit interaction. *Proc Natl Acad Sci USA* 90:7706–7710
 105. Simonds WF, Manji HK, Garritsen A (1993) G proteins and β ARK: a new twist for the coiled coil. *Trends J Biochem Sci* 18:315–317
 106. Bubis J, Khorana HG (1990) Sites of interaction in the complex between β - and γ -subunits of transducin. *J Biol Chem* 265:12995–12999
 107. Pronin AN, Gautam N (1992) Interaction between G protein β and γ subunit types is selective. *Proc Natl Acad Sci USA* 89:6220–6224

108. Higgins JB, Casey PJ (1994) In vitro processing of recombinant G protein γ subunits. *J Biol Chem* 269:9067–9073
109. Fukada Y, Matsuda T, Kokame K, Takao T, Shimonishi Y, Akino T, Yoshizawa T (1994) Effects of carboxyl methylation of photoreceptor G protein γ -subunit in visual transduction. *J Biol Chem* 269:5163–5170
110. Dietrich A, Meister M, Brazil D, Camps M, Gierschik P (1994) Stimulation of phospholipase C- β 2 by recombinant guanine-nucleotide-binding protein $\beta\gamma$ -dimers produced in a baculovirus/insect cell expression system. Requirement of γ -subunit isoprenylation for stimulation of phospholipase C. *Eur J Biochem* 219:171–178
111. Yamada M, Jahangir A, Hosoya Y, Inanobe A, Katada T, Kurachi Y (1994) G_K^* and brain $G_{\beta\gamma}$ activate muscarinic K^+ channel through the same mechanism. *J Biol Chem* 268:24551–24554
112. Thomason PA, James SR, Casey PJ, Downes CP (1994) A G protein $\beta\gamma$ -subunit-responsive phosphoinositide 3-kinase activity in human platelet cytosol. *J Biol Chem* 269:16525–16528
113. Lefkowitz RJ (1993) G protein-coupled receptor kinases. *Cell* 74:409–412
114. Crespo P, Xu N, Simonds WF, Gutkind JS (1994) Ras-dependent activation of MAP kinase pathway mediated by G protein $\beta\gamma$ subunits. *Nature (London)* 369:418–420
115. Faure M, Voyno-Yasenetskaya TA, Bourne HR (1994) cAMP and $\beta\gamma$ subunits of heterotrimeric G proteins stimulate the mitogen-activated protein kinase pathway in COS-7 cells. *J Biol Chem* 269:7851–7854
116. Musacchio A, Gibson T, Rice P, Thompson J, Saraste M (1993) The PH domain: a common piece in the structural patchwork of signalling proteins. *Trends Biochem Sci* 18:343–348
117. Gibson T, Hyvönen M, Musacchio A, Saraste M, Birney E (1994) PH domain: the first anniversary. *Trends Biochem Sci* 19:349–353
118. Parker PJ, Hemmings BA, Gierschik P (1994) PH domains and phospholipases – a meaningful relationship? *Trends Biochem Sci* 18:343–348
119. Touhara K, Inglese J, Pitcher JA, Shaw G, Lefkowitz RJ (1994) Binding of G protein $\beta\gamma$ -subunits to pleckstrin homology domains. *J Biol Chem* 269:10217–10220
120. Ismailov II, McDuffie JH, Benos DJ (1994) Protein kinase A phosphorylation and G protein regulation of purified renal Na^+ channels in planar bilayer membranes. *J Biol Chem* 269:10235–10241
121. Bubien JK, Jope RS, Warnock DG (1994) G proteins modulate amiloride-sensitive sodium channels. *J Biol Chem* 269:17780–17783
122. Bokoch GM, Gilman AG (1984) Inhibition of receptor-mediated release of arachidonic acid by pertussis toxin. *Cell* 39:301–308
123. Okajima F, Ui M (1984) ADP-ribosylation of the specific membrane proteins by islet-activating protein, pertussis toxin, associated with inhibition of a chemotactic peptide-induced arachidonate release in neutrophils. *J Biol Chem* 259:13863–13871
124. Exton JH (1990) Signaling through phosphatidylcholine breakdown. *J Biol Chem* 265:1–4
125. Fasolato C, Innocenti B, Pozzan T (1994) Receptor-activated Ca^{2+} influx: how many mechanisms for how many channels? *Trends Pharmacol. Sci.* 15:77–83