

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/261185183>

# Synthesis of substituted pyrimidines as corticotropin releasing factor (CRF) receptor ligands

ARTICLE *in* EUROPEAN JOURNAL OF MEDICINAL CHEMISTRY · MARCH 2014

Impact Factor: 3.45 · DOI: 10.1016/j.ejmech.2014.03.040 · Source: PubMed

---

CITATIONS

3

---

READS

75

## 4 AUTHORS, INCLUDING:



**Bhimanna Kuppast**

South Dakota State University

10 PUBLICATIONS 14 CITATIONS

SEE PROFILE



**George Liapakis**

University of Crete

53 PUBLICATIONS 1,862 CITATIONS

SEE PROFILE



## Original article

## Synthesis of substituted pyrimidines as corticotropin releasing factor (CRF) receptor ligands

Bhimanna Kuppast<sup>a</sup>, Katerina Spyridaki<sup>b</sup>, George Liapakis<sup>b</sup>, Hesham Fahmy<sup>a,\*</sup><sup>a</sup> Department of Pharmaceutical Sciences, College of Pharmacy, South Dakota State University, Brookings, SD 57007, USA<sup>b</sup> Department of Pharmacology, Faculty of Medicine, University of Crete, Voutes, Heraklion 71003, Crete, Greece

## ARTICLE INFO

## Article history:

Received 17 September 2013

Received in revised form

11 March 2014

Accepted 13 March 2014

Available online 15 March 2014

## Keywords:

Pyrimidines

Thiazolo[4,5-d]pyrimidines

Corticotropin releasing factor receptor ligands

CRF

## ABSTRACT

Corticotropin releasing factor (CRF) is a neuropeptide hormone produced from the hypothalamus that controls the secretion of corticotropin (ACTH) from the anterior pituitary gland that, in turn, prompts the adrenal glands to secrete glucocorticoids. This involvement in the hypothalamic–pituitary–adrenal axis (HPA) in response to stress and also playing a key role in behavioral, cardiovascular, immune and gastrointestinal systems made CRF binding to its receptors an important target in drug discovery aiming to develop lead compounds with the potential to treat various stress-related disorders including depression, anxiety and addictive disorders. Several non-peptide CRF<sub>1</sub> receptor antagonists were developed by pharmaceutical companies and are currently in clinical trials with the aim of improving the health consequences of chronic stress and for use in the clinical management of anxiety and stress. Many showed promising results not only in treatment of anxiety and depression but also in treatment of CRF-induced hypertension, as well as in treatment of arthritis, irritable bowel syndrome and peptic ulcers. In this manuscript, we describe the synthesis of substituted pyrimidines with close structural similarities to reported lead compounds with promising CRF<sub>1</sub> receptor affinities and carrying groups known to be associated with optimum affinity to CRF<sub>1</sub> receptors. The affinity of the newly prepared compounds in comparison to antalarmin, a potent CRF<sub>1</sub> receptor antagonist in clinical trials as a standard, is also described. Four compounds from the new series showed promising CRF<sub>1</sub> receptor affinity.

© 2014 Elsevier Masson SAS. All rights reserved.

## 1. Introduction

Corticotropin releasing factor (CRF), also called corticotropin releasing hormone (CRH), is a hypothalamic hormone that plays a key role in the maintenance of homeostasis. CRF is secreted by the paraventricular nucleus of the hypothalamus in response to stress and is transported via the portal vein to the anterior lobe of the pituitary gland where it causes the release of corticotropin (also called adrenocorticotrophic hormone ACTH). The released ACTH is then transported by the blood to the adrenal glands where it stimulates the release of endogenous glucocorticoids [1,2]. In addition to the regulation of the hypothalamus–pituitary–adrenal axis (HPA), CRF plays an important role in stress as well as in many physiological and pathological processes such as the cardiovascular, gastrointestinal, behavioral, immune and reproductive systems [3–17].

CRF is a 41 amino acid neuropeptide and belongs to a family of structurally related highly homologous peptides from several species such as rat h/rCRF, human h/rCRF, goat, cow, pig and xenopus CRF [1,18–22]. In addition to the previously mentioned CRF peptides, the CRF family also includes peptides from various species such as sauvagine, urotensin and urocortins. These peptides are closely related to CRF and are considered CRF-like peptides. Sauvagine (SVG) and urotensin (URO) have been isolated from the frog *Phyllomedusa sauvagei* and the sucker fish *Catostomus commersoni*, respectively, whereas urocortin (Ucn), urocortin II (UcnII) and urocortin III (UcnIII) have been isolated from mammals. These are natural ligands that bind to and activate CRF receptors [23–27].

CRF and its analogs exert their actions by interacting with two types of plasma membrane receptors, CRF<sub>1</sub> and CRF<sub>2</sub>. Both CRF receptor types belong to the superfamily of G protein-coupled receptors (GPCR) [28–30]. Several receptor sub-types exist for CRF<sub>1</sub> and CRF<sub>2</sub> [31]. The main functional receptor subtype of CRF<sub>1</sub> is CRF<sub>1α</sub> that mediates most of the action of CRF family peptides. CRF<sub>1α</sub> will be mentioned in this manuscript as CRF<sub>1</sub> for simplicity.

\* Corresponding author.

E-mail address: [Hesham.Fahmy@sdstate.edu](mailto:Hesham.Fahmy@sdstate.edu) (H. Fahmy).

Because of important role and involvement of CRF and its receptors in CNS, behavioral, cardiovascular, gastrointestinal, immune and reproductive systems, several non-peptide CRF antagonists were developed as new leads in drug discovery to treat various stress-related disorders including depression, anxiety [32,33] and addictive disorders [34–36]. Most non-peptide CRF antagonists discovered to date are substituted five or six membered heterocyclic or bicyclic rings [37,38]. These non-peptide antagonists bind selectively to CRF<sub>1</sub> and offer advantages over the peptide congeners in terms of stability, ease of preparation, ease of structural modification to enhance the pharmacokinetic profile and better blood–brain barrier penetration.

Several non-peptide CRF receptor antagonists were developed by pharmaceutical companies such as Antalarmin, CP-154526, DMP-904, DMP-696 and NBI-27914 (Fig. 1) and are currently in clinical trials with the aim of improving the health consequences of chronic stress and for use in the clinical management of anxiety and stress [38]. Antalarmin showed promising results in treatment of CRF-induced hypertension [39–42]. Promising results for antalarmin and other CRF<sub>1</sub> antagonists were also observed in the area of drug addiction disorders. Antalarmin also showed anti-inflammatory effects and has been suggested as having potential uses in treatment of arthritis [43], irritable bowel syndrome [44,45] and peptic ulcers [46].

In this manuscript we describe the synthesis and characterization of several substituted pyrimidines with close structural similarities to NBI-27914, a prominent CRF receptor antagonist with excellent binding affinity. The binding affinities of the synthesized compounds were evaluated in comparison to antalarmin as a standard. Four of the newly synthesized compounds showed promising binding affinities and were selected as lead compounds for further structural modifications and for in vivo testing in animal models of anxiety.

## 2. Results and discussion

### 2.1. Chemistry

The general synthetic scheme for synthesis of the target compounds (1–28) is described in Fig. 2. The selected substituted phenyl isothiocyanate was treated with cynoacetamide and sulfur in presence of triethylamine following a Gewald type of reaction to give intermediate 4-amino-5-carboxamido-3-substituted phenylthiazol-2-thiones (I a,b) in excellent yields according to reported methods [47–51]. The thiazoles were then treated with acetic anhydride at reflux temperature to yield 6-methylthiazolo[4,5-d]pyrimidin-6-ones (II a,b) [50] or with a 1:1 mixture of triethyl orthoformate and acetic anhydride to yield thiazolo[4,5-d]pyrimidin-6-one unsubstituted at position 6 (II c). It is worth mentioning that the selected isothiocyanates used were 2,4,6-trimethylphenylisothiocyanate and 2-bromo-4-isopropylphenyl isothiocyanate since previous studies have shown that this

particular substitutions at that particular position is required for optimum CRF<sub>1</sub> binding antagonist activity [37,49].

The thiazolo[4,5-d]pyrimidin-6-ones (II a–c) were then subjected to thiazole ring opening using 4 N sodium hydroxide solution at reflux temperature in presence of positive pressure of nitrogen or argon gas. If the reaction is carried out in presence of air, oxidation reaction takes place where two molecules of the resulting pyrimidin-6-one-5-thiols are oxidized to form a dimer linked through an S–S bond [52].

Alkylation of the thiol group was spontaneously carried out in the same reaction mixture using methyl iodide or ethyl iodide in 1,4-dioxane, in presence of nitrogen or argon gas to give 5-alkylthio-pyrimidin-6-ones (III a–f) in excellent yields. Chlorination of III a–f with phosphorous oxychloride yielded the 6-chloro pyrimidine derivatives (IV a–f) in excellent yields. Because the chloro compounds (IV a–f) have electron donating groups (methyl, aminoaryl and thioalkyl at positions 2, 4 and 5), they are less reactive towards nucleophilic substitution reaction of the 6-chloro group with secondary amines. Thus synthesis of compounds (1–28) was carried out by a modification of the method reported by Beletskaya [53]. The chloro derivatives (IV a–f) were treated with secondary amines using tris(dibenzylideneacetone) dipalladium(0), Xphos and cesium carbonate, with the secondary amines as both nucleophile and solvent. Five secondary amines were selected for this reaction: *N,N*-diethylamine, *N,N*-di(*n*-propyl)amine, *N*-ethyl, *N*-(*n*-butyl)amine, *N,N*-bis(2-methoxyethyl)amine and *N*-[2-(cyclopropyl)ethyl], *N*-(*n*-propyl)amine. This selection was based on previous reports showing an optimum CRF<sub>1</sub> receptor antagonist activity for those particular amino groups in the target compounds [49].

The structures of the all intermediates were confirmed using <sup>1</sup>H NMR and Mass spectroscopy and the structures of the compounds (1–28) were elucidated using <sup>1</sup>H NMR and <sup>13</sup>C NMR and confirmed with Mass spectroscopy and elemental analysis. The detailed structural variations of the individual compounds (1–28) are shown in Table 1.

### 2.2. Receptor binding study

The series of the synthesized pyrimidines (1–28) were evaluated for their binding affinity to CRF<sub>1</sub> receptors. First, the ability of compounds (1–28) at a single concentration of 500 nM to inhibit the specific binding of [<sup>125</sup>I]-Tyr<sup>0</sup>sauvagine to membranes from HEK 293 cells stably expressing the CRF<sub>1</sub> receptor in binding experiments performed under equilibrium conditions was evaluated. In this initial screening experiment, compounds showing low affinity (1, 2, 4, 5, 7, 10–28) as measured by their inability to inhibit close to or more than 50% of [<sup>125</sup>I]-Tyr<sup>0</sup>-sauvagine specific binding (Fig. 3) were excluded from further pharmacological evaluation. The other compounds were further pharmacologically characterized by determining their binding affinities (IC<sub>50</sub>) for CRF<sub>1</sub> receptors in competition experiments performed under equilibrium




Fig. 1. Structures of CRF<sub>1</sub> receptor antagonists.




Fig. 2. Synthetic scheme of the compounds (1–28).

conditions in membranes from HEK 293 cells stably expressing the CRF<sub>1</sub> receptor. Four lead compounds (**3**, **6**, **8** and **9**) were identified (Fig. 3). These lead compounds were found to bind to CRF<sub>1</sub> receptors in a concentration dependent manner (Fig. 4) and with promising binding affinities (Table 2).

### 2.3. SAR study

Four compounds showed promising CRF receptor binding affinities and were identified as lead compounds (**3**, **6**, **8** and **9**). All the lead compounds have a methyl group at position 2. Compounds having a 2,4,6-trimethylphenyl group at –position 4 showed better binding affinities than compounds having a 2-bromo-4-isopropylphenyl group. Both the methyl and ethyl groups of the alkylthiol group at position 5 showed promising affinity. At position 6, only the *N,N*-bis(methoxyethyl)amino, the *N*-cyclopropyl-*N*-methylpropylamino and the *N,N*-di-(*n*-propyl)amino displayed promising CRF receptor binding affinity.

### 3. Conclusion

A series of novel substituted pyrimidines were synthesized as CRF<sub>1</sub> receptor ligands. The newly synthesized compounds have the groups known for optimum binding affinity for CRF<sub>1</sub> receptors, particularly a methyl group at position 2, a substituted phenyl at position 4 and dialkylamino group at position 6 of the pyrimidine ring while having different thioalkyl groups at position 5. The synthesized compounds were evaluated for their binding affinity for CRF<sub>1</sub> receptor in comparison with antalarmin, a known non-peptide CRF<sub>1</sub> receptor antagonist, as a standard. Four lead compounds of the new series (**3**, **6**, **8** and **9**) were identified. These four

compounds showed a promising CRF<sub>1</sub> receptor binding affinity. These compounds showed receptor binding affinity with (–log IC<sub>50</sub>) values of 6.23–6.61 compared to 7.73 for antalarmin. This indicates that further structural modifications of these compounds may lead to compounds with superior CRF<sub>1</sub> receptor binding affinity and a potential for development as new anti-anxiety and antidepressant agents.


## 4. Experimental

### 4.1. Chemistry

#### 4.1.1. General information


All chemicals were purchased from commercial sources. The substituted phenyl isothiocyanates were purchased from Oakwood Products, Inc. SC, USA. Tris(dibenzylideneacetone)dipalladium(0), (±)-2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl, Copper(I) iodide, and 2-(dicyclohexylphosphino) 2',4',6'-triisopropylbiphenyl were purchased from Acros organics, NJ, USA. Palladium (II) acetate was purchased from Tokyo chemical industry, Japan. cesium carbonate was purchased from Alfa Aesar, MA. All chemical were stored in desiccator. All laboratory grade, secondary amines and dry solvents were purchased from Sigma Aldrich USA, Acros organics, Fisher scientific USA and used without further purification. Flash column chromatography separation was performed using Acros organics silica gel 40–60 μm, 60 Å using combination of ethyl acetate and hexanes. Whatman TLC plates were used for thin layer chromatography and visualization was done using UV fluorescence at 254 nm. Melting points were recorded on a Mel-Temp, Laboratory devices, Inc and are uncorrected. %CHN Analyzer by combustion/TCD and %S by O flask combustion/IC were used for elemental

**Table 1**  
5-thioalkyl-6-*N,N*-diaminoalkyl-4-substituted phenylaminopyrimidines (**1–28**).




No	R	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	X
1	CH <sub>3</sub>	CH <sub>3</sub>	C <sub>3</sub> H <sub>7</sub>	C <sub>3</sub> H <sub>7</sub>	2,4,6-triCH <sub>3</sub>
2	CH <sub>3</sub>	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	2,4,6-triCH <sub>3</sub>
3	CH <sub>3</sub>	CH <sub>3</sub>	C <sub>2</sub> H <sub>4</sub> OCH <sub>3</sub>	C <sub>2</sub> H <sub>4</sub> OCH <sub>3</sub>	2,4,6-triCH <sub>3</sub>
4	CH <sub>3</sub>	CH <sub>3</sub>	<i>c</i> -propylCH <sub>2</sub>	C <sub>3</sub> H <sub>7</sub>	2,4,6-triCH <sub>3</sub>
5	CH <sub>3</sub>	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>4</sub> H <sub>9</sub>	2,4,6-triCH <sub>3</sub>
6	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>3</sub> H <sub>7</sub>	C <sub>3</sub> H <sub>7</sub>	2,4,6-triCH <sub>3</sub>
7	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	2,4,6-triCH <sub>3</sub>
8	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>4</sub> OCH <sub>3</sub>	C <sub>2</sub> H <sub>4</sub> OCH <sub>3</sub>	2,4,6-triCH <sub>3</sub>
9	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	<i>c</i> -propylCH <sub>2</sub>	C <sub>3</sub> H <sub>7</sub>	2,4,6-triCH <sub>3</sub>
10	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>4</sub> H <sub>9</sub>	2,4,6-triCH <sub>3</sub>
11	CH <sub>3</sub>	CH <sub>3</sub>	C <sub>3</sub> H <sub>7</sub>	C <sub>3</sub> H <sub>7</sub>	2-Br-4-CH(CH <sub>3</sub> ) <sub>2</sub>
12	CH <sub>3</sub>	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	2-Br-4-CH(CH <sub>3</sub> ) <sub>2</sub>
13	CH <sub>3</sub>	CH <sub>3</sub>	C <sub>2</sub> H <sub>4</sub> OCH <sub>3</sub>	C <sub>2</sub> H <sub>4</sub> OCH <sub>3</sub>	2-Br-4-CH(CH <sub>3</sub> ) <sub>2</sub>
14	CH <sub>3</sub>	CH <sub>3</sub>	<i>c</i> -propylCH <sub>2</sub>	C <sub>3</sub> H <sub>7</sub>	2-Br-4-CH(CH <sub>3</sub> ) <sub>2</sub>
15	CH <sub>3</sub>	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>4</sub> H <sub>9</sub>	2-Br-4-CH(CH <sub>3</sub> ) <sub>2</sub>
16	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>3</sub> H <sub>7</sub>	C <sub>3</sub> H <sub>7</sub>	2-Br-4-CH(CH <sub>3</sub> ) <sub>2</sub>
17	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	2-Br-4-CH(CH <sub>3</sub> ) <sub>2</sub>
18	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>4</sub> OCH <sub>3</sub>	C <sub>2</sub> H <sub>4</sub> OCH <sub>3</sub>	2-Br-4-CH(CH <sub>3</sub> ) <sub>2</sub>
19	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	<i>c</i> -propylCH <sub>2</sub>	C <sub>3</sub> H <sub>7</sub>	2-Br-4-CH(CH <sub>3</sub> ) <sub>2</sub>
20	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>4</sub> H <sub>9</sub>	2-Br-4-CH(CH <sub>3</sub> ) <sub>2</sub>
21	H	CH <sub>3</sub>	C <sub>3</sub> H <sub>7</sub>	C <sub>3</sub> H <sub>7</sub>	2,4,6-triCH <sub>3</sub>
22	H	CH <sub>3</sub>	C <sub>2</sub> H <sub>4</sub> OCH <sub>3</sub>	C <sub>2</sub> H <sub>4</sub> OCH <sub>3</sub>	2,4,6-triCH <sub>3</sub>
23	H	CH <sub>3</sub>	<i>c</i> -propylCH <sub>2</sub>	C <sub>3</sub> H <sub>7</sub>	2,4,6-triCH <sub>3</sub>
24	H	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>4</sub> H <sub>9</sub>	2,4,6-triCH <sub>3</sub>
25	H	C <sub>2</sub> H <sub>5</sub>	C <sub>3</sub> H <sub>7</sub>	C <sub>3</sub> H <sub>7</sub>	2,4,6-triCH <sub>3</sub>
26	H	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	2,4,6-triCH <sub>3</sub>
27	H	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>4</sub> OCH <sub>3</sub>	C <sub>2</sub> H <sub>4</sub> OCH <sub>3</sub>	2,4,6-triCH <sub>3</sub>
28	H	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>4</sub> H <sub>9</sub>	2,4,6-triCH <sub>3</sub>

analysis of final compounds and performed by Micro Analysis Inc., Wilmington DE, USA and are within  $\pm 0.4\%$ .  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were obtained on a Bruker Avance 400 MHz instrument using  $\text{CDCl}_3$  as solvent unless otherwise stated.  $^1\text{H}$  NMR Spectra are



**Fig. 3.** Screening of compounds for binding to human CRF<sub>1</sub> receptor. Inhibition of [ $^{125}\text{I}$ ]-Tyr<sup>0</sup>-sauvagine specific binding by 500 nM of compounds **1–28** on membranes from HEK 293 cells stably expressing the human CRF<sub>1</sub> receptor. The bars represent the % inhibition of radioligand specific binding by the compounds, determined from 2 to 5 experiments (with their means and S.E.). The dashed line represents the 50% inhibition of radioligand specific binding.



**Fig. 4.** Competition binding isotherms of compounds **3**, **6**, **8** and **9** to human CRF<sub>1</sub> receptor. Competition of [ $^{125}\text{I}$ ]-Tyr<sup>0</sup>-sauvagine specific binding by increasing concentrations of the 4 lead compounds on membranes from HEK 293 cells stably expressing the human CRF<sub>1</sub> receptor. The means and S.E. (duplicate determination) are shown and the data were fit to a one-site competition model by nonlinear regression.

reported in order; multiplicity, number of protons and signals were characterized as s (singlet), dd (doublet of doublet), t (triplet), m (multiplet), br s (broad signal), q (quartet), quin (quintet), tquin (triplet of quintet), sxt (sextet), spt (septet). Chemical shifts are relative to TMS as an internal standard. Mass spectra were recorded on Finnigan LCQTM DECA by Thermo Quest San Jose, CA, using electrospray ionization method. All the reactions were carried out in flame dried glassware under an atmosphere of nitrogen unless otherwise stated.

#### 4.1.2. 4-Amino-5-carboxamido-3-(substituted phenyl)-2,3-dihydrothiazole-2-thiones (**1a,b**) [50]

To a stirred solution of triethylamine (1.0 mmol) and cyn-oacetamide (1.0 mmol) in dimethyl sulfoxide (2.5 mL), sulfur was added (1.0 mmol) and the reaction mixture was stirred for 10 min at ambient temperature, followed by addition of the selected substituted phenyl isothiocyanate (1.0 mmol). The reaction mixture was stirred at ambient temperature for 4 h and completion of the reaction was monitored by TLC using 9:1 dichloromethane and methanol. The reaction mixture was then diluted with water and extracted with ethyl acetate. The combined organic layer was washed with water and brine solution and dried over anhydrous sodium sulfate and concentrated under vacuum to give the product as a solid residue **1a** and **1b**.

**4.1.2.1. 4-amino-3-mesityl-2-thioxo-2,3-dihydrothiazole-5-carboxamide (**1a**).** Yield: yellow solid (73%); MP: 203 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.08 (s, 2H), 5.85 (br. s., 2H), 5.10 (br. s., 2H), 2.37 (s, 3H), 2.11 (s, 6H); ESI-MS  $m/z$ : 294.95 ( $\text{MH}^+$ ).

**Table 2**

The IC<sub>50</sub> values of the lead compounds compared to antalarmin.

Compound	–LogIC <sub>50</sub> $\pm$ S.E
Antalarmin	7.73 $\pm$ 0.09
Compound <b>3</b>	6.56 $\pm$ 0.12
Compound <b>6</b>	6.27 $\pm$ 0.26
Compound <b>8</b>	6.23 $\pm$ 0.06
Compound <b>9</b>	6.61 $\pm$ 0.06



4.1.2.2. 4-Amino-3-(2-bromo-4-isopropylphenyl)-2-thioxo-2,3-dihydrothiazole-5-carboxamide (**I b**). Yield: yellow solid (71%); MP: 206 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.68 (d,  $J = 1.8$  Hz, 1H), 7.42 (dd,  $J = 1.8, 8.2$  Hz, 1H), 7.30 (d,  $J = 8.2$  Hz, 1H), 5.86 (br. s., 2H), 5.09 (br. s., 2H), 3.01 (spt,  $J = 7.1$  Hz, 1H), 1.32 (d,  $J = 7.1$  Hz, 6H); ESI–MS  $m/z$ : 371.94 ( $\text{MH}^+$ ), 373.93 ( $\text{MH} + 2^+$ ).

#### 4.1.3. 5-Substituted-2-thioxo-3-(substitutedphenyl)-2,3-dihydrothiazolo[4,5-d]pyrimidin-7(6H)-one (**II a–c**) [50]

A mixture of **I a,b** (1.0 mmol) in acetic anhydride (for **II a,b**) (3 mL) or 1:1 mixture of acetic anhydride and triethyl orthoformate (for **II c**) (3 mL) was stirred at reflux temperature for 3–4 h and completion of the reaction was monitored by TLC using 9:1 dichloromethane and methanol. The reaction mixture was cooled to room temperature. The separated product was filtered and washed with ethanol and dried to give **II a–c**.

4.1.3.1. 3-Mesityl-5-methyl-2-thioxo-2,3-dihydrothiazolo[4,5-d]pyrimidin-7(6H)-one (**II a**). Yield: brown crystals (69.7%); MP: 340 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  13.13 (br. s., 1H), 7.06 (s, 2H), 2.49 (s, 3H), 2.39 (s, 3H), 2.02 (s, 6H); ESI–MS  $m/z$ : 318.17 ( $\text{MH}^+$ ).

4.1.3.2. 3-(2-Bromo-4-isopropylphenyl)-5-methyl-2-thioxo-2,3-dihydrothiazolo[4,5-d]pyrimidin-7(6H)-one (**II b**). Yield: light yellow solid (70.4%); MP: 237 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  12.94 (br. s., 1H), 7.65 (d,  $J = 1.8$  Hz, 1H), 7.39 (dd,  $J = 1.8, 8.2$  Hz, 1H), 7.25 (d,  $J = 8.2$  Hz, 1H), 3.02 (spt,  $J = 7.1$  Hz, 1H), 2.50 (s, 3H), 1.34 (d,  $J = 7.1$  Hz, 6H); ESI–MS  $m/z$ : 396.08 ( $\text{MH}^+$ ), 398.06 ( $\text{MH} + 2^+$ ).

4.1.3.3. 3-Mesityl-2-thioxo-2,3-dihydrothiazolo[4,5-d]pyrimidin-7(6H)-one (**II c**). Yield: brown color solid (84%); MP: 315 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  13.07 (br. s., 1H), 8.08 (s, 1H), 7.09 (s, 2H), 2.38 (s, 3H), 2.04 (s, 6H); ESI–MS  $m/z$ : 304.14 ( $\text{MH}^+$ ).

#### 4.1.4. 6-(Substituted phenyl)-2-methyl-5-(alkylthio)pyrimidin-4(3H)-ones (**III a–f**)

A mixture of **II a–c** (1.0 mmol) in 4 N NaOH (10 mL) was stirred at 100 °C for 3.5–4 h under the positive pressure of nitrogen. The reaction mixture was cooled to ambient temperature, and a solution of the selected alkyl iodide (1.5 mmol) in 1,4-dioxane (2 mL) was added slowly at ambient temperature and stirred for 2 h under positive atmosphere of nitrogen. The reaction mixture was cooled to 0 °C, and pH was adjusted to 4 pH using conc. HCl. The precipitated solid was filtered and washed with ice cold water and dried to give a crude residue. The pure compound was isolated by flash chromatography using gradient ethyl acetate in hexanes to give **III a–f**.

4.1.4.1. 6-(Mesitylamino)-2-methyl-5-(methylthio)pyrimidin-4(3H)-one (**III a**). Yield: whitish solid (87%); MP: 335 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  12.54 (br. s., 1H), 7.48 (s, 1H), 6.93 (s, 2H), 2.31 (s, 6H), 2.29 (s, 3H), 2.17 (s, 3H), 2.14 (s, 3H); ESI–MS  $m/z$ : 290.12 ( $\text{MH}^+$ ).

4.1.4.2. 5-(Ethylthio)-6-(mesitylamino)-2-methylpyrimidin-4(3H)-one (**III b**). Yield: whitish solid (87%); MP: 244 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  13.03 (br. s., 1H), 7.49 (s, 1H), 6.93 (s, 2H), 2.82 (q,  $J = 7.4$  Hz, 2H), 2.34 (s, 3H), 2.29 (s, 3H), 2.19 (s, 6H), 1.30 (t,  $J = 7.4$  Hz, 3H); ESI–MS  $m/z$ : 304.07 ( $\text{MH}^+$ ).

4.1.4.3. 6-((2-Bromo-4-isopropylphenyl)amino)-2-methyl-5-(methylthio)pyrimidin-4(3H)-one (**III c**). Yield: light yellow solid (87%); MP: 238 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  13.05 (br. s., 1H), 8.64 (s, 1H), 8.21 (d,  $J = 8.3$  Hz, 1H), 7.41 (d,  $J = 2.0$  Hz, 1H), 7.21 (dd,  $J = 2.0,$

8.3 Hz, 1H), 2.92 (spt,  $J = 6.9$  Hz, 1H), 2.55 (s, 3H), 2.40 (s, 3H), 1.24 (d,  $J = 6.9$  Hz, 6H); ESI–MS  $m/z$ : 368.1 ( $\text{MH}^+$ ), 370.07 ( $\text{MH} + 2^+$ ).

4.1.4.4. 6-((2-Bromo-4-isopropylphenyl)amino)-5-(ethylthio)-2-methylpyrimidin-4(3H)-one (**III d**). Yield: light yellow solid (87%); MP: 207 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  13.07 (br. s., 1H), 8.68 (s, 1H), 8.19 (d,  $J = 8.3$  Hz, 1H), 7.44 (d,  $J = 1.8$  Hz, 1H), 7.19 (dd,  $J = 1.8, 8.3$  Hz, 1H), 2.95 (spt,  $J = 6.8$  Hz, 1H), 2.76 (q,  $J = 7.4$  Hz, 2H), 2.49 (s, 3H), 1.30 (t,  $J = 7.4$  Hz, 3H), 1.25 (d,  $J = 6.8$  Hz, 6H); ESI–MS  $m/z$ : 382.07 ( $\text{MH}^+$ ), 384.07 ( $\text{MH} + 2^+$ ).

4.1.4.5. 6-(Mesitylamino)-5-(methylthio)pyrimidin-4(3H)-one (**III e**). Yield: white solid (90%); MP: 247 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  13.52 (br. s., 1H), 7.95 (s, 1H), 7.52 (s, 1H), 6.95 (s, 2H), 2.34 (s, 3H), 2.30 (s, 3H), 2.17 (s, 6H); ESI–MS  $m/z$ : 276.25 ( $\text{MH}^+$ ).

4.1.4.6. 5-(Ethylthio)-6-(mesitylamino)pyrimidin-4(3H)-one (**III f**). Yield: white solid (90%); MP: 218 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  13.51 (br. s., 1H), 7.92 (s, 1H), 7.50 (s, 1H), 6.92 (s, 2H), 2.73 (q,  $J = 7.3$  Hz, 2H), 2.29 (s, 3H), 2.19 (s, 6H), 1.24 (t,  $J = 7.3$  Hz, 3H); ESI–MS  $m/z$ : 290.14 ( $\text{MH}^+$ ).

#### 4.1.5. 6-Chloro-2-substituted-4-(substitutedphenylamino)-5-alkylthiopyrimidines (**IV a–f**)

A mixture of **III a–f** (1.0 mmol) and phosphorous oxychloride (2.5 mL) were stirred at reflux temperature for 3–4 h and completion of the reaction was monitored by TLC using ethyl acetate and hexanes mixture. The reaction mixture was cooled to room temperature and carefully diluted with ice cold water, extracted with ethyl acetate. The combined organic layer was washed with water, brine and dried over anhydrous sodium sulfate and concentrated under vacuum to get residue. The pure compound was isolated by flash chromatography using gradient ethyl acetate in hexanes to give **IV a–f**.

4.1.5.1. 6-Chloro-*N*-mesityl-2-methyl-5-(methylthio)pyrimidin-4-amine (**IV a**). Yield: off white solid (78%); MP: 114 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.66 (br. s., 1H), 6.95 (s, 2H), 2.39 (s, 3H), 2.38 (s, 3H), 2.32 (s, 3H), 2.16 (s, 6H); ESI–MS  $m/z$ : 308.17 ( $\text{MH}^+$ ), 310.15 ( $\text{MH} + 2^+$ ).

4.1.5.2. 6-Chloro-5-(ethylthio)-*N*-mesityl-2-methylpyrimidin-4-amine (**IV b**). Yield: off white solid (81%); MP: 77 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.66 (br. s., 1H), 6.95 (s, 2H), 2.90 (q,  $J = 7.4$  Hz, 2H), 2.38 (s, 3H), 2.32 (s, 3H), 2.16 (s, 6H), 1.34 (t,  $J = 7.4$  Hz, 3H); ESI–MS  $m/z$ : 322.13 ( $\text{MH}^+$ ), 324.14 ( $\text{MH} + 2^+$ ).

4.1.5.3. *N*-(2-Bromo-4-isopropylphenyl)-6-chloro-2-methyl-5-(methylthio)pyrimidin-4-amine (**IV c**). Yield: off white solid (81%); MP: 94 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.91 (s, 1H), 8.44 (d,  $J = 8.6$  Hz, 1H), 7.46 (d,  $J = 2.0$  Hz, 1H), 7.24 (dd,  $J = 2.0, 8.6$  Hz, 1H), 2.90 (spt,  $J = 7.1$  Hz, 1H), 2.57 (s, 3H), 2.40 (s, 3H), 1.26 (d,  $J = 7.1$  Hz, 6H); ESI–MS  $m/z$ : 386.04 ( $\text{MH}^+$ ), 388.03 ( $\text{MH} + 2^+$ ), 390.03 ( $\text{MH} + 4^+$ ).

4.1.5.4. *N*-(2-bromo-4-isopropylphenyl)-6-chloro-5-(ethylthio)-2-methylpyrimidin-4-amine (**IV d**). Yield: off white solid (65%); MP: 65 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.94 (s, 1H), 8.48 (d,  $J = 8.6$  Hz, 1H), 7.46 (d,  $J = 2.0$  Hz, 1H), 7.24 (dd,  $J = 2.0, 8.6$  Hz, 1H), 2.95 (spt,  $J = 6.8$  Hz, 1H), 2.84 (q,  $J = 7.4$  Hz, 2H), 2.53 (s, 3H), 1.32 (t,  $J = 7.4$  Hz, 3H), 1.26 (d,  $J = 6.8$  Hz, 6H); ESI–MS  $m/z$ : 400.15 ( $\text{MH}^+$ ), 402.11 ( $\text{MH} + 2^+$ ), 404.08 ( $\text{MH} + 4^+$ ).

4.1.5.5. 6-Chloro-*N*-mesityl-5-(methylthio)pyrimidin-4-amine (**IV e**). Yield: off white solid (70%); MP: 194 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.26 (s, 1H), 7.69 (br. s., 1H), 6.97 (s, 2H), 2.43 (s, 3H), 2.31 (s, 3H), 2.16 (s, 6H); ESI–MS  $m/z$ : 294.48 ( $\text{MH}^+$ ), 296.44 ( $\text{MH} + 2^+$ ).

**4.1.5.6. 6-Chloro-5-(ethylthio)-N-mesitylpyrimidin-4-amine (IV f).** Yield: off white solid (94%); MP: 115 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.10 (s, 1H), 7.55 (s, 1H), 6.94 (s, 2H), 2.75 (q,  $J = 7.3$  Hz, 2H), 2.29 (s, 3H), 2.18 (s, 6H), 1.23 (t,  $J = 7.3$  Hz, 3H); ESI–MS  $m/z$ : 308.47 ( $\text{MH}^+$ ), 310.56 ( $\text{MH} + 2^+$ ).

**4.1.6. 2-Substituted-6-dialkylamino-4-(substitutedphenylamino)-5-alkylthiopyrimidines (1–28)**

An oven-dried 3 necked flask equipped with a magnetic stir bar and fitted with a condenser and a rubber septum was evacuated and backfilled with nitrogen. This step was repeated two times with a total of three times. The chloro compound (**IV a–f**) (1.0 mmol) and cesium carbonate (1.5 mmol) were added and flask was evacuated and backfilled with nitrogen. *tris*(dibenzylideneacetone)dipalladium(0) (0.1 mmol, 10 mol%) and 2-dicyclohexyl phosphine-2',4',6'-triisopropyl biphenyl (0.1 mmol, 10 mol%) were added and flask was evacuated and backfilled with nitrogen. 10 mL of the selected secondary amine was added to the reaction mixture via a syringe and the flask was evacuated and backfilled with nitrogen for 3 times. The flask was placed in an oil bath and the temperature was maintained at 100 °C under positive pressure of nitrogen and the reaction was monitored by TLC. After the reaction completion, the reaction mixture was cooled to room temperature, filtered through celite and washed with dichloromethane ( $3 \times 10$  mL). The combined dichloromethane layer was washed with 0.1 N HCl solution until the pH was neutral, followed by water ( $3 \times 10$  mL) and brine solution ( $1 \times 10$  mL). The organic layer was dried over anhydrous sodium sulfate and evaporated under reduced pressure to semisolid residue. The residue was purified by flash chromatography with a gradient eluant of 0–10% ethyl acetate in hexanes to give the desired product (**1–28**).

**4.1.6.1.  $N^4$ -mesityl-2-methyl-5-(methylthio)- $N^6$ , $N^6$ -dipropylpyrimidine-4,6-diamine (1).** Yield: semisolid (61%); MP: 206 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.57 (s, 1H), 6.91 (s, 2H), 3.56 (t,  $J = 7.6$  Hz, 4H), 2.29 (s, 3H), 2.20 (s, 3H), 2.19 (s, 3H), 2.18 (s, 6H), 1.67 (sxt,  $J = 7.4$  Hz, 4H), 0.90 (t,  $J = 7.4$  Hz, 6H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  165.4, 163.9, 163.4, 135.5, 134.1, 128.7, 88.9, 52.6, 26.3, 21.5, 21.0, 18.9, 18.4, 11.4; ESI–MS  $m/z$ : 373.38 ( $\text{MH}^+$ ).

**4.1.6.2.  $N^4$ , $N^4$ -diethyl- $N^6$ -mesityl-2-methyl-5-(methylthio)pyrimidine-4,6-diamine (2).** Yield: light yellow solid (55%); MP: 68 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.55 (s, 1H), 6.91 (s, 2H), 3.66 (q,  $J = 6.9$  Hz, 4H), 2.29 (s, 3H), 2.21 (s, 3H), 2.20 (s, 3H), 2.18 (s, 6H), 1.23 (t,  $J = 6.9$  Hz, 6H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  165.5, 163.8, 163.4, 135.5, 134.1, 128.7, 88.9, 44.5, 26.3, 21.0, 18.8, 18.3, 13.6; ESI–MS  $m/z$ : 345.25 ( $\text{MH}^+$ ).

**4.1.6.3.  $N^4$ -mesityl- $N^6$ , $N^6$ -bis(2-methoxyethyl)-2-methyl-5-(methylthio)pyrimidine-4,6-diamine (3).** Yield: off white semisolid (61%); MP: 60 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.59 (s, 1H), 6.92 (s, 2H), 3.93 (t,  $J = 6.0$  Hz, 4H), 3.67 (t,  $J = 6.0$  Hz, 4H), 3.38 (s, 6H), 2.31 (s, 3H), 2.22 (s, 3H), 2.22 (s, 3H), 2.18 (s, 6H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  165.5, 163.7, 163.6, 135.7, 135.5, 133.9, 128.7, 89.2, 71.5, 58.8, 50.8, 26.2, 21.0, 18.8, 18.2; ESI–MS  $m/z$ : 405.26 ( $\text{MH}^+$ ).

**4.1.6.4.  $N^4$ -(cyclopropylmethyl)- $N^6$ -mesityl-2-methyl-5-(methylthio)- $N^4$ -propylpyrimidine-4,6-diamine (4).** Yield: Yellow semisolid (69%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.57 (s, 1H), 6.93 (s, 2H), 3.69 (t,  $J = 7.3$  Hz, 2H), 3.52 (d,  $J = 6.5$  Hz, 2H), 2.32 (s, 3H), 2.24 (s, 3H), 2.23 (s, 3H), 2.20 (s, 6H), 1.71 (sxt,  $J = 7.3$  Hz, 2H), 1.19 (tquin,  $J = 6.8$  Hz, 1H), 0.93 (t,  $J = 7.4$  Hz, 3H), 0.52 (dt,  $J = 6.5$  Hz, 2H), 0.26 (dt,  $J = 6.5$ , 12.6 Hz, 2H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  165.4, 164.5, 163.4, 135.5, 134.1, 128.7, 89.6, 55.1, 52.2, 26.2, 21.4, 21.0, 18.8, 18.1, 11.4, 10.3, 3.7; ESI–MS  $m/z$ : 385.28 ( $\text{MH}^+$ ).

**4.1.6.5.  $N^4$ -butyl- $N^4$ -ethyl- $N^6$ -mesityl-2-methyl-5-(methylthio)pyrimidine-4,6-diamine (5).** Yield: light yellow semisolid (66%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.58 (s, 1H), 6.93 (s, 2H), 3.70 (q,  $J = 7.1$  Hz, 2H), 3.60 (t,  $J = 7.5$  Hz, 2H), 2.31 (s, 3H), 2.23 (s, 3H), 2.22 (s, 3H), 2.20 (s, 6H), 1.67 (quin,  $J = 7.5$  Hz, 2H), 1.36 (sxt,  $J = 7.5$  Hz, 2H), 1.24 (t,  $J = 7.1$  Hz, 3H), 0.97 (t,  $J = 7.4$  Hz, 3H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  165.4, 163.9, 163.4, 135.5, 134.1, 128.7, 88.9, 49.8, 45.3, 30.5, 26.2, 21.0, 20.3, 18.8, 18.3, 14.0, 13.6; ESI–MS  $m/z$ : 373.35 ( $\text{MH}^+$ ).

**4.1.6.6. 5-(Ethylthio)- $N^4$ -mesityl-2-methyl- $N^6$ , $N^6$ -dipropylpyrimidine-4,6-diamine (6).** Yield: Light yellow solid (59.0%); MP: 85 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.55 (s, 1H), 6.92 (s, 2H), 3.57 (t,  $J = 7.4$  Hz, 4H), 2.67 (q,  $J = 7.5$  Hz, 2H), 2.30 (s, 3H), 2.23 (s, 3H), 2.18 (s, 6H), 1.66 (sxt,  $J = 7.6$  Hz, 4H), 1.21 (t,  $J = 7.4$  Hz, 3H), 0.91 (t,  $J = 7.4$  Hz, 6H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  165.3, 164.5, 163.7, 135.6, 134.1, 128.7, 87.3, 52.7, 28.9, 26.2, 21.5, 20.9, 18.9, 14.3, 11.4; ESI–MS  $m/z$ : 387.27 ( $\text{MH}^+$ ).

**4.1.6.7.  $N^4$ , $N^4$ -diethyl-5-(ethylthio)- $N^6$ -mesityl-2-methylpyrimidine-4,6-diamine (7).** Yield: off white semisolid (44.5%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.54 (s, 1H), 6.92 (s, 2H), 3.67 (q,  $J = 6.9$  Hz, 4H), 2.68 (q,  $J = 7.4$  Hz, 2H), 2.31 (s, 3H), 2.24 (s, 3H), 2.18 (s, 6H), 1.28 (t,  $J = 6.9$  Hz, 6H), 1.24 (t,  $J = 7.4$  Hz, 3H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  165.4, 164.7, 163.7, 135.6, 134.1, 128.7, 87.5, 44.6, 28.8, 26.2, 20.9, 18.9, 14.3, 13.6; ESI–MS  $m/z$ : 359.22 ( $\text{MH}^+$ ).

**4.1.6.8. 5-(Ethylthio)- $N^4$ -mesityl- $N^6$ , $N^6$ -bis(2-methoxyethyl)-2-methylpyrimidine-4,6-diamine (8).** Yield: semisolid (60%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.55 (s, 1H), 6.92 (s, 2H), 3.92 (t,  $J = 6.2$  Hz, 4H), 3.64 (t,  $J = 6.1$  Hz, 4H), 3.37 (s, 6H), 2.69 (q,  $J = 7.5$  Hz, 2H), 2.30 (s, 3H), 2.23 (s, 3H), 2.18 (s, 6H), 1.22 (t,  $J = 7.4$  Hz, 3H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  165.4, 164.4, 163.9, 135.5, 133.9, 128.7, 87.8, 71.4, 58.8, 50.8, 28.7, 26.2, 20.9, 18.8, 14.3; ESI–MS  $m/z$ : 419.27 ( $\text{MH}^+$ ).

**4.1.6.9.  $N^4$ -(cyclopropylmethyl)-5-(ethylthio)- $N^6$ -mesityl-2-methyl- $N^4$ -propylpyrimidine-4,6-diamine (9).** Yield: off white solid (72.7%); MP: 70 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.53 (s, 1H), 6.92 (s, 2H), 3.67 (t,  $J = 7.3$  Hz, 2H), 3.50 (d,  $J = 6.5$  Hz, 2H), 2.69 (q,  $J = 7.4$  Hz, 2H), 2.30 (s, 3H), 2.24 (s, 3H), 2.18 (s, 6H), 1.69 (sxt,  $J = 7.3$  Hz, 2H), 1.21 (t,  $J = 7.4$  Hz, 3H), 1.16 (tquin,  $J = 5.0$ , 6.5 Hz 1H), 0.91 (t,  $J = 7.3$  Hz, 3H), 0.50 (dt,  $J = 5.0$  2H), 0.24 (dt,  $J = 5.0$  Hz, 2H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  166.1, 165.2, 163.8, 135.6, 134.1, 128.7, 88.0, 55.1, 52.3, 28.7, 26.2, 21.3, 21.0, 18.9, 14.4, 11.4, 10.3, 3.7; ESI–MS  $m/z$ : 399.27 ( $\text{MH}^+$ ).

**4.1.6.10.  $N^4$ -butyl- $N^4$ -ethyl-5-(ethylthio)- $N^6$ -mesityl-2-methylpyrimidine-4,6-diamine (10).** Yield: semisolid (66.7%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.55 (s, 1H), 6.93 (s, 2H), 3.69 (q,  $J = 7.1$  Hz, 2H), 3.62 (t,  $J = 7.1$  Hz, 2H), 2.68 (q,  $J = 7.6$  Hz, 2H), 2.31 (s, 3H), 2.24 (s, 3H), 2.19 (s, 6H), 1.65 (quin,  $J = 7.6$  Hz, 2H), 1.37 (sxt,  $J = 7.5$  Hz, 2H), 1.22 (t,  $J = 7.3$  Hz, 6H), 0.97 (t,  $J = 7.4$  Hz, 3H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  165.3, 164.6, 163.7, 135.6, 134.1, 128.7, 87.4, 49.9, 45.3, 30.4, 28.8, 26.2, 20.9, 20.3, 18.9, 14.3, 14.0, 13.5; ESI–MS  $m/z$ : 387.27 ( $\text{MH}^+$ ).

**4.1.6.11.  $N^4$ -(2-bromo-4-isopropylphenyl)-2-methyl-5-(methylthio)- $N^6$ , $N^6$ -dipropylpyrimidine-4,6-diamine (11).** Yield: semisolid (72.9%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.91 (s, 1H), 8.56 (d,  $J = 8.6$  Hz, 1H), 7.41 (d,  $J = 2.0$  Hz, 1H), 7.19 (dd,  $J = 2.0$ , 8.6 Hz, 1H), 3.59 (t,  $J = 7.4$  Hz, 4H), 2.86 (spt,  $J = 7.1$  Hz, 1H), 2.41 (s, 3H), 2.19 (s, 3H), 1.67 (sxt,  $J = 7.4$  Hz, 4H), 1.25 (d,  $J = 7.1$  Hz, 6H), 0.91 (t,  $J = 7.4$  Hz, 6H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  164.9, 161.9, 143.8, 135.7, 130.0, 126.0, 121.5, 114.0, 91.3, 52.7, 33.3, 26.1, 24.0, 21.5, 18.7, 11.3; ESI–MS  $m/z$ : 451.24 ( $\text{MH}^+$ ), 453.20 ( $\text{MH} + 2^+$ ).

4.1.6.12.  $N^4$ -(2-bromo-4-isopropylphenyl)- $N^6,N^6$ -diethyl-2-methyl-5-(methylthio)pyrimidine-4,6-diamine (**12**). Yield: semisolid (76.7%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.88 (s, 1H), 8.55 (d,  $J$  = 8.6 Hz, 1H), 7.41 (d,  $J$  = 2.0 Hz, 1H), 7.19 (dd,  $J$  = 2.0, 8.6 Hz, 1H), 3.69 (q,  $J$  = 7.1 Hz, 4H), 2.86 (spt,  $J$  = 6.9 Hz, 1H), 2.42 (s, 3H), 2.20 (s, 3H), 1.26 (t,  $J$  = 7.0 Hz, 6H), 1.20 (t,  $J$  = 7.1 Hz, 6H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  164.9, 164.0, 161.8, 143.9, 135.7, 130.0, 126.0, 121.5, 114.1, 91.3, 44.7, 33.3, 26.1, 24.0, 18.6, 13.6; ESI-MS  $m/z$ : 423.19 ( $\text{MH}^+$ ), 425.17 ( $\text{MH} + 2^+$ ).

4.1.6.13.  $N^4$ -(2-bromo-4-isopropylphenyl)- $N^6,N^6$ -bis(2-methoxyethyl)-2-methyl-5-(methylthio)pyrimidine-4,6-diamine (**13**). Yield: semisolid (56%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.90 (s, 1H), 8.53 (d,  $J$  = 8.6 Hz, 1H), 7.41 (d,  $J$  = 2.0 Hz, 1H), 7.19 (dd,  $J$  = 2.0, 8.6 Hz, 1H), 3.96 (t,  $J$  = 6.0 Hz, 4H), 3.65 (t,  $J$  = 6.0 Hz, 4H), 3.37 (s, 6H), 2.87 (spt,  $J$  = 6.9 Hz, 1H), 2.41 (s, 3H), 2.21 (s, 3H), 1.25 (d,  $J$  = 6.9 Hz, 6H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  165.0, 163.9, 162.0, 144.1, 135.5, 130.0, 126.0, 121.6, 114.1, 91.7, 71.4, 58.8, 51.0, 33.3, 26.1, 23.9, 18.3; ESI-MS  $m/z$ : 483.2 ( $\text{MH}^+$ ), 485.17 ( $\text{MH} + 2^+$ ).

4.1.6.14.  $N^4$ -(2-bromo-4-isopropylphenyl)- $N^6$ -(cyclopropylmethyl)-2-methyl-5-(methylthio)- $N^6$ -propylpyrimidine-4,6-diamine (**14**). Yield: semisolid (68.4%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.88 (s, 1H), 8.56 (d,  $J$  = 8.6 Hz, 1H), 7.41 (d,  $J$  = 2.0 Hz, 1H), 7.19 (dd,  $J$  = 2.0, 8.6 Hz, 1H), 3.70 (t,  $J$  = 7.5 Hz, 2H), 3.53 (d,  $J$  = 6.8 Hz, 2H), 2.87 (spt,  $J$  = 6.9 Hz, 1H), 2.42 (s, 3H), 2.21 (s, 3H), 1.68 (sxt,  $J$  = 7.5 Hz, 2H), 1.25 (d,  $J$  = 6.9 Hz, 6H), 1.17 (tquin,  $J$  = 5.0, 6.8 Hz, 1H), 0.91 (t,  $J$  = 7.4 Hz, 3H), 0.51 (dt,  $J$  = 2.0, 5.0 Hz, 2H), 0.25 (dt,  $J$  = 2.0, 5.0 Hz, 2H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  164.9, 164.7, 161.9, 143.9, 135.7, 130.0, 126.0, 121.5, 114.0, 92.0, 55.2, 52.3, 33.3, 26.1, 23.9, 21.4, 18.4, 11.3, 10.3, 3.7; ESI-MS  $m/z$ : 463.23 ( $\text{MH}^+$ ), 465.19 ( $\text{MH} + 2^+$ ).

4.1.6.15.  $N^4$ -(2-bromo-4-isopropylphenyl)- $N^6$ -butyl- $N^6$ -ethyl-2-methyl-5-(methylthio)pyrimidine-4,6-diamine (**15**). Yield: semisolid (72%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.88 (s, 1H), 8.54 (d,  $J$  = 8.6 Hz, 1H), 7.40 (d,  $J$  = 2.0 Hz, 1H), 7.18 (dd,  $J$  = 2.0, 8.6 Hz, 1H), 3.69 (q,  $J$  = 7.0 Hz, 2H), 3.60 (t,  $J$  = 7.2 Hz, 2H), 2.85 (spt,  $J$  = 6.9 Hz, 1H), 2.40 (s, 3H), 2.18 (s, 3H), 1.63 (quin,  $J$  = 7.4 Hz, 2H), 1.34 (sxt,  $J$  = 7.3 Hz, 2H), 1.25 (d,  $J$  = 6.9 Hz, 6H), 1.18 (t,  $J$  = 7.0 Hz, 3H), 0.95 (t,  $J$  = 7.3 Hz, 3H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  164.9, 164.1, 161.9, 143.8, 135.7, 130.0, 126.0, 121.5, 114.0, 91.3, 49.9, 45.4, 33.3, 30.5, 26.1, 24.0, 20.2, 18.6, 14.0, 13.6; ESI-MS  $m/z$ : 451.23 ( $\text{MH}^+$ ), 453.20 ( $\text{MH} + 2^+$ ).

4.1.6.16.  $N^4$ -(2-bromo-4-isopropylphenyl)-5-(ethylthio)-2-methyl- $N^6,N^6$ -dipropylpyrimidine-4,6-diamine (**16**). Yield: semisolid (66.4%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.93 (s, 1H), 8.58 (d,  $J$  = 8.6 Hz, 1H), 7.40 (d,  $J$  = 2.0 Hz, 1H), 7.18 (dd,  $J$  = 2.0, 8.6 Hz, 1H), 3.59 (t,  $J$  = 7.3 Hz, 4H), 2.86 (spt,  $J$  = 6.9 Hz, 1H), 2.64 (q,  $J$  = 7.5 Hz, 2H), 2.42 (s, 3H), 1.66 (sxt,  $J$  = 7.3 Hz, 4H), 1.24 (d,  $J$  = 6.9 Hz, 6H), 1.18 (t,  $J$  = 7.4 Hz, 3H), 0.90 (t,  $J$  = 7.3 Hz, 6H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  164.8, 164.8, 162.3, 143.7, 135.8, 130.0, 126.0, 121.2, 113.8, 89.6, 52.8, 33.3, 29.5, 26.2, 24.0, 21.4, 14.1, 11.3; ESI-MS  $m/z$ : 465.24 ( $\text{MH}^+$ ), 467.20 ( $\text{MH} + 2^+$ ).

4.1.6.17.  $N^4$ -(2-bromo-4-isopropylphenyl)- $N^6,N^6$ -diethyl-5-(ethylthio)-2-methylpyrimidine-4,6-diamine (**17**). Yield: semisolid (54%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.91 (s, 1H), 8.57 (d,  $J$  = 8.6 Hz, 1H), 7.41 (d,  $J$  = 2.0 Hz, 1H), 7.19 (dd,  $J$  = 2.0, 8.6 Hz, 1H), 3.68 (q,  $J$  = 7.1 Hz, 4H), 2.86 (spt,  $J$  = 6.9 Hz, 1H), 2.65 (q,  $J$  = 7.4 Hz, 2H), 2.43 (s, 3H), 1.28 (d,  $J$  = 6.9 Hz, 6H), 1.26 (t,  $J$  = 7.1 Hz, 6H), 1.18 (t,  $J$  = 7.4 Hz, 3H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  164.9, 162.3, 143.8, 135.7, 130.0, 126.0, 121.3, 113.9, 89.7, 44.8, 33.3, 29.4, 26.2, 24.0, 14.2, 13.5; ESI-MS  $m/z$ : 437.4 ( $\text{MH}^+$ ), 439.20 ( $\text{MH} + 2^+$ ).

4.1.6.18.  $N^4$ -(2-bromo-4-isopropylphenyl)-5-(ethylthio)- $N^6,N^6$ -bis(2-methoxyethyl)-2-methylpyrimidine-4,6-diamine (**18**). Yield: semisolid (69.4%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.92 (s, 1H), 8.56 (d,

$J$  = 8.6 Hz, 1H), 7.41 (d,  $J$  = 2.0 Hz, 1H), 7.19 (dd,  $J$  = 2.0, 8.6 Hz, 1H), 3.95 (t,  $J$  = 6.0 Hz, 4H), 3.63 (t,  $J$  = 6.0 Hz, 4H), 3.36 (s, 6H), 2.86 (spt,  $J$  = 6.8 Hz, 1H), 2.67 (q,  $J$  = 7.4 Hz, 2H), 2.42 (s, 3H), 1.24 (d,  $J$  = 6.8 Hz, 6H), 1.19 (t,  $J$  = 7.4 Hz, 3H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  164.9, 164.6, 162.4, 144.0, 135.5, 130.0, 126.0, 121.4, 113.9, 90.1, 71.2, 58.8, 51.0, 33.3, 29.2, 26.1, 23.9, 14.1; ESI-MS  $m/z$ : 497.23 ( $\text{MH}^+$ ), 499.19 ( $\text{MH} + 2^+$ ).

4.1.6.19.  $N^4$ -(2-bromo-4-isopropylphenyl)- $N^6$ -(cyclopropylmethyl)-5-(ethylthio)-2-methyl- $N^6$ -propylpyrimidine-4,6-diamine (**19**). Yield: semisolid (59.6%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.91 (s, 1H), 8.59 (d,  $J$  = 8.6 Hz, 1H), 7.41 (d,  $J$  = 2.0 Hz, 1H), 7.19 (dd,  $J$  = 1.8, 8.6 Hz, 1H), 3.74 (t,  $J$  = 7.5 Hz, 2H), 3.53 (d,  $J$  = 6.5 Hz, 2H), 2.86 (spt,  $J$  = 6.9 Hz, 1H), 2.66 (q,  $J$  = 7.3 Hz, 2H), 2.43 (s, 3H), 1.67 (sxt,  $J$  = 7.5 Hz, 2H), 1.25 (d,  $J$  = 7.1 Hz, 6H), 1.17 (tquin,  $J$  = 5.0, 6.8 Hz, 1H), 1.19 (t,  $J$  = 7.4 Hz, 3H), 0.90 (t,  $J$  = 7.3 Hz, 3H), 0.51 (dt,  $J$  = 2.0, 5.0 Hz, 2H), 0.25 (dt,  $J$  = 2.0, 5.0 Hz, 2H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  165.4, 164.8, 162.3, 143.8, 135.7, 130.0, 126.0, 121.3, 113.8, 90.2, 55.2, 52.4, 33.3, 29.2, 26.2, 24.0, 21.3, 14.2, 11.3, 10.2, 3.7; ESI-MS  $m/z$ : 477.27 ( $\text{MH}^+$ ), 479.21 ( $\text{MH} + 2^+$ ).

4.1.6.20.  $N^4$ -(2-bromo-4-isopropylphenyl)- $N^6$ -butyl- $N^6$ -ethyl-5-(ethylthio)-2-methylpyrimidine-4,6-diamine (**20**). Yield: semisolid (59.6%);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.92 (s, 1H), 8.58 (d,  $J$  = 8.6 Hz, 1H), 7.41 (d,  $J$  = 2.0 Hz, 1H), 7.19 (dd,  $J$  = 2.0, 8.6 Hz, 1H), 3.70 (q,  $J$  = 7.1 Hz, 2H), 3.60 (t,  $J$  = 7.2 Hz, 2H), 2.86 (spt,  $J$  = 6.9 Hz, 1H), 2.64 (q,  $J$  = 7.3 Hz, 2H), 2.42 (s, 3H), 1.63 (quin,  $J$  = 7.2 Hz, 2H), 1.34 (sxt,  $J$  = 7.3 Hz, 2H), 1.25 (d,  $J$  = 6.9 Hz, 6H), 1.20 (t,  $J$  = 7.1 Hz, 3H), 0.96 (m, 6H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  164.8, 164.8, 162.3, 143.7, 135.7, 130.0, 126.0, 121.3, 113.8, 89.7, 50.0, 45.5, 33.3, 30.4, 29.4, 26.2, 24.0, 20.2, 14.2, 14.0, 13.5; ESI-MS  $m/z$ : 465.25 ( $\text{MH}^+$ ), 467.21 ( $\text{MH} + 2^+$ ).

4.1.6.21.  $N^4$ -mesityl-5-(methylthio)- $N^6,N^6$ -dipropylpyrimidine-4,6-diamine (**21**). Yield: off white solid (67.2%); MP: 83 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.09 (s, 1H), 7.58 (s, 1H), 6.94 (s, 2H), 3.63 (t,  $J$  = 7.4 Hz, 4H), 2.29 (s, 3H), 2.26 (s, 3H), 2.19 (s, 6H), 1.70 (sxt,  $J$  = 7.5 Hz, 4H), 0.93 (t,  $J$  = 7.4 Hz, 6H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  163.8, 156.6, 136.4, 135.9, 133.6, 128.9, 91.6, 52.7, 21.4, 21.0, 18.6, 18.2, 11.3; ESI-MS  $m/z$ : 459.25 ( $\text{MH}^+$ ).

4.1.6.22.  $N^4$ -mesityl- $N^6,N^6$ -bis(2-methoxyethyl)-5-(methylthio)pyrimidine-4,6-diamine (**22**). Yield: light yellow solid (60.9%); MP: 89 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.08 (s, 1H), 7.59 (s, 1H), 6.94 (s, 2H), 3.96 (t,  $J$  = 6.0 Hz, 4H), 3.66 (t,  $J$  = 6.0 Hz, 4H), 3.37 (s, 6H), 2.29 (s, 3H), 2.27 (s, 3H), 2.18 (s, 6H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  163.9, 163.6, 156.6, 136.6, 135.9, 129.0, 92.0, 71.4, 58.9, 51.0, 21.0, 18.5; ESI-MS  $m/z$ : 391.54 ( $\text{MH}^+$ ).

4.1.6.23.  $N^4$ -(cyclopropylmethyl)- $N^6$ -mesityl-5-(methylthio)- $N^4$ -propylpyrimidine-4,6-diamine (**23**). Yield: white solid (75.4%); MP: 85 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.12 (s, 1H), 7.56 (s, 1H), 6.94 (s, 2H), 3.70 (t,  $J$  = 7.3 Hz, 2H), 3.53 (d,  $J$  = 6.5 Hz, 2H), 2.30 (s, 3H), 2.27 (s, 3H), 2.20 (s, 6H), 1.71 (sxt,  $J$  = 7.4 Hz, 2H), 1.20 (tquin,  $J$  = 12.5, 6.5 Hz, 1H), 0.94 (t,  $J$  = 7.3 Hz, 3H), 0.52 (dt,  $J$  = 6.5, 12.6 Hz, 2H), 0.26 (dt,  $J$  = 6.5, 12.6 Hz, 2H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  164.4, 163.8, 156.6, 136.5, 135.9, 133.5, 129.0, 92.3, 55.2, 52.2, 21.3, 21.0, 18.6, 17.9, 11.3, 10.2, 3.7; ESI-MS  $m/z$ : 371.29 ( $\text{MH}^+$ ).

4.1.6.24.  $N^4$ -butyl- $N^4$ -ethyl- $N^6$ -mesityl-5-(methylthio)pyrimidine-4,6-diamine (**24**). Yield: off white solid (73.8%); MP: 57 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.10 (s, 1H), 7.57 (s, 1H), 6.94 (s, 2H), 3.70 (q,  $J$  = 7.1 Hz, 2H), 3.60 (t,  $J$  = 7.5 Hz, 2H), 2.31 (s, 3H), 2.23 (s, 3H), 2.20 (s, 6H), 1.67 (quin,  $J$  = 7.5 Hz, 2H), 1.36 (sxt,  $J$  = 7.5 Hz, 2H), 1.24 (t,  $J$  = 7.1 Hz, 3H), 0.97 (t,  $J$  = 7.4 Hz, 3H);  $^{13}\text{C}$  NMR (400 MHz,  $\text{CDCl}_3$ )



$\delta$  163.8, 163.7, 156.7, 136.4, 135.9, 133.6, 129.0, 91.5, 50.0, 45.4, 30.5, 21.0, 20.3, 18.6, 18.1, 14.0, 13.5; ESI–MS  $m/z$ : 359.27 ( $MH^+$ ).

**4.1.6.25.** 5-(Ethylthio)- $N^4$ -mesityl- $N^6$ , $N^6$ -dipropylpyrimidine-4,6-diamine (**25**). Yield: light yellow semisolid (76.4%);  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.10 (s, 1H), 7.55 (s, 1H), 6.94 (s, 2H), 3.60 (t,  $J = 7.4$  Hz, 4H), 2.72 (q,  $J = 7.4$  Hz, 2H), 2.29 (s, 3H), 2.18 (s, 6H), 1.68 (sxt,  $J = 7.6$  Hz, 4H), 1.23 (t,  $J = 7.4$  Hz, 3H), 0.92 (t,  $J = 7.3$  Hz, 6H);  $^{13}C$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  164.4, 164.1, 156.5, 136.4, 135.9, 129.0, 90.0, 52.8, 28.7, 21.4, 21.0, 18.6, 14.3, 11.3; ESI–MS  $m/z$ : 373.29 ( $MH^+$ ).

**4.1.6.26.**  $N^4$ , $N^4$ -diethyl-5-(ethylthio)- $N^6$ -mesitylpyrimidine-4,6-diamine (**26**). Yield: (42.03%); MP: 45 °C;  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.11 (s, 1H), 7.55 (s, 1H), 6.94 (s, 2H), 3.69 (q,  $J = 6.8$  Hz, 4H), 2.73 (q,  $J = 7.3$  Hz, 2H), 2.29 (s, 3H), 2.19 (s, 6H), 1.28 (t,  $J = 6.9$  Hz, 6H), 1.24 (t,  $J = 6.9$  Hz, 3H);  $^{13}C$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  164.4, 164.1, 156.6, 135.9, 133.5, 128.9, 90.0, 44.8, 28.7, 21.0, 18.6, 14.3, 13.5; ESI–MS  $m/z$ : 345.27 ( $MH^+$ ).

**4.1.6.27.** 5-(Ethylthio)- $N^4$ -mesityl- $N^6$ , $N^6$ -bis(2-methoxyethyl)pyrimidine-4,6-diamine (**27**). Yield: light yellow semisolid (61.5%);  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.10 (s, 1H), 7.55 (s, 1H), 6.94 (s, 2H), 3.95 (t,  $J = 6.0$  Hz, 4H), 3.64 (t,  $J = 6.0$  Hz, 4H), 3.36 (s, 6H), 2.75 (q,  $J = 7.3$  Hz, 2H), 2.29 (s, 3H), 2.18 (s, 6H), 1.23 (t,  $J = 7.4$  Hz, 3H);  $^{13}C$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  164.2, 156.5, 136.5, 135.8, 133.4, 129.0, 90.5, 71.2, 58.8, 50.9, 28.5, 21.0, 18.6, 14.3; ESI–MS  $m/z$ : 405.28 ( $MH^+$ ).

**4.1.6.28.**  $N^4$ -butyl- $N^4$ -ethyl-5-(ethylthio)- $N^6$ -mesitylpyrimidine-4,6-diamine (**28**). Yield: off white semisolid (76%);  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.10 (s, 1H), 7.54 (s, 1H), 6.94 (s, 2H), 3.69 (q,  $J = 7.0$  Hz, 2H), 3.60 (t,  $J = 7.1$  Hz, 2H), 2.72 (q,  $J = 7.4$  Hz, 2H), 2.29 (s, 3H), 2.18 (s, 6H), 1.61 (quin,  $J = 7.6$  Hz, 2H), 1.34 (sxt,  $J = 7.5$ , 2H), 1.23 (t,  $J = 7.3$  Hz, 6H), 0.96 (t,  $J = 7.4$  Hz, 3H);  $^{13}C$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  164.4, 164.1, 156.6, 136.4, 135.9, 133.6, 129.0, 90.1, 50.1, 45.5, 30.4, 28.7, 21.0, 20.2, 18.6, 14.3, 14.0, 13.4; ESI–MS  $m/z$ : 373.32 ( $MH^+$ ).

## 4.2. CRF<sub>1</sub> receptor binding study

Binding studies were performed in membrane homogenates from human embryonic kidney cells (HEK 293) stably expressing CRF<sub>1</sub> receptors and using [ $^{125}I$ ]-Tyr<sup>0</sup>-sauvagine as radioligand. Membrane homogenates were prepared according to the method of Gkountelias [54] CRF<sub>1</sub>-expressing HEK 293 cells, grown in DMEM/F12 (1:1) containing 3.15 g/L glucose, 10% bovine calf serum and 300  $\mu$ g/ml of the antibiotic, Geneticin at 37 °C and 5% CO<sub>2</sub>, were washed with phosphate-buffered saline (PBS) (4.3 mM Na<sub>2</sub>HPO<sub>4</sub>·7H<sub>2</sub>O, 1.4 mM KH<sub>2</sub>PO<sub>4</sub>, 137 mM NaCl, 2.7 mM KCl, pH 7.2–7.3 at R.T). Then the cells were briefly treated with PBS containing 2 mM EDTA (PBS/EDTA), and then dissociated in PBS/EDTA. Cells suspensions were centrifuged at 1000× g for 5 min at room temperature, and the pellets were homogenized in 1.5 mL of buffer H (20 mM HEPES, containing 10 mM MgCl<sub>2</sub>, 2 mM EGTA, 0.2 mg/ml bacitracin and 0.93  $\mu$ g/ml aprotinin pH 7.2 at 4 °C) using a Janke & Kunkel IKA Ultra Turrax T25 homogenizer, at setting ~20, for 10–15 s, at 4 °C. The homogenates were centrifuged at 16,000× g, for 10 min, at 4 °C. The membrane pellets were re-suspended by homogenization, as described above, in 1 mL buffer B (buffer H containing 0.1% BSA, pH 7.2 at 20 °C). The membrane suspensions were then diluted in buffer B and aliquots of suspensions (50  $\mu$ l) were added into tubes containing buffer B and 20–25 pM [ $^{125}I$ ]-Tyr<sup>0</sup>-sauvagine without or with compounds **1–28** at a single concentration of 500 nM or with increasing concentrations of these compounds (competition binding experiments) in a final volume of 0.2 mL. The mixtures were incubated at 20–21 °C for 120 min and

then filtered through Whatman 934AH filters, presoaked for 1 h in 0.3% polyethylenimine at 4 °C. The filters were washed 3 times with 0.5 mL of ice-cold PBS, pH 7.1 containing 0.01% Triton X-100 and assessed for radioactivity in a gamma counter. The amount of membranes used was adjusted to insure that the specific binding was always equal to or less than 10% of the total concentration of the added radioligand. Specific [ $^{125}I$ ]-Tyr<sup>0</sup>-sauvagine binding was defined as total binding less nonspecific binding in the presence of 1000 nM antalarmin. Data for competition binding were analyzed by nonlinear regression analysis, using Prism 4.0 (GraphPad Software, San Diego, CA). IC<sub>50</sub> values were obtained by fitting the data from competition studies to a one-site competition model.

## 4.3. Statistical analysis

Prism 4.0 was used for all statistical analysis (GraphPad Software, San Diego, CA). Competition binding was analyzed by nonlinear regression analysis, and IC<sub>50</sub> values were obtained by fitting the data from competition studies to a one-site competition model by nonlinear regression.

## Acknowledgments

This research project was funded by Department of Pharmaceutical Sciences, South Dakota State University, Brookings, South Dakota. CRF receptor binding evaluation was performed at the Department of Pharmacology, Faculty of Medicine, University of Crete, Crete, Greece.

## References

- [1] W. Vale, J. Spiess, C. Rivier, J. Rivier, Characterization of a 41-residue ovine hypothalamic peptide that stimulates secretion of corticotropin and beta-endorphin, *Science* 213 (1981) 1394–1397.
- [2] G.P. Chrousos, The hypothalamic-pituitary-adrenal axis and immune-mediated inflammation, *The New England Journal of Medicine* 332 (1995) 1351–1362.
- [3] M.J. Owens, C.B. Nemeroff, Physiology and pharmacology of corticotropin-releasing factor, *Pharmacological Reviews* 43 (1991) 425–473.
- [4] P.W. Gold, G.P. Chrousos, Organization of the stress system and its dysregulation in melancholic and atypical depression: high vs low CRH/NE states, *Molecular Psychiatry* 7 (2002) 254–275.
- [5] M.E. Keck, F. Holsboer, Hyperactivity of CRH neuronal circuits as a target for therapeutic interventions in affective disorders, *Peptides* 22 (2001) 835–844.
- [6] J.M. Reul, F. Holsboer, Corticotropin-releasing factor receptors 1 and 2 in anxiety and depression, *Current Opinion in Pharmacology* 2 (2002) 23–33.
- [7] F.M. Dautzenberg, G.J. Kilpatrick, R.L. Hauger, J. Moreau, Molecular biology of the CRH receptors— in the mood, *Peptides* 22 (2001) 753–760.
- [8] D.P. Behan, S.C. Heinrichs, J.C. Troncoso, X.J. Liu, C.H. Kawan, N. Ling, E.B. De Souza, Displacement of corticotropin releasing factor from its binding protein as a possible treatment for Alzheimer's disease, *Nature* 378 (1995) 284–287.
- [9] E.B. De Souza, P.J. Whitehouse, M.J. Kuhar, D.L. Price, W.W. Vale, Reciprocal changes in corticotropin-releasing factor (CRF)-like immunoreactivity and CRF receptors in cerebral cortex of Alzheimer's disease, *Nature* 319 (1986) 593–595.
- [10] A. Makriganakis, E. Zoumakis, S. Kalantaridou, C. Coutifaris, A.N. Margioris, G. Coukos, K.C. Rice, A. Gravanis, G.P. Chrousos, Corticotropin-releasing hormone promotes blastocyst implantation and early maternal tolerance, *Nature Immunology* 2 (2001) 1018–1024.
- [11] G.P. Chrousos, Interactions between the hypothalamic-pituitary-adrenal axis and the female reproductive system: clinical implications, *Annals of Internal Medicine* 129 (1998) 229.
- [12] V. Martinez, J. Rivier, L. Wang, Y. Tache, Central injection of a new corticotropin-releasing factor (CRF) antagonist, astressin, blocks CRF- and stress-related alterations of gastric and colonic motor function, *The Journal of Pharmacology and Experimental Therapeutics* 280 (1997) 754–760.
- [13] D.N. Orth, Corticotropin-releasing hormone in humans, *Endocrine Reviews* 13 (1992) 164–191.
- [14] D.G. Parkes, R.S. Weisinger, C.N. May, Cardiovascular actions of CRH and urocortin: an update, *Peptides* 22 (2001) 821–827.
- [15] M. Venihaki, P. Dikkes, A. Carrigan, K.P. Karalis, Corticotropin-releasing hormone regulates IL-6 expression during inflammation, *The Journal of Clinical Investigation* 108 (2001) 1159–1166.
- [16] M. Venihaki, J.A. Majzoub, Animal models of CRH deficiency, *Frontiers in Neuroendocrinology* 20 (1999) 122–145.

- [17] E. Zoumakis, A.N. Margioris, A. Makrigiannakis, C. Stournaras, A. Gravanis, Human endometrium as a neuroendocrine tissue: expression, regulation and biological roles of endometrial corticotropin-releasing hormone (CRH) and opioid peptides, *Journal of Endocrinological Investigation* 20 (1997) 158–167.
- [18] J. Rivier, J. Spiess, W. Vale, Characterization of rat hypothalamic corticotropin-releasing factor, *Proceedings of the National Academy of Sciences of the United States of America* 80 (1983) 4851–4855.
- [19] S. Shibahara, Y. Morimoto, Y. Furutani, M. Notake, H. Takahashi, S. Shimizu, S. Horikawa, S. Numa, Isolation and sequence analysis of the human corticotropin-releasing factor precursor gene, *The EMBO Journal* 2 (1983) 775–779.
- [20] N. Ling, F. Esch, P. Bohlen, A. Baird, R. Guillemin, Isolation and characterization of caprine corticotropin-releasing factor, *Biochemical and Biophysical Research Communications* 122 (1984) 1218–1224.
- [21] M. Pathy, J. Horvath, M. Mason-Garcia, B. Szoke, D.H. Schlesinger, A.V. Schally, Isolation and amino acid sequence of corticotropin-releasing factor from pig hypothalamus, *Proceedings of the National Academy of Sciences of the United States of America* 82 (1985) 8762–8766.
- [22] M.P. Stenzel-Poore, K.A. Heldwein, P. Stenzel, S. Lee, W.W. Vale, Characterization of the genomic corticotropin-releasing factor (CRF) gene from *Xenopus laevis*: two members of the CRF family exist in amphibians, *Molecular Endocrinology* 6 (1992) 1716–1724.
- [23] P.C. Montecucchi, A. Henschen, Amino acid composition and sequence analysis of sauvagine, a new active peptide from the skin of *Phyllomedusa sauvagei*, *International Journal of Peptide and Protein Research* 18 (1981) 113–120.
- [24] K. Lederis, A. Letter, D. McMaster, G. Moore, D. Schlesinger, Complete amino acid sequence of urotensin I, a hypotensive and corticotropin-releasing neuropeptide from *Catostomus*, *Science* 218 (1982) 162–165.
- [25] T.M. Reyes, K. Lewis, M.H. Perrin, K.S. Kunitake, J. Vaughan, C.A. Arias, J.B. Hogenesch, J. Gulyas, J. Rivier, W.W. Vale, P.E. Sawchenko, Urocortin II: a member of the corticotropin-releasing factor (CRF) neuropeptide family that is selectively bound by type 2 CRF receptors, *Proceedings of the National Academy of Sciences of the United States of America* 98 (2001) 2843–2848.
- [26] K. Lewis, C. Li, M.H. Perrin, A. Blount, K. Kunitake, C. Donaldson, J. Vaughan, T.M. Reyes, J. Gulyas, W. Fischer, L. Bilezikjian, J. Rivier, P.E. Sawchenko, W.W. Vale, Identification of urocortin III, an additional member of the corticotropin-releasing factor (CRF) family with high affinity for the CRF2 receptor, *Proceedings of the National Academy of Sciences of the United States of America* 98 (2001) 7570–7575.
- [27] P.V. Pallai, M. Mabilia, M. Goodman, W. Vale, J. Rivier, Structural homology of corticotropin-releasing factor, sauvagine, and urotensin I: circular dichroism and prediction studies, *Proceedings of the National Academy of Sciences of the United States of America* 80 (1983) 6770–6774.
- [28] T.L. Bale, W.W. Vale, CRF and CRF receptors: role in stress responsivity and other behaviors, *Annual Review of Pharmacology and Toxicology* 44 (2004) 525–557.
- [29] E.B. De Souza, Corticotropin-releasing factor receptors: physiology, pharmacology, biochemistry and role in central nervous system and immune disorders, *Psychoneuroendocrinology* 20 (1995) 789–819.
- [30] F.M. Dautzenberg, R.L. Hauger, The CRF peptide family and their receptors: yet more partners discovered, *Trends in Pharmacological Sciences* 23 (2002) 71–77.
- [31] J. Saunders, J. Williams, Antagonists of the corticotropin releasing factor receptor, *Progress in Medicinal Chemistry* 41 (2003) 195–247.
- [32] D.E. Grigoriadis, The corticotropin-releasing factor receptor: a novel target for the treatment of depression and anxiety-related disorders, *Expert Opinion on Therapeutic Targets* 9 (2005) 651–684.
- [33] B.J. Kolber, M.S. Roberts, M.P. Howell, D.F. Wozniak, M.S. Sands, L.J. Muglia, Central amygdala glucocorticoid receptor action promotes fear-associated CRH activation and conditioning, *Proceedings of the National Academy of Sciences of the United States of America* 105 (2008) 12004–12009.
- [34] M.A. Watanabe, S. Kucenas, T.A. Bowman, M. Ruhlman, M.M. Knuepfer, Angiotensin II and CRF receptors in the central nucleus of the amygdala mediate hemodynamic response variability to cocaine in conscious rats, *Brain Research* 1309 (2010) 53–65.
- [35] Z. Sarnyai, É. Biró, J. Gardi, M. Vecsernyés, J. Julesz, G. Telegdy, Brain corticotropin-releasing factor mediates 'anxiety-like' behavior induced by cocaine withdrawal in rats, *Brain Research* 675 (1995) 89–97.
- [36] O. George, S. Ghazland, M.R. Azar, P. Cottone, E.P. Zorrilla, L.H. Parsons, L.E. O'Dell, H.N. Richardson, G.F. Koob, CRF-CRF1 system activation mediates withdrawal-induced increases in nicotine self-administration in nicotine-dependent rats, *Proceedings of the National Academy of Sciences of the United States of America* 104 (2007) 17198–17203.
- [37] P.J. Gilligan, D.W. Robertson, R. Zaczek, Corticotropin releasing factor (CRF) receptor modulators: progress and opportunities for new therapeutic agents, *Journal of Medicinal Chemistry* 43 (2000) 1641–1660.
- [38] E.P. Zorrilla, G.F. Koob, Progress in corticotropin-releasing factor-1 antagonist development, *Drug Discovery Today* 15 (2010) 371–383.
- [39] E.P. Zorrilla, G.R. Valdez, J. Nozulak, G.F. Koob, A. Markou, Effects of antalarmin, a CRF type 1 receptor antagonist, on anxiety-like behavior and motor activation in the rat, *Brain Research* 952 (2002) 188–199.
- [40] E.L. Webster, D.B. Lewis, D.J. Torpy, E.K. Zachman, K.C. Rice, G.P. Chrousos, In vivo and in vitro characterization of antalarmin, a nonpeptide corticotropin-releasing hormone (CRH) receptor antagonist: suppression of pituitary ACTH release and peripheral inflammation, *Endocrinology* 137 (1996) 5747–5750.
- [41] T. Deak, K.T. Nguyen, A.L. Ehrlich, L.R. Watkins, R.L. Spencer, S.F. Maier, J. Licinio, M.L. Wong, G.P. Chrousos, E. Webster, P.W. Gold, The impact of the nonpeptide corticotropin-releasing hormone antagonist antalarmin on behavioral and endocrine responses to stress, *Endocrinology* 140 (1999) 79–86.
- [42] R.J. Briscoe, C.L. Cabrera, T.J. Baird, K.C. Rice, J.H. Woods, Antalarmin blockade of corticotropin releasing hormone-induced hypertension in rats, *Brain Research* 881 (2000) 204–207.
- [43] E.L. Webster, R.M. Barrientos, C. Contoreggi, M.G. Isaac, S. Ligier, K.E. Gabry, G.P. Chrousos, E.F. McCarthy, K.C. Rice, P.W. Gold, E.M. Sternberg, Corticotropin releasing hormone (CRH) antagonist attenuates adjuvant induced arthritis: role of CRH in peripheral inflammation, *The Journal of Rheumatology* 29 (2002) 1252–1261.
- [44] B. Greenwood-Van Meerveld, A.C. Johnson, S. Cochrane, J. Schulkin, D.A. Myers, Corticotropin-releasing factor 1 receptor-mediated mechanisms inhibit colonic hypersensitivity in rats, *Neurogastroenterology and Motility: The Official Journal of the European Gastrointestinal Motility Society* 17 (2005) 415–422.
- [45] V. Martinez, Y. Tache, CRF1 receptors as a therapeutic target for irritable bowel syndrome, *Current Pharmaceutical Design* 12 (2006) 4071–4088.
- [46] K.E. Gabry, G.P. Chrousos, K.C. Rice, R.M. Mostafa, E. Sternberg, A.B. Negrao, E.L. Webster, S.M. McCann, P.W. Gold, Marked suppression of gastric ulcerogenesis and intestinal responses to stress by a novel class of drugs, *Molecular Psychiatry* 7 (2002) 474–483.
- [47] K. Gewald, U. Hain, P. Hartung, Zur Chemie der 4-aminothiazolin-2-thione, *Monatshefte für Chemie* 112 (1981) 1393–1404.
- [48] E. Badawey, S.M. Rida, A.A. Hazza, H.T.Y. Fahmy, Y.M. Gohar, Potential antimicrobials. II. Synthesis and in vitro anti-microbial evaluation of some thiazolo[4,5-d]pyrimidines, *European Journal of Medicinal Chemistry* 28 (1993) 97–101.
- [49] J.P. Beck, M.A. Curry, R.J. Chorvat, L.W. Fitzgerald, P.J. Gilligan, R. Zaczek, G.L. Trainor, Thiazolo[4,5-d]pyrimidine thiones and -ones as corticotropin-releasing hormone (CRH-R1) receptor antagonists, *Bioorganic & Medicinal Chemistry Letters* 9 (1999) 1185–1188.
- [50] J. Beck, Thiazolo[4, 5-d] pyrimidines and pyridines as corticotropin releasing factor (CRF) antagonists, in, *WO Patent 1,999,051,608*, 1999.
- [51] M.A. El Sherbeny, F.A. Badria, S.A. Kheira, synthesis of some new thiazolo[4,5-d]pyrimidine derivatives and evaluation of their antifungal, antiviral and cytotoxic activities, *Medicinal Chemistry Research* 6 (1996) 28–39.
- [52] K. Gewald, U. Hain, R. Schindler, M. Gruner, Zur Chemie der 4-Amino-thiazolin-2-thione, 2. Mitt, *Monatshefte für Chemie* 125 (1994) 1129–1143.
- [53] G.A. Artamkina, M.V. Ermolina, I.P. Beletskaya, Solvent-free Pd-catalysed N-arylation of amines, amides and diaza-18-crown-6, *Mendeleev Communications* (2003) 158–160.
- [54] K. Gkountelias, T. Tselios, M. Venihaki, G. Deraos, I. Lazaridis, O. Rassouli, A. Gravanis, G. Liapakis, Alanine scanning mutagenesis of the second extracellular loop of type 1 corticotropin-releasing factor receptor revealed residues critical for peptide binding, *Molecular Pharmacology* 75 (2009) 793–800.