



Molecular characteristics of community-associated methicillin-resistant *Staphylococcus aureus* colonizing surgical patients in Greece☆☆☆

Linus Hadjihannas^a, Mina Psychogiou^a, Joanna Empel^c, Chris Kosmidis^a, Dimitrios Goukos^a, Jina Bouzala^a, Sotirios Georgopoulos^b, Surbhi Malhotra-Kumar^d, Stephan Harbarth^e, George L. Daikos^{a,*} and in collaboration with the MOSAR-WP no. 1 and 4 study teams

^a First Department of Propaedeutic Medicine, University of Athens, Athens, Greece

^b First Department of Surgery, University of Athens, Athens, Greece

^c Department of Molecular Microbiology, National Medicine's Institute, Warsaw, Poland

^d Department of Medical Microbiology, Vaccine and Infectious Disease Institute, Universiteit Antwerpen, Antwerp, Belgium

^e Infection Control Program, Geneva University Hospitals and Medical School, Geneva, Switzerland

ARTICLE INFO

Article history:

Received 29 April 2012

Received in revised form 8 August 2012

Accepted 19 August 2012

Available online 26 September 2012

Keywords:

CA-MRSA

Colonization

Epidemiology

Surgical patients

Multidrug resistance

ABSTRACT

Fifty-one of 925 patients screened for methicillin-resistant *Staphylococcus aureus* (MRSA) upon admission to a surgical unit were MRSA carriers; 15 were classified as community- and 36 as hospital-associated-MRSA. Fourteen of 22 isolates typed by molecular methods belonged to the European clone ST80-IVc, 3 of which exhibited resistance to ≥ 3 non- β -lactam antibiotics.

© 2012 Elsevier Inc. All rights reserved.

Community-associated methicillin-resistant *S. aureus* (CA-MRSA) is an important cause of recurrent skin infections and severe necrotizing pneumonia in healthy individuals in the community (David and Daum, 2010). Although CA-MRSA strains are usually susceptible to many non- β -lactam antibiotics, emergence of resistance to multiple classes of antibiotics has become increasingly recognized in several community-associated clones, such as those of ST8, ST80, ST59, and ST772 genetic lineages, adding to the overall burden of infection caused by this pathogen (Ellington et al., 2009; Francois et al., 2008; Hanssen et al., 2005).

☆ This work was partially supported by funding from the European Community (MOSAR network contract LSHP-CT-2007-037941) and an unrestricted educational grant, Antibacterial Europe ASPIRE Research award 2010, from Pfizer International Operations (WS813738). JE was partly supported by a grant (no. 934/6. PR UE/2009/7) from the Ministry of Science and Higher Education, Poland. The work performed at NMI was partially supported by the Mikrobank 2 Programme from the Ministry of Science and Higher Education, Poland. GLD was partially supported by Pfizer Hellas for MRSA molecular typing (WS588755).

☆☆ Conflicts of interest: GLD has served as an advisory board member for Pfizer Hellas. SH is a member of the speakers' bureau of bioMérieux and Pfizer, a member of the advisory board of Destiny Pharma, DaVolterra, and bioMérieux, and has received financial support for MRSA research activities from B. Braun, Pfizer, and the European Commission.

* Corresponding author. Tel.: +30-2107462636; fax: +30-2107462635.

E-mail address: gdaikos@med.uoa.gr (G.L. Daikos).

Since its first identification in Greece in 1998 (Aires de Sousa et al., 2003), the so-called European clone (ST80-IVc, PVL-positive) has been recognized with increasing frequency as a cause of CA-MRSA infections both in children and in adults (Katopodis et al., 2010; Niniou et al., 2008; Vourli et al., 2009). More importantly, MRSA clones of community origin could become endemic in the healthcare setting causing severe infections in hospitalized patients (Popovich et al., 2008). According to a previous study, PVL-positive community-associated clones account already for 45% of health-care-associated (HA) MRSA infections at several hospitals in Greece (Chini et al., 2006).

It is well established that the colonizing strains of *S. aureus* are the ones that usually cause infection in surgical patients. Therefore, it is important to know what proportion of MRSA colonizing surgical patients in our geographic region belongs to community clones and what are the characteristics of the strains colonizing this patient population. Herein, we report the molecular typing and the antimicrobial susceptibility patterns of MRSA isolates from surgical patients screened upon admission to surgical wards.

Consecutive patients admitted to a surgical unit of a tertiary care hospital in Athens, Greece, between January and July 2009, were screened within 48 h of admission for MRSA carriage. The study was approved by the Institutional Review Board of Laiko General Hospital. The nares and inguinal area were sampled with a sterile swab, and

samples were inoculated onto BBL CHROMagar MRSA II medium. Isolates were classified as CA-MRSA if patients did not have surgery, hospitalization, or hemodialysis within the previous year; did not reside in a chronic-care facility during the previous year; did not have indwelling catheters at the time of sampling; and had no previous isolation of MRSA, according to Centers for Diseases Control and Prevention epidemiologic definition (Klevens et al., 2007). Antimicrobial susceptibility to ciprofloxacin, erythromycin, clindamycin, fusidic acid, co-trimoxazole, gentamicin, tetracycline, and rifampin was determined for all CA-MRSA isolates, by disk-diffusion testing, using Kirby–Bauer disks and interpreted according to Clinical and Laboratory Standards Institute criteria (CLSI, 2011). Genomic DNA for polymerase chain reaction analysis was prepared using the QIAamp DNA Mini and Blood Mini kit (Qiagen, Hilden, Germany). All MRSA isolates were examined for the presence of Panton–Valentine leukocidin (PVL) and *mecA* genes as previously described (Krziwanek et al., 2007). SCCmec typing and subtyping, *spa* typing, and multilocus sequence typing (MLST) were performed for all CA-MRSA isolates and for a selected number of HA-MRSA as described previously (Enright et al., 2000; Harmsen et al., 2003; Milheirico et al., 2007; Zhang et al., 2005). The amplicons were purified, and sequence analysis was performed with the BigDye Terminator kit v.3.1 (Applied Biosystems, CA, USA) using the 3730 DNA Analyser (Applied Biosystems, CA, USA), Biogenomica, Athens, Greece. DNA alignment was performed with the BLAST software (<http://www.ncbi.nlm.nih.gov>). The sequence types (ST) of MRSA isolates were assigned through the MLST database (<http://saureus.mlst.net/sql/multiplelocus.asp>), and the *spa* types were determined using the BioNumerics Demo Web Server (<http://bnas.applied-maths.com/spaupload.aspx>). All MRSA-colonized patients were followed prospectively for development of infection. The infection was presumed to be due to the same strain if the colonizing and infecting isolates had the same ST, *spa* type, and SCCmec type or subtype.

A total of 925 patients (51% males; median age, 62.8 years) were screened, of whom 51 (5.5%) were positive for MRSA at the time of admission. Fifteen (29.4%) of 51 MRSA isolates fulfilled the epidemiologic criteria for CA-MRSA of which 9 belonged to community MRSA clones and 6 to hospital MRSA clones. The community clones were ST80-IVc ($n = 7$) and ST30-IVc ($n = 2$), whereas the hospital clones were ST239-III ($n = 2$), ST5-II ($n = 2$), ST105-II ($n = 1$), and ST22-IVc ($n = 1$) as shown in the Table 1. Of note, one of the PVL-negative

community MRSA clones (ST30-IVc) was of a novel *spa* type, t8767 (<http://spa.ridom.de/spatypes.shtml>). Of 36 HA-MRSA isolates, as designated by epidemiologic criteria, 7 of the 8 PVL-positive isolates that were analyzed belonged to the community MRSA clone, ST80-IVc (Table 1). The remaining 28 PVL-negative HA-MRSA isolates were examined only for the presence of SCCmec elements: 4 isolates carried the SCCmec II element; 6, the SCCmec III; 13, the SCCmec IV; 2, the SCCmec V; 2, the SCCmec VI; and 1 was not typable.

Similarly with a previous report (Katopodis et al., 2010), the most common antimicrobial resistance pattern observed in the ST80-IVc clone was resistance to fusidic acid and tetracycline. Three of the 14 PVL-positive isolates of the ST80-IVc genetic lineage exhibited resistance to 3 or more non- β -lactam classes of antibiotics (Table 1; isolate nos. 6, 7, and 21). As expected, the isolates belonging to hospital MRSA clones (Table 1; isolates nos. 11, 15, 17–19) exhibited more extensive antimicrobial resistance profile. Two of the patients colonized with the ST80-IVc community clone at the time of admission (Table 1; patient nos. 1 and 24) developed clinical infection by the same MRSA strain during their hospital stay.

This work provides important information on the molecular characteristics of MRSA that had colonized patients prior to admission to a surgical unit of a Greek tertiary care hospital. At least one-third of MRSA-colonizing isolates belonged to the ST80-IVc clone, indicating that this clone is the most predominant among the community MRSA clones circulating in our region, as it seems to be in most European countries (Otter and French, 2010). Moreover, 2 patients colonized with this particular clone developed clinical infection by the same strains, as was evidenced by the identical ST, *spa* type, and SCCmec of the respective colonizing and infecting isolates. These findings suggest that the ST80-IVc clone not only is an epidemiologically successful clone but also has the potential to become a nosocomial pathogen. In view of the spread of community MRSA clones in the healthcare setting and the probable diffusion of hospital clones to the community, the molecular typing of MRSA isolates appears to be not so useful a method for designating MRSA isolates as community or hospital acquired.

More worrisome is the emergence of multidrug resistance in several community MRSA clones and particularly in the ST80-IVc clone. This is in accordance with other studies that have observed multiresistant isolates belonging to the ST80 lineage (Hanssen et al.,

Table 1
Genetic characteristics and phenotypes of MRSA colonizing surgical patients.

No.	PVL	ST	SCCmec	<i>spa</i>	Ciprofloxacin	Erythromycin	Clindamycin	Fusidic acid	Co-trimoxazole	Gentamicin	Tetracycline	Rifampin
<i>CA-MRSA isolates</i>												
1	+	80	IVc	t044	S	S	S	R	S	S	R	S
2	+	80	IVc	t044	S	S	S	R	S	S	R	S
3	+	80	IVc	t044	S	S	S	R	S	S	R	S
4	+	80	IVc	t044	S	S	S	R	S	S	R	S
5	+	80	IVc	t044	S	S	S	R	S	S	R	S
6	+	80	IVc	t044	S	R	R	R	S	S	R	S
7	+	80	IVc	t131	R	R	R	R	S	S	R	S
8	+	30	IVc	t8026	S	S	S	S	S	S	S	S
11	–	239	III	t421	R	R	R	R	R	S	R	S
12	–	22	IVc	t005	S	S	S	S	S	S	S	S
13	–	30	IVc	t8767	S	S	S	S	S	S	S	S
15	–	105	II	t002	R	R	R	R	S	S	S	S
17	–	5	II	t002	R	R	R	S	R	S	S	S
18	–	5	II	t002	R	R	R	S	R	S	S	S
19	–	239	III	t1152	R	R	R	R	R	R	R	S
<i>HA-MRSA isolates</i>												
20	+	80	IVc	t044	S	S	S	R	S	S	R	S
21	+	80	IVc	t044	S	S	S	R	S	R	R	S
22	+	80	IVc	t044	S	S	S	R	R	S	R	S
23	+	80	IVc	t131	S	S	S	R	S	S	R	S
24	+	80	IVc	t044	S	S	S	R	S	S	R	S
25	+	80	IVc	t044	S	S	S	R	S	S	R	S
26	+	80	IVc	t044	S	S	S	R	S	S	R	S

PVL = Panton–Valentine leukocidin; ST = multilocus sequence type; S = susceptible; R = resistant.

2005; Katopodis et al., 2010). In Greece, one of the most common resistance patterns exhibited by the ST80-IV clone is resistance to fusidic acid, tetracycline, erythromycin, and clindamycin (Katopodis et al., 2010). Notably, in the present report, 1 isolate belonging to the ST80-IVc clone and to a unique *spa* type (t131) expressed a further increase in its resistance repertoire including 5 different classes of non- β -lactam antibiotics, namely, quinolones, macrolides, clindamycin, fusidic acid, and tetracyclines.

In conclusion, a significant proportion of the MRSA colonizing surgical patients in our geographic region belongs to the epidemiologically successful ST80-IVc European clone which has the potential to become multidrug resistant and cause nosocomial infections. These findings raise serious concerns and cause new challenges in infection control practitioners and clinicians.

Acknowledgment

The authors would like to acknowledge the help of Andie Lee for the setup of the study.

References

- Aires de Sousa M, Bartzavali C, Spiliopoulou I, Santos Sanches I, Crisóstomo MI, de Lencastre H. Two international methicillin-resistant *Staphylococcus aureus* clones endemic in a university hospital in Patras, Greece. *J Clin Microbiol* 2003;41:2027–32.
- Chini V, Petinaki E, Foka A, Paratiras S, Dimitracopoulos G, Spiliopoulou I. Spread of *Staphylococcus aureus* clinical isolates carrying Panton–Valentine leukocidin genes during a 3-year period in Greece. *Clin Microbiol Infect* 2006;12:29–34.
- Clinical and Laboratory Standards Institute (CLSI). Performance standards for antimicrobial susceptibility testing: 21st informational supplement (CLSI document M100-S21). Wayne, PA: CLSI; 2011.
- David MZ, Daum RS. Community-associated methicillin-resistant *Staphylococcus aureus*: epidemiology and clinical consequences of an emerging epidemic. *Clin Microbiol Rev* 2010;23:616–87.
- Ellington MJ, Ganner M, Warner M, Cookson BD, Kearns AM. Polyclonal multiply antibiotic-resistant methicillin-resistant *Staphylococcus aureus* with Panton–Valentine leukocidin in England. *J Antimicrob Chemother* 2009;65:46–50.
- Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J Clin Microbiol* 2000;38:1008–15.
- Francois P, Harbarth S, Huyghe A, et al. Methicillin-resistant *Staphylococcus aureus*, Geneva, Switzerland, 1993–2005. *Emerg Infect Dis* 2008;14:304–7.
- Hanssen AM, Fossum A, Mikalsen J, Halvorsen DS, Bukholm G, Ericson Sollid JU. Dissemination of community-acquired methicillin-resistant *Staphylococcus aureus* clones in northern Norway: sequence types 8 and 80 predominate. *J Clin Microbiol* 2005;43:2118–24.
- Harmsen D, Claus H, Witte W, Rothganger J, Turnwald D, Vogel U. Typing of methicillin-resistant *Staphylococcus aureus* in a university hospital setting by using novel software for *spa* repeat determination and database management. *J Clin Microbiol* 2003;41:5442–8.
- Katopodis GD, Grivea IN, Tsantsaridou AJ, Pournaras S, Petinaki E, Syrogiannopoulos GA. Fusidic acid and clindamycin resistance in community-associated, methicillin-resistant *Staphylococcus aureus* infections in children of central Greece. *BMC Infect Dis* 2010;10:351–9.
- Klevens RM, Morrison MA, Nadle J, et al. Invasive methicillin-resistant *Staphylococcus aureus* infections in the United States. *JAMA* 2007;298:1763–71.
- Krzywanek K, Luger C, Sammer B, et al. Pvl-positive MRSA in Austria. *Eur J Clin Microbiol Infect Dis* 2007;26:931–5.
- Milheirico C, Oliveira DC, de Lencastre H. Update to the multiplex PCR strategy for assignment of *mec* element types in *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2007;51:3374–7.
- Niniou I, Vourli S, Lebesse E, et al. Clinical and molecular epidemiology of community-acquired, methicillin-resistant *Staphylococcus aureus* infections in children in central Greece. *Eur J Clin Microbiol Infect Dis* 2008;27:831–7.
- Otter JA, French GL. Molecular epidemiology of community-associated methicillin-resistant *Staphylococcus aureus* in Europe. *Lancet Infect Dis* 2010;10:227–39.
- Popovich KJ, Weinstein RA, Hota B. Are community-associated methicillin-resistant *Staphylococcus aureus* (MRSA) strains replacing traditional nosocomial MRSA strains? *Clin Infect Dis* 2008;46:787–94.
- Vourli S, Vagiakou H, Ganteris G, et al. High rates of community-acquired, Panton–Valentine leukocidin (PVL)-positive methicillin-resistant *S. aureus* (MRSA) infections in adult outpatients in Greece. *Euro Surveill* 2009;14.
- Zhang K, McClure JA, Elsayed S, Louie T, Conly JM. Novel multiplex PCR assay for characterization and concomitant subtyping of staphylococcal cassette chromosome *mec* types I to V in methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 2005;43:5026–33.