

Rare genetic variant register

A rare genetic variant of the T cell receptor gamma joining segment TRGJ1

M. Weill, X.M. Zhang, F. Bernard, G. Lefranc, and M.-P. Lefranc

Laboratoire d'Immunogénétique Moléculaire, LIGM, URA CNRS 1191, Université Montpellier II, UFR Sciences, CC012, Place E. Bataillon, F-34095 Montpellier Cedex 5, France

Received June 11, 1991 / Revised July 5, 1991

Summary. A rare *Hind*III restriction fragment variant of the human T cell receptor gamma joining segment TRGJ1 has been identified, in addition to two previously described alleles.

Source/description. The TRG1.4 probe (Bensmana et al. 1991) is a genomic 1450-bp *Eco*RI-*Hind*III fragment cloned in pUC18, and located between the TRGJ1 joining segment and the exon 1 of TRGC1 (Lefranc and Rabbitts 1985; Lefranc et al. 1986). This probe cross-hybridizes to the corresponding region located between TRGJ2 and TRGC2.

Polymorphism. *Hind*III (A/AGCTT) identifies three allelic restriction fragments of 2.1 kb (*A1), 5.0 kb (*A2) and 3.7 kb (*A3). Constant band: 2.1 kb.

Frequency. *A1: 0.64, *A2: 0.35, *A3: 0.01, as studied in 52 unrelated Caucasoids; the variant was not polymorphic for *Eco*RI (5 kb + 7 kb), *Bam*HI (20 kb + 13 kb), *Kpn*I (9 kb + 16 kb), *Sac*I (6.4 kb + 7.0 kb) studied in 25 unrelated individuals. The two constant bands observed with these four enzymes correspond to the TRGC1 and TRGC2 regions, respectively.

Chromosomal localisation. The human TRG locus has been mapped on chromosome 7 (Rabbitts et al. 1985) at band p14-p15 (Murre et al. 1985; Bensmana et al. 1991).

Mendelian Inheritance. Co-dominant segregation was demonstrated in five families.

Probe availability. Available from Dr. M.-P. Lefranc.

Offprint requests to: M.-P. Lefranc

Other comments. The two alleles, *A1 (2.1 kb) and *A2 (5.0 kb), have been previously described using the TRGJ1 probe, pH60 (Lefranc and Rabbitts 1985; Lefranc et al. 1986; Buresi et al. 1989). In this report, we describe a third allele, *A3 (3.7 kb), corresponding to the absence of the *Hind*III site located 160 bp upstream of TRGJ1. The location of this polymorphic *Hind*III site 5' of TRGJ1 does not interfere with the sizes of the V γ -J γ rearranged bands and therefore with the assignment of the given V γ genes and J γ segments, as described by Forster et al. (1987).

References

- Bensmana M, Mattei M-G, Lefranc M-P (1991) Localization of the human T-cell receptor gamma locus (TCRG) to 7p14-p15 by in situ hybridization. *Cytogenet Cell Genet* 56:31-32
- Buresi C, Ghanem N, Huck S, Lefranc G, Lefranc M-P (1989) Exon duplication and triplication in the human T-cell receptor gamma constant region genes and RFLP in French, Lebanese, Tunisian, and Black African populations. *Immunogenetics* 29:161-172
- Forster A, Huck S, Ghanem N, Lefranc M-P, Rabbitts TH (1987) New subgroups in the human T-cell rearranging V γ gene locus. *EMBO J* 6:945-950
- Lefranc M-P, Rabbitts TH (1985) Two tandemly organized human genes encoding the T-cell gamma constant region sequences show multiple rearrangements in different T-cell types. *Nature* 316:464-466
- Lefranc M-P, Forster A, Rabbitts TH (1986) Genetic polymorphism and exon changes of the constant regions of the human T-cell rearranging gene γ . *Proc Natl Acad Sci USA* 83:9596-9600
- Murre C, Waldmann RA, Morton CC, Bongiovanni KF, Waldmann TA, Shows TB, Seidman JG (1985) Human γ -chain genes are rearranged in leukaemic T-cells and map to the short arm of chromosome 7. *Nature* 316:549-552
- Rabbitts TH, Lefranc M-P, Stinson MA, Sims JE, Schroder J, Steinmetz M, Spurr NL, Solomon E, Goodfellow PN (1985) The chromosomal location of the T-cell receptor genes and a T-cell rearranging gene: possible correlation with specific translocations in human T-cell leukaemia. *EMBO J* 4:1461-1465