

Proclivity of Activated Neutrophils to Cause Postischemic Cardiac Dysfunction: Participation in Stunning?

Kevin Mullane and Robert Engler

*Gensia Pharmaceuticals, Inc., San Diego, CA, and VAMC,
Department of Medicine, University of California San Diego,
La Jolla, CA, USA*

Summary. Myocardial stunning is a reversible defect in contractile function provoked by brief episodes of ischemia followed by reperfusion. Many studies have demonstrated the potential involvement of free radicals in the etiology of myocardial stunning. While activated neutrophils have the capacity to release free radicals and evoke contractile dysfunction, it is not clear that this potential is realized in the absence of myocellular damage. Attempts to define the contribution of activated neutrophils to myocardial stunning by removing the cells from the bloodstream are contradictory, and the apparent simplicity of this seemingly logical approach is an illusion. For example, it is not known how many neutrophils are required to induce contractile failure, the site of action within the heart, the mechanisms that may be responsible, or even the time course or process of neutrophil activation. The production of free radicals and endothelial dysfunction may create conditions propitious for neutrophil recruitment. However, because activated neutrophils synthesize and release various mediators that are potentially toxic to myocardium, once the stage is reached for leukocyte accumulation, it may herald the progression from reversible to irreversible cardiac injury.

Cardiovasc Drugs Ther 1991;5:915-924

Key Words. myocardial stunning, neutrophils, free radicals, endothelial dysfunction

The advent of new techniques to restore coronary blood flow during acute myocardial ischemia has led to the recognition that the recovery of pump function is sluggish, even if the myocardium is still viable, a situation termed the *stunned myocardium* [1]. This review will adhere to the definition of myocardial "stunning," as outlined by Bolli [2,3], as a reversible defect in contractile function observed after a period of ischemia, in the absence of overt myocellular damage or necrosis. The distinction between sublethal or reversible injury and irrevocable cell death is important because, while both situations are associated with mechanical derangements, the mechanisms of contractile dysfunction are probably quite different. The

background to the elucidation and recognition of the stunned myocardium has recently been reviewed [3].

Evidence for an important (but not exclusive) role of oxygen-derived free radicals and metabolites in the pathogenesis of myocardial stunning has been elegantly presented by Bolli and colleagues and other investigators [1-3]. It is natural to consider the potential for neutrophil activation as a contributory factor to myocardial stunning, because on a quantitative basis neutrophils are a major potential source of oxygen metabolites. However, free-radical-mediated injury and neutrophil-mediated injury are not necessarily synonymous, and it may be erroneous to equate the two events.

Postischemic contractile abnormalities have been described in isolated buffer-perfused hearts subjected to global ischemia [5-8]. Although it is not clear that this in vitro response mimics myocardial stunning in vivo where the time for contractile function to recover generally takes much longer [9]; nevertheless, it corresponds to a temporary postischemic mechanical abnormality. The in vitro model of stunning is accompanied by a burst of free-radical/oxygen metabolite formation within the first few seconds to minutes of reperfusion [7,10,11], and is ameliorated by various free-radical scavengers [6,7]. Consequently, this response shares many features of myocardial stunning in vivo, and it indicates that neutrophils are not an essential component of some aspects of the contractile defect because it occurs in the absence of any blood elements. Thus a more pertinent issue is whether the activation of neutrophils can exacerbate postischemic cardiac dysfunction in vivo, rather than account for the complete response. Clearly there are other

Author for correspondence and reprint requests: Kevin Mullane, Gensia Pharmaceuticals, Inc., 11025 Roselle Street, San Diego, CA 92121, USA.

sources of free radicals, such as the vascular endothelium [12–14] or myocyte mitochondria [15,16] and additional mechanisms that also contribute to the ultimate response [3].

In reviewing the contribution of activated neutrophils to myocardial stunning, it is apparent that the evidence is fragmentary and incomplete. Many studies have relied upon techniques to deplete neutrophils from the blood and compare the postischemic contractile failure to a control group with neutrophils. Unfortunately, the apparent simplicity of this logic is illusory, and the results are inconsistent, perhaps due, in part, to the inherent difficulty in producing true tissue neutropenia. Therefore it is necessary to examine the issue in a broader context of whether neutrophils have the capacity to influence contractile function, the mechanisms involved, and whether there is a basis to consider these mechanisms as contributing to myocardial stunning. Moreover, myocardial stunning has been described in a variety of settings; after ischemia induced by a single brief occlusion, multiple occlusions or global ischemia during cardioplegic arrest, and during exercise or demand ischemia [3]. It is not inconceivable that differences between the various models may reflect the interplay of different components of injury such that generalized conclusions could be misleading. Recognition of a component of injury attributable to activated neutrophils or other inflammatory cells may be important because this response can become self-perpetuating. Recruitment of inflammatory cells may herald the potential for a reversible injury to be “pushed over the edge” in a transition to lethal damage, due to the variety of injurious mediators produced by activated neutrophils.

Can Neutrophils Impair Contractile Function?

Quiescent or inactivated neutrophils do not release enzymes and mediators, and apparently do not influence resting cardiac function. Perfusion of isolated hearts with diluted blood or buffer neutrophils does not alter baseline contractile function [8,17,18], despite the long transit times of leukocytes through the microcirculation [19] and the potential for diminished perfusion. This is probably because of substantial capillary reserve in the heart, where it is suggested that only 50–60% capillaries are perfused during normal cardiac function [20,21]. Moreover, removal of leukocytes from blood perfusing hearts *in situ* is also without effect on baseline cardiac function [22–25]. Changes in contractile function elicited by neutrophils could be secondary to intravascular se-

questration or entrapment, thereby causing perfusion defects and local ischemia, or by neutrophil-derived products directly depressing contractile function. Consequently, Kraemer and coworkers [26,27] studied the interaction of neutrophils with papillary muscles mounted in organ baths and devoid of a vascular component. In this setting, where the direct influences of neutrophils on contractility can be assessed, nonstimulated cells do not impair cardiac function. Thus, resting neutrophils do not alter myocardial contractility *in vitro* or *in vivo*.

Activation of neutrophils by chemotactic factors leads to the production and/or release of a host of different mediators, many of which could influence myocardial contractility. These mediators include lipid peroxidation products of arachidonic acid [28]; 1-*O*-alkyl-2-aryl-sn-glycerol-3-phosphorylcholine, or platelet activating factor (PAF), which promotes contractile dysfunction either directly [29] or through a mechanism dependent upon the presence of blood elements [30]; metabolites of oxygen including superoxide anions (O_2^-), hydrogen peroxide (H_2O_2), and hypochlorite anions ($HOCl^-$), each of which has been identified as a negative inotropic species [7,26,31,32]; and proteases such as collagenase and elastase, which may compromise structural and cellular integrity [33,34] to impair cardiac function.

Although activated neutrophils elaborate multiple mediators capable of compromising contractility, the question remains as to whether this potential is actually realized in a dynamic setting of cell-tissue interactions. This was addressed directly by the addition of neutrophils to isolated papillary muscles and monitoring changes in contractile function [26,27]. Activated neutrophils provoked a concentration-dependent decline in papillary muscle function, which developed over 5–10 minutes (Figure 1). PAF, leukotrienes, superoxide anions, or hypochlorite anions were excluded from mediating this response. The decrease in function was attributed to the release of hydrogen peroxide because catalase attenuated the neutrophil-induced contractile derangement while exogenous hydrogen peroxide mimicked the effect [26]. Monoclonal antibodies to the CD18 adhesion complex of the neutrophil abolished contractile failure provoked by activated neutrophils [27]. Adherent neutrophils also produce substantially more H_2O_2 , and for longer periods of time, than cells in suspension [35]. Prevention of adhesion with monoclonal antibodies to the CD11/CD18 adhesion complex suppressed this augmented H_2O_2 release [36] and could account for the beneficial effects of the antiadhesion antibodies.

Activated neutrophils also adhere to isolated myocytes that have been primed with a cytokine such as




Fig. 1. A: Time-dependent reduction in rabbit papillary muscle function by human neutrophils (PMNs) activated with human recombinant (hr) C5a (●---●, $n = 18$), while either PMNs (□---□) or hrC5a (○---○) alone were without effect. Data from Kraemer et al., [27]. ** $p < 0.01$, *** $p < 0.001$, repeated measures ANOVA. **B:** Concentration-dependent re-

duction in rabbit papillary muscle function by the cumulative addition of rabbit neutrophils (□---□, $n = 18$) versus time-related control papillary muscles in the absence of PMNs (○---○, $n = 10$). The response was allowed to develop over 10 minutes before the next concentration of cells was added. Data from Kraemer et al. [26]. * $p < 0.01$

interleukin-1 or tumor necrosis factor, and express intercellular adhesion molecule-1 (ICAM-1) [36]. Adhesion is mediated by an interaction between CD18 and ICAM-1, and results in the generation and release of H_2O_2 by the neutrophils, which is associated with contracture of the myocytes [36]. Thus, neutrophils have the capacity to interact directly with myocytes. These studies support the suggestion that activated neutrophils can release mediators in amounts sufficient to provoke cardiac dysfunction. The nature of this functional defect—whether it reflects overt damage of the endocardial covering of the muscle, as has been described with activated eosinophils [37]; oxidant-induced injury to the myocytes, perhaps interfering with calcium transport [38], or by compromising either cellular integrity or the contractile apparatus by interactions between oxidants and neutrophil-derived proteases [33]—remains to be elucidated. Importantly, it is not clear if the functional derangement reflects a reversible phenomenon, which is mandatory for the definition of stunning, or it represents irreversible overt damage to the muscle, and thus, a different phenomenon. In this context it should be noted that isolated myocytes with attached neutrophils acquired the rounded configuration of damaged cells [36].

Neutrophils and Coronary Vascular Resistance

Another means by which neutrophil activation could impair myocardial contractility is indirectly via an effect on coronary perfusion. Isolated hearts perfused with a buffer containing neutrophils exhibit vasoconstriction accompanied by a decrease in left ventricular developed pressure when the neutrophils are stimulated with phorbol myristate acetate [39]. The vasoconstriction was attributed to the production of oxygen metabolites by the neutrophils [39]. Subjecting the neutrophil-perfused heart to a period of hypoxia followed by reoxygenation leads to neutrophil activation and tissue sequestration [8,17], accompanied by vasoconstriction during hypoxia and compromised recovery of contractile function upon reoxygenation [8]. Isolated rat hearts perfused with blood diluted 1:1 with Krebs' buffer and subjected to 30 minute isch-

emia showed a 62% decrease in capillary perfusion and a 76% increase in coronary vascular resistance at reperfusion, whereas perfusion with blood depleted of leukocytes gave a 33% reduction in perfused capillaries and only a 5% increase in vascular resistance [18]. The increased vascular resistance associated with whole blood perfusion was attributed to microvascular plugging and/or vasoconstrictor production by the neutrophils [18]. Thus, various studies using isolated perfused hearts concluded that activated neutrophils could impair coronary perfusion, and the diminished perfusion was associated with a decline in contractile function. However, these studies do not permit differentiation between contractile failure secondary to inadequate perfusion and a direct negative inotropic effect of neutrophils or neutrophil-derived mediator(s).

Capillary plugging

Neutrophil entrapment within the microcirculation has been observed in the reperfused myocardium *in vivo* and is suggested to account for the capillary no-reflow phenomenon, where flow is not restored uniformly to the postischemic region [40], while leukocyte depletion reduces the extent of no reflow [41,42]. However, no reflow only occurs within regions of the myocardium that are irreversibly injured [43], so by definition is unlikely to be a significant factor in the stunned myocardium. Furthermore, 12 minutes of coronary occlusion followed by 1 hour of reperfusion was associated with a paradoxical 40% decrease in myocardial neutrophil content [44], indicating capillary entrapment does not occur in this setting.

Neutrophil-mediated vasoconstriction

The ability of neutrophils to influence vascular tone directly has been addressed by adding neutrophils to organ baths containing rings of vascular tissue [45–50]. The neutrophil-induced vasoconstriction that is observed upon activation of the leukocytes has been variously attributed to oxygen metabolites, in particular, superoxide anions, which degrade endothelial-derived relaxing factor [EDRF: 47], the transcellular metabolism of neutrophil-derived leukotriene A_4 to leukotriene C_4 by the endothelium [49]; or the release of an unidentified peptide [50]. Stimulated neutrophils release a vasoconstrictor principle that is heat stable and produces contraction of vascular smooth muscle in the absence of neutrophils [50]. This material also augments platelet-mediated contractions [50], attributed primarily to serotonin and thromboxane A_2 [51,52]. Isolated, coronary arteries perfused *ex vivo* after being subjected to 15 minutes hypertension *in vivo* to induce neutrophil and platelet adhesion to the endothelium, exhibited a fourfold increase in sensitiv-

ity to the vasoconstrictor effects of serotonin that was reversed by a combination of thromboxane and leukotriene receptor blockade [56]. Thus, activated neutrophils can promote vasoconstriction directly by the release of vasoactive mediators, and indirectly by potentiating the activities of other vasoconstrictor agents. The significance of these neutrophil-mediated changes in vascular tone are unknown at this time, but if occurring locally in the reperfused myocardium might create areas of hypoperfusion, which, in turn, would promote regional wall-motion abnormalities.

Vascular stunning

While dramatic vasoconstriction is not observed in stunned myocardium, resting subendocardial blood flow is diminished after a single 15-minute ischemic episode followed by reperfusion [9,53,54]. In this setting, the reative hyperemic response following a 40-second occlusion and the maximal vasodilator responses to adenosine or papaverine are also impaired, giving rise to the term *microvascular stunning* [54]. These vascular derangements do not correlate with the contractile dysfunction [54], and it is unlikely that changes in coronary tone contribute to the functional derangement. Moreover, changes in vascular function are not apparent after 10 minutes of ischemia and reperfusion [55], suggesting that different models of stunning, while sharing a common defect, may reflect different events that culminate in injury. Vascular stunning and a diminished coronary vascular reserve is apparently associated with more “severe” forms of myocardial stunning and may represent the beginning of a transition into lethal injury. The possibility that defects in subendocardial perfusion and vascular reserve are linked to neutrophil recruitment has not been addressed.

Neutrophils and Contractile Failure in Vivo

Few studies have addressed directly whether the local activation of neutrophils *in vivo* is accompanied by contractile dysfunction. Based on the apparent importance of components of complement as neutrophil chemoattractants in the setting of myocardial ischemia and reperfusion [57,58], the chemotactic complement fragment C5a was injected into the coronary artery of anesthetized open-chest pigs [55–58]. C5a transiently evoked coronary vasoconstriction, compromised contractile function, and promoted myocardial leukocyte sequestration [59,60]. The derangements in cardiac function were attributed to the peptido leukotrienes, such as leukotriene D_4 , and to thromboxane A_2 , mediators released by C5a, because

the response was ameliorated by a combination of antagonists to these eicosanoids [61]. These antagonists did not attenuate the leukocyte accumulation in the heart [61], and the leukocytes and/or other blood elements have been proposed as the source of the vasoconstrictor material [61,62]. However, myocardial leukocyte uptake could be an epiphenomenon of unknown significance. Leukotriene B₄ is another chemotactic agent that induced equivalent leukocyte sequestration to C5a when injected into a coronary artery, but did not cause vasoconstriction or contractile dysfunction [63]. Moreover, while neutrophil depletion with filters prevented vasoconstriction in response to C5a, hearts previously perfused with filtered blood remained refractory to the C5a-induced thromboxane A₂ production and vasoconstriction, even when perfused with whole blood never exposed to filters [64]. Pigs rendered severely neutropenic with cyclophosphamide were equisensitive to C5a-induced ischemia as normal pigs [65]. Therefore cells within the heart, rather than circulating blood elements, appear to be responsible for the response to C5a, and neutrophil trapping is an epiphenomenon for the acute response. The source(s) of the vasoconstrictor material elicited by C5a remain to be defined and could be tissue leukocytes (i.e., macrophages) that synthesize both leukotriene D₄ and thromboxane A₂, in contrast to the circulating neutrophils [66]. Because the predominant effect of C5a is on tissue cells, rather than the neutrophils, it remains to be determined if activation of leukocytes in blood perfusing the heart can promote contractile failure.

Does transient ischemia and reperfusion create conditions propitious for neutrophil activation?

The adhesion of neutrophils to the vascular endothelium is a prerequisite for their recruitment and accumulation in tissues. The site specificity of neutrophil adhesion implies that it results from a change in the status or function of the endothelium. Normally, endothelial cells elaborate and release various mediators that oppose neutrophil adhesion, including EDRF or nitric oxide [66], prostacyclin, and adenosine [67–69]. Although endothelial cells are relatively resistant to ischemic damage [70], endothelial dysfunction expressed as an increased permeability to macromolecules and impaired relaxations to agonists that operate through the release of EDRF, has been described after 15 minutes of ischemia followed by reperfusion [71], a classical setting of myocardial stunning. The formation of oxygen metabolites at reperfusion may account, at least in part, for the impaired

endothelial-dependent relaxations, because they can be reproduced with exogenous free-radical generating systems such as xanthine-xanthine oxidase [50]. In addition, oxygen metabolites inhibit prostacyclin production [72] and deplete endothelial adenosine triphosphate [73], a major source of adenosine, while the administration of superoxide dismutase prior to reperfusion preserves endothelial function after longer periods of ischemia [74–76]. Thus, brief periods of ischemia followed by reperfusion promote endothelial dysfunction, with impaired production of various autacoids that normally attenuate neutrophil attachment.

Oxygen metabolites, in particular H₂O₂, not only provoke a loss of EDRF, PGI₂, and adenosine, but also induce endothelial cells to rapidly express GMP-140 [77] and PAF [78], two molecules that are intimately involved in neutrophil adhesion. Cooperative interactions between these adhesion molecules could lead to the early recruitment of neutrophils. In addition, H₂O₂ activates CD11b/CD18-dependent cell adhesion [79].

The significance of these functional changes of the endothelium to neutrophil activation and accumulation, or indeed to myocardial stunning, is not clear, but serves to highlight changes that could favor neutrophil recruitment. However, these changes alone may not be sufficient to promote neutrophil accumulation, and the production of an agent to activate the leukocytes may also be required.

To date, there is a paucity of direct evidence demonstrating the formation of a chemotactic agent after brief ischemic episodes. Preliminary studies in patients undergoing coronary-artery balloon-angioplasty induced ischemia for a duration of only 1.5 minutes was associated with neutrophil activation attributed to the local release of an undefined chemotactic substance(s) [80]. Using accumulation of Clq, a subunit of the first component of complement, in ischemic canine myocardium as a measure of complement activation and equating Clq localization with severity of ischemia, Rossen and colleagues [57] showed some local complement activation after 15 minutes of ischemia that did not achieve statistical significance ($p = 0.09$).

Electron paramagnetic spin resonance studies of either myocardial tissue [9] or spin-trap agents perfused through the heart [11,81,82] demonstrate free-radical production at reperfusion that is suppressed by the administration of superoxide dismutase and catalase [7,83]. Because some of the spin-trap agents used remain confined to the vascular compartment [82,83], it appears that oxygen metabolites such as superoxide anions may be released into the bloodstream. Petrone et al. [84] described the production of a neutrophil

chemotactic factor in plasma exposed to superoxide anions. This factor did not cause neutrophil degranulation or free-radical generation, but was suggested to account for the ability of superoxide dismutase to attenuate neutrophil accumulation at sites of inflammation [84]. The identity of this factor, or whether it is actually formed in situ, remains unknown. While these observations suggest that elements necessary for neutrophil activation and sequestration may be present in some situations that provoke myocardial stunning, assessments of III-indium-labeled neutrophils, as mentioned above, failed to show neutrophil accumulation after 12 minutes of ischemia and reperfusion in the anesthetized dog [44]. Indeed, a paradoxical decrease in myocardial neutrophil content was observed for reasons that are not clear.

Neutrophil Depletion and Myocardial Stunning

Four studies have addressed the influence of neutrophil depletion on myocardial stunning in the anesthetized dog, using either filters to remove cells from blood entering the coronary circulation [22,24,25] or antiserum to canine neutrophils to effect a systemic neutropenia [23]. A further study used $f(ab')_2$ fragments of a monoclonal antibody to the CD11b/CD18 adhesion complex on the leukocyte to prevent neutrophil accumulation [85]. In some cases a reduction in the degree of stunning was noted [22,25], while no benefit was observed in other studies [23,24,85]. While it can be argued that differences in dogs, diet, severity of ischemia, anesthesia, specificity of cell depletion etc. could account for some of these differences, two other points may be particularly noteworthy. The first pertains to the number of activated neutrophils that are required to provoke cardiac dysfunction. In vitro experiments [8,26] and the in vivo study of Engler and Covell [22] suggest that activation of only a few neutrophils may be necessary to evoke contractile dysfunction. Since the techniques to remove neutrophils are incomplete and probably activate the cells, the remaining cells may exert a greater influence than would be anticipated on a quantitative basis, and obscure interpretation of the studies. It should be noted that the absolute number of neutrophils in the tissue may not relate to leukocyte-mediated injury. Rather, the microvascular location and the state of activation may be more important determinants [86,87].

The second issue is that both studies that reported a beneficial effect of neutrophil depletion used Leukopak® filters to remove the cells [22,25], while Jeremy and Becker [24] used Imugard IG500 filters. Therefore it is appropriate to ask if the Leukopak® filters

somehow confer a protective effect on the postischemic myocardium independent of their ability to remove neutrophils. This has not and cannot now be addressed since Leukopak® filters are no longer available. However, it has been noted that leukocyte depletion with Pall filters desensitizes the heart to the coronary vasoconstrictor and negative inotropic actions of C5a, which persist during subsequent perfusion with normal blood, and are thus independent of leukocyte depletion [64]. These findings indicate that filtered blood confers some ill-defined protective effect on the heart. This is potentially important because it suggests that there is an endogenous factor(s) that can be "activated" or "formed" within the tissue to provide persistent protection against the ischemia provoked by C5a. Moreover, because filtering blood solely at reperfusion attenuates stunning [25], it indicates that any endogenous protectant can prevent injury from occurring after an ischemic event. This is an area that warrants further investigation.

Summary

The participation of neutrophils in myocardial stunning remains uncertain. While it appears that neutrophils have the potential to elicit contractile dysfunction, it is not evident that this potential is realized after transient ischemic periods not associated with myocellular necrosis. Other mechanisms may account for the beneficial effects of filtering the blood of neutrophils on the stunned myocardium. However, this review is filled with disclaimers and qualifiers indicating large gaps in our knowledge concerning the time course of neutrophil activation, number of cells necessary to elicit functional derangements, the site of action of neutrophils within the heart, the mechanisms responsible for neutrophil-induced dysfunction, etc. Merely depleting circulating blood of neutrophils is not an adequate means to define their contribution to postischemic contractile dysfunction, and different approaches to this issue are mandated.

References

1. Braunwald E. and Kloner RA. The stunned myocardium: Prolonged, postischemic ventricular dysfunction. *Circulation* 1982;66:1146-1149.
2. Bolli R. Oxygen-derived free radicals and postischemic myocardial dysfunction ("stunned myocardium"). *J Am Coll Cardiol* 1988;12:239-249.
3. Bolli R. Mechanism of myocardial "stunning." *Circulation* 1990;82:723-738.
4. Mullane KM. Oxygen-derived free radicals and reperfusion injury of the heart. In: Sonnecklick, Larach, Lesch, eds. *New frontiers in cardiovascular therapy: Focus on ACE inhibition*. 1989:234-270.
5. Ambrosio G, Jacobus WE, Bergman CA, et al. Preserved

- high energy phosphate metabolic reserve in globally "stunned" hearts despite reduction of basal ATP content and contractility. *J Mol Cell Cardiol* 1987;19:953-964.
6. Ambrosio G, Zweier JL, Jacobus WE, et al. Improvement of postischemic myocardial function and metabolism induced by administration of desferrioxamine at the time of reflow: The role of iron in the pathogenesis of reperfusion injury. *Circulation* 1987;76:906-915.
 7. Zweier JL. Measurement of superoxide-derived free radicals in the reperfused heart: Evidence for a free radical mechanism of reperfusion injury. *J Biol Chem* 1988;263:1353-1357.
 8. Kraemer R, Mullane KM. Neutrophils delay functional recovery of the posthypoxic heart of the rabbit. *J Pharm Exp Ther* 1989;251:620-626.
 9. Bolli R, Zhu WX, Thoraby JJ, et al. Time course and determinants of recovery of function after reversible ischemia in conscious dogs. *Am J Physiol* 1988;254:H102-H114.
 10. Zweier JL, Flaherty JT, Weisfeldt ML. Direct measurement of free radical generation following reperfusion of ischemic myocardium. *Proc Natl Acad Sci USA* 1987;84:1404-1407.
 11. Garlick PB, Davies MJ, Hearse DJ, Slater TF. Direct detection of free radicals in the reperfused rat heart using electron spin resonance spectroscopy. *Circ Res* 1987;61:757-760.
 12. Ratych RE, Chuknyiska RS, Bulkley GB. The primary localization of free radical generation after anoxia/reoxygenation in isolated endothelial cells. *Surgery* 1987;102:122-131.
 13. Zweier JL, Kuppusamy P, Luty GA. Measurement of endothelial cell free radical generation: Evidence for a central mechanism of free radical injury in postischemic tissues. *Proc Natl Acad Sci USA* 1988;85:4046-4050.
 14. Schinetti ML, Sbarbati R, Scarlattini M. Superoxide production by human umbilical vein endothelial cells in an anoxia-reoxygenation model. *Cardiovasc Res* 1989;23:76-80.
 15. Shlafer M, Myers CL, Adkins S. Mitochondrial hydrogen peroxide generation and activities of glutathione peroxidase and superoxide dismutase following global ischemia. *J Mol Cell Cardiol* 1987;19:1195-1206.
 16. Vandeplasche G, Hermans C, Thone F, Borgers M. Mitochondrial hydrogen peroxide generation by NADH-oxidase activity following regional myocardial ischemia in the dog. *J Mol Cell Cardiol* 1989;21:383-392.
 17. Gillespie MN, Kojima S, Owasojo JO, et al. Hypoxia provokes leukotriene-dependent neutrophil sequestration in perfused rabbit hearts. *J Pharm Exp Ther* 1987;241:812-816.
 18. Reynolds JM, McDonagh PF. Early in reperfusion, leukocytes alter perfused coronary capillary and vascular resistance. *Am J Physiol* 1989;256:HJ982-HJ989.
 19. Schmid-Schonbein GW, Engler RL. Granulocytes as active participants in acute myocardial ischemia and infarction. *Am J Cardiol Pathol* 1986;11:15-30.
 20. Martini J, Honig CR. Direct measurements of intercapillary distance in beating rat heart in situ under various conditions of O₂ supply. *Microvasc Res* 1969;1:244-256.
 21. Duran WN, Marsicano TH, Anderson RW. Capillary reserve in isometrically contracting dog hearts. *Am J Physiol* 1977;232:H276-H281.
 22. Engler R, Covell JW. Granulocytes cause reperfusion ventricular dysfunction after 15-minute ischemia in the dog. *Circ Res* 1987;61:20-28.
 23. O'Neill PG, Charlat ML, Michael LH, et al. Influence of neutrophil depletion on myocardial function and flow after reversible ischemia. *Am J Physiol* 1989;256:H341-H351.
 24. Jeremy RW, Becker LC. Neutrophil depletion does not prevent myocardial dysfunction after brief coronary occlusion. *J Am Coll Cardiol* 1989;13:1155-1163.
 25. Westlin W, Mullane KM. Alleviation of myocardial stunning by leukocyte and platelet depletion. *Circulation* 1989;80:1828-1836.
 26. Kraemer R, Seligmann B, Mullane KM. Polymorphonuclear leukocytes reduce cardiac function in vitro by release of H₂O₂. *Am J Physiol* 1990;258:H1847-H1855.
 27. Kraemer R, Smith CW, Mullane KM. Activated human polymorphonuclear leukocytes reduce rabbit papillary muscle function: Role of the CD18 glycoprotein adhesion complex. *Cardiovasc Res* 1991;25:172-175.
 28. Mullane KM, Westlin W, Kraemer R. Activated neutrophils release mediators that may contribute to myocardial injury and dysfunction associated with ischemia and reperfusion. *Ann NY Acad Sci* 1988;524:103-121.
 29. Levi R, Burke JA, Guo ZG, et al. Acetyl glyceryl ether phosphorylcholine (AGEPC) a putative mediator of cardiac anaphylaxis in the guinea pig. *Circ Res* 1984;54:117-124.
 30. Kenzora JL, Perez JE, Bergmann SR, Lange LG. Effects of acetyl glyceryl ether of phosphorylcholine (platelet activating factor) on ventricular preload, afterload, and contractility in dogs. *J Clin Invest* 1984;74:1193-1203.
 31. Schrier GM, Hess ML. Quantitative identification of superoxide anion as a negative inotropic species. *Am J Physiol* 1988;255:H138-H143.
 32. Myers CL, Weiss SJ, Kirsch MM, Shlafer M. Involvement of hydrogen peroxide and hydroxyl radical in the "oxygen paradox": Reduction of creatine kinase release by catalase, allopurinol or deferoxamine, but not by superoxide dismutase. *J Mol Cell Cardiol* 1985;17:675-684.
 33. Weiss SJ. Tissue destruction by neutrophils. *N Engl J Med* 1989;320:365-376.
 34. Henson PM, Johnston RB. Tissue injury in inflammation: Oxidants, proteinases, and cationic proteins. *J Clin Invest* 1987;79:669-674.
 35. Nathan CF. Neutrophil activation on biological surfaces: Massive secretion of hydrogen peroxide in response to products of macrophages and lymphocytes. *J Clin Invest* 1987;80:1550-1560.
 36. Entman ML, Youker K, Shappell SB, et al. Neutrophil adherence to isolated adult canine myocytes: Evidence for a CD18-dependent mechanism. *J Clin Invest* 1990;85:1497-1506.
 37. Shah AM, Brutsaert DL, Meulemans AL, et al. Eosinophils from hypereosinophilic patients damage endocardium of isolated feline heart muscle preparations. *Circulation* 1990;81:1081-1088.
 38. Hearse DJ. Stunning: A radical re-view. *Cardiovasc Drugs Ther* 1991;5:853-876.
 39. Gillespie MN, Kojima S, Kunitomo M, Jay M. Coronary and myocardial effects of activated neutrophils in perfused rabbit hearts. *J Pharmacol Exp Ther* 1986;239:836-840.
 40. Engler R, Schmid-Schonbein GW, Pavelec RS. Leukocyte capillary plugging in myocardial ischemia and reperfusion in the dog. *Am J Pathol* 1983;111:98-111.
 41. Engler RL, Dahlgren MD, Morris DD, et al. Role of leukocytes in response to acute myocardial ischemia and reflow in dogs. *Am J Physiol* 1986;251:H314-H322.
 42. Litt MR, Jeremy RW, Weisman HF, et al. Neutrophil depletion limited to reperfusion reduces myocardial

- infarct size after 90 minutes of ischemia: Evidence for neutrophil-mediated reperfusion injury. *Circulation* 1989;80:1816-1827.
43. Ambrosio G, Weisman HF, Mannisi JA, Becker LC. Progressive impairment of regional myocardial perfusion after initial restoration of postischemic blood flow. *Circulation* 1989;80:1846-1861.
 44. Go LO, Murry CE, Richard VJ, et al. Myocardial neutrophil accumulation during reperfusion after reversible or irreversible ischemic injury. *Am J Physiol* 1988;255:H1188-H1198.
 45. Rimeie TJ, Sturm RJ, Adams LM, et al. Interaction of neutrophils with vascular smooth muscle: Identification of a neutrophil-derived relaxing factor. *J Pharmacol Exp Ther* 1988;245:102-111.
 46. Mullane KM, Sessa W. Neutrophil-induced modulation of local blood flow: Association with myocardial ischemia and reperfusion injury. In: Bevan JA, Majeurski H, Maxwell RA, Story DF, eds. *Vascular neuroeffector mechanisms* Washington D.C.: IRL Press, 1988:2889-298.
 47. Ohlstein EH, Nichols AJ. Rabbit polymorphonuclear neutrophils elicit endothelium-dependent contraction in vascular smooth muscle. *Circ Res* 1989;65:917-924.
 48. Mehta JL, Lawson DL, Nicholas WW, Mehta P. Modulation of vascular tone by neutrophils: Dependence on endothelial integrity. *Am J Physiol* 1989;257:H1315-H1320.
 49. Nishida M, Kuzuya T, Hoshida S, et al. Polymorphonuclear leukocytes induced vasoconstriction in isolated canine coronary arteries. *Circ Res* 1990;66:253-258.
 50. Sessa WC, Mullane KM. Release of a neutrophil-derived vasoconstrictor agent which augments platelet-induced contractions of blood vessels in vitro. *Br J Pharmacol* 1990;99:553-559.
 51. Cohen R, Shepherd JT, Vanhoutte PM. Inhibitory role of the endothelium in the response of isolated coronary arteries to platelets. *Science* 1983;221:273-274.
 52. Houston DS, Shepherd JT, Vanhoutte PM. Aggregating human platelets cause direct contraction and endothelium-dependent relaxation of isolated canine coronary arteries: Role of serotonin, thromboxane A₂, and adenosine nucleotides. *J Clin Invest* 1986;78:539-544.
 53. Heyndrickx GR, Baig H, Nellens P, et al. Depression of regional blood flow and wall thickening after brief coronary occlusions. *Am J Physiol* 1978;234:H653-H659.
 54. Bolli R, Triana JF, Jeroudi MO. Prolonged impairment of coronary vasodilation after reversible ischemia: Evidence for microvascular "stunning." *Circ Res* 1990;67:332-343.
 55. Jeremy RW, Stahl L, Gillinov M, et al. Preservation of coronary flow reserve in stunned myocardium. *Am J Physiol* 1989;256:H1303-H1310.
 56. Cappelli-Bigazzi M, Lamping KG, Nuno DW, Harrison DG. Leukocyte and platelet-derived factors augment canine coronary constriction to serotonin. *Am J Physiol* 1990;259:H1161-H1170.
 57. Rossen RD, Swain JL, Michael LH, et al. Selective accumulation of the first component of complement and leukocytes in ischemic canine heart muscle: A possible initiator of an extra myocardial mechanism of ischemic injury. *Circ Res* 1985;57:119-130.
 58. Weisman HF, Bartow T, Leppo MK, et al. Soluble human complement receptor type 1: In vivo inhibitor of complement suppressing post-ischemic myocardial inflammation and necrosis. *Science* 1990;249:146-151.
 59. Martin SE, Chenoweth DE, Engler RL, et al. Complement C5a decreases regional coronary blood flow and myocardial function in pigs: Implications for a granulocyte mechanism. *Circ Res* 1988;63:483-491.
 60. Ito BR, Roth DM, Chenoweth DE, et al. Thromboxane is produced in response to intracoronary infusions of complement C5a in pigs: Cyclooxygenase blockade does not reduce the myocardial ischemia and leukocyte accumulation. *Circ Res* 1989;65:1220-1232.
 61. Ito BR, Roth DM, Engler RL. Thromboxane A₂ and peptidoleukotrienes contribute to the myocardial ischemia and contractile dysfunction in response to intracoronary infusion of complement C5a in pigs. *Circ Res* 1990;66:596-607.
 62. Ito BR, Roth DM, Chenoweth D, Engler RL. Mechanisms of complement C5a induced myocardial ischemia: Formed blood elements are required/depressed contractile function is secondary to blood flow reduction (Abst). *Circulation* 1988;78 (Suppl II):II484.
 63. Engler RL, Ito BR, Roth DL, Chenoweth D. Dissociation of leukocyte accumulation from myocardial dysfunction: LTB₄ versus C5a induced PMN activation (Abstr). *Circulation* 1988;78 (Suppl II):II-651.
 64. Engler RL, del Balzo U, Nork SE, Ito BR. Leukocyte filtered blood attenuates the complement C5a induced myocardial ischemia and thromboxane production: Leukocyte dependent or independent? *Circulation* 1990;82 (Suppl III):III702.
 65. del Balzo U, Ito BR, Nork SE, Engler RL. Leukocyte depletion does not attenuate the myocardial ischemia and thromboxane production in response to complement C5a. *Circulation* 1990;82 (Suppl III):III276.
 66. Lewis RA, Austen KF. The biologically active leukotrienes. *J Clin Invest* 1984;73:889-897.
 67. Gryglewski RJ, Botting RM, Vane JR. Mediators produced by the endothelial cell. *Hypertension* 1988;12:530-548.
 68. Mullane K. Neutrophil and endothelial changes in reperfusion injury. *Trends Cardiovasc Med* 1991, in press.
 69. Cronstein BN, Levin RI, Belanoff J, et al. Adenosine: An endogenous inhibitor of neutrophil-mediated injury to endothelial cells. *J Clin Invest* 1986;78:760-770.
 70. Piper JM, Buderus S, Krutzfeldt A, et al. Sensitivity of the endothelium to hypoxia and reoxygenation. In: Piper HM, ed. *Pathophysiology of severe ischemic myocardial injury*. Dordrecht: Kluwer Academic Publishers, 1990:359-379.
 71. Dauber IM, Van Benthuyzen KM, McMurty IF, et al. Functional coronary microvascular injury evident as increased permeability due to brief ischemia and reperfusion. *Circ Res* 1990;66:986-998.
 72. Whorton AR, Montgomery ME, Kent RS. Effect of hydrogen peroxide on prostaglandin production and cellular integrity in cultured porcine aortic endothelial cells. *J Clin Invest* 1985;76:295-302.
 73. Andreoli SP, Liechty EA, Mallett C. Exogenous adenosine nucleotides replete endothelial cell adenosine triphosphate after oxidant injury by adenosine uptake. *J Lab Clin Med* 1990;115:304-313.
 74. Tsao PS, Aoki N, Lefer DJ, et al. Time course of endothelial dysfunction and myocardial injury during myocardial ischemia and reperfusion in the cat. *Circulation* 1990;82:1402-1412.
 75. Mehta JL, Nichols WW, Donnelly WH, et al. Protection by superoxide dismutase from myocardial dysfunction and attenuation of vasodilator reserve after coronary occlusion and reperfusion in dog. *Circ Res* 1989;65:1283-1295.
 76. Dauber IM, Lesnefsky EJ, VanBenthuyzen KM, et al. Re-

- active oxygen metabolite scavengers decrease function coronary microvascular injury due to ischemia-reperfusion. *Am J Physiol* 1991;260:H42-H49.
77. Patel KD, Zimmerman GA, Prescott SM, et al. Oxygen radicals induce human endothelial cells to express GMP-140 and bind neutrophils. *J Cell Biol* 1991, in press.
 78. Lewis MS, Whatley RE, Cain P, et al. Hydrogen peroxide stimulates the synthesis of platelet-activating factor by endothelium and induces endothelial cell-dependent neutrophil adhesion. *J Clin Invest* 1988;82:2045-2055.
 79. Skogund G, Cotgreave I, Rincon J, et al. H_2O_2 activates CD11b—CD18-dependent cell adhesion. *Biochem Biophys Res Comm* 1988;157:443-449.
 80. Neumann FJ, Rauch B, Richard G, et al. Local alteration of neutrophil function in short term myocardial ischemia. *Circulation* 1990;82 (Suppl III):III-702.
 81. Kramer JH, Arroyo CM, Dickens BF, et al. Spin-trapping evidence that graded myocardial ischemia alters post-ischemic superoxide production. *Free Rad Biol Med* 1987;3: 153-159.
 82. Bolli R, Patel BS, Jeroudi MO, et al. Demonstration of free radical generation in "stunned" myocardium of intact dogs with the use of the spin trap α -phenyl N-tert-butyl nitron. *J Clin Invest* 1988;82:476-485.
 83. Bolli R, Jeroudi MO, Patel BS, et al. Direct evidence that oxygen-derived free radicals contribute to postischemic myocardial dysfunction in the intact dog. *Proc Natl Acad Sci USA* 1989;86:4695-4699.
 84. Petrone WF, English DK, Wong K, McCord JM. Free radicals and inflammation: Superoxide-depressed activation of a neutrophil chemotactic factor in plasma. *Proc Natl Acad Sci USA* 1980;77:1159-1163.
 85. Schott RJ, Nao BS, McClanahan TB, et al. F(ab')₂ fragments of anti-Mo1 (904) monoclonal antibodies do not prevent myocardial stunning. *Circ Res* 1989;65:1112-1124.
 86. Barroso-Aranda J, Schmid-Schonbein J, Zweifach GW, Engler RL. Granulocytes and no-reflow phenomenon in irreversible hemorrhagic shock. *Circ Res* 1988;63:437-447.
 87. Barroso-Aranda J, Schmid-Schonbein GW. Transformation of neutrophils as indicator of irreversibility in hemorrhagic shock. *Am J Physiol* 1989;257:H846-H852.