

To each graded algebra  $R$  with a finite number of generators we associate the series  $T(R, z) = \sum d_n z^n$ , where  $d_n$  is the dimension of the homogeneous component of  $R$ . It is proved that if the dimensions  $d_n$  have polynomial growth, then the Krull dimension of  $R$  cannot exceed the order of the pole of the series  $T(R, z)$  for  $z = 1$  by more than 1.

We shall mainly consider the graded algebra  $R = \sum_{i=0}^{\infty} R_i$  over an arbitrary field  $k$  having a finite number of generators and a finite number of defining relations. However, we shall require a somewhat more general formulation of the main hypothesis.

Let  $M = \sum_{i=0}^{\infty} M_i$  be a graded bimodule over a free algebra  $k(x_1, \dots, x_s)$  having a finite number of generators and a finite number of defining relations. Let the symbol  $d_i(M)$ , or simply  $d_i$ , denote the dimension of  $M_i$  as a vector space over  $k$ . The main hypothesis has three equivalent formulations.

1. There exists a linear recurrence relation with integral coefficients  $\lambda_i$ :

$$d_n = \sum_{i=1}^k \lambda_i d_{n-i}, \quad (1)$$

which is valid for all sufficiently large  $n$ . The polynomial  $Q(M, z) = z^k - \sum \lambda_i z^{k-i}$  will be called the characteristic polynomial of the module  $M$ .

2. For all sufficiently large  $n$

$$d_n = \sum P_i(n) \alpha_i^n, \quad (2)$$

where the  $\alpha_i$  are algebraic integers and the  $P_i(n)$  are polynomials with rational coefficients.

3. The series

$$T(M, z) = \sum_{n=0}^{\infty} d_n z^n \quad (3)$$

is a rational function with integral coefficients and lowest term in the denominator equal to 1.

The proof that these formulations are equivalent can be found in [1], Chap. 3.

The foregoing quantities are related as follows. The  $\alpha_i$  are the roots of the characteristic polynomial  $Q(M, z)$ ;  $P_i(n)$  is a polynomial of degree  $k_i$ , where  $k_i + 1$  is the multiplicity of  $\alpha_i$ ;  $T(M, z) = f(z)g^{-1}(z)$ , where  $g(z) = z^k Q(M, z^{-1})$ , and the degree of  $f(z)$  is less than the number starting with which the recurrence relation holds for all following numbers.

In the general case the main hypothesis remains unproved. A proof exists in the following special cases: modules over commutative algebras ([2], Theorem 15.2); algebras defined by a finite collection of words ([3], Theorem 2); algebras having global dimension less than 3 ([3], Theorem 3). Theorem 1 below proves the main hypothesis for a rather narrow class of algebras; however, it is interesting because it establishes a connection between the coefficients of recurrence relation (1) and the Möbius function for a monoid (see [4]) or a partially ordered set (see [1], 2.2).

Moscow Institute of Electronic Machine Building. Translated from *Matematicheskie Zametki*, Vol. 14, No. 2, pp. 209-216, August, 1973. Original article submitted January 3, 1972.

© 1974 Consultants Bureau, a division of Plenum Publishing Corporation, 227 West 17th Street, New York, N. Y. 10011. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, microfilming, recording or otherwise, without written permission of the publisher. A copy of this article is available from the publisher for \$15.00.

Following Cartier and Foata [4], we introduce the following definitions. Let  $M$  be a monoid and  $M^*$  the set of nonidentity elements of  $M$ . By a factorization of  $x$  in  $M$  we mean any sequence  $(x_1, x_2, \dots, x_n)$  of elements of  $M^*$  such that  $x = x_1 x_2 \dots x_n$ . The number  $n$  is called the length of the factorization. We consider a monoid in which any element admits only a finite number of distinct factorizations. The number of factorizations of  $x$  is denoted by  $d(x)$ . The number of factorizations of even length is denoted by  $d_+(x)$  and of odd length by  $d_-(x)$ . It is obvious that  $d(x) = d_+(x) + d_-(x)$ . The function  $\mu(x) = d_+(x) - d_-(x)$  is called the Möbius function of  $M$ .

Let  $A$  be the set of functions on the monoid  $M$ . We introduce a multiplication operation on  $A$  as follows:

$$fg(x) = \sum f(x_1) g(x_2), \quad (4)$$

which turns  $A$  into a monoid. The function  $\varepsilon(x)$ ,  $\varepsilon(1) = 1$ , and  $\varepsilon(x) = 0$  for  $x \neq 1$  is the identity of  $A$ . The sum in (4) can be extended to all pairs  $(x_1, x_2)$  such that  $x = x_1 x_2$ ; among these pairs are  $(1, x)$  and  $(x, 1)$ . Let  $\zeta(x) = 1$  for all  $x \in M$ . Then (see [4])

$$\zeta u = \mu \zeta = \varepsilon.$$

**THEOREM 1.** Let  $M$  be a graded monoid without zero and with a finite number of generators. Let  $\mu(x)$  be such that  $\mu(x) \neq 0$  for only a finite number of elements of  $M$ . Then the semigroup algebra  $k(M)$  satisfies the main hypothesis.

Proof. Consider the following series product:

$$\sum_{x \in M} \mu(x) x \sum_{y \in M} \zeta(y) y = \sum_{z \in M} \varepsilon(z) z = 1.$$

Since there is no zero in  $M$ ,

$$\sum \mu(x) t^{|x|} \sum \zeta(y) t^{|y|} = 1,$$

where  $|x|$  is the degree of  $x$ . Collecting the coefficients of  $t^N$ , we obtain

$$\sum \mu(x) \zeta(y) = 0, \quad (5)$$

where  $|x| + |y| = N > 0$ .

Let  $\lambda_i = \sum_{|x|=i} \mu(x)$  and let  $d_j = \sum_{|y|=j} \zeta(y)$  be the number of elements of  $M$  having degree  $j$ . These elements can be taken as a basis for the  $j$ -th homogeneous component of the semigroup algebra  $k(M)$ . Therefore  $d_j$  coincides with  $d_j(k(M))$ .

Equality (5) can now be rewritten in the form

$$\sum_{|x|+|y|=N} \mu(x) \zeta(y) = \sum_{|x|<N} \mu(x) \sum_{|y|=N-|x|} \zeta(y) = \sum_{i=0}^N \lambda_i d_{N-i} = 0. \quad (6)$$

Since almost all  $\lambda_i = 0$ , by the condition of the theorem, equality (6) becomes recurrence relation (1).

**COROLLARY.** Let an algebra  $R$  be defined by a finite number of generators  $x_1, x_2, \dots, x_s$  and relations of the form  $x_i x_j - x_j x_i$  for some  $i$  and  $j$ . Then  $R$  satisfies the main hypothesis.

Proof. The algebra  $R$  satisfies all the conditions of Theorem 1 (see [4]).

The basic ring-theoretic operations, when applied to algebras satisfying the main hypothesis, again result in algebras satisfying it.

a) For any exact sequence

$$0 \rightarrow M_1 \rightarrow M \rightarrow M_2 \rightarrow 0, \\ T(M, z) = T(M_1, z) + T(M_2, z).$$

b) It is obvious that

$$T(M_1 \oplus {}_k M_2, z) = T(M_1, z) \cdot T(M_2, z).$$

c) If  $R_1$  and  $R_2$  are graded algebras and  $R_1 * R_2$  is their free product, then

$$T(R_1 * R_2, z) = [T^{-1}(R_1, z) + T^{-1}(R_2, z) - 1]^{-1}. \quad (7)$$

Viewing  $R_1$ ,  $R_2$ , and  $R_1 * R_2$  as  $k$ -modules, we obtain

$$R_1 * R_2 = k + I_1 + I_2 + I_1 \otimes I_2 + I_2 \otimes I_1 + I_1 \otimes I_2 \otimes I_1 + \dots,$$

where  $I_1(I_2)$  is the augmentation ideal of the algebra  $R_1(R_2)$ . Let  $T(I_1, z) = a_1$  and  $T(I_2, z) = a_2$ . Then

$$\begin{aligned} T(R_1 * R_2, z) &= 1 + a_1 + a_2 + 2a_1a_2 + a_1^2a_2 + a_1a_2^2 + \dots = \\ &= 1 + (a_1 + a_2 + 2a_1a_2)(1 + a_1a_2 + a_1^2a_2^2 + \dots) = 1 + (a_1 + a_2 + 2a_1a_2)(1 - a_1a_2)^{-1}. \end{aligned}$$

Substituting  $T(R_1, z) = a_1 + 1$  and  $T(R_2, z) = a_2 + 1$  into the resulting equation, we obtain Eq. (7).

If we know the expressions for  $T(R_1 * R_2, z)$ , it is easy to obtain the characteristic polynomial for  $R_1 * R_2$ . An especially simple formula is obtained when recurrence relation (1) is valid for all numbers starting with the first. In this case

$$T(R_i, z) = Q^{-1}(R_i, z^{-1}) z^{-k}$$

and

$$T(R_1 * R_2, z) = [Q(R_1, z^{-1}) z^k + Q(R_2, z^{-1}) z^k - 1]^{-1},$$

i.e.,

$$Q(R_1 * R_2, z) = Q(R_1, z) + Q(R_2, z) - z^k.$$

This means that the right sides of the recurrence relations can be added.

d) For the algebra of polynomials over the algebra  $A$  we obtain

$$T(A[x], z) = T(A)(1 - z)^{-1}. \quad (8)$$

Indeed, the algebra  $A[x]$ , viewed as a  $k$ -module, can be decomposed into the direct sum  $A[x] = A + Ax + Ax^2 + \dots$ . Therefore

$$T(A[x], z) = T(A) + zT(A) + z^2T(A) + \dots = T(A)(1 - z)^{-1}.$$

**LEMMA 1.** If  $r$  is the radius of convergence of the series  $T(M, z)$ , then  $T(M, z)$  has a pole at the point  $z = r$ , and all the poles of this series on the circle  $|z| = r$  have order not exceeding the order of the pole at  $z = r$ .

**Proof.** Since the terms of the series are positive, the existence of a pole at  $z = r$  is obvious. The comparison of the poles is obtained from the inequality

$$\left| \sum d_i u^i \right| \leq \sum d_i |u|^i.$$

**LEMMA 2.** If the series  $T(M, z)$  has radius of convergence equal to 1, then all of its poles are situated at roots of 1.

**Proof.** Let  $T(M, z) = f(z)g^{-1}(z)$ , where  $g(z) = 1 + a_1z + \dots + a_kz^k$ . Since the product of the roots of  $g(z)$  is equal to 1 and the smallest has modulus equal to unity, all the roots of  $g(z)$  are equal to 1 in modulus. If all the conjugate algebraic integers are equal to 1 in modulus, then they are roots of 1 of integral degree [5].

All modules that satisfy the main hypothesis can be divided into two classes:

- a) modules of polynomial type, i.e.,  $T(M, z)$  has poles only on the circle  $|z| = 1$ ;
- b) modules of exponential type, i.e.,  $T(M, z)$  has at least one pole for  $z = r < 1$ .

It is obvious that for any module  $M$  of polynomial type

$$T(M, z) = f(z) \prod_i (1 - z^{m_i}).$$

Let  $R$  be a graded algebra and  $R[x]$  the algebra of polynomials over  $R$  in the commuting variable  $x$ . Let us establish a correspondence between ideals of the ring  $R$  and homogeneous ideals of the ring  $R[x]$ . This correspondence is established as in [6], Chap. 7, § 5. Since we need to perform the same construction in a more general case, we shall verify these properties.

Let  $a = a_k + a_{k+1} + \dots + a_n \in R$ , where  $a_i$  is a homogeneous element of degree  $i$  in  $R$  and  $a_k \neq 0$ . Set  $\varphi(a) = a_k x^{n-k} + a_{k+1} x^{n-k-1} + \dots + a_n$ , and define  $\psi: R[x] \rightarrow R$ , by setting  $\psi(x) = 1$  and extending it further as a ring homomorphism. The mapping  $\varphi$  is not a homomorphism; the mapping  $\psi$  is a homomorphism, but we shall consider it only on homogeneous elements. Obviously,

$$\psi\varphi(a) = a, \quad \psi\varphi(b) = x^{-s}b, \quad (9)$$

where  $x^s$  is the highest degree of  $s$  dividing  $b$ .

These mappings can be extended to ideals. Small Gothic letters will denote ideals of  $R$ , and capital letters ideals of  $R[x]$ . If  $\mathfrak{a} \in R$ , then  $\varphi(\mathfrak{a})$  is not an ideal in  $R[x]$ , since it does not contain elements divisible by  $x$ . However, it is easy to verify that the set of elements of the form  $x^s\varphi(a)$ , where  $a \in \mathfrak{a}$ , and  $s$  is any natural number, is an ideal in  $R[x]$ , which we shall henceforth denote by  $\varphi(\mathfrak{a})$ .

Equalities (9) extended to ideals become the inclusions

$$\psi\varphi(\mathfrak{a}) = \mathfrak{a}, \quad \varphi\psi(\mathfrak{A}) \supset \mathfrak{A}. \quad (10)$$

Since the ideal  $\varphi(\mathfrak{a})$  is generated by the elements  $\varphi(a)$ , where  $a \in \mathfrak{a}$ , we have  $\varphi(ab) = \varphi(a)\varphi(b)$ , and if  $\mathfrak{a} \subset \mathfrak{b}$ , then  $\varphi(\mathfrak{a}) \subset \varphi(\mathfrak{b})$ . Let us prove that prime ideals are carried to prime ideals. Indeed, let  $\mathfrak{A}\mathfrak{B} \subset \varphi(\mathfrak{p})$  where  $\mathfrak{p}$  is prime. Then  $\psi(\mathfrak{A})\psi(\mathfrak{B}) \subset \psi\varphi(\mathfrak{p}) = (\mathfrak{p})$ . Let  $\psi(\mathfrak{A}) \subset \mathfrak{p}$ . Then  $\varphi\psi(\mathfrak{A}) \subset \varphi(\mathfrak{p})$ . Using (10), we obtain  $\mathfrak{A} \subset \varphi\psi(\mathfrak{A}) \subset \varphi(\mathfrak{p})$ , i.e.,  $\varphi(\mathfrak{p})$  is prime.

**THEOREM 2.** Let  $R$  be a graded algebra of polynomial type, and let the series  $T(R, z)$  have a pole of order  $s$  at the point  $z = 1$ . Then the Krull dimension of  $R$  does not exceed  $s + 1$ .

**Proof.** Let  $\mathfrak{q}_0 \subset \mathfrak{q}_1$  be homogeneous prime ideals of  $R$ . Replacing  $R/\mathfrak{q}_0$  by  $A$ , we can assume that  $\mathfrak{q}_0 = 0$ . For any left ideal  $\mathfrak{q}$  of  $A$  we have  $0 \neq \mathfrak{q}_1 \mathfrak{q} \in \mathfrak{q}_1 \cap \mathfrak{q}$ , whence  $A$  is an essential extension of  $\mathfrak{q}_1$ . Therefore,  $\mathfrak{q}_1$  contains an element  $a$  which is not a divisor of zero (see [7]). The left  $A$ -modules  $A$  and  $Aa$  are isomorphic, and so  $T(A, z) = z^k T(Aa, z)$ , where  $k$  is the degree of  $a$ . Hence we obtain the coefficient inequality  $T(\mathfrak{q}_1, z) \geq z^k T(A, z)$ , which remains valid upon comparing the values of the series for positive values of  $z$  not exceeding 1. Therefore

$$T(A/\mathfrak{q}_1, z) = T(A, z) - T(\mathfrak{q}_1, z) \leq T(A, z) - z^k T(A, z) = (1 - z)h(z)T(A, z),$$

where  $h(z)$  is a polynomial. It is apparent from the resulting inequality that the series  $T(A/\mathfrak{q}_1, z)$  has a pole of smaller order than the series  $T(A, z)$ . If the series  $T(A, z)$  does not have  $z = 1$  as a singular point then it is a polynomial. In this case the algebra  $A$  is finite-dimensional over the field  $k$ , and so it cannot contain a chain of prime ideals of nonzero length. Using induction, we obtain that the algebra  $R$  cannot contain a chain of homogeneous prime ideals of length bigger than  $s$ .

The series  $T(R[x], z)$  according to formula (8), has a pole of order  $s + 1$  at the point  $z = 1$ . Using the correspondence  $\varphi$  between prime ideals of  $R$  and homogeneous prime ideals of  $R[x]$ , we obtain the assertion of the theorem.

Commutative algebras with a finite number of generators are of polynomial type and are noetherian. It is apparent from the proof of Theorem 2 that the ascending chain condition on ideals in a ring of polynomial type holds if there exist nondivisors of zero in a sufficient number of factors of this series. The question arises of whether all algebras of polynomial type are noetherian. It turns out that this is false. The algebra  $R = k(xy)/(x^2, yx, yxy)$  serves as an example. If  $\varphi: k(x, y) \rightarrow R$  is the canonical homomorphism and  $\varphi(x) = a$ ,  $\varphi(y) = b$ , then as a basis for the homogeneous component  $R_n$  we can take the four elements  $b^n, ab^{n-1}, b^{n-1}a$ , and  $ab^{n-2}a$ . It is easy to compute  $T(R, z) = (1 + z + z^2 + z^3)(1 - z)^{-1}$ ; however,  $R$  contains an ideal  $\mathfrak{a}$ , generated by the infinite collection of elements  $ab^n a$  ( $n = 1, 2, \dots$ ).

The ideal  $\mathfrak{a}$  splits into an infinite direct sum, since  $Rab^n aR = kab^n ak$ , i.e., has infinite dimension in the sense of Goldie; therefore (see [8]) it has an infinite Krull dimension in the sense of Rentschler and Gabriel [9].

#### LITERATURE CITED

1. M. Hall, Combinatorics [Russian translation], Moscow (1970).
2. Yu. I. Manin, Lectures on Algebraic Geometry [in Russian], Moscow (1968).
3. V. E. Govorov, "Graded algebras," *Matem. Zametki*, **12**, No. 2, 197-204 (1972).

4. B. Cartier and D. Foata, Problemes Combinatoires de Commutation et Rearrangements, Lecture Notes in Math., No. 85 (1969).
5. L. Bieberbach, "Über einen Satz Polyascher," Art. Arch. Math., 4, 23-28 (1953).
6. O. Zariski and P. Samuel, Commutative Algebra, Van Nostrand, Princeton, Vol. I (1958), Vol. II (1960).
7. A. W. Goldie, "Semi-prime rings with maximum conditions," Proc. London Math. Soc., 10, 201-220 (1960).
8. P. Lemonnier, "Quelques application de la dimension de Krull," C. R. Acad. Sci., A1395-A1397, 270 (1970).
9. R. Rentschler and P. Gabriel, "Sur la dimension des anneaux et ensembles ordonnés" C. R. Acad. Sci., A712-A715, 265 (1970).