

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/226005099>

ChemInform Abstract: Microwave Assisted Synthesis and Antibacterial Activity of New Quinolone Derivatives.

ARTICLE *in* MONATSHFTE FUER CHEMIE/CHEMICAL MONTHLY · AUGUST 1998

Impact Factor: 1.22 · DOI: 10.1007/PL00013504

CITATIONS

17

READS

22

6 AUTHORS, INCLUDING:



Rajesh Kumar

Indian Agricultural Research Institute

87 PUBLICATIONS 414 CITATIONS

SEE PROFILE

Microwave Assisted Synthesis and Antibacterial Activity of New Quinolone Derivatives

Mazaahir Kidwai^{1,*}, Preeti Misra¹, Rajesh Kumar¹, Rajendra K. Saxena², Rani Gupta², and Sapna Bradoo²

¹ Department of Chemistry, University of Delhi, Delhi 110007, India

² Department of Microbiology, University of Delhi, South Campus, Delhi 110021, India

Summary. A series of novel 6-fluoro-7-(5-alkyl-1,3,4-thiadiazol/oxadiazol-2-ylsulfanyl)-4-quinolone-3-carboxylic acids were synthesized using microwave irradiation. The compounds were tested for their *in vitro* antibacterial activity. All compounds containing the 1,3,4-thiadiazole/oxadiazole moiety at position 7 showed promising antibacterial activity.

Keywords. Microwave irradiation; Quinolones; Antibacterial activity.

Mikrowellenunterstützte Synthese und antibakterielle Aktivität von neuen Chinolonderivaten

Zusammenfassung. Einige neue 6-Fluoro-7-(5-alkyl-1,3,4-thiadiazol/oxadiazol-2-sulfanyl)-4-chinolon-3-carbonsäuren wurden unter Bestrahlung mit Mikrowellen synthetisiert. Die Verbindungen wurden hinsichtlich ihrer antibakteriellen Aktivität *in vitro* untersucht. Sämtliche Verbindungen mit 7-ständigen 1,3,4-Thiadiazol/Oxadiazol-Einheiten zeigten vielversprechende Aktivität.

Introduction

In recent years, three quinolones [1] (ciprofloxacin [2], ofloxacin [3], and norfloxacin [4]) have become commercially available. Quinolones inhibit the replication of DNA [5] by inhibition of the DNA gyrase. The latter is probably a metalloenzyme and interacts with the DNA molecule *via* an intermediate (iron complex) that carries the quinolone drug. It has been reported that quinolone antibacterial agents do not bind to the gyrase subunit, but rather to DNA, and the mechanism of action is based on drug DNA complex formation [6–7]. Quinolones act as bidentate chelating agents.

Quinolone derivatives have been associated with antibacterial [8], antimicrobial [9], and antistaphylococcal [10] activities. Thiadiazoles and oxadiazoles are pharmacologically important [11]. Keeping in view the importance of MORE (Microwave induced Organic Reaction Enhancement [12–17] chemistry and the biological importance of the above mentioned moieties, it was thought worthwhile to develop a method for the rapid synthesis of 6-fluoro-7-(5-alkyl-1,3,4-thiadiazol/oxadiazol-2-ylsulfanyl)-4-quinolone-3-carboxylic acids and to screen them for bactericidal activity.

* Corresponding author

Results and Discussion

To our knowledge, this is the first report on quinolone synthesis in which microwave technique has been used and thiadiazole/oxadiazole rings are incorporated.

Diethyl-3-chloro-4-fluoroanilinemethylene malonate (**1**) was synthesized by condensation of 3-chloro-4-fluoroaniline with diethylethoxymethylene malonate. Cyclization of **1** in *PPA* under microwave irradiation for 3 min yielded the cyclized ester which upon alkaline hydrolysis afforded **2** as evidenced by the disappearance of the ester group signals in the ^1H NMR spectrum at $\delta = 1.5$ and 4.2 ppm. Formal nucleophilic substitution of the chlorine in **2** with mercapto substituted 1,3,4-thiadiazoles/oxadiazoles was achieved under microwave irradiation for 4–5 min furnishing **3a–f** (Scheme 1). The structures of the compounds were confirmed by their analytical and spectroscopic data given in the experimental part.




Table 1. Antibacterial activity of compounds **3a–f**

Organism	3a	3b	3c	3d	3e	3f	Norfloxacin
2253- <i>Corynebacterium rubrum</i> NCL	+++	+	+++	++++	+++	+++	++++
2281- <i>Klebsiella aerogens</i> NCL	+	–	–	+++	++	+	++++
2491- <i>Erwinia herbicola</i> NCL	++	+	++	+++	+++	+++	+++
2715- <i>Bacillus lichenformis</i> NCL	+	+	++	++	+	–	++++
2323- <i>Bacillus coagulans</i> NCL	+	–	+	+	++	++	+++
K ₁₂ - <i>E. coli</i>	++	+	+	+++	++	++	++++
2689- <i>Bacillus lichenformis</i> NCL	+++	+	–	++	+++	++	+++

–: No measurable activity; +: 3–9 mm; ++: 10–12 mm; +++: 13–16 mm; ++++: 17–21 mm

Compounds **3a–f** were tested (Table 1) for their *in vitro* antibacterial activity against 6 bacterial strains by the cup-plate agar diffusion method [18]. They were dissolved in *DMF* at a concentration of 50 µg/ml. All compounds were found active against *Corynebacterium rubrum*, *Escherichia coli*, and *Erwinia herbicola*. Compounds with a 1,3,4-oxadiazole ring at position 7 possess better antibacterial activity than those with a 1,3,4-thiadiazole ring. Among all synthesized quinolones, **3d** provided the best antibacterial activity and compared well with the activities of norfloxacin. **3a**, **3e**, and **3f** also possess promising antibacterial activity.

Experimental

Melting points were taken on an electrothermal apparatus and are uncorrected. IR spectra were recorded on a 1710 Perkin Elmer FTIR spectrophotometer using KBr discs. ^1H NMR spectra were recorded on a FTNMR Hitachi R-600 spectrometer operating at 90 MHz using *TMS* as internal standard (δ in ppm). Elemental analysis data agreed satisfactorily with the calculated ones. A Padmini Essentia microwave oven, Model Brownie, at 2450 MHz was used. For TLC analysis, silica gel coated Al plates (Merck) were used.

Diethyl-3-chloro-4-fluoroanilinomethylene malonate (1; C₁₄H₁₅O₄ NCIF)

A stirred mixture containing an equimolar ratio of 3-chloro-4-fluoro aniline and diethylethoxymethylene malonate was heated at 120–140°C until the evolution of ethanol ceased. The disappearance of aniline was confirmed by TLC using petrol ether:CHCl₃:acetone (80:15:5) as eluent. The yield was almost quantitative, and the crude product (oil) was pure enough for the next step.

Yield: 91%; IR (KBr): 3320 (N-H), 1725 (C=O) cm⁻¹; ^1H NMR (CDCl₃, δ): 1.3 (t, 6H, 2CH₂CH₃), 4.1 (q, 4H, 2CH₂CH₃), 7.0–7.3 (m, 3H, Ar-H), 9.1 (br, 1H, NH), 9.4 (s, 1H, C₂-H) ppm; MS: *m/z* calculated for C₁₄H₁₅O₄ NCIF (M⁺) 315.18, found 315.03.

7-Chloro-6-fluoro-4-quinolone-3-carboxylic acid (2; C₁₀H₅O₃ NCIF)

A mixture containing 1.5 g **1** and 7 ml polyphosphoric acid (*PPA*) was subjected to microwave irradiation (MWI) for 3 min. The reaction was monitored by TLC using petrol ether:ethylacetate (9:1) as eluent. The resulting mixture was alkalized with an excess of NaOH and subjected to MWI for 4–5 min. The resulting solution was neutralized with 10% HCl to adjust the *pH* value to 4 and extracted with ethyl acetate. After drying over Na₂SO₄, the extract was evaporated to dryness. **2** was isolated as slightly yellow crystalline product.

Yield: 78%; m.p.: >300°C; IR (KBr): 3400 (N-H), 1700 (C=O) cm⁻¹; ^1H NMR (CDCl₃, δ): 8.7 (d, 1H, *J* = 9 Hz, C₅-H), 9.0 (d, 1H, *J* = 5 Hz, C₈-H), 9.6 (s, 1H, C₂-H), 9.2 (br, 1H, NH) ppm; MS: *m/z* calculated for C₁₀H₅O₃ NCIF (M⁺) 241.51, found 241.43.

General procedure for the preparation of 6-fluoro-7-(5-alkyl-1,3,4-thiadiazol/oxadiazol-2-ylsulfanyl)-4-quinolone-3-carboxylic acids (3a–f)

A mixture of **2** (0.05 mol), 1, 3, 4-thiadiazole/oxadiazole (0.05 mol), and 4 g K₂CO₃ in 5 ml *DMF* was subjected to MWI for 4–5 min. The inorganic salt was filtered off, and the resulting filtrate was poured into an excess of ice cold water (100 ml) and extracted with ethyl acetate. The organic layer was washed with water (2×50 ml), dried over Na₂SO₄, and evaporated to afford the corresponding products which were recrystallized from *DMF*–EtOH.

6-Fluoro-7-(5-methyl-1,3,4-thiadiazol-2-ylsulfanyl)-4-quinolone-3-carboxylic acid
(**3a**; C₁₃H₈O₃N₃FS₂)

Yield: 50%; m.p.: >300°C; IR (KBr): 1680 (C=O), 1580 (C=N) cm⁻¹; ¹H NMR (CDCl₃, δ): 2.7 (s, 3H, CH₃), 8.65 (d, 1H, *J* = 9 Hz, C₅-H), 9.13 (d, 1H, *J* = 5 Hz, C₈-H), 9.62 (s, 1H, C₂-H), 9.27 (br, 1H, NH) ppm; MS: *m/z* calculated for C₁₃H₈O₃N₃FS₂ (M⁺) 337.12, found 337.02.

6-Fluoro-7-(5-nonyl-1,3,4-thiadiazol-2-ylsulfanyl)-4-quinolone-3-carboxylic acid
(**3b**; C₂₁H₂₄O₃N₃FS₂)

Yield: 72%; m.p.: 278°C; IR (KBr): 1685 (C=O), 1575 (C=N) cm⁻¹; ¹H NMR (CDCl₃, δ): 0.88 (t, 3H, 9'CH₃), 1.27 (m, 12H, 6CH₂), 1.6 (m, 2H, 2'CH₂), 2.34 (t, 2H, 1'CH₂), 8.67 (d, 1H, *J* = 9 Hz, C₅-H), 9.15 (d, 1H, *J* = 5 Hz, C₈-H), 9.65 (s, 1H, C₂-H), 9.25 (br, 1H, NH) ppm; MS: *m/z* calculated for C₂₁H₂₄O₃N₃FS₂ (M⁺) 449.29, found 449.14.

6-Fluoro-7-(5-undecyl-1,3,4-thiadiazol-2-ylsulfanyl)-4-quinolone-3-carboxylic acid
(**3c**; C₂₃H₂₈O₃N₃FS₂)

Yield: 67%; m.p.: 290°C; IR (KBr): 1680 (C=O), 1575 (C=N) cm⁻¹; ¹H NMR (CDCl₃, δ): 0.88 (t, 3H, 11'CH₃), 1.26 (m, 16H, 8CH₂), 1.62 (m, 2H, 2'CH₂), 2.34 (t, 2H, 1'CH₂), 8.63 (d, 1H, *J* = 9 Hz, C₅-H), 9.10 (d, 1H, *J* = 5 Hz, C₈-H), 9.58 (s, 1H, C₂-H), 9.20 (br, 1H, NH) ppm; MS: *m/z* calculated for C₂₃H₂₈O₃N₃FS₂ (M⁺) 477.31, found 477.18.

6-Fluoro-7-(5-methyl-1,3,4-oxadiazol-2-ylsulfanyl)-4-quinolone-3-carboxylic acid
(**3d**; C₁₃H₈O₄N₃FS)

Yield: 58%; m.p.: >300°C; IR (KBr): 1675 (C=O), 1530 (C=N) cm⁻¹; ¹H NMR (CDCl₃, δ): 2.76 (s, 3H, CH₃), 8.70 (d, 1H, *J* = 9 Hz, C₅-H), 9.10 (d, 1H, *J* = 5 Hz, C₈-H), 9.76 (s, 1H, C₂-H), 9.32 (br, 1H, NH) ppm; MS: *m/z* calculated for C₁₃H₈O₄N₃FS (M⁺) 321.14, found 321.03.

6-Fluoro-7-(5-nonyl-1,3,4-oxadiazol-2-ylsulfanyl)-4-quinolone-3-carboxylic acid
(**3e**; C₂₁H₂₄O₄N₃FS)

Yield: 69%; m.p.: 280°C; IR (KBr): 1680 (C=O), 1525 (C=N) cm⁻¹; ¹H NMR (CDCl₃, δ): 0.89 (t, 3H, 9'CH₃), 1.26 (m, 12H, 6CH₂), 1.62 (m, 2H, 2'CH₂), 2.34 (t, 2H, 1'CH₂), 8.67 (d, 1H, *J* = 9 Hz, C₅-H), 9.15 (d, 1H, *J* = 5 Hz, C₈-H), 9.65 (s, 1H, C₂-H), 9.27 (br, 1H, NH) ppm; MS: *m/z* calculated for C₂₁H₂₄O₄N₃FS (M⁺) 433.22, found 433.20.

6-Fluoro-7-(5-undecyl-1,3,4-oxadiazol-2-ylsulfanyl)-4-quinolone-3-carboxylic acid
(**3f**; C₂₃H₂₈O₄N₃FS)

Yield: 63%; m.p.: 286°C; IR (KBr): 1685 (C=O), 1520 (C=N) cm⁻¹; ¹H NMR (CDCl₃, δ): 0.88 (t, 3H, 11'-CH₃), 1.25 (m, 16H, 8CH₂), 1.6 (m, 2H, 2'-CH₂), 2.34 (t, 2H, 1'-CH₂), 8.65 (d, 1H, *J* = 9 Hz, C₅-H), 9.15 (d, 1H, *J* = 5 Hz, C₈-H), 9.60 (s, 1H, C₂-H), 9.27 (br, 1H, NH) ppm; MS: *m/z* calculated for C₂₃H₂₈O₄N₃FS (M⁺) 461.24, found 461.10.

References

- [1] Davies BI, Maesen FPV (1987) Pharmaceutisch Weekblad Scientific Edition **9**: 53
- [2] Wise R, Andrews JM, Edwards LJ (1983) Antimicrob Agents Chemother **23**: 559

- [3] Hayakawa I, Hiramitsu T, Tanaka Y (1984) Chem Pharm Bull **32**: 4907
- [4] Koga H, Itoh A, Murayama S, Suzue S, Irikura T (1980) J Med Chem **23**: 1358
- [5] Cozarelli NR (1980) Science **207**: 953
- [6] Shen LL, Pernet AG (1985) Proc Natl Acad Sci USA **82**: 307
- [7] Sharma RK, Joseph S, Kidwai M (1997) J Indian Chem Soc **74**: 11
- [8] Zhang MQ, Haemers A, Berghe DV, Pattyn SR, Bollaert W (1991) J Heterocycl Chem **28**: 685
- [9] Roy K, Srivastava RP, Tekwani BL, Pandey VC, Bhaduri AP (1996) Bioorg Med Chem Lett **6**: 121
- [10] Reuman M, Daum SJ, Singh B, Wentland MP, Perni B, Pennock P, Carabates PM, Gruett MD, Saindane MT, Droff PH, Coughlin SA, Sedlock DM, Rake JB, Leshner GY (1995) J Med Chem **38**: 2531
- [11] Mullican MD, Wilson MW, Connor DT, Kostlan CR, Schrier DJ, Dyer RD (1993) J Med Chem **36**: 1090
- [12] Kidwai M, Kumar P (1997) J Chem Res(S) **5**: 178
- [13] Kidwai M, Kumar P (1996) J Chem Res(S) **5**: 254
- [14] Kidwai M, Kumar P, Kohli S (1997) J Chem Res(S) **24**
- [15] Kidwai M, Kumar R, Goel Y (1997) Main Gp Met Chem **20**: 367
- [16] Kidwai M, Goel Y (1996) Polyhedron **15**: 2819
- [17] Kidwai M, Bhushan KR, Kumar P, Kumar R (1997) Monatsh Chem **128**: 1291
- [18] British Pharmacopaea (1953) Pharmaceutical Press, London, p 796

Received January 19, 1998. Accepted (revised) February 10, 1998