

See discussions, stats, and author profiles for this publication at:
<https://www.researchgate.net/publication/226216890>

ChemInform Abstract: A General Survey and Recent Advances in the Activation of Unreactive Chemical Bonds by Metal Complexes

CHAPTER *in* CHEMINFORM · APRIL 1999

Impact Factor: 0.74 · DOI: 10.1007/3-540-68525-1_1 · Source: OAI

CITATIONS

4

READS

34

2 AUTHORS, INCLUDING:



Gerard van koten

Utrecht University

1,113 PUBLICATIONS **27,997**

CITATIONS

SEE PROFILE

A General Survey and Recent Advances in the Activation of Unreactive Chemical Bonds by Metal Complexes

Robert A. Gossage^a, Gerard van Koten*

Debye Institute, Department of Metal-Mediated Synthesis, Utrecht University,
Padualaan 8, 3584 CH, Utrecht, The Netherlands

E-mail: g.vankoten@chem.uu.nl

^aAnorMED Inc., #100 20353–64th Ave., Langley, British Columbia, Canada, V2Y 1N5

| | | |
|---|---|---|
| 1 | General Introduction | 1 |
| 2 | The Activation of C–H Bonds | 2 |
| 3 | The Activation of C–Cl and C–F Bonds | 3 |
| 4 | The Activation of C–C and Si–Si Bonds | 4 |
| 5 | The Activation of C–O Bonds | 4 |
| 6 | The Activation of Molecular Nitrogen | 5 |
| 7 | Concluding Remark | 5 |
| | References | 6 |

1 General Introduction

The activation of chemical bonds is perhaps the most important area of modern chemistry. In our never ending search for cheap raw materials, the use of abundant but traditionally unreactive molecules as synthetic precursors is becoming increasingly more attractive. A major goal of this endeavor is to devise not only better methods to produce bulk (commodity) chemicals but also to synthesize new materials for industry, medicine and research. Unreactive chemicals include compounds such as hydrocarbons and other media which, under normal circumstances, do not react with other substrates (or themselves). Two primary examples are molecular nitrogen and saturated alkanes. These substances represent very inexpensive potential sources of nitrogen and carbon, respectively. Hydrocarbons (i.e., oil and petroleum products) are the largest fraction of world primary energy production and are thus readily available starting materials. The same is true for dinitrogen as it is a major component of the earth's atmosphere. In addition, the activation of general classes of inert bonds, such as the C–Cl, C–F

or C–O bonds, has importance in the destruction of certain man-made environmental toxins (PCB's, CFC's, etc.) and in the potential application of much cheaper chlorinated compounds as reagents (e.g., in Grignard reactions). The activation of specific C–C bonds has great potential in speciality chemical synthesis as does, to a lesser extent, the Si–Si bond in materials science.

This volume of Topics in Organometallic Chemistry is devoted to recent advances in the activation, by metal complexes, of what are termed unreactive chemical bonds. In this introduction, a brief overview of each topic will be presented. This is certainly not intended as a comprehensive review of each subject but merely as a stepping stone to the more detailed chapters that follow within.

2

The Activation of C–H Bonds

The synthetic utility of activating C–H bonds has long been recognized in chemistry. For example, the combustion of hydrocarbons as an energy source with the concomitant formation of water and carbon dioxide is a fundamental reaction in most machines. The *selective* activation of specific types of C–H bonds is however, by no means a simple problem. The strength of the C–H bond in for example, methane (440 kJ/mol; 105 kcal/mol) or benzene (461 kJ/mol; 110 kcal/mol), is a factor which alone does not dominate the chemistry, but it does make activation of this type of bond relative to weaker bonds present in a molecule more difficult [1]. Catalytic activation by metal containing compounds has therefore been an area of intense research.

The study of the reactivity of aryl C–H bonds began in the last century. However, direct bond activation was only observed in the last 35 years, despite the quite early successes of scientists like Charles Friedel and James Crafts who discovered (1877) that aluminum chloride catalysts can promote the alkylation of benzene. Specifically, *orthometallation* of aryl groups attached to coordinated donor atoms was among the first examples of direct C–H bond activation in a transition metal complex. This work demonstrated that an M–C(aryl) fragment and a metal hydride could be formed directly by oxidative addition of an *ortho* H–C(aryl) group [2–8]. An early example of this is the orthometallation of a coordinated triphenylphosphine ligand bound to an Ir(I) metal center [2–6]. Intermediates in this reaction have been isolated and all show close interatomic contact distances between the metal and a hydrogen nucleus. Examples include the very early crystallographic work of Laplaca and Ibers [9] of the Ru(II) complex $\text{RuCl}_2(\text{PPh}_3)_3$. Related chemistry includes examples of hydrogen atoms of alkyl groups in close proximity to a metal centre, such as the early disclosure by Maitlis and coworkers concerning a Pt phosphine compound with a close intramolecular $\text{Pt}\cdots\text{H}(\text{alkyl})$ distance [10]. Crabtree and others have coined the phrase *agostic* to describe this type of interaction and have used these complexes as models of the intimate first stages of C–H bond activation [11, 12]. However, the direct use of aromatic compounds in synthesis is generally restricted to the activation of groups other than the C–H fragment (e.g., C–X bonds: X=Br, Cl).

Since aryl halides are fairly cheap reagents, there has been less recent emphasis (see Section 3 below) on the development of aryl relative to that of alkyl C–H bond activation [13–17]. However, the manufacture of aryl halides is not an environmentally friendly process and thus the future of bulk aromatic synthesis may lie in the direct activation of C–H bonds. For example, the formation of benzaldehyde from the insertion of CO into a C–H bond of benzene is a recent development in this area [17].

Bergman has referred to the selective activation of C–H bonds of saturated alkanes as one of last remaining Holy Grails of synthetic chemistry [18–20]. The initial (perceived) breakthrough in alkyl C–H bond activation came from the simultaneous [21, 22] but independent work of Graham and coworkers (University of Alberta, Canada) and by Bergman et al. (University of California at Berkeley, U.S.A.). Both of these groups discovered that cyclopentadienyl complexes of Ir and Rh can, under photochemical conditions, oxidatively add alkanes to yield hydridometal alkyls. Although none of these systems have yet been shown to operate catalytically, detailed study of the fundamental aspects of this chemistry have continued in earnest [23–29]. Kinetic analysis has revealed that initial agostic interactions of the alkane with the metal center is a key (reversible) intermediate step and that later oxidative addition likely occurs via a simple three-center transition state [30]. Further work in this important area of research will be disclosed in the later chapters of this text by Prof. W. D. Jones (Stoichiometric Activation of C–H Bonds), Prof. F. Kakiuchi and Prof. S. Murai (Catalytic Activation of C–H Bonds) and the chapter by Prof. A. Sen (Catalytic Activation of Methane and Ethane by Metal Compounds).

3

The Activation of C–Cl and C–F Bonds

The activation of chlorinated hydrocarbons is an area of study that has direct environmental consequences in relation to the facile destruction of polychlorinated biphenyls (PCB's C–Cl bond strength 402 kJ/mol; 96 kcal/mol for C_6H_5-Cl) and other pollutants. The high reactivity of simple polyhalogenated alkanes such as tetrachloromethane [31–33], is generally facilitated by the ease of formation of the trichloromethane radical. However, other chlorocarbons are not so easily activated. There are a few examples of metal complexes which contain coordinated alkyl halides [34–37] and these may be considered as agostic interactions (i.e., $M \cdots X-R$) between the metal and an alkyl halide atom. Direct oxidative addition of unreactive haloalkanes such as dichloromethane has been reported [38, 39]. The activation of C–Cl bonds will be described in the chapter by Dr. V. V. Grushin and Prof. H. Alper (Activation of Otherwise Unreactive C–Cl Bonds) which details the use of chlorocarbons for a variety of applications in synthesis. The environmental (e.g., ozone depletion) and potential pharmaceutical relevance of fluorocarbons has made C–F bond activation an area of quite active research. Despite the high energy of C–F bonds (e.g., 644 kJ/mol; 154 kcal/mol for the C–F bond in C_6F_6), the activation of this fragment by a

number of complexes has been reported [40, 41]. Many metal complexes can activate C–F bonds in a stoichiometric fashion and the list includes reagents containing Ti [42], Fe [43], Ni [44], Ru [45, 46], Rh [47–50], Ir [47–50], W [51], U [52] and Yb [53]. Recently, this area has expanded to the catalytic activation of C–F bonds as reported by Aizenberg and Milstein [54, 55]. The use of electron-rich Rh phosphine compounds enables the activation of fluorobenzenes, although turnover numbers are low (<1 turnover per hour at 94°C) [54]. Further details on this chemistry can be found in the relevant chapter by Prof. T. G. Richmond (Metal Reagents for Activation and Functionalization of C–F Bonds).

4

The Activation of C–C and Si–Si Bonds

The activation of C–C bonds (specifically cleavage reactions) by metal compounds has been extensively studied and there are many reagents that can be used to perform the selective manipulation of alkynes and alkenes to produce a plethora of new compounds [56–59]. Transition metal and lanthanide complexes are playing an ever increasing role in the selective formation and cleavage of C–C bonds [60]. Considerable effort has been applied to the application of metal compounds in such diverse areas as ring forming reactions [61–64], regio- and enantio-selective addition reactions [65–71], direct C–C bond formation [72–76] and other processes [56, 57, 77]. This represents a very large cross section of chemical reactivity and the review in this text by Dr. M. Murakami and Prof. Y. Ito (Cleavage of C–C Single Bonds by Transition Metals) will emphasize this selected area of organometallic chemistry and (homogeneous) catalysis.

A related area of this chemistry is the selective reactivity of Si–Si bonds [78]. The use of disilanes in organic synthesis for the creation of new (and reactive) Si–C bonds is an area of current interest and application [79–81]. This work has stemmed from earlier studies which have demonstrated that disilanes readily oxidatively add to a variety of metal centres [82] to produce transition metal silyl complexes [83]. The investigation of this area will be detailed by Prof. M. Sugimone and Prof. Y. Ito (Activation of Si–Si Bonds: Stoichiometric and Catalytic Reactions).

5

The Activation of C–O Bonds

The cleavage of C–O bonds has significant applications in a number of important areas of synthetic organic and organometallic chemistry [56, 57, 83, 84]. The activation of carbon-oxygen single, double and triple bonds has been an area of active research in relation to catalysis and to many fundamental chemical reactions [85]. The reactions of carbon monoxide or dioxide has long been recognized in organometallic chemistry and thus can not be considered unreactive substrates by the definition of this book. Other organic compounds which contain the C=O functionality such as aldehydes and ketones can also not be con-

sidered unreactive as per the definition of this manuscript [84, 85]. However, asymmetric reactions at this functional group is an area of great current interest. Specifically, Prof. Y.-S. Lin and Prof. A. Yamamoto will describe recent advances in the activation of the important allylic C–O bond with special detail paid to allyl metal complexes. The allylic functionality has been studied extensively because of its importance in a number of specialty organic syntheses [86–92]. In addition, the chapter “Activation of C–O Bonds. Stoichiometric and Catalytic Reactions” will review recent applications of transition metal compounds to the cleavage of C–O bonds of esters, ethers [91, 92] and anhydrides. The rupture of the C–O multiple bonds is also detailed.

6

The Activation of Molecular Dinitrogen

Catalytic dinitrogen activation has been one of the most difficult challenges in the field of organometallic chemistry. Nitrogen gas is used as the N-containing feedstock in the (high temperature and pressure) Haber process for the production of ammonia. The drastic conditions that are necessary for efficient catalysis has lead to a vast amount of study directed towards the design of a mild ammonia synthesis from N_2 . The very high strength of the N–N bond (945 kJ/mol; 226 kcal/mol) and the differences in energy distribution between the bonds of N_2 compared with acetylene for example, lead to few similarities in the reactivity of these compounds [93–96]. Dinitrogen is a very weak base and hence does not easily (or strongly in most cases) coordinate to metals. The first example of an N_2 complex was not isolated until 1965 by Allen and Senoff [97]. Since that time, dinitrogen complexes have been isolated for almost all of the transition metals and lanthanides. However, simple coordination of N_2 did not immediately lead to any activation of the molecule. It took several years before a simple N_2 -compound could even be stoichiometrically converted to hydrazine [98]. The most promising results in this area have been reported by Laplaza and Cummins [99]. Their 1995 disclosure was the first example of a mild, catalytic conversion of N_2 to ammonia using a high valent Mo complex as catalyst [100]. The chapter presented by Prof. M. Hidai and Prof. Y. Mizobe entitled “Activation of the N–N Triple Bond in Molecular Nitrogen: Toward its Chemical Transformation into Organo-nitrogen Compounds” will discuss recent work concerning direct organonitrogen synthesis using transition metal catalysts.

7

Concluding Remark

The use of metal compounds in the activation of chemical bonds is an area of useful and fascinating research [101]. The following chapters will detail many of the recent advances in this area of chemistry with the emphasis on the use of traditionally unreactive substrates.

References

1. CRC Handbook of Chemistry and Physics. 72nd edn. (1991). CRC, Boston, Sect. D
2. Duff JM, Shaw BL (1972) *J Chem Soc Dalton Trans* 2219
3. Valentine JS (1973) *J Chem Soc Chem Commun* 857
4. Ryabov AD (1990) *Chem Rev* 90:403 and references cited therein
5. Parshall GW (1970) *Acc Chem Res* 3:139 and references cited therein
6. Omae I (1986) *Organometallic intramolecular-coordination compounds*. Elsevier, Amsterdam (and references cited therein)
7. Steenwinkel P, Gossage RA, van Koten G (1998) *Chem Eur J* 4:159
8. Bruce MI (1977) *Angew Chem* 89:75, *Angew Chem Int Ed Engl* 16:73
9. La Placa SJ, Ibers JA (1965) *Inorg Chem* 4:778
10. Roe DM, Bailey PM, Moseley K, Maitlis PM (1972) *Chem Commun* 1273
11. Crabtree RH, Hamilton DG (1988) *Adv Organomet Chem* 28:299
12. Richardson TB, Koetzle TF, Crabtree RH (1996) *Inorg Chim Acta* 250:69
13. Arndtsen BA, Bergman RG (1995) *Science* 270:1970
14. Gutiérrez E, Monge A, Nicasio MC, Poveda ML, Carmona E (1994) *J Am Chem Soc* 116:791 and references cited therein
15. Fukuyama T, Chatani N, Kakiuchi F, Murai S (1997) *J Org Chem* 62:5647
16. Murai S, Chatani N, Kakiuchi F (1997) *Catalysis Surveys from Japan* 1:35 and references cited therein
17. Sakakura T, Sodeyama T, Sasaki K, Wada K, Tanaka M (1990) *J Am Chem Soc* 112:7221 and references cited therein
18. Arndtsen BA, Bergman RG, Mobley TA, Peterson TH (1995) *Acc Chem Res* 28:154
19. Jones WD (1989) Alkane activation by cyclopentadienyl complexes of rhodium, iridium and related species. In: Hill CL (ed) *Activation and functionalization of alkanes*. John Wiley, NY, chap 4
20. Shilov AE (1994) *Activation of saturated hydrocarbons by transition metal complexes*. Reidel, Boston
21. Hoyano JK, Graham WG (1982) *J Am Chem Soc* 104:3723
22. Janowicz AH, Bergman RG (1982) *J Am Chem Soc* 104:352
23. Lohrenz JCW, Jacobsen H (1996) *Angew Chem* 108:1403, *Angew Chem Int Ed Engl* 35:1305
24. Pérez PJ, Poveda ML, Carmona E (1995) *Angew Chem* 107:242, *Angew Chem Int Ed Engl* 34:231
25. Schneider JJ (1996) *Angew Chem* 108:1132, *Angew Chem Int Ed Engl* 35:1068
26. Cundai TR (1994) *J Am Chem Soc* 116:340
27. Hovey AH, Morken JP (1996) *Angew Chem* 108:1378, *Angew Chem Int Ed Engl* 35:1262
28. Jonas RT, Stack TDP (1997) *J Am Chem Soc* 119:8566
29. Bromberg SE, Yang H, Asplund AT, McNamara BK, Kotz KT, Yeston JS, Wilkens M, Frei H, Bergman RG, Harris CD (1997) *Science* 278:260
30. Bengali AA, Arndtsen BA, Burger PM, Schultz RH, Weiller BH, Kyle KR, Bradley S, Moore C, Bergman RG (1995) *Pure Appl Chem* 67:281
31. Knapen JWJ, van der Made AW, de Wilde JC, van Leeuwen PWNM, Wijkens P, Grove DM, van Koten G (1994) *Nature* 372:359
32. van de Kuil LA, Grove DM, Gossage RA, Zwikker JW, Jenneskens LW, Drenth W, van Koten G (1997) *Organometallics* 16:4985
33. Gossage RA, van de Kuil LA, van Koten G (1998) *Acc Chem Res* 31:423
34. Friedrich HB, Moss JR (1991) *Adv Organomet Chem* 33:235
35. Kulawiec RJ, Crabtree RH (1990) *Coord Chem Rev* 99:89
36. Peng T-S, Winter CH, Gladysz JA (1994) *Inorg Chem* 33:2534
37. Pathak DD, Adams H, White C (1994) *J Chem Soc Chem Commun* 733

38. Huser M, Youinou M-T, Osborn J (1989) *Angew Chem* 101:1427
39. Olivèn M, Caulton KG (1997) *Chem Commun* 1733
40. Kiplinger JL, Richmond TG, Osterberg CE (1994) *Chem Rev* 94:373
41. Hughes RP (1990) *Adv Organomet Chem* 31:183
42. Kiplinger JL, Richmond TG (1996) *J Am Chem Soc* 118:1805
43. Kiplinger JL, Richmond TG (1993) *J Am Chem Soc* 115:5303
44. Bach I, Pöschke K-R, Goddard R, Kopiske C, Krüger C, Rufinska A, Seevogel K (1996) *Organometallics* 15:4959
45. Whittlesey MK, Perutz RN, Greener B, Moore MH (1997) *Chem Commun* 187
46. Whittlesey MK, Perutz RN, Moore MH (1996) *Chem Commun* 787
47. Edelbach BL, Jones WD (1997) *J Am Chem Soc* 119:7734
48. Jones WD, Partridge MG, Perutz RN (1991) *J Chem Soc Chem Commun* 264
49. Belt ST, Helliwell M, Jones WD, Partridge MG, Perutz RN (1993) *J Am Chem Soc* 115:1429
50. Blum O, Frolov F, Milstein D (1991) *J Chem Soc Chem Commun* 258
51. Kiplinger JL, King MA, Fechtenkotter A, Arif AM, Richmond TG (1996) *Organometallics* 15:5292
52. Weydert M, Anderson RA, Bergman RG (1993) *J Am Chem Soc* 115:8837
53. Deacon GB, Forsyth CM, Sun J (1994) *Tetrahedron Lett* 35:1095
54. Aizenberg M, Milstein D (1995) *Science* 265:359
55. Aizenberg M, Milstein D (1995) *J Am Chem Soc* 117:359
56. Schlosser M (ed, 1994) *Organometallics in synthesis: a manual*. John Wiley, New York
57. Davies SG (1982) *Organotransition metal chemistry: applications to organic synthesis*. Pergamon, Frankfurt
58. Keim W (1994) *New J Chem* 18:93
59. Schwarz H (1989) *Acc Chem Res* 22:282
60. Trost BM (1983) *Science* 219:245
61. Trost BM, Parquette JR, Marquart AL (1995) *J Am Chem Soc* 117:3284
62. Trost BM, Higuchi RI (1996) *J Am Chem Soc* 118:10094
63. Negishi E-I, Takahashi T (1994) *Acc Chem Res* 27:124
64. Trost BM, Grese TA (1991) *J Am Chem Soc* 113:7363
65. van Koten G, Gossage RA, Grove DM, Jastrzebski JTBH (1998) Selective product formation with organometallic radicals of nickel and zinc. In: Matyjaszewski K (ed) *Controlled radical polymerization*. ACS Pub, Washington, chap 5
66. Wissing E, van der Linden S, Rijnberg E, Boersma J, Smeets WJJ, Spek AL, van Koten G (1994) *Organometallics* 13:2602
67. Trost BM, Indolese AF, Miller TJJ, Treptow B (1995) *J Am Chem Soc* 117:615
68. Trost BM, Imi K, Davies IW (1995) *J Am Chem Soc* 117:5371
69. Trost BM, Toste FD (1996) *J Am Chem Soc* 118:6305
70. van der Boom ME, Kraatz H-B, Ben-David Y, Milstein D (1996) *Chem Commun* 2167 and references cited therein
71. Donkervoort JG, Vicario JL, Jastrzebski JTBH, Gossage RA, Cahiez G, van Koten G (1998) *J Organomet Chem* 558:61
72. de Vries AHM, Meetsma A, Feringa BL (1996) *Angew Chem* 108:2526, *Angew Chem Int Ed Engl* 35:2374
73. RajanBabu TV, Casalnuovo AL (1994) *Pure Appl Chem* 66:1535
74. Rijnberg E, Hovestad NJ, Kleij AW, Jastrzebski JTBH, Boersma J, Janssen MD, Spek AL, van Koten G (1997) *Organometallics* 16:2847
75. Hoveya AH, Moreken JP (1996) *Angew Chem* 108:1378, *Angew Chem Int Ed Engl* 35:1263
76. van Koten G (1994) *Pure Appl Chem* 66: 1455
77. Tsuji J (1975) *Organic synthesis by means of transition metal complexes*. Springer, Berlin Heidelberg New York (and references cited therein)

78. Sharma HK, Pannell KH (1995) *Chem Rev* 95:1351
79. Tamao K, Okazaki S, Kumada M (1978) *J Organomet Chem* 146: 87
80. Hayashi T, Matsumoto Y, Ito Y (1988) *J Am Chem Soc* 110:5579
81. Ito H, Ishizuka T, Tateiwa J, Sonoda M, Hosomi A (1998) *J Am Chem Soc* 120:11196
82. Sugimone M, Ito Y (1998) *J Chem Soc Dalton Trans* 1925
83. See for example: Gossage RA, McLennan G, Stobart SR (1996) *Inorg Chem* 35:1729
84. van Leeuwen PWNM, van Koten G (1993) Rhodium Catalyzed Hydroformylation. In: Moulijn JA, van Leeuwen PWNM, van Santen RA (eds) *Catalysis: an integrated approach to homogeneous, heterogeneous and industrial catalysis*. Elsevier, Amsterdam, Sect. 6.2
85. Elschenbroich Ch, Salzer A (1992) *Organometallics: a concise introduction*. VCH, Basel, chap 17
86. Mandai T, Matsumoto T, Kawada M, Tsuji J (1994) *Tetrahedron* 50:475
87. Shimizu I, Ishii H (1994) *Tetrahedron* 50:487
88. Yamamoto A, Ozawa F, Osakada K, Huang L (1991) *Pure Appl Chem* 63:687
89. Trost B (1981) *Pure Appl Chem* 53:2357
90. Trost B (1980) *Acc Chem Res* 13:385
91. Djakovich L, Moulines F, Astruc D (1996) *New J Chem* 20:1071
92. Maercher A (1987) *Angew Chem* 99:1002, *Angew Chem Int Ed Engl* 26:972
93. Bazhenova TA, Shilov AE (1995) *Coord Chem Rev* 144:69
94. Leigh GJ (1992) *Acc Chem Res* 25:177
95. Dilworth JR (1996) *Coord Chem Rev* 154:163
96. Richards RL (1996) *Coord Chem Rev* 154:83
97. Allen AD, Senoff CV (1965) *J Chem Soc Chem Commun* 621
98. Hidai M, Mizobe Y (1995) *Chem Rev* 95:1115
99. Laplaza CE, Cummins CC (1995) *Science* 268:861
100. For a critical evaluation of the Laplaza/Cummins system [99] see: Leigh GJ (1995) *Science* 268:827.
101. Herrmann WA, Cornils B (1997) *Angew Chem* 109:1074, *Angew Chem Int Ed Engl* 36:1048