

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/6184111>

A Computationally Designed Rh(I)-Catalyzed Two-Component [5+2+1] Cycloaddition of Ene-vinylcyclopropanes and CO for the Synthesis of Cyclooctenones

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · SEPTEMBER 2007

Impact Factor: 12.11 · DOI: 10.1021/ja072505w · Source: PubMed

CITATIONS

93

READS

44

10 AUTHORS, INCLUDING:



Jingxin Wang

Johns Hopkins University

24 PUBLICATIONS 494 CITATIONS

SEE PROFILE



Lei Jiao

Technische Universität München

29 PUBLICATIONS 1,248 CITATIONS

SEE PROFILE



Yong Liang

Peking University

42 PUBLICATIONS 1,446 CITATIONS

SEE PROFILE



Zhi-Xiang Yu

Peking University

94 PUBLICATIONS 2,678 CITATIONS

SEE PROFILE

A Computationally Designed Rh(I)-Catalyzed Two-Component [5+2+1] Cycloaddition of Ene-vinylcyclopropanes and CO for the Synthesis of Cyclooctenones

Yuanyuan Wang,[†] Jingxin Wang,[†] Jiachun Su,[†] Feng Huang,[†] Lei Jiao,[†] Yong Liang,[†] Dazhi Yang,[†] Shiwei Zhang,[†] Paul A. Wender,[‡] and Zhi-Xiang Yu^{*,†}

Beijing National Laboratory for Molecular Sciences (BNLMS), Key Laboratory of Bioorganic Chemistry and Molecular Engineering of Ministry of Education, College of Chemistry, Peking University, Beijing 100871, China, and Departments of Chemistry and of Chemical and Systems Biology, Stanford University, Stanford, California 94305-5080

Received April 10, 2007; E-mail: yuzx@pku.edu.cn


The design or discovery of new reactions for the synthesis of eight-membered rings has been stimulated by their occurrence in molecules of biomedical importance (e.g., taxol), ligands for catalysis (e.g., COT, COD), and precursors for natural products and novel materials (e.g., polyacetylenes).¹ The main difficulty encountered in the synthesis of eight-membered rings is the co-occurrence of unfavorable entropic and enthalpic factors associated with ring closure.¹ Several transition-metal-catalyzed cycloadditions² have been introduced in recent years to address this problem including Ni-, Rh-, or Ru-complex catalyzed [4+4], [6+2], [5+2+1], and [4+2+2] cycloadditions.³ Herein we report a computationally designed and experimentally verified, new two-component [5+2+1] cycloaddition of tethered ene-vinylcyclopropanes (ene-VCPs) and CO that provides for the synthesis of ring-fused cyclooctenones, a ring system encountered in numerous natural products, non-natural targets, and their precursors (Scheme 1).

Our method is based on work from the Wender laboratory leading to the first metal-catalyzed [5+2] cycloadditions between vinylcyclopropanes and 2- π components (such as alkynes, alkenes, and allenes) which afford seven-membered ring cycloadducts.⁴ [Rh(CO)₂Cl]₂ was found to be an especially effective and versatile catalyst for this and many related cycloadditions.⁵ Importantly, however, the intramolecular [5+2] cycloaddition of ene-VCPs has not been realized using [Rh(CO)₂Cl]₂ as catalyst,^{4b} even though this transformation can be achieved with RhCl(PPh₃)₃ or [(arene)Rh(COD)]-SbF₆ catalyst.^{4c}

We speculated that the failure of [Rh(CO)₂Cl]₂ to catalyze the [5+2] cycloaddition of ene-VCPs (relative to yne-VCPs and allenyl-VCPs) could be attributed to the reductive elimination (RE) step in the catalytic cycle (Scheme 1).⁶ The RE step in the intramolecular [5+2] cycloaddition of ene-VCP would lead to the formation of an (sp³)C–C(sp³) bond. However, it is known that this kind of RE reaction is not facile when compared to the migratory reductive elimination (MRE) of an (sp²)C–M–C(sp³) subunit for the formation of an (sp²)C–C(sp³) bond.^{6,7} We hypothesized that raising the reaction temperature or extending the reaction time of the [Rh(CO)₂Cl]₂ catalyzed intramolecular [5+2] cycloaddition of an ene-VCP could enable this reaction to occur, if no side reactions compete (hypothesis A). We further hypothesized that the putative intermediate **I** (Scheme 1), whose RE to the [5+2] product is disfavored, would in the presence of CO be converted to intermediate **II**, whose MRE to a [5+2+1] cycloadduct would be favored by the formation of an (sp²)C–C(sp³) bond (hypothesis B).

Before testing our hypotheses experimentally, we computed the energies for the RE, the CO insertion, and MRE steps shown in


Scheme 1. Rationale for the Two-Component [5+2+1] Reaction



Scheme 1 (X = CH₂) using the B3LYP method.⁸ The activation energies of the RE steps are about 25–30 kcal/mol, whereas the CO insertion and MRE steps have activation energies of about 13–14 and 23–24 kcal/mol, respectively, suggesting that the [5+2+1] path would be favored over the [5+2] path and the dominant product would be the [5+2+1] cycloadducts.⁸

Encouraged by the above preliminary calculations, we began our experimental test. We observed that, in the absence of CO, the [Rh(CO)₂Cl]₂ catalyzed (10 mol %) reaction of ene-VCP substrate **1** at 110 °C did indeed give a double-bond-isomerized [5+2] cycloadduct **3** in 59% isolated yield, together with 10% of the

Table 1. Optimization Studies of the [5+2+1] Cycloadditions

							
entry	CO [atm]	catalyst [mol %]	T [°C]	solvent	conc [M]	t [h]	yield [%]
1	0	10% [Rh(CO) ₂ Cl] ₂	110	toluene	0.05	24	10 ^a
2	1	5% [Rh(CO) ₂ Cl] ₂	80	dioxane	0.05	5	44 ^b
3	4	5% [Rh(CO) ₂ Cl] ₂	80	dioxane	0.05	24	8
4	0.2 ^c	5% [Rh(CO) ₂ Cl] ₂	80	dioxane	0.05	5	70 ^d
5	0.2	5% [Rh(CO) ₂ Cl] ₂	60	dioxane	0.05	48	17
6	0.2	5% [Rh(CO) ₂ Cl] ₂	90	dioxane	0.05	5	70
7	0.2	5% [Rh(CO) ₂ Cl] ₂	100	dioxane	0.05	5	61
8	0.2	5% [Rh(CO) ₂ Cl] ₂	80	DCE	0.05	5	62 ^e
9	0.2	5% [Rh(CO) ₂ Cl] ₂	80	toluene	0.05	12	14
10	0.2	5% [Rh(CO) ₂ Cl] ₂	80	dioxane	0.01	5	68
11	0.2	5% [Rh(CO) ₂ Cl] ₂	80	dioxane	0.20	5	34
12	0.2	10% [Rh(CO) ₂ Cl] ₂	80	dioxane	0.05	5	72
13	1	10% RhCl(PPh ₃) ₃	80	dioxane	0.05	17	N.R.
14	1	10% RhCl(PPh ₃) ₃ + 10% AgOTf	80	dioxane	0.05	18	23 ^f
15	1	5% [Rh(CO) ₂ Cl] ₂ + 10% AgOTf	80	dioxane	0.05	13	7

^a Accompanied with a [5+2] cycloadduct **3** (59%); see Supporting Information for details. ^b Cis/trans = 5:1. ^c Conditions: 0.2 atm CO + 0.8 atm N₂. ^d Cis/trans > 20:1. ^e Cis/trans = 4:1. ^f Cis/trans = 1:1.


[†] Peking University.

[‡] Stanford University.

Table 2. Rh(I)-Catalyzed [5+2+1] Cycloaddition Reactions of Ene-vinylcyclopropane Substrates and CO^a

Ene-VCP Substrates	1	4	6	8	10	12	14	16	18	20
Cyclo-adducts										
	70%	81%	90% ^b	29%	71%	78%	92%	73%	83% ^d	90% ^e

^a E = CO₂Me. Isolated yields were reported unless otherwise indicated. ^b GC yield. Isolated yield is 44% owing to the volatility of the product. ^c Confirmed by X-ray analysis. ^d A [5+2] product was obtained in 11% yield. ^e Combined yield of diastereomers (trans/cis = 5:1).

Scheme 2

^a Confirmed by X-ray analysis.

[5+2+1] cycloadduct **2** in which the carbonyl group comes from the CO ligand of the [Rh(CO)₂Cl]₂ catalyst (Table 1, entry 1). This result supported our hypothesis A that [5+2] ene-VCP cycloadditions can be achieved when the reaction temperature is increased. We then turned our attention to promoting the [5+2+1] cycloaddition. Gratifyingly, substrate **1** in the presence of CO (balloon, 1 atm) and 5 mol % [Rh(CO)₂Cl]₂ catalyst in dioxane, gave after 5 h at 80 °C the [5+2+1] cycloadduct, cyclooctenone **2** (cis/trans = 5:1), in 44% isolated yield (entry 2). We then systematically optimized the reaction conditions for this new [5+2+1] cycloaddition using substrate **1**. Comparison of the reactions in entries 2 to 12 suggested that the optimal conditions for the [5+2+1] cycloaddition were the use of a substrate concentration of 0.05 M, 5 mol % [Rh(CO)₂Cl]₂ catalyst, CO (delivered by balloon admixed with N₂ in the ratio of 1:4 at a pressure of 1 atm),⁹ dioxane as solvent, and a reaction temperature of 80–90 °C (entries 4 and 6). Under these conditions, substrate **1** gave the cis-fused [5+2+1] cycloaddition product **2** as a single diastereomer. Other Rh(I) catalysts such as Wilkinson's complex and the cationic rhodium(I) catalysts were found ineffective for the [5+2+1] cycloaddition (entries 13, 14, and 15).

With the above optimal conditions in hand, we studied the Rh(I)-catalyzed [5+2+1] cycloadditions of diverse substrates (Table 2). The results indicate that the cycloaddition reactions are tolerant of tethers incorporating geminal diester, sulfonamide, and ether functionalities. Comparison of reaction yields of substrates with different tethers shows that heteroatom substitution in the tether results in higher yields of [5+2+1] cycloadducts. This [5+2+1] cycloaddition also tolerates methyl substitution on the alkene (substrates **8**, **10**, and **12**) and the VCP (substrates **18**, **22**, **24**, **25**, and **27**). Phenyl substitution of the cyclopropane leads efficiently to the [5+2+1] cycloadducts **13**, **15**, **17**, and **19**. In addition to the 5/8-ring system, the trans-fused 6/8-ring system can also be efficiently established in the cycloaddition reaction from **20** to **21**, albeit with decreased diastereoselectivity. It is noteworthy as well that the E/Z geometry of the C=C bonds in the VCP moieties of ene-VCPs affects the cis/trans stereochemistry of the bicyclic products (Scheme 2). In addition to its synthetic merit, this has

mechanistic implications suggesting that the starting alkene geometry is conserved through the multistep pathway.

In conclusion, a new Rh(I)-catalyzed [5+2+1] cycloaddition has been designed computationally and verified experimentally. This study provides a flexible, convenient, and efficient method for constructing bicyclic cyclooctenones. The reaction proceeds in good to excellent yields with a variety of tether types and substitution patterns and allows for the preparation of 5/8- and 6/8-fused ring systems even those containing quaternary centers. Further studies of this reaction (scope, mechanism, and stereochemistry) and its synthetic applications to natural products are being pursued.

Acknowledgment. We are indebted to generous financial support from Peking University, the Natural Science Foundation of China (9800445, 0240203 and 20672005), and the National Science Foundation USA (Grant CHE-040638). Professor K. N. Houk of UCLA is highly appreciated for his support to Z.X.Y. and the theoretic and synthetic organic chemistry lab at PKU.

Supporting Information Available: Computational and experimental details. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (a) Yet, L. *Chem. Rev.* **2000**, *100*, 2963. (b) Michaut, A.; Rodriguez, J. *Angew. Chem., Int. Ed.* **2006**, *45*, 5740. (c) Sieburth, S. M.; Cunard, N. T. *Tetrahedron* **1996**, *52*, 6251. (d) Mehta, G.; Singh, V. *Chem. Rev.* **1999**, *99*, 881. (e) Wender, P. A.; Croatt, M. P.; Deschamps, N. M. in *Comprehensive Organometallic Chemistry III*; Ojima, I., Ed.; Elsevier Ltd: 2007; Vol. 10, pp 603–648. (f) Korshak, Y. V.; Korshak, V. V.; Kanischka, G.; Hocker, H. *Makromol. Chem., Rapid Commun.* **1985**, *6*, 685. (g) Scherman, O. A.; Rutenberg, I. M.; Grubbs, R. H. *J. Am. Chem. Soc.* **2003**, *125*, 8515.
- (2) Recent reviews: (a) Lautens, M.; Klute, W.; Tam, W. *Chem. Rev.* **1996**, *96*, 49. (b) Saito, S.; Yamamoto, Y. *Chem. Rev.* **2000**, *100*, 2901. (c) Murakami, M. *Angew. Chem., Int. Ed.* **2003**, *42*, 718.
- (3) (a) Wender, P. A.; Ihle, N. C. *J. Am. Chem. Soc.* **1986**, *108*, 4678. (b) Wender, P. A.; Christy, J. P. *J. Am. Chem. Soc.* **2006**, *128*, 5354. (c) Wender, P. A.; Correa, A. G.; Sato, Y.; Sun, R. *J. Am. Chem. Soc.* **2000**, *122*, 7815. (d) Wender, P. A.; Gamber, G. G.; Hubbard, R. D.; Zhang, L. *J. Am. Chem. Soc.* **2002**, *124*, 2876. (e) Wegner, H. A.; de Meijere, A.; Wender, P. A. *J. Am. Chem. Soc.* **2005**, *127*, 6530. (f) Murakami, M.; Ashida, S. R.; Matsuda, S. *J. Am. Chem. Soc.* **2006**, *128*, 2166. (g) Gilbertson, S.; DeBoef, B. *J. Am. Chem. Soc.* **2002**, *124*, 8784. (h) Evans, P. A.; Baum, E. W.; Fazal, A. N.; Pink, M. *Chem. Commun.* **2005**, 63.
- (4) (a) Wender, P. A.; Takahashi, H.; Witulski, B. *J. Am. Chem. Soc.* **1995**, *117*, 4720. (b) Wender, P. A.; Sperandio, D. *J. Org. Chem.* **1998**, *63*, 4164. (c) Wender, P. A.; Williams, T. *J. Angew. Chem., Int. Ed.* **2002**, *41*, 4550.
- (5) (a) Wender, P. A.; Pedersen, T. M.; Scano, M. J. C. *J. Am. Chem. Soc.* **2002**, *124*, 15154. (b) Jeong, N.; Sung, B. K.; Choi, Y. K. *J. Am. Chem. Soc.* **2000**, *122*, 6771. (c) Brummond, K. M.; Chen, H.; Sill, P.; You, L. *J. Am. Chem. Soc.* **2002**, *124*, 15186. (d) Wender, P. A.; Paxton, T. J.; Williams, T. J. *J. Am. Chem. Soc.* **2006**, *128*, 14814.
- (6) For a theoretical study on the Wender [5+2] cycloaddition, see: Yu, Z.-X.; Wender, P. A.; Houk, K. N. *J. Am. Chem. Soc.* **2004**, *126*, 9154.
- (7) Ozawa, F.; Mori, T. *Organometallics* **2003**, *22*, 3593 and references therein.
- (8) Computational details, references, and results are given in the Supporting Information.
- (9) It was found that the Pauson–Khand reaction catalyzed by [Rh(CO)₂Cl]₂ was very effective under a low CO pressure, see: Kobayashi, T.; Koga, Y.; Narasaka, K. *J. Organomet. Chem.* **2001**, *624*, 73.

JA072505W