

David B. Weinreb

Re: Detection of JC virus sequences in colorectal cancers in Japan

Received: 18 October 2005 / Accepted: 26 October 2005 / Published online: 20 December 2005
© Springer-Verlag 2005

To the Editors:

We read “Detection of JC virus DNA sequences in colorectal cancers in Japan” by Hori et al. with great excitement. These authors offer convincing evidence that polyoma T antigen may be integrated into the cellular genome of colon carcinoma tissues, although the T-antigen protein is not expressed [4]. We enthusiastically agree with their conclusions and would like to offer additional data to advance this discussion.

First, we have investigated the presence of polyomavirus (PV) T antigen in tumors by immunohistochemistry (IHC), using a primary antibody that reacts with the large T antigen of both the JC and BK viruses [1, 5]. T antigen was not detected in any of 50 cases of colon carcinoma, 18 cases of gastric cancer, or 15 cases of bladder carcinoma.

In addition, we examined by IHC the presence of T antigen in six cases of polyoma virus nephropathy (PVN) in renal transplant patients. In all of these six patients, robust nuclear staining specific for T antigen was observed in scattered tubular cells. The infected tubular cells were characterized by distinct cytopathological features, including anisonucleosis, hyperchromasia with clumped chromatin, and intranuclear inclusions. Such cellular changes represent a “permissive” infection, i.e., a state of viral replication. These histopathological features are absent in our 83 cases of carcinoma.

We agree with these authors that if PV is present in tumors, it is likely to be integrated into the cellular genome, representing a “nonpermissive” infection in which viral replication does not occur.

We conclude that future IHC studies of tumors are likely to be uninformative because we fail to detect viral antigens in nonpermissive infections, which are precisely the category of infections hypothesized to drive oncogenesis. Although recent studies have detected T antigen by IHC in esophageal [2] and colon [3] carcinoma, it remains to be seen whether these data truly represent evidence for a viral etiology in these tumors, or, alternatively, a sequelae of the patient’s immunocompromised state.

We look forward to hearing the results of future studies from Dr. Hori’s team.

References

1. Del Valle L, Gordon J, Assimakopoulou M, Enam S, Geddes JF, Varakis JN, Katsetos CD, Croul S, Khalili K (2001) Detection of JC virus DNA sequences and expression of the viral regulatory protein T-antigen in tumors of the central nervous system. *Cancer Res* 61:4287–4293
2. Del Valle L, White MK, Enam S, Oviedo SP, Bromer MQ, Thomas RM, Parkman HP, Khalili K (2005) Detection of JC virus DNA sequences and expression of viral T antigen and agnoprotein in esophageal carcinoma. *Cancer* 103(3):516–527
3. Enam S, Del Valle L, Lara C, Gan DD, Ortiz-Hidalgo C, Palazzo JP, Khalili K (2002) Association of human polyomavirus JCV with colon cancer: evidence for interaction of viral T-antigen and beta-catenin. *Cancer Res* 62(23):7093–7101
4. Hori R, Murai Y, Tsuneyama K, Abdel-Aziz HO, Nomoto K, Takahashi H, Cheng CM, Kuchina T, Harman BV, Takano Y (2005) Detection of JC virus DNA sequences in colorectal cancers in Japan. *Virchows Arch* 14:1–8
5. Mann RS, Carroll RB (1984) Cross-reaction of BK virus large T antigen with monoclonal antibodies directed against SV40 large T antigen. *Virology* 138:379

D. B. Weinreb (✉)
School of Medicine,
Mount Sinai Hospital,
Room 11C1, 50 East 98th Street,
New York, NY 10029, USA
e-mail: David.weinreb@mssm.edu
Tel.: +1-917-8339472