

New Pharmacologic Agents for Diabetes

Clifford J. Bailey, PhD, FRCP, FRCPath

Address

Department of Pharmaceutical Sciences, Aston University,
Aston Triangle, Birmingham B4 7ET, UK.

Current Diabetes Reports 2001, 1:119–126

Current Science Inc. ISSN 1534-4827

Copyright © 2001 by Current Science Inc.

New agents are being developed to address the underlying endocrinopathies and metabolic disturbances of type 2 diabetes. Stimulants of the nuclear peroxisome proliferator-activated receptor γ (PPAR γ) are being identified to selectively improve insulin actions, and dual agonists of PPAR γ and PPAR α are being evaluated for enhanced control of hyperglycemia and dyslipidemia. Novel activators of insulin receptor phosphorylation and inhibitors of receptor dephosphorylation are offering encouraging leads for new agents. Analogues of glucagon-like peptide-1 that increase glucose-induced insulin secretion may additionally increase β -cell neogenesis from progenitor duct cells. The amylin analogue pramlintide, which suppresses glucagon secretion and reduces weight, is advancing in clinical trial. Direct stimulants of glucose utilization and partial inhibitors of gluconeogenesis are providing useful new drug templates. Thus, new pharmacologic approaches are emerging to treat the multiple lesions of type 2 diabetes.

Introduction

Established and recently introduced oral agents for the treatment of type 2 diabetes offer substantial efficacy through a range of actions (Table 1). However they seldom reinstate and sustain normal glycemic control [1], leaving an urgent need for improved therapies. This article evaluates strategies for the future drug treatment of type 2 diabetes with agents other than insulins. It examines proposed sites of intervention and potential new pharmacologic approaches to improve glycemic control.

Current Treatment Strategies

It is now abundantly clear that improved glycemic control translates into benefits against the chronic vascular complications of type 2 diabetes. Hence, current treatment strategies are designed to achieve the best possible glycemic control, consonant with the circumstances of the individual patient [2]. Approaches to the treatment process have been shaped by an appreciation that type 2 diabetes typically emerges through the composite effects of insulin

resistance and β -cell dysfunction, often aggravated by the impositions of dyslipidemia [3]. Thus, early and intensive intervention against each of these underlying pathogenic features is advisably taken into account in the selection of drug therapy, often requiring combinations of two and possibly three differently acting agents. The choice and timing of therapies can also provide a balanced control of basal and postprandial components of the hyperglycemia. The latter component is presently receiving added attention following reports that it is strongly predictive of fatal vascular events [4].

Hyperglycemia is one of a constellation of risk factors for the macrovascular complications of type 2 diabetes, collectively encompassed within the insulin resistance (or metabolic) syndrome. All of these risk factors, which include body weight, lipids, blood pressure, and clotting status should be considered in the treatment process. In this context the concept of “treating to target” has emerged as a valuable, if somewhat arbitrary guide to the ideal, or least damaging, values to strive for [2].

Future Treatment Strategies


Future treatment strategies for type 2 diabetes are expected to integrate existing and new agents against insulin resistance, β -cell failure, and dyslipidemia to improve the prospect of attaining and consolidating treatment targets (Fig. 1) [5,6•]. The treatment algorithm might therefore reflect more closely the natural history of type 2 diabetes, which presents different therapeutic challenges at different stages throughout its progression. Because insulin resistance has been implicated in the early metabolic disturbances of type 2 diabetes as well as many accompanying vascular risk factors, it may be appropriate to underpin the treatment process from the outset with agents that will improve or mimic the metabolic actions of insulin. As the various intracellular defects of insulin action become better characterized, additional agents to counter insulin resistance by new cellular mechanisms can be anticipated.

Incessant deterioration of β -cell function and loss of β -cell mass are crucial determinants of antidiabetic therapy, because a severe decline in insulin concentrations eventually precludes the effective use of current oral agents. It is still uncertain whether β -cell function or mass can be preserved by early and aggressive intervention with current agents. Intermittent (*eg*, prandial) rather than continual stimulation of β -cell function, early use of insulin therapy or agents to relieve insulin resistance, and unique β -cell

Table 1. Oral agents for the treatment of type 2 diabetes

Agents	Examples of agents	Mode of action	Time of action
Alpha-glucosidase inhibitors Sulfonylureas	Acarbose, miglitol, voglibose Glimepiride, glyburide,* glipizide, gliclazide, tolbutamide	Slow rate of carbohydrate digestion Stimulate insulin secretion	Mainly postprandial Basal and postprandial
Meglitinides	Repaglinide, nateglinide	Stimulate insulin secretion: prandial (short-acting)	Mainly postprandial
Biguanide Thiazolidinediones	Metformin Rosiglitazone, pioglitazone	Improve insulin action Improve insulin action (PPAR γ agonists)	Basal and postprandial Basal and postprandial

*Glyburide is also known as glibenclamide. Fixed-dose combination tablets of glyburide with metformin are available.

**Figure 1.** Potential intervention sites for new agents to treat type 2 diabetes.

effects of thiazolidinediones have all been mooted in this respect. An important future focus of antidiabetic therapy must be to prevent the apoptosis and promote replacement neogenesis of β cells. From a functional perspective it will be advantageous for new insulin secretagogues to act in a glucose-dependent manner to strongly counter postprandial hyperglycemia and obviate hypoglycemia. β -cell function would also be improved if such agents stimulated insulin biosynthesis commensurate with their stimulation of secretion. For the interim, substantial benefits can be gained by mixing oral agents with insulin, and taking advantage of the expanding range of insulin preparations, delivery routes, and devices.

In the future, dyslipidemia might be considered for primary intervention in type 2 diabetes. The lipotoxic effects of chronically raised fatty acid and triglyceride concentrations involve detrimental influences on insulin action and β -cell function, and independently increase vascular risk. Glucotoxicity also exacerbates insulin resistance and β -cell failure while promoting glycation and its clinical sequelae. Thus, agents that reduce blood glucose concentrations by any safe means could, in principle, serve a useful therapeutic purpose. Because the hyperglycemia of type 2 diabetes derives in part from excess hepatic glucose output, agents that suppress gluconeogenesis and glycogenolysis are suitable candidates, but they must not cause

Table 2. Examples of PPAR γ agonists and dual PPAR γ and α agonists under investigation as potential new oral antidiabetic agents

PPAR γ agonists	Manufacturer
Thiazolidinediones	
BM-13.1246 and BM-15.2054	Boehringer Mannheim
CS-011/CI-1037*	Sankyo
MCC-555	Mitsubishi Chemical
NC-2100	Nipon Chemipharm
NIP-221* and NIP-223	Nissan Chemical
T-174	Tanabe
TZD-300512*	Takeda and Lilly
Nonthiazolidinediones	
BM-17.0744	Boehringer Mannheim
GI-262570*	GlaxoSmithKline
GW-1929	GlaxoSmithKline
L-764406	Merck
Dual PPAR γ and α agonists	
Thiazolidinediones	
DRF-2189	Dr. Reddy's Research Foundation
KRP-297	Kyorin Pharmaceutical
Nonthiazolidinediones	
JTT-501	Japan Tobacco and Pharmacia
BMS-298585	Bristol-Myers Squibb
AZ-242	AstraZeneca
NN-622	NovoNordisk

*Peroxisome proliferator-activated receptor γ (PPAR γ) agonists reported to significantly lower plasma triglycerides may be acting in part via weak PPAR α agonism. Dual PPAR γ and α agonists are selected for plasma triglyceride-lowering activity as well as improved insulin sensitivity and glucose-lowering.

extensive or irreversible inhibition of hepatic glucose output if hypoglycemia is to be avoided. Impaired glucose uptake and utilization by tissues that are acutely insulin-dependent could be addressed, at least in part, by agents that directly stimulate these functions.

Weight loss in overweight type 2 diabetic patients can effectively reduce hyperglycemia [2]. There is an opportunity for greater use of safe antiobesity agents as initial intervention. In anticipation that rigorous control of hypertension and other modifiable cardiovascular risk factors will be routine in type 2 patients, new antidiabetic agents should be compatible with existing standard treatments and prophylactic measures against these risks.

Addressing Insulin Resistance

Several agents are in development to improve or mimic insulin action. Particular research interest is presently focused on stimulants of the nuclear peroxisome proliferator-activated receptor γ (PPAR γ). However, clinical studies with most of these compounds are in their early stages and a new product is not imminently poised to enter the market.

PPAR γ agonists

Stimulating PPAR γ is now an established route to improve insulin sensitivity in type 2 diabetes, as illustrated by the thiazolidinediones in present clinical use, namely rosiglitazone and pioglitazone [7•]. PPAR γ is expressed mainly in adipose tissue where it promotes adipogenesis, lipogenesis, and glucose uptake. Stimulation of PPAR γ causes transcription of certain insulin-sensitive genes such as lipoprotein lipase, fatty acid transporter protein, adipocyte fatty acid-binding protein (aP2), fatty acyl-CoA synthase, malic enzyme, glucokinase, phosphoenolpyruvate carboxykinase, and glucose transporter isoform-4 (Glut-4) [8,9]. The blood glucose-lowering effect of PPAR γ agonists is at least partly due to increased adipocyte lipogenesis. This decreases circulating concentrations of non-esterified fatty acids, which resets the glucose-fatty acid (Randle) cycle to favor glucose utilization by muscle [10]. PPAR γ agonists also improve insulin action by reducing adipocyte production of the cytokine tumor necrosis factor α (TNF α) and the hormone resistin, as well as activating phosphatidylinositol 3-kinase (PI3K) and possibly reducing leptin levels [10–12]. However, PPAR γ agonists remain effective in animal models that are almost devoid of adipose tissue, emphasizing the importance of effects on other tissues [13]. PPAR γ is expressed in skeletal muscle and liver, and direct actions of PPAR γ agonists in these tissues could contribute to the blood glucose-lowering effect [14,15]. However, the actions of PPAR γ in these tissues have yet to be resolved [16•]. Hypomagnesemia is believed to aggravate insulin resistance in some type 2 diabetic individuals, and PPAR γ agonists have been reported to address this problem [17]. Also, some actions of thiazolidinedione PPAR γ agonists occur too rapidly to be explained by a genomic action, suggesting that some effects are independent of the PPAR γ receptor [10].

Many new PPAR γ agonists are reported to be under investigation as potential antidiabetic agents (Table 2). Some of these are thiazolidinediones [14,17–23], but there are several nonthiazolidinedione PPAR γ agonists [24–27]. The thiazolidinediones have previously shown a close positive correlation between their binding affinity for PPAR γ and their blood glucose-lowering efficacy in animal models of type 2 diabetes [28]. However, several new thiazolidinediones with weak PPAR γ agonism, such as MCC-555 and NC-2100, have shown potent glucose-lowering activity in glucose intolerant and type 2 diabetic animal models [19,20]. This indicates that variations in the nonthiazolidinedione moiety make an important contribution to the glucose-lowering efficacy. This could partly reflect the structure of the ligand-binding domain of PPAR γ , which is sufficiently large to allow the different PPAR γ agonists to produce slightly different conformational changes to the receptor. This in turn will alter the recruitment of coactivators that will modify the selection of genes that are transcribed [29,30]. Thus, the actions of PPAR γ agonists can be modified by structural changes to the nonthiazolidinedione part of the molecule for customized selection of the required therapeutic profile.

Although PPAR γ is expressed by pancreatic β cells, most studies have shown little or no direct effect of thiazolidinediones on total insulin secretion [7•,10]. However, thiazolidinediones may improve the pattern of insulin secretion and the morphology of the pancreatic islets in diabetic states, and defer or prevent β -cell hyperplasia while restoring granulation in animal models [31,32]. Interestingly, the peroxisome proliferator response element has been identified in the promoter of the *GLUT2* gene, which is expressed in β cells [33]. The prospect that PPAR γ agonists might limit detrimental changes to β cells in patients with impaired glucose tolerance raises the possibility of a preventative application for these agents.

Some further potentially advantageous effects of thiazolidinedione PPAR γ agonists include vascular protection through decreased platelet aggregation, increased fibrinolysis, and possibly an indirect decrease in plasminogen activator inhibitor-1 [7•]. Independent effects on glomerular structure, which reduce nephropathic symptoms, have also been reported [32,34]. Additionally, thiazolidinediones have been ascribed with putative effects on inflammation, atherogenesis, and tumor proliferation [35]. It is pertinent to note that the idiosyncratic liver toxicity that occurred during use of the first clinically available thiazolidinedione, troglitazone, has not been observed with later thiazolidinediones in clinical use, but preclinical tests may not discriminate the potential for hepatotoxicity in humans [36]. PPAR γ agonists are prone to cause fluid retention [7•,10], and newer agents should be chosen to minimize this effect. Weight gain is another unwanted effect of PPAR γ stimulation [7•,10], but weaker agonists, which also increase the expression of uncoupling proteins (and combination with a PPAR α or a retinoid X receptor [RXR] agonist), could be helpful in this respect [20].

Thiazolidinedione PPAR γ agonists have consistently reduced circulating nonesterified fatty acids, but effects on other lipid parameters are variable. Limited evidence from the clinical use of thiazolidinediones suggests that the diversity of lipid changes probably reflects the structure of the nonthiazolidinedione part of the molecule and the pretreatment lipid profile of the patient [7•,8–10]. In hyperlipidemic patients, circulating total triglycerides can decline, and cell culture studies indicate increased lipid oxidation, implying reduced fat deposition in muscle [16•]. Low-density lipoprotein (LDL) cholesterol and high-density lipoprotein cholesterol may increase during initial treatment with a thiazolidinedione, although the ratio of these fractions is little changed. Preliminary reports indicate that LDL cholesterol may return to pretreatment levels during chronic use.

Dual PPAR γ and PPAR α agonists

Peroxisome proliferator-activated receptor α is expressed in muscle, liver, and various other tissues where it promotes the oxidation of fatty acids through increased transcription of enzymes such as acyl-CoA synthase, acyl-CoA oxidase,

and thiolase [9]. Stimulation of PPAR α (eg, by fibrates) is an established mechanism to lower circulating lipid concentrations. Several established PPAR γ agonists are known to bind PPAR α with low affinity, and enhanced stimulation of PPAR α is associated with a greater lipid-lowering effect. Hence, dual agonists of PPAR γ and PPAR α are now seen as an opportunity to achieve both glucose- and lipid-lowering effects (Table 2). Several thiazolidinedione and nonthiazolidinedione compounds are receiving particular attention in this respect [37–39]. It will be important for such agents to selectively balance the two agonistic effects for optimal therapeutic efficacy at the same circulating concentration.

Dual PPAR γ and RXR agonists

Peroxisome proliferator-activated receptor γ exists as a heterodimer with the RXR, and coactivation of PPAR γ and RXR has been reported to cause an additive or synergistic improvement of insulin action in skeletal muscle compared with stimulation of PPAR γ alone [16•]. Stimulating RXR alone may be ineffective or produce variable effects such as reduced food intake, reduced body weight gain, increased expression of uncoupling proteins, and increased expression of the p85a subunit of PI3K [40–42]. Each of these effects could facilitate insulin action, but they may depend on structural components of the RXR agonist that do not actually bind to the RXR ligand-binding site of the RXR receptor. Thus, binding of an RXR agonist to the RXR may be accompanied by interactions of the agonist with other regions of the receptor, and this will alter the profile of genes transcribed. Thus, there is a potential for selective RXR agonists and dual PPAR γ -RXR agonists.

Mineral enhancers of insulin action

Several mineral supplements, notably magnesium and chromium, will improve insulin action in patients who are deficient in these minerals. The roles of these minerals in insulin signaling and energy metabolism have been reviewed previously [5].

Salts of vanadium, molybdenum, selenium, and tungsten can improve or mimic some metabolic actions of insulin in isolated cells and tissues. They also improve glycemic control in insulin-deficient and hyperinsulinemic animal models of diabetes, suggesting a possible therapeutic use [43–47]. Vanadium salts appear to prolong insulin receptor signaling by inhibiting phosphatases that dephosphorylate and deactivate the tyrosine kinase activity of the β subunit. They also improve muscle glucose uptake and suppress hepatic glucose output by other uncharacterized mechanisms that bypass the known pathways of insulin action—hence, attracting the label of insulin mimetic agents [44]. Concerns about toxicity and bioavailability of vanadium salts may be allayed by highly potent peroxovanadium salts and other organic vanadium complexes [6•,43,44]. Molybdenum and tungsten salts act predominantly to decrease hepatic glucose output, but the mecha-




Figure 2. Actions of glucagon-like peptide-1 (GLP-1) analogues to promote neogenesis and proliferation of pancreatic β cells.

nism of action of these salts and selenium salts remains to be established [45–47].

Other insulin action enhancers

One approach to overcome the multiple postreceptor signaling defects of insulin resistance [48] has been to interrupt the intracellular negative feedback effects of downstream signals such as PI3K and protein kinase B, which normally cause a temporary suppression of subsequent insulin signaling [49]. For example, like vanadium salts, various compounds such as benzonaphthofurans and thiophenes inhibit protein tyrosine phosphatase 1B, improving insulin action in insulin-resistant obese-diabetic animals [50]. The tyrosine kinase activity of the insulin receptor and insulin receptor substrates is inhibited by serine-threonine phosphorylation. Certain isoforms of protein kinase C (PKC) have been implicated in this process, providing a potential intervention target [51].

Direct stimulation of insulin receptor tyrosine kinase activity has recently been reported using metabolite (LY783281) from a *Pseudomassaria* fungus [52] and a synthetic molecule (TLK16998) [53•]. Both compounds appear to promote phosphorylation of the intracellular segment of the β subunit of the insulin receptor. Other agents that enhance or partially mimic insulin-signaling intermediates by undefined mechanisms could also provide leads for the development of new antidiabetic agents, as considered elsewhere [5,6•,54–56].

Addressing β -cell Dysfunction

Although current insulin-releasing agents make an important contribution to the treatment of type 2 diabetes, they do not actually reinstate normality of the glucose-sensing process or the pathways of insulin production by individual β cells. Also, they do not restore the diminishing β -cell mass. Several agents in development promise to address these issues.

GLP-1, exendin-4, and DPP-IV inhibitors

The intestinal hormone glucagon-like peptide-1 (GLP-1) is released during meal digestion and potentiates nutrient-induced insulin secretion and proinsulin biosynthesis

[57•]. It acts on the β cell via specific receptors coupled to the adenylate cyclase-cAMP protein kinase A (PKA) pathway and other intracellular mediators. Release of endogenous GLP-1 is reduced in diabetic states [58], and the ability of GLP-1 to enhance predominantly glucose-stimulated insulin secretion facilitates the control of postprandial hyperglycemia without increased risk of hypoglycemia. GLP-1 increases both the acute and chronic phases of glucose-stimulated insulin secretion, and partly restores the normal pulsatile pattern of insulin secretion [57•,59]. This is complemented by increased β -cell granulation, resulting from increased stimulation of proinsulin biosynthesis through PKA-dependent and independent mechanisms [60]. (It is noteworthy that meglitinide insulin secretagogues are also used to target postprandial hyperglycemia, but they act on the β cell in a similar manner to sulfonylureas, which “induce” rather than “potentiate” insulin secretion [5]).

The main biologically active form of GLP-1 (the truncated 7-36 amide) and various analogues are being assessed as potential antidiabetic drugs. Receptors for GLP-1 are widely distributed, and GLP-1 exerts a satiety-inducing effect, reduces glucagon secretion, and slows gastric emptying [57•]. Animal studies have suggested that GLP-1 has further extra-pancreatic effects that improve insulin action, but this has not been a consistent finding in human type 2 diabetes [61]. A recent study in type 1 patients indicated that GLP-1 improves insulin-mediated peripheral glucose disposal but may reduce insulin-induced suppression of hepatic glucose output [62].

A particularly attractive and novel feature of GLP-1 and its analogues is increased β -cell neogenesis, growth, and differentiation [63•,64,65]. GLP-1 increases expression of the transcription factor PDX-1 (IDX-1) that is involved in the early differentiation of β cells from ductal progenitor cells, and the proliferation of established β cells (Fig. 2). PDX-1 is also involved in glucose-mediated gene transcription; thus, GLP-1 offers a possible means to recruit progenitor cells into β cells and to expand the β -cell mass while improving glucose-mediated proinsulin biosynthesis and glucose-stimulated insulin secretion.

Glucagon-like peptide-1 is a short-acting peptide. In clinical trials it has been delivered by either intravenous infusion or repeated subcutaneous injection. The short biological half-life (about 1 minute) is largely due to inactivation by the circulating enzyme dipeptidylpeptidase-IV (DPP-IV), which cleaves the N-terminal dipeptide. Orally effective inhibitors of this enzyme are now under consideration to enhance the activity of endogenous and exogenously administered GLP-1 [66], as well as analogues that are resistant to DPP-IV [67]. A long-acting analogue of GLP-1 is exendin-4, a 39 amino acid peptide from the venom of the Gila monster (*Heloderma suspectum*), having 52% sequence homology with GLP-1. Exendin-4 activates GLP-1 receptors and produces similar biological effects, making it an interesting prototype drug. To circumvent the problem of repeated parenteral delivery, depot implants of GLP-1 analogues are being considered, and buccal and oral delivery of GLP-1 analogues have recently been reported [67].

Other insulin releasers

Stimulation of insulin release using phosphodiesterase inhibitors, α -2 adrenoceptor antagonists, or succinate esters has been difficult to exploit therapeutically, because the actions of these agents are not specific to the β cell [5,6•]. Recognition that imidazoline compounds can stimulate insulin secretion by closure of Kir 6.2 channels and activation of intracellular calcium mobilization has prompted exploration of the therapeutic potential of these types of compounds. A recently described imidazoline, BL11282, enhanced glucose-induced insulin secretion without affecting Kir 6.2 channels, possibly by activating PKA and PKC [68].

Direct modulators of metabolism

A selection of substances that directly stimulate glucose uptake and metabolism or suppress gluconeogenesis, and might be considered as potential leads for the development of new antidiabetic agents, is reviewed elsewhere [5]. Among these substances aminoimidazole carbox-amide ribonucleotide (AICAR), a ribofuranoside activator of AMP-activated protein kinase, is interesting because it inhibits (by inhibiting fructose-1, 6-bisphosphatase) hepatic gluconeogenesis and increases the expression of Glut-4 and hexokinase in muscle [69]. Inhibitors of glucose-6 phosphatase (G-6-Pase) translocase and G-6-Pase catalytic protein are under investigation as novel routes for the suppression of excess hepatic glucose output in type 2 diabetes [70,71]. Glycogen phosphorylase inhibitors provide another potentially useful means to suppress hepatic glucose output [72].

Reactive oxygen species are believed to contribute to the development of insulin resistance, but a beneficial effect of antioxidants in the treatment of type 2 diabetes has not been demonstrated. Recently, the antioxidant lipoic acid was shown to prevent insulin resistance caused by oxidative stress in cultured muscle cells [73]. Lipoic acid has been reported

to improve insulin sensitivity in type 2 diabetic patients, although this could be due to other actions: for example, lipoic acid is a co-factor for mitochondrial dehydrogenases and could therefore facilitate glucose oxidation. Lipoic acid has also been reported to enhance postreceptor insulin signaling [73]. Interestingly, lipoic acid is being considered as a treatment for diabetic neuropathy.

Manipulations of lipid metabolism provide opportunities to improve the circulating lipid profile and reduce hyperglycemia [5]. Agents that suppress fatty acid oxidation have received considerable interest in this respect. By reducing the supply of energy for gluconeogenesis and by facilitating peripheral glucose utilization, inhibitors of fatty acid oxidation can effectively lower blood glucose concentrations. However, it has been difficult to titrate the dosages of drugs acting in this manner to avoid episodes of hypoglycemia [5]. The concept of suppressing fatty acid oxidation has also been called into question by evidence that this causes intracellular accumulation of lipids, which then aggravates insulin resistance [74].

Antiobesity agents

The antiobesity agent sibutramine, which induces satiety via serotonin norepinephrine reuptake inhibition (SNRI), may improve glycemic control by mechanisms other than weight loss in obese type 2 diabetic patients. The primary amine metabolite of sibutramine increases basal and insulin-stimulated glucose uptake by cultured muscle cells via a mechanism that is independent of PI3K and unrelated to SNRI activity [75].

Developing a potent and highly specific β_3 -adrenoceptor agonist as an antiobesity agent has been hampered by differences in receptor structure between rodents and humans. β_3 agonists promote lipolysis and increase energy expenditure by thermogenesis through increased expression of uncoupling proteins (UCPs), principally UCP-1, in brown and white adipose tissue. The β_3 agonist AJ-9677 has recently been shown to decrease expression of TNF α and increase expression of Glut-4 in adipose tissue associated with reduced hyperglycemia and improved insulin action in obese-diabetic animals [76].

The amylin analogue pramlintide produces a range of beneficial actions to improve glycemic control in type 1 and type 2 diabetic patients [77]. These include reduced appetite and weight reduction, suppression of glucagon secretion, and slowing of gastric emptying. Trials have delivered the peptide by injection immediately before meals, but it should be possible to devise preparations for administration by other routes.

Conclusions

The multiple endocrine and metabolic disturbances that constitute type 2 diabetes provide many potential opportunities for intervention by new oral antidiabetic agents.

Although it would be advantageous to address insulin resistance and β -cell dysfunction as the primary endocrinopathies, any safe interventions that will improve glycemic control deserve consideration in the quest for future generations of antidiabetic drugs.

References and Recommended Reading

Papers of particular interest, published recently, have been highlighted as:

- Of importance
- Of major importance

1. Turner RC, Cull CA, Frighi V, Holman RR: Glycemic control with diet, sulfonylurea, metformin, or insulin in patients with type 2 diabetes mellitus. *JAMA* 1999, 281:2005–2012.
 2. Krentz AJ, Bailey CJ: *Type 2 Diabetes*. London: Royal Society Medicine Press; 2001.
 3. DeFronzo RA: Pathogenesis of type 2 diabetes: metabolic and molecular implications for identifying diabetes genes. *Diabetes Rev* 1997, 5:177–269.
 4. The DECODE Study Group: Glucose tolerance and mortality: comparison of WHO and American Diabetes Association diagnostic criteria. *Lancet* 1999, 354:617–621.
 5. Bailey CJ: New pharmacological approaches to glycemic control. *Diabetes Rev* 1999, 7:94–113.
 6. Bailey CJ: Potential new treatments for type 2 diabetes. *Trends Pharmacol Sci* 2000, 21:259–265.
- Provides an overview of the mechanisms of action of potential new agents for the treatment of hyperglycemia in type 2 diabetes. It indicates how these agents might be integrated into future treatment strategies.
7. Mudaliar S, Henry RR: New oral therapies for type 2 diabetes mellitus: the glitazones or insulin sensitizers. *Annu Rev Med* 2001, 52:239–257.
- Current appraisal of thiazolidinediones in clinical use, showing diverse range of effects achievable by stimulation of PPAR γ .
8. Auwerx J: PPAR γ , the ultimate thrifty gene. *Diabetologia* 1999, 42:1033–1049.
 9. Desvergne B, Wahli W: Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocr Rev* 1999, 20:649–688.
 10. Day C: Thiazolidinediones: a new class of antidiabetic drugs. *Diabetic Med* 1999, 16:179–192.
 11. Claire M, Bailey ST, Bhat S, et al.: The hormone resistin links obesity to diabetes. *Nature* 2001, 409:307–312.
 12. Katsuki A, Sumida Y, Murata K, et al.: Troglitazone reduces plasma levels of tumour necrosis factor- α in obese patients with type 2 diabetes. *Diabetes Obesity Metab* 2000, 2:189–191.
 13. Burant CF, Sreenan S, Hirano K, et al.: Troglitazone action is independent of adipose tissue. *J Clin Invest* 1998, 100:2900–2908.
 14. Furnsinn C, Brunmair B, Meyer M, et al.: Chronic and acute effects of thiazolidinediones BM 13.1258 and BM 15.2054 on rat skeletal muscle glucose metabolism. *Br J Pharmacol* 1999, 128:1141–1148.
 15. Loviscach M, Rehman N, Carter L, et al.: Distribution of peroxisome proliferator-activated receptors (PPARs) in human skeletal muscle and adipose tissues: relation to insulin action. *Diabetologia* 2000, 43:304–311.
 16. Cha BS, Ciaraldi TP, Carter L, et al.: Peroxisome proliferator-activated receptor (PPAR) γ and retinoid X receptor (RXR) agonists have complementary effects on glucose and lipid metabolism in human skeletal muscle. *Diabetologia* 2001, 44:444–452.

Evidence of the potential advantages of dual PPAR γ and RXR activation in human tissue. Also provides a useful debate on current appreciation of the role of PPAR γ in muscle.

17. Arakawa K, Ishihara T, Aoto M, et al.: Actions of novel antidiabetic thiazolidinedione, T-174, in animal models of non-insulin-dependent diabetes mellitus (NIDDM) and in cultured muscle cells. *Br J Pharmacol* 1998, 125:429–436.

18. Araki K, Yachi M, Haggisawa Y, et al.: Antidiabetic characterization of CS-011: a new thiazolidinedione with potent insulin-sensitizing activity. *Diabetes* 2000, 49(suppl 1):A425.
19. Upton R, Widdowson PS, Ishii S, et al.: Improved metabolic status and insulin sensitivity in obese fatty (fa/fa) Zucker rats and Zucker Diabetic Fatty (ZDF) rats treated with the thiazolidinedione MCC-555. *Br J Pharmacol* 1998, 125:1708–1714.
20. Fukui Y, Masui S, Osada S, et al.: A new thiazolidinedione, NC-2100 which is a weak PPAR γ activator, exhibits potent antidiabetic effects and induces uncoupling protein 1 in white adipose tissue of KKAY obese mice. *Diabetes* 2000, 49:759–767.
21. Naitoh T, Nakabeppu H, Kamon J, et al.: NIP-221 is a novel and potent thiazolidinedione-based peroxisome proliferator-activated receptor (PPAR) γ ligand with a prolonged biological half-time. *Diabetes* 2000, 49(suppl 1):A480.
22. Sohda T, Mizuno K, Momose Y, et al.: Studies on antidiabetic agents. Novel thiazolidinedione derivatives as potent hypoglycemic and hypolipidemic agents. *J Med Chem* 1992, 35:2617–2626.
23. Yakubu-Madus FE, Stephens TW, Johnson WT: Lipid lowering explains the insulin sensitivity enhancing effects of a thiazolidinedione, 5-(4-(2-(2-phenyl-4-oxazolyl)ethoxy)benzyl)-2,4 thiazolidinedione. *Diabetes Obesity Metab* 2000, 2:155–163.
24. Fiedorek F, Wilson GG, Frith L, et al.: Monotherapy with GI262570, a tyrosine-based non-thiazolidinedione PPAR γ agonist, improves metabolic control in type 2 diabetes mellitus patients. *Diabetes* 2000, 49(suppl 1):A157.
25. Brown KK, Henke BR, Blanchard SG, et al.: A novel N-aryl tyrosine activator of peroxisome proliferator-activated receptor- γ reverses the diabetic phenotype of the Zucker diabetic fatty rat. *Diabetes* 2000, 49:1415–1424.
26. Way JM, Harrington WW, Brown KK, et al.: Comprehensive messenger ribonucleic acid profiling reveals that peroxisome proliferator-activated receptor gamma activation has coordinate effects on gene expression in multiple insulin-sensitive tissues. *Endocrinology* 2001, 142:1269–1277.
27. Elbrecht A, Chen Y, Adams A, et al.: L-764406 is a partial agonist of human peroxisome proliferator-activated receptor gamma. The role of Cys313 in ligand binding. *J Biol Chem* 1999, 274:7913–7922.
28. Berger J, Bailey P, Biswas C, et al.: Thiazolidinediones produce a conformational change in peroxisome proliferator-activated receptor-gamma: binding and activation correlate with antidiabetic actions in db/db mice. *Endocrinology* 1996, 137:4189–4195.
29. Nolte RT, Wisely GB, Westin S, et al.: Ligand binding and co-activator assembly of the peroxisome proliferator-activated receptor- γ . *Nature* 1998, 395:137–143.
30. Camp HS, Li O, Wise SC, et al.: Differential activation of peroxisome proliferator-activated receptor- γ by troglitazone. *Diabetes* 2000, 49:539–547.
31. Dubois M, Pattou F, Kerr-Conte J, et al.: Expression of peroxisome proliferator-activated receptor γ (PPAR γ) in normal human pancreatic islet cells. *Diabetologia* 2000, 43:1165–1169.
32. Buckingham RE, Al-Barazanji KA, Toseland CD, et al.: Peroxisome proliferator-activated receptor- γ agonist, rosiglitazone, protects against nephropathy and pancreatic islet abnormalities in Zucker fatty rats. *Diabetes* 1998, 47:1326–1334.
33. Kim H, Kim J, Kim S, et al.: Identification and functional characterization of the peroxisomal proliferator response element in rat GLUT2 promoter. *Diabetes* 2000, 49:1517–1524.
34. Isshiki K, Haneda M, Koya D, et al.: Thiazolidinedione compounds ameliorate glomerular dysfunction independent of their insulin-sensitizing action in diabetic rats. *Diabetes* 2000, 49:1022–1032.
35. Murphy GJ, Holder JC: PPAR- γ agonists: therapeutic role in diabetes, inflammation and cancer. *Trends Pharmacol Sci* 2000, 21:469–474.
36. Bailey CJ: The rise and fall of troglitazone. *Diabetic Med* 2000, 17:414–415.
37. Shibata T, Takeuchi S, Yokota S, et al.: Effects of peroxisome proliferator-activated receptor- α and - γ agonist, JTT-501, on diabetic complications in Zucker diabetic fatty rats. *Br J Pharmacol* 2000, 130:495–504.

38. Lohray BB, Bhushan V, Rao BP, *et al.*: Novel euglycemic and hypolipidemic agents. *J Med Chem* 1998, 41:1619–1630.
 39. Murakami K, Tobe K, Ide T, *et al.*: A novel insulin sensitizer acts as a coligand for peroxisome proliferator-activated receptor- α (PPAR α) and PPAR- γ . *Diabetes* 1998, 47:1841–1847.
 40. Mukherjee R, Davies PJ, Crombie D, *et al.*: Sensitization of diabetic and obese mice to insulin by retinoid x receptor agonists. *Nature* 1997, 386:407–410.
 41. Emilsson V, O'Dowd J, Wang S, *et al.*: The effects of rexinoids and rosiglitazone on body weight and uncoupling protein isoform expression in the Zucker fa/fa rat. *Metabolism* 2000, 49:1610–1615.
 42. Liu Y, Sennitt MV, Hislop DC, *et al.*: Retinoid x receptor agonists have anti-obesity effects and improve insulin sensitivity in Zucker fa/fa rats. *Int J Obesity* 2000, 24:997–1004.
 43. Reul BA, Amin SS, Buchet JP, *et al.*: Effects of vanadium complexes with organic ligands on glucose metabolism: a comparison study in diabetic rats. *Br J Pharmacol* 1999, 126:467–477.
 44. Tsiani E, Bogdanovic E, Sorisky A, *et al.*: Tyrosine phosphatase inhibitors, vanadate and pervanadate, stimulate glucose transport and GLUT translocation in muscle cells by a mechanism independent of phosphatidylinositol 3-kinase and protein kinase C. *Diabetes* 1998, 47:1676–1686.
 45. Reul BA, Becker DJ, Ongemba LN, *et al.*: Improvement of glucose homeostasis and hepatic insulin resistance in ob/ob mice given oral molybdate. *J Endocrinol* 1997, 155:55–64.
 46. Stapleton SR: Selenium: an insulin-mimetic. *Cell Mol Life Sci* 2000, 57:1874–1879.
 47. Munoz MC, Barbera A, Dominguez J, *et al.*: Effects of tungstate, a new potential oral antidiabetic agent, in Zucker diabetic fatty rats. *Diabetes* 2001, 50:131–138.
 48. Zierath JR, Krook A, Wallberg-Henriksson H: Insulin action and insulin resistance in human skeletal muscle. *Diabetologia* 2000, 43:821–835.
 49. Pederson TM, Kramer DL, Rondinone CM: Serine threonine phosphorylation of IRS-1 triggers its degradation. Possible regulation by tyrosine phosphorylation. *Diabetes* 2001, 50:24–31.
 50. Wrobel J, Sredy J, Moxham C, *et al.*: PTP1B inhibition and antihyperglycemic activity in the ob/ob mouse model of novel 11-arylbenzo(b)-naphtho(2,3-d)furans and 11-arylbenzo(b)-naphtho(2,3-d)thiophenes. *J Med Chem* 1999, 42:3199–3202.
 51. Itani S, Zhou Q, Pories WJ, *et al.*: Involvement of protein kinase C in human skeletal muscle insulin resistance and obesity. *Diabetes* 2000, 49:1353–1358.
 52. Zhang B, Salituro G, Szalkowski D, *et al.*: Discovery of a small molecule insulin mimetic with antidiabetic activity in mice. *Science* 1999, 284:974–977.
 53. Manchem VP, Goldfine ID, Kohanski RA, *et al.*: A novel small molecule that directly sensitises the insulin receptor in vitro and in vivo. *Diabetes* 2001, 50:824–830.
- Example of recently reported compound that offers a novel structural template for the design of agents to enhance signaling activity of the insulin receptor.
54. Bates SH, Jones RB, Bailey CJ: Insulin-like effect of pinitol. *Br J Pharmacol* 2000, 130:1944–1948.
 55. Meyer K, Voss E, Neidlein R, *et al.*: Omega-substituted alkyl carboxylic acids as antidiabetic and lipid-lowering agents. *Eur J Med Chem* 1998, 33:775–787.
 56. Day C: Traditional plant treatments for diabetes mellitus. *Actualities de Chimie Ther* 2000, 26:131–150.
 57. Drucker DJ: Glucagon-like peptides. *Diabetes* 1998, 47:159–169.
- Comprehensive review of the potential advantages and challenges for the development of GLP-1 as a treatment for type 2 diabetes.
58. Vilsboll T, Krarup T, Deacon CE, *et al.*: Reduced postprandial concentrations of intact biologically active glucagon-like peptide 1 in type 2 diabetic patients. *Diabetes* 2001, 50:609–613.
 59. Juhl CB, Schmitz O, Pincus S, *et al.*: Short-term treatment with GLP-1 increases pulsatile insulin secretion in type II diabetes with no effect on orderliness. *Diabetologia* 2000, 43:583–588.
 60. Skoglund G, Hussain MA, Holz GG: Glucagon-like peptide 1 stimulates insulin gene promoter activity by protein kinase A independent activation of the rat insulin I gene cAMP response element. *Diabetes* 2000, 49:1156–1164.
 61. Vella A, Shah P, Basu R, *et al.*: Effect of glucagon-like peptide-1 (7-36) amide on glucose effectiveness and insulin action in people with type 2 diabetes. *Diabetes* 2000, 49:611–617.
 62. Vella A, Shah P, Basu R, *et al.*: Effect of glucagon-like peptide-1 (7-36) amide on initial splanchnic glucose uptake and insulin action in humans with type 1 diabetes. *Diabetes* 2001, 50:565–572.
 63. Xu W, Stoffers DA, Habener JE, *et al.*: Exendin-4 stimulates both beta-cell replication and neogenesis, resulting in increased beta-cell mass and improved glucose tolerance in diabetes rats. *Diabetes* 1999, 48:2270–2276.
- Early indication of the potential for GLP-1 analogues to stimulate β -cell neogenesis and replication, providing a new approach to the replacement of β -cell mass in diabetic states.
64. Stoffers DA, Kieffer TJ, Hussain MA, *et al.*: Insulinotropic glucagon-like peptide 1 agonists stimulate expression of homeodomain protein IDX-1 and increase islet size in mouse pancreas. *Diabetes* 2000, 49:741–748.
 65. Hui H, Wright C, Perfetti R: Glucagon-like peptide 1 induces differentiation of islet duodenal homeobox-1 positive pancreatic ductal cells into insulin-secreting cells. *Diabetes* 2001, 50:785–796.
 66. Holst JJ, Deacon CF: Inhibition of the activity of dipeptidyl-peptidase IV as a treatment for type 2 diabetes. *Diabetes* 1998, 47:1663–1670.
 67. Joseph JW, Kalitsky J, Brubaker PL: Oral delivery of glucagon-like peptide-1 in a modified polymer preparation normalizes basal glycaemia in diabetic db/db mice. *Diabetologia* 2000, 43:1319–1328.
 68. Efanou AM, Zaitsev SU, Mest HJ, *et al.*: The novel imidazoline compound BL11282 potentiates glucose-induced insulin secretion in pancreatic beta-cells in the absence of modulation of K-ATP channel activity. *Diabetes* 2001, 50:797–802.
 69. Bahl ES, Jessen N, Schmitz O, *et al.*: Chronic treatment with 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranoside increases insulin-stimulated glucose uptake and Glut4 translocation in rat skeletal muscles in a fiber type-specific manner. *Diabetes* 2001, 50:12–17.
 70. Parker JC, Van Volkenburg MA, Levy CB, *et al.*: Plasma glucose levels are reduced in rats and mice with an inhibitor of glucose-6-phosphate translocase. *Diabetes* 1998, 47:1630–1636.
 71. Madsen P, Lundbeck JM, Jakobsen P, *et al.*: Glucose-6-phosphatase catalytic enzyme inhibitors: synthesis and in vitro evaluation of novel 4,5,6,7-tetrahydrothienol[3,2-c]- and -[2,3-c]pyridines. *Bioorg Med Chem* 2000, 8:2277–2289.
 72. Treadway JL, Mendys P, Hoover DJ: Glycogen phosphorylase inhibitors for treatment of type 2 diabetes mellitus. *Expert Opin Invest Drugs* 2001, 10:439–454.
 73. Maddux BA, See W, Lawrence JC, *et al.*: Protection against oxidative stress-induced insulin resistance in rat L6 muscle cells by micromolar concentrations of α -lipoic acid. *Diabetes* 2001, 50:404–410.
 74. Dobbins RL, Szczepaniak LS, Bentley B, *et al.*: Prolonged inhibition of muscle carnitine palmitoyltransferase-1 promotes intramyocellular lipid accumulation and insulin resistance in rats. *Diabetes* 2001, 50:123–130.
 75. Bailey CJ, Turner SL, Bates SH, *et al.*: Sibutramine metabolites increase glucose transport by cultured rat muscle cells. *Int J Obesity* 2001, 25:478–485.
 76. Kato H, Ohue M, Kato K, *et al.*: Mechanism of amelioration of insulin resistance by B_3 -adrenoceptor agonist AJ-9677 in the KK-A y /Ta diabetic obese mouse model. *Diabetes* 2001, 50:113–122.
 77. Young AA: Amylin's physiology and its role in diabetes. *Curr Opin Endocrinol Diabetes* 1997, 4:282–290.