

See discussions, stats, and author profiles for this publication at:  
<https://www.researchgate.net/publication/257638123>

# Lapachol and its congeners as anticancer agents: A review

ARTICLE *in* PHYTOCHEMISTRY REVIEWS · MARCH 2014

Impact Factor: 2.41 · DOI: 10.1007/s11101-013-9289-1

CITATIONS

9

READS

228

5 AUTHORS, INCLUDING:



**Francesco Epifano**

Università degli Studi G. d'Annunzio C...

230 PUBLICATIONS 2,552 CITATIONS

SEE PROFILE



**Salvatore Genovese**

Università degli Studi G. d'Annunzio C...

134 PUBLICATIONS 1,095 CITATIONS

SEE PROFILE



**Serena Fiorito**

Università degli Studi G. d'Annunzio C...

34 PUBLICATIONS 48 CITATIONS

SEE PROFILE



**Robert Kiss**

Fonds de la Recherche Scientifique (F...

521 PUBLICATIONS 12,373 CITATIONS

SEE PROFILE

# Lapachol and its congeners as anticancer agents: a review

Francesco Epifano · Salvatore Genovese ·  
Serena Fiorito · Véronique Mathieu ·  
Robert Kiss



Received: 14 November 2012 / Accepted: 5 April 2013 / Published online: 13 April 2013  
© Springer Science+Business Media Dordrecht 2013

**Abstract** Lapachol is a naturally occurring 1,4-naphthoquinone originally isolated by the Italian phytochemist E. Paterno from *Tabebuia avellanedae* (Bignoniaceae) in 1882 and subsequently found in several other genera belonging to the families of Leguminosae, Malvaceae, Plumbaginaceae, Lamiaceae, Arecaceae, Scrophulariaceae, Verbenaceae, Celastraceae, Avicenniaceae, Caesalpiniaceae, Rubiaceae, and Proteaceae. A wide range of pharmacological activities have been observed for lapachol and its semi-synthetic derivatives in the literature, such as antileishmanial, anticarcinomic, anti-inflammatory, antimalarial, antiseptic, antitumor, antiviral, bactericidal, fungicidal, insectifugal, pesticidal, schistosomicidal, termiticidal, and viricidal effects. The aim of this review is to discuss in detail the phytochemical properties and pharmacological effects of the title compound that have been reported thus far,

highlighting its potential therapeutic benefits for the future.



**Keywords** Bignoniaceae · Lapachol ·  
1,4-Naphthoquinones · Anticancer activity

## Historical introduction of lapachol

1,4-Naphthoquinones represent a large class of natural compounds and are found in a wide range of plant families as well as in fungi and bacteria. Notable examples of naturally occurring 1,4-naphthoquinones include the K vitamins, juglone (isolated from the black walnut, *Juglans nigra* L. [Juglandaceae]), and plumbagin (obtained from *Plumbago*, *Drosera*, and *Nepenthes* spp.). Naphthoquinone derivatives have valuable pharmacological effects, acting as cytotoxic, antibacterial, antifungal, antiviral, antiprotozoal, insecticidal, anti-inflammatory, and antipyretic agents (Grolig and Wagner 2005). Atovaquone® (2), a derivative of lapachol (1), has been approved for the treatment of *Pneumocystis pneumonia*, toxoplasmosis, and malaria (Eyong et al. 2008). This latter was also tested as anti-tumor agent by the National Cancer Institute (Bethesda, MD, USA) in the 60-cell-line panel under the NSC-759582 NCI code, but displaying a poor activity ranging from a minimum percentage of inhibition of 75.5 % against UO-31 to a maximum of 122.4 % against RXF 393 renal cancer cell lines.

F. Epifano (✉) · S. Genovese · S. Fiorito  
Dipartimento di Farmacia, Università “G. D’Annunzio”  
Chieti-Pescara, Via dei Vestini 31, 66100 Chieti Scalo,  
CH, Italy  
e-mail: fepifano@unich.it

V. Mathieu · R. Kiss  
Laboratoire de Toxicologie, Faculté de Pharmacie,  
Université Libre de Bruxelles (ULB), Campus de la  
Plaine, CP205/1, Boulevard du Triomphe, 1050 Brussels,  
Belgium



The mechanisms of action underlying the observed effects of naphthoquinone derivatives are mainly due to their capacity to interact with topoisomerases and to generate semiquinone radicals and reactive oxygen species (ROS) inside the cell (da Silva et al. 2003). Plants containing naphthoquinones are widely used in Southeast Asia and South America to treat malignant and parasitic diseases. In this context, lapachol **1** (2-hydroxy-3-isopentenyl-1,4-naphthoquinone) represents one of the best examples of a research trend in phytochemistry.

Lapachol was first isolated in 1882 by the Italian chemist Emanuele Paternò from the tree *Tabebuia avellanedae* Lorentz ex Griseb. (Bignoniaceae) (Paternò 1882). Its name is derived from the term “lapacho,” the name by which this plant is commonly known in Brazil, where it is also called “pau d’arco” and “taheebo.” Extracts of the inner bark of *T. avellanedae*, from which lapachol was first purified, are used by local populations as analgesic, anti-inflammatory, anti-neoplastic, antimicrobial, and diuretic agents (Miranda et al. 2001). The compound was first named “lapachic acid” by Paternò because of its fairly acidic properties, and the term “lapachol” was assigned by Hooker (1892). In 1927, Fieser definitively confirmed the structure of **1** by comparing an authentic sample extracted from the bark of *T. avellanedae* with one obtained by chemical synthesis (Fieser 1927).

### Natural sources of lapachol

Several plants in the Bignoniaceae family are the richest sources of lapachol. However, this natural naphthoquinone has been also extracted from genera of the families

Leguminosae, Malvaceae, Plumbaginaceae, Lamiales, Arecaceae, Scrophulariaceae, Verbenaceae, Celastraceae, Avicenniaceae, Caesalpiniaceae, Rubiaceae, and Proteaceae, as outlined in Table 1.

### Screening of lapachol in plant extracts

Several methods are currently available to detect the presence of lapachol in vegetable extracts. The absorption spectra of the title compound were first fully described in Cooke et al. (1939). In the early 1960s, experimentation with the use of lapachol for therapeutic purposes prompted the search for alternative and more sensitive methodologies to accurately measure the concentration of compound **1** not only in plant extracts but also in biological fluids. In 1969, a fluorimetric method for the determination of lapachol in serum was developed by Finkel and Harrison (1969). These authors exploited the capacity of compound **1** to fluoresce at 410 nm after reduction with  $\text{NaHSO}_3$ , leading to a highly sensitive analytical technique. Then, Dawson et al. (1989) provided the complete NMR assignment and interpretation of lapachol, using one- and two-dimensional techniques. The IR behavior of **1** was well studied by Farfan et al. (2006). These authors were able to accurately calculate and assign all vibration frequencies of lapachol, even in enriched plant extracts, using the density functional theory. Currently, the most widespread and frequently used technique for the qualitative and quantitative analysis of compound **1** in complex matrices is high-performance liquid chromatography (HPLC). The first report of the use of this technique in the literature was provided by Awang et al. (1986). These authors

**Table 1** Main natural sources of lapachol

Sources	References
<i>Abutilon pakistanicum</i> Jafri and Ali in Jafri (Malvaceae)	Bakhat et al. (2010)
<i>Acacia jacquemontii</i> Benth. (Leguminosae)	Singh et al. (2010)
<i>Acacia nilotica</i> (Linn.) Del. (Leguminosae)	Prakash and Garg (1981)
<i>Austroplenckia populnea</i> var. <i>Ovata</i> (Reissek ex Mart.) Lundell (Celastraceae)	De Sousa et al. (2006)
<i>Avicenna alba</i> Blume (Avicenniaceae)	Ito et al. (2000)
<i>Avicennia tomentosa</i> Jacq. (Avicenniaceae)	Bournot (1914)
<i>Bauhinia guianensis</i> Aubl. (Caesalpiniaceae)	Viana et al. (1999)
<i>Bignonia gracilis</i> Lodd. (Bignoniaceae)	Prakash and Singh (1980a, b, c)
<i>Bignonia leucoxydon</i> L. (Bignoniaceae)	Oosterde (1917) and Oosterde (1916)
<i>Bignonia unguis-cati</i> Bureau and K.Schum. (Bignoniaceae)	Joshi et al. (1985)
<i>Capraria biflora</i> L. (Scrophulariaceae)	Lemos et al. (2007)
<i>Catalpa longissima</i> Sims (Bignoniaceae)	Chauhan et al. (1988)
<i>Conospermum</i> spp. (Proteaceae)	Cannon et al. (1975)
<i>Desmodium pulchellum</i> (L.) Benth. (Fabaceae)	Joshi et al. (1975)
<i>Diphysa robinoides</i> Benth. (Fabaceae)	Sagrero-Nievese (1986)
<i>Dolichandrone crispa</i> Seem (Bignoniaceae)	Prakash and Singh (1980a, b, c)
<i>Euterpe precatoria</i> Mart. (Arecaceae)	Galotta et al. (2008)
<i>Heterophragma adenophyllum</i> Seem ex Benth. and Hook. F. (Bignoniaceae)	Singh et al. (1972, 2006) and Joshi et al. (1979)
<i>Hibiscus tiliaceus</i> L. (Malvaceae)	Ali et al. (1980)
<i>Jacaranda mimosaeifolia</i> D. Don (Fabaceae)	Joshi et al. (1975)
<i>Kigelia tipulat</i> (Lam.) Benth. (Bignoniaceae)	Khan and Mlungwana (1999)
<i>Kigelia pinnata</i> (Jacq.) DC (Bignoniaceae)	Singh et al. (2010), Joshi et al. (1981) and Govindachari et al. (1971)
<i>Lippia microphylla</i> Cham. (Verbenaceae)	Lemos et al. (2007)
<i>Lippia sidoides</i> Cham. (Verbenaceae)	Lemos et al. (2007)
<i>Lomatia illicifolia</i> R.Br. (Proteaceae)	Rennie (1895)
<i>Lomatia longifolia</i> R.Br. (Proteaceae)	Rennie (1895)
<i>Macfadyena unguis-cati</i> (L.) A.H.Gentry (Bignoniaceae)	Duarte et al. (2000)
<i>Mansoa</i> spp. (Bignoniaceae)	Zoghbi et al. (2009)
<i>Markhamia hildebrandtii</i> (Baker) Sprague (Bignoniaceae)	Chen and Lee (1986)
<i>Markhamia platycalyx</i> Sprague (Bignoniaceae)	Joshi et al. (1985)
<i>Markhamia tipulate</i> Wall (Bignoniaceae)	Singh and Singh (1980) and Joshi et al. (1978)
<i>Markhamia zanzibarica</i> K. Schum. (Bignoniaceae)	Khan and Mlungwana (1999)
<i>Melloa quadrivalvis</i> (Jacq.) A. H. Gentry (Bignoniaceae)	Lima et al. (2005)
<i>Millingtonia hortensis</i> Linn. (Bignoniaceae)	Prakash and Garg (1981) and Singh et al. (1972)
<i>Newboldia laevis</i> (P. Beauv.) Seemann ex Bureau (Bignoniaceae)	Eyong et al. (2005, 2006) and Houghton et al. (1994)
<i>Oroxylum indicum</i> (L.) Benth. Ex Kurz (Bignoniaceae)	Joshi et al. (1977a, b, c, d)
<i>Paratecoma peroba</i> (Record) Kuhlman (Bignoniaceae)	Sanderman et al. (1968)
<i>Paulownia kawakamii</i> T. Ito (Scrophulariaceae)	Huang et al. (2004)
<i>Phyllarthron comorense</i> DC (Bignoniaceae)	Joshi et al. (1973a, b, 1975, 1976)
<i>Plumbago scandens</i> L. (Plumbaginaceae)	Rodrigues et al. 2006.
<i>Plumbago zeylanica</i> L. (Plumbaginaceae)	Kishore et al. (2010)
<i>Radermachera sinica</i> Hemsl. (Bignoniaceae)	Inoue et al. (1981)

**Table 1** continued

Sources	References
<i>Radermachera xylocarpa</i> K. Schum. (Bignoniaceae)	Shetgiri et al. (2001)
<i>Randia dumatorium</i> L. (Rubiaceae)	Joshi et al. (1981)
<i>Rubia tinctorum</i> L. (Rubiaceae)	Kawasaki et al. (1992)
<i>Salvia sahendica</i> Boiss and Buhse (Lamiaceae)	Jassabi et al. (2006)
<i>Stereospermum kunthianum</i> Cham. (Bignoniaceae)	Ghogomu-Tih et al. (1986)
<i>Stereospermum personatum</i> (Hassk.) Chatterjee (Bignoniaceae)	Kumar et al. (2005)
<i>Stereospermum suaveolens</i> DC. (Bignoniaceae)	Joshi et al. (1977a, b, c, d)
<i>Stereospermum tetragonum</i> DC (Bignoniaceae)	Purushothaman and Natarajan (1974)
<i>Tabebuia avellanadae</i> Lorentz ex Griseb. (Bignoniaceae)	Viana et al. (2003), Krustrak (2001), Wagner et al. (1989), Burnett and Thomson (1967) and de Lima et al. (1962)
<i>Tabebuia chrysantha</i> (Jacq.) G.Nicholson (Bignoniaceae)	Burnett and Thomson (1968)
<i>Tabebuia chrysotricha</i> (Mart. Ex DC.) Standl. (Bignoniaceae)	Grazziotin et al. (1992)
<i>Tabebuia coralibe</i> Standl. (Bignoniaceae)	Rodriguez, et al. (1999)
<i>Tabebuia flavescens</i> Benth. and Hook.f. ex Griseb. (Bignoniaceae)	Orth (1960)
<i>Tabebuia guayacan</i> Hemsl. (Bignoniaceae)	Manners and Jurd (1976)
<i>Tabebuia heptaphylla</i> (Vell.) Toledo (Bignoniaceae)	Schmeda-Hirschmann and Papastergiou (2003)
<i>Tabebuia impetiginosa</i> (Mart. Ex DC.) Standl. (Bignoniaceae)	Gomez Castellanos et al. (2009)
<i>Tabebuia ochracea</i> (Cham.) Standl. (Bignoniaceae)	Zani et al. (1991)
<i>Tabebuia palmeri</i> Rose (Bignoniaceae)	Villegas et al. (1995)
<i>Tabebuia pentaphylla</i> Hemsl. (Bignoniaceae)	Rohatgi et al. (1983), and Prakash and Singh (1980a, b, c, 1981)
<i>Tabebuia rosea</i> DC (Bignoniaceae)	Girard et al. (1988) and Joshi et al. (1973a, b, 1976, 1977a, b, c, d)
<i>Tabebuia serratifolia</i> Vahl. Nicholson (Bignoniaceae)	Lemos et al. (2007), Velasquez et al. (2004) and Vidal-Tessier et al. (1988)
<i>Tecoma stans</i> Juss. (Bignoniaceae)	Dohnal (1976)
<i>Tecomella undulata</i> Seem (Bignoniaceae)	Singh et al. (2008a, b), Joshi et al (1977a, b, c, d, 1986), Joshi and Singh (1974) and Gupta et al. (1969)
<i>Tectona grandis</i> L.f. (Lamiaceae)	Lukmandaru et al. (2009), Singh et al. (1989, 2008a, b), Moreira et al. (2006), Windeisen et al. (2003), Khan and Mlungwana (1998) and Sandermann and Simatupang (1964, 1965)
<i>Zeyheria digitalis</i> (Vell.) L.B. Sm. and Sandwith (Bignoniaceae)	da Silveira et al. (1975)
<i>Zeyheria montana</i> Mart. (Bignoniaceae)	Jacome et al. (1999)

developed a sensitive screening technique for lapachol using a reverse-phase HPLC with a Brownlee RP-18 Sheri-10 column and MeCN (aq) 0.25 %–AcOH 1:1 as the mobile phase. The HPLC separation of **1** in mixtures containing structurally similar products was reported about 10 years later by Steinert et al. (1996) using an isocratic elution and by Novotna et al. (1999). An alternative reversed-phase methodology, using a LiChrosorb RP-10 column and MeOH (aq) 5 %–

AcOH 8:2 as the elution mixture, has been recently described by Fonseca et al. (2004). Mass spectrometry represents the most recently described method for the analysis of lapachol in plant extracts. The fragmentation study and the electrospray ionization mass spectrometry of lapachol were thoroughly reported in Vessecchi et al. (2010). These authors studied the occurrence of protonated and carbocationic species in the positive mode and deprotonated species in the

negative ones using collision-induced dissociation experiments. Another very sensitive technique, full-scan high resolution mass spectrometry, was also used for the analysis of lapachol in food, feed, and general botanicals (Mol et al. 2011). Finally, alternative methodologies, such as voltammetry (Ngameni et al. 2000) and thermal analysis (Santan et al. 2008), have also been recently used for the same purposes.

### Physicochemical properties of lapachol and synthetic methodologies

Lapachol is a light to dark yellow crystalline powder that typically forms prisms with a pKa of 6.15 (Ossowski et al. 2008) and that has a melting point of 141–143 °C. Lapachol is fairly soluble in water, although its solubility is greatly affected by pH; the solubility ranges from 1.5 µg/mL at pH 4.0–5 mg/mL at pH 10. The UV absorption maxima of lapachol are 251, 278, and 331 nm. As a bidentate ligand, lapachol tends to form stable complexes with transition metals such as copper, iron (Sawhney et al. 1983), zinc, cobalt (Martinez et al. 2005), nickel (Farfan et al. 2009), bismuth, antimony (de Oliveira et al. 2011), and several others (Bhatia and Vohra 1982). The previous and renewed interest in lapachol from the phytochemical and pharmacognostic points of view prompted the development of several synthetic processes to synthesize this biologically active naphthoquinone. As stated above, in 1892, Hooker described the first method to obtain lapachol by the reaction of isovaleraldehyde and 1,4-naphthoquinone (Hooker 1892). Subsequently, in 1927, Fieser obtained the title compound by reacting the silver salt of 2-hydroxy-1,4-naphthoquinone (lawsone) with 3,3-dimethylallyl bromide as the alkylating agent, producing **1** with a very low yield (5 %) (Fieser 1927). Approximately 40 years later, Pettit and Houghton described a synthetic scheme by which lapachol was obtained using a six-step process starting from lawsone and succinyl peroxide followed by a series of reductions and Grignard reactions (Pettit and Houghton 1971). At almost the same time, Jacobsen and Torssell applied the alkylation of quinones with radicals obtained from the decarboxylation of carboxylic acid promoted by peroxydisulfate ions to lawsone (Jacobsen and Torssell 1973). A Williamson etherification followed by a heat-promoted Claisen

rearrangement was employed to synthesize **1** from lawsone and 3,3-dimethylallyl bromide. In Sun et al. (1998) alkylated the lithium salt of the lawsone nucleophile with 3,3-dimethylallyl bromide, providing lapachol in 40 % yield. Finally, in recent years, Kazantzi et al. (2007) used the Pd-catalyzed allylation of lawsone with isopentenyl alcohol or isopentenyl methyl ether to synthesize **1** with good yield, and Ferreira et al. (2011) similarly used the formic acid–water mediated one-pot reduction of *o*-quinone methides. Lapachol is easily oxidized both in vitro and in vivo. The redox transformations of the title compound are currently of great interest because they are believed to be partly responsible for the observed biological effects of lapachol. For these reasons, simple in vitro methods to mimic the biooxidation process of **1** have recently been established. A Mn(III) complex with H<sub>2</sub>O<sub>2</sub> and hypervalent iodine species has been used to investigate and structurally characterize the chemical entities derived from the total or partial oxidation of lapachol (Niehues et al. 2012a, b; Pires et al. 2011; Bodini and Arancibia 1989).

### In vitro pharmacological and toxicological analyses of the antitumor activity of lapachol

Numerous lapachol derivatives have already been synthesized and have been shown to exert significant in vitro growth inhibitory effects in various cancer cell lines. The antitumor activity of lapachol was first described in the early 1970s (Block et al. 1974), and this compound was tested by the National Cancer Institute (Bethesda, MD, USA) in the 60-cell-line panel under the NSC-11905 NCI code (Fig. 1).

The mean in vitro growth inhibitory concentrations shown in Fig. 1 indicate that the 60 cancer cell lines analyzed show only small differences in lapachol sensitivity at the micromolar concentrations tested. Lapachol has also been demonstrated to reduce the number of tumors caused by doxorubicin in *Drosophila melanogaster* heterozygous for the tumor suppressor gene *wts* (Costa et al. 2011).


In addition to its antiproliferative activity in cancer cells, lapachol also decreases the invasion of HeLa cells and could therefore represent an interesting scaffold with which to generate novel anti-metastatic compounds (Balassiano et al. 2005). Lapachol-

**Fig. 1** The data illustrate the average 50 % growth inhibitory concentrations obtained for lapachol in a panel of 60 cancer cell lines. The testing was performed by the Developmental Therapeutics Program, Division of Cancer Treatment and Diagnosis, National Cancer Institute. The URL to the Program's website is <http://dtp.cancer.gov>. These data are presented here with the permission of the NCI

### GI<sub>50</sub> Mean Graph for Compound 11905

NCI Cancer Screen Current Data

Average GI<sub>50</sub> over all cell lines is 2.41E-5 M





induced anti-metastatic effects have also been demonstrated by Maeda et al. (2008) as detailed in the next section on the *in vivo* effects of lapachol.

Lapachol also displays anti-angiogenic properties, especially when it is coordinated to bismuth (III) (Parrilha et al. 2012). Furanonaphthoquinones are also phytoconstituents of the Bignoniaceae family, and 2-acetylfuranonaphthoquinone (**3**; Table 2) displays higher *in vitro* antitumor activity than lapachol (Eyong et al. 2008). Compound **3** is available in minute amounts from nature, whereas exposure of lapachol to ozonolysis conditions results in the formation of the expected aldehyde **4** (Table 2) at 70 % yield along with this unusual compound **3** at 30 % yield (Eyong et al. 2008). Pentacyclic derivatives of lapachol such as ( $\pm$ )1,4-naphthoquinones (compound **5** in Table 2) display ten times higher *in vitro* antitumor activity than lapachol (da Silva et al. 2002). These pentacyclic derivatives can also be considered pterocarpan (**6**; Table 2) derivatives where the A-ring is substituted by the 1,4-naphthoquinone nucleus (da Silva et al. 2002). Pentacyclic 1,4-naphthoquinone derivatives such as **7–9** (Salustiano et al. 2010) are active in single micromolar ranges against oxidative stress-resistant K562 and multidrug-resistant (MDR) Lucena-1 human leukemia cells, whereas lapachol **1** and  $\alpha$ -lapachol (**10**, Table 2) are not (Salustiano et al. 2010). In addition, these pentacyclic 1,4-naphthoquinone derivatives display low toxicity in lymphocytes activated by phytohemagglutinin, a feature that emphasizes their bioselectivity for cancer cells over normal cells (Salustiano et al. 2010). The cytotoxic effects of these compounds on cells from leukemia patients are related to the activation of apoptosis (Salustiano et al. 2010).

As emphasized by Maia et al. (2011), despite the relevant therapeutic progress obtained with imatinib, clinical resistance to this drug has emerged and reemerged after cytogenetic remission in a group of patients with chronic myeloid leukemia (CML). Maia et al. (2011) have thus evaluated the anti-CML activity and mechanisms of action of LQB-118 (**11**, Table 2), a pterocarpanquinone structurally related to lapachol **1**. LQB-118 treatment resulted in a substantial reduction in viability of two cell lines derived from CML, the vincristine-sensitive K562 cell line and the vincristine-resistant K562-Lucena (a cell line overexpressing P-glycoprotein). These authors observed, in agreement with these results, that the induction of caspase-3

activation by this compound indicated a significant rate of apoptosis, and apoptosis induced by LQB-118 **11** was accompanied by a reduction of P-glycoprotein, survivin, and XIAP expression (Maia et al. 2011).

Although, as stated above, the anti-malarial agent atovaquone was not seen to be an effective antitumor agent, its 2-piperazinyl and 2-piperidinyl analogues have been shown to significantly exert *in vitro* growth inhibitory effects favourably comparing to the activity of the known cancer chemotherapeutic doxorubicine (Zhou et al. 2009).

### In vivo pharmacological and toxicological analyses of the antitumor activity of lapachol





The pharmacokinetic of lapachol in humans has been studied by Niehues et al. (2012a, b). The maximum plasma concentration of this naphthoquinone derivative was seen to be 30  $\mu\text{g/mL}$ . Several reports have demonstrated that lapachol displays *in vivo* anticancer activity. Indeed, da Consolação et al. (1975) showed that the acetylglucosylation of lapachol (**9**, Table 2) resulted in a compound that extended lapachol activity and that was effective against mouse lymphocytic leukemia P-388. These authors reported that mice inoculated with 1 million leukemic cells and treated with the drug for 9 days had an 80 % increase in lifespan compared with control animals (da Consolação et al. 1975). Maeda et al. (2008) reported that lapachol can act as a vitamin K antagonist with *in vivo* anticancer activity. These authors used the B16BL6 mouse metastatic melanoma model and observed that a single oral administration of a highly toxic dose of lapachol (80–100 mg/kg) 6 h before an intravenous injection of tumor cells in the mice dramatically promoted metastasis (Maeda et al. 2008). This increase in metastasis was also observed in T cell-deficient mice and NK-suppressed mice, and *in vitro* treatment of B16BL6 cells with lapachol promoted metastasis only slightly, indicating that lapachol promotes metastasis primarily by affecting host factors other than T and NK cells (Maeda et al. 2008). The increase in metastasis caused by lapachol was almost completely suppressed by administering vitamin K3 before and 6 h after oral administration of lapachol, and the protein C level was reduced maximally





without an increase in prothrombin time (Maeda et al. 2008). These authors thus suggested that a high dose of lapachol promotes metastasis by inducing a hypercoagulable state as a result of the inhibition of a vitamin

K-dependent pathway (Maeda et al. 2008). In contrast, chronic oral administration of low, non-toxic doses of lapachol (5–20 mg/kg) weakly but significantly suppressed metastasis by an unknown mechanism,

**Table 2** Illustration of lapachol congeners with in vitro and/or in vivo antitumor activity

Compounds	Reference
 <p style="text-align: right;"><b>3</b></p>	Eyong et al. (2008)
 <p style="text-align: right;"><b>4</b></p>	Eyong et al. (2008)
 <p style="text-align: right;"><b>5</b></p>	da Silva et al. (2002)
 <p style="text-align: right;"><b>6</b></p>	Salustiano et al. (2010)

**Table 2** continued

Compounds	Reference
 <p><b>7</b> <math>R^1 = OH, R^2 = OCH_3</math>; <b>8</b> <math>R^1 = OCH_3, R^2 = OH</math>;  <b>9</b> <math>R^1 = R^2 = Obn</math>; <b>10</b> <math>R^1 = R^2 = H</math></p>	Salustiano et al. (2010) and Maia et al. (2011)
 <p style="text-align: center;"><b>11</b></p>	da Consolação et al. (1975)

nevertheless suggesting the possible use of lapachol as an anti-metastatic agent (Maeda et al. 2008).

## Conclusions

Lapachol is a compound with a relatively simple chemical structure that can be easily derivatized into various subgroups of compounds with increasing levels of anticancer activity both in vitro and in vivo. Lapachol can be obtained by extraction from a large variety of plants; it can also be synthesized in six chemical steps. This compound displays not only antiproliferative activity but also antimetastatic effects, again both in vitro and in vivo. While lapachol has been identified for several decades, it merits further chemical and pharmacological analyses because it has higher antiproliferative and antimigratory effects in cancer cells than in normal cells. Lapachol could therefore lead to the discovery of novel anticancer agents with marked anti-metastatic effects, and the development of such agents is

especially important because 90 % of cancer patients currently die from their metastases and not from their primary cancer.

## References

- Ali S, Singh P, Thomson RH (1980) Naturally occurring quinones. Part 28. Sesquiterpenoid quinones and related compounds from *Hibiscus tiliaceus*. J Chem Soc Perkin Trans 1:257–259
- Awang DVC, Kindack D, Dawson BA (1986) Tandem high-performance liquid chromatography methods for resolution of lapachol and related naphthaquinones. J Chromat 368:439–443
- Bakhat A, Itrat F, Abdul M et al (2010) New glycosidic constituents of *Abutilon pakistanicum*. Helv Chim Acta 93:2245–2250
- Balassiano IT, De Paulo SA, Henriques Silva N et al (2005) Demonstration of the lapachol as a potential drug for reducing cancer metastasis. Oncol Rep 13:329–333
- Bhatia BML, Vohra N (1982) Kinetic studies of the nonisothermal decomposition of metal chelates of lapachol with calcium (II), barium (II), and lead (II). Thermochim Acta 53:361–364

- Block JB, Serpick AA, Miller VV et al (1974) Early clinical studies with lapachol (NSC-11905). *Cancer Chemother Rep* 4:27–28
- Bodini ME, Arancibia V (1989) Manganese complexes with 2-hydroxy-3-(3-methyl-2-butenyl)-1,4-naphthoquinone (lapachol). Redox chemistry and spectroscopy in dimethyl sulfoxide. *Polyhedron* 8:1407–1412
- Bournot K (1914) Separation of lapachol from the heartwood of *Avicennia tomentosa*. *Archiv Pharm* 251:351–354
- Burnett AR, Thomson RH (1967) Naturally occurring quinones. XII. Extractives from *Tabebuia avellanedae*. *J Chem Soc C* 2100–2104
- Burnett AR, Thomson RH (1968) Naturally occurring quinones. XII. Extractives from *Tabebuia chrysantha* and other Bignoniaceae. *J Chem Soc C* 850–853
- Cannon JR, Joshi KR, McDonald IA et al (1975) Structures of nine quinones isolated from two *Conospermum* species. *Tetrahedron Lett* 32:2795–2798
- Chauhan AK, Dobhal MP, Uniyal PN (1988) Phytochemical investigation of *Catalpa longissima*. *Herba Pol* 34:3–5
- Chen CC, Lee MH (1986) Constituents of *Markhamia hildebrandtii* (Baker) Sprague and their antitumor activity. *Huaxue* 44:61–64
- Cooke RG, Macbeth AK, Winzer FL (1939) Absorption spectra of some naturally occurring naphthoquinones and their derivatives. *J Chem Soc* 878–884
- Costa WF, Oliveira AB, Nepomuceno JC (2011) Lapachol as an epithelial tumor inhibitor agent in *Drosophila melanogaster* heterozygote for tumor suppressor gene wts. *Genet Mol Res* 10:3236–3245
- da Consolação M, Linardi F, de Oliveira MM et al (1975) A lapachol derivative active against mouse lymphocytic leukemia P-388. *J Med Chem* 18:1159–1161
- da Silva AJ, Buarque CD, Brito FV et al (2002) Synthesis and preliminary pharmacological evaluation of new ( $\pm$ )-1,4-naphthoquinones structurally related to lapachol. *Bioorg Med Chem* 10:2731–2738
- da Silva MN, da Souza MCBV, Ferreira VF et al (2003) Synthesis of new aldehyde derivatives from  $\beta$ -lapachone and nor- $\beta$ -lapachone. *Arkivoc* 156
- da Silveira JC, Gottlieb OR, de Oliveira GG (1975) Chemistry of Brazilian Bignoniaceae. 1. Zeyherol, a dilignol from *Zeyheria digitalis*. *Phytochemistry* 14:1829–1830
- Dawson BA, Girard M, Kindack D et al (1989) Carbon-13 NMR of lapachol and some related naphthoquinone. *Magn Res Chem* 27:1176–1177
- de Lima OG, D'Albuquerque IL, de Lima CG et al (1962) Antibiotic substances in higher plants. XX. Antimicrobial activity of some derivatives of lapachol as compared with xyloidone, a new natural o-naphthoquinone isolated from extracts of heartwood of *Tabebuia avellanedae*. *Rev Inst Antibiot Univ Recife* 4:3–17
- de Oliveira LG, Silva MM, de Paula FCS et al (2011) Antimony(V) and bismuth(V) complexes of lapachol: synthesis, crystal structure and cytotoxic activity. *Molecules* 16:10314–10323
- de Sousa JR, Silva GDF, Miyakoshi T et al (2006) Constituents of the root wood of *Austroplenckia populnea* var. *ovata*. *J Nat Prod* 69:1225–1227
- Dohnal B (1976) Investigations of some metabolites of *Tecoma stans* Juss. Callus tissue II. Chromatographic analysis of alkaloid and quinone compounds. *Acta Soc Bot Pol* 45:369–381
- Duarte DS, Dolabela MF, Salas CE et al (2000) Chemical characterization and biological activity of *Macfadyena unguis-cati* (Bignoniaceae). *J Pharm Pharmacol* 52:347–352
- Eyong KO, Krohn K, Hussain H et al (2005) Newbouldiaquinone and newbouldiamide: a new naphthoquinone-anthraquinone coupled pigment and a new ceramide from *Newbouldia laevis*. *Chem Pharm Bull* 53:616–619
- Eyong KO, Folefoc GN, Kuete V et al (2006) Newbouldiaquinone A: a naphthoquinone-anthraquinone ether coupled pigment, as a potential antimicrobial and antimalarial agent from *Newbouldia laevis*. *Phytochemistry* 67:605–609
- Eyong KO, Kumar PS, Kuete V et al (2008) Semisynthesis and antitumoral activity of 2-acetyl-furanonaphthoquinone and other naphthoquinone derivatives from lapachol. *Bioorg Med Chem Lett* 18:5387–5390
- Farfan RA, Molina JR, Ottaviani E et al (2006) Infrared band assignment of lapachol by means of comparative theoretical and experimental studies. *Inf Technol* 17:63–66
- Farfan RA, Espindola JA, Martinez MA et al (2009) Synthesis and crystal structure of a new lapacholate complex with nickel(II), [Ni(Lap)<sub>2</sub>(DMF)(H<sub>2</sub>O)]. *J Coord Chem* 62:3738–3744
- Ferreira SB, Rodrigues da Rocha D, Carneiro JWM et al (2011) A new method to prepare 3-alkyl-2-hydroxy-1,4-naphthoquinones: synthesis of lapachol and phthiocol. *Synlett* 1551–1554
- Fieser LF (1927) the alkylation of hydroxynaphthoquinone. A synthesis of lapachol. *J Am Chem Soc* 49:857–864
- Finkel JM, Harrison SD Jr (1969) Fluorimetric method for the determination of lapachol in serum. *Anal Chem* 41:1854–1855
- Fonseca SGC, da Silva LBL, Castro RF et al (2004) Validation of the analytical methodology for the determination of lapachol in solutions by HPLC. *Quim Nova* 27:157–159
- Galotta ALQA, Bonaventura MAD, Lima LARS (2008) Antioxidant and cytotoxic activities of “acai” (*Euterpe precatoria* Mart.). *Quim Nova* 31:1427–1430
- Ghogomu-Tih R, Nyasse B, Tsamo E et al (1986) Chemical constituents of the stem heart wood of *Stereospermum kunthianum*. *Planta Med* 1986:342
- Girard M, Kindack D, Dawson BA et al (1988) Naphthoquinones constituents from *Tabebuia* spp. *J Nat Prod* 51:1023–1024
- Gomez Castellanos JR, Prieto JM, Heinrich M (2009) Red lapacho (*Tabebuia impetiginosa*). A global ethnopharmacologically commodity? *J Ethnopharmacol* 121:1–13
- Govindachari TR, Patankar SJ, Viswanathan N (1971) Isolation and structure of two new dihydroisocoumarins from *Kigelia pinnata*. *Phytochemistry* 10:1603–1606
- Grazziotin JD, Schapoval EES, Chaves CG et al (1992) Phytochemical and analgesic investigation of *Tabebuia chrysotricha*. *J Ethnopharmacol* 36:249–251
- Grolig J, Wagner R (2005) Naphthoquinones. In: Ullmann's encyclopedia of industrial chemistry. Wiley-VCH, Weinheim
- Gupta SR, Malik KK, Seshadri TR (1969) Lapachol from the heartwood of *Tecoma undulata* and a note on its reaction. *Ind J Chem* 7:457–459

- Hooker SC (1892) The constitution of lapachic acid (lapachol) and its derivatives. *J Chem Soc* 61:611–650
- Houghton P, Pandev R, Hawkes JE (1994) Naphthoquinones and an alkaloid from the roots of *Newboldia laevis*. *Phytochemistry* 35:1602–1603
- Huang P, Karagianis G, Wei S et al (2004)  $\alpha$ -Lapachone and other naphthoquinones from the heartwood of *Paulownia kawakamii*. *Biochem Syst Ecol* 32:1047–1049
- Inoue K, Chen CC, Inouye H et al (1981) Quinones and related compounds in higher plants. Part 16. Naphthoquinones from *Radermachera sinica* Hemsl. (Bignoniaceae). *J Chem Soc Perkin Trans 1*:2764–2770
- Ito C, Katsuno S, Kondo Y et al (2000) Chemical constituents of *Avicenna alba*. Isolation and structural elucidation of new naphthoquinones and their analogues. *Chem Pharm Bull* 48:339–343
- Jacobsen N, Torssell K (1973) Synthesis of naturally occurring quinones. Alkylation with the silver ion-peroxydisulfate-carboxylic acid system. *Acta Chem Scand* 27:3211–3216
- Jacome RLRP, de Oliveira AB, Raslan DS et al (1999) Analysis of naphthoquinones in *Zeyheria montana* crude extracts. *Quim Nova* 22:175–177
- Jassabi AR, Mehrdad M, Eghtesadi F et al (2006) Novel rearranged abietane diterpenoids from the roots of *Salvia sahendica*. *Chem Biodiv* 3:916–922
- Joshi KC, Singh LB (1974) Quinonoid and other constituents from the heartwood of *Tecomella undulate*. *Phytochemistry* 13:663–664
- Joshi KC, Prakash L, Singh P (1973a) Quinones and other constituents from *Phyllarthron comorense*. *Phytochemistry* 12:469–470
- Joshi KC, Prakash L, Singh P (1973b) Quinones and other constituents from *Tabebuia rosea*. *Phytochemistry* 12:942–943
- Joshi KC, Bansal RK, Singh P et al (1975) Components of the stem barks of *Phyllarthron comorense* and *Jacaranda mimosaeifolia* and the roots of *Desmodium pulchellum*. *Ind J Chem* 13:869–870
- Joshi KC, Singh P, Singh G (1976) Crystalline components of roots of *Phyllarthron comorense* DC and of stem bark of *Tabebuia rosea* DC. *Ind J Chem B* 14:637–638
- Joshi KC, Bansal RK, Patni R (1977a) Chemical examination of the roots of *Stereospermum suaveolens* DC. *J Ind Chem Soc* 54:648–649
- Joshi KC, Prakash L, Shah RK (1977b) Chemical examination of the roots of *Tabebuia rosea* and *Oroxylum indicum*. *Planta Med* 257–258
- Joshi KC, Singh P, Pardasani RT (1977c) Quinones and other constituents from the roots of *Tecomella undulate*. *Planta Med* 14–16
- Joshi KC, Singh P, Singh G (1977d) Chemical examination of *Tecomella undulata* Seem. *Curr Sci* 46:145–146
- Joshi KC, Singh P, Pardasani RT (1978) Chemical constituents from the stem heartwood of *Markhamia stipulata*. *Planta Med* 219–221
- Joshi KC, Singh P, Pardasani RT et al (1979) Quinones and other constituents from *Heterophragma adenophyllum*. *Planta Med* 1979:60–63
- Joshi KC, Singh P, Taneja S (1981) Crystalline components of the stem heartwood of *Randia dumatorium* and *Kigelia pinnata*. *J Ind Chem Soc* 58:825–826
- Joshi KC, Singh P, Sharma MC (1985) Quinones and other constituents of *Markhamia platycalyx* and *Bignonia unguiscati*. *J Nat Prod* 48:145
- Joshi KC, Sharma AK, Singh P (1986) A new ferulic ester from *Tecomella undulata*. *Planta Med* 71–72
- Kawasaki Y, Goda Y, Yoshihira K (1992) The mutagenic constituents of *Rubia tinctorum*. *Chem Pharm Bull* 40:1504–1509
- Kazantzi G, Malamidou-Xenikaki E, Spyroudis S (2007) Palladium-catalyzed allylation of 2-hydroxy-1,4-naphthoquinone. Application to the preparation of lapachol. *Synlett* 427–430
- Khan RM, Mlungwana SM (1998) 5-Hydroxylapachol: a cytotoxic agent from *Tectona grandis*. *Phytochemistry* 50:439–442
- Khan MR, Mlungwana SM (1999)  $\gamma$ -Sitosterol, a cytotoxic sterol from *Markhamia zanzibarica* and *Kigelia Africana*. *Fitoterapia* 70:111–112
- Kishore N, Mishra BB, Tiwari VK et al (2010) Difuranonaphthoquinones from *Plumbago zeylanica* roots. *Phytochem Lett* 3:62–65
- Krustrak D (2001) Taheebo-Lapacho-Tabebuia impetiginosa. *Farm Glasn* 57:215–222
- Kumar US, Tiwari AK, Reddy SV et al (2005) Free-radical scavenging and xanthine oxidase inhibitory constituents from *Stereospermum personatum*. *J Nat Prod* 68:1615–1621
- Lemos TLG, Monte FJQ, Kellen A et al (2007) Quinones from plants of northeastern Brazil: structural diversity, chemical transformations, NMR data and biological activities. *Nat Prod Res* 21:529–550
- Lima CSA, de Amorim ELC, Nascimento SC et al (2005) Cytotoxic pyranonaphthoquinones from *Melloa quadrivalvis* (Bignoniaceae). *Nat Prod Res* 19:217–222
- Lukmandaru G, Ashitani T, Takahashi K (2009) Color and chemical characterization of partially black-streaked heartwood in teak (*Tectona grandis*). *J Forestry Res* 20:377–380
- Maeda M, Murakami M, Takegami T et al (2008) Promotion or suppression of experimental metastasis of B16 melanoma cells after oral administration of lapachol. *Toxicol Appl Pharmacol* 229:232–238
- Maia RC, Vasconcelos FC, de Sá Bacelar T et al (2011) LQB-118, a pterocarpanquinone structurally related to lapachol [2-hydroxy-3-(3-methyl-2-butenyl)-1,4-naphthoquinone]: a novel class of agent with high apoptotic effect in chronic myeloid leukemia cells. *Invest New Drugs* 29:1143–1155
- Manners GD, Jurd L (1976) A new naphthoquinone from *Tabebuia guayacan*. *Phytochemistry* 15:225–226
- Martinez MA, de Jimenez MCL, Castellano EE et al (2005) Two isostructural complexes of Co (II) and Zn (II) with lapacholate, dimethylformamide and water. *J Arg Chem Soc* 93:185–193
- Miranda FGG, Vilar JC, Alves IAN et al (2001) Antinociceptive and antiedematogenic properties and acute toxicity of *Tabebuia avellanadae* Lor. Ex Griseb. Inner bark aqueous extract. *BMC Pharmacol* 1:6
- Mol HGJ, van Dam RCJ, Zomer P et al (2011) Screening of plant toxins in food, feed, and botanicals using full-scan high-resolution (Orbitrap) mass spectrometry. *Food Addit Contam A* 28:1405–1423

- Moreira RYO, Arruda MSP, Arruda AC et al (2006) Anthraquinones and naphthoquinones from the bark of *Tectona grandis* reforestation specimen. *Rev Bras Farmac* 16:392–396
- Ngameni E, Tonle IK, Nanseu et al (2000) Voltammetry study of 2-hydroxy-3-isoprenyl-1,4-naphthoquinone using a carbon paste electrode. *Electroanalysis* 12:847–852
- Niehues M, Barros VP, Emery FS et al (2012a) Biomimetic in vitro oxidation of lapachol: a model to predict and analyse the in vivo phase I metabolism of bioactive compounds. *Eur J Med Chem* 54:804–812
- Niehues M, Barros VP, Emery FS et al (2012b) From in vitro to in vivo: preclinical data on the oxidative metabolism and pharmacokinetics of lapachol. *Planta Med* 78:PH9
- Novotna P, Pacakova V, Bosakova Z et al (1999) High-performance liquid chromatographic determination of some anthraquinone and naphthoquinone dyes occurring in historical textiles. *J Chromat A* 863:234–241
- Oesterde OA (1916) Substance accompanying lapachol in greenheart wood. *Archiv Pharm* 254:346–348
- Oesterde OA (1917) Substance accompanying lapachol in greenheart wood. *J Chem Soc* 112:505
- Orth H (1960) Isolation of extraneous materials from *Tabebuia flavescens* by extraction. *Holzforsch* 14:89–91
- Ossowski T, Goulart MOF, de Abreu FC et al (2008) Determination of the pK<sub>a</sub> values of some biologically active and inactive hydroxyquinones. *J Braz Chem Soc* 19:175–183
- Parrilha GL, Vieira RP, Campos PP et al (2012) Coordination of lapachol to bismuth(III) improves its anti-inflammatory and anti-angiogenic activities. *Biometals* 25:55–62
- Paternò E (1882) Ricerche sull'acido lapachico. *Gazz Chim It* 12:337–392
- Pettit GR, Houghton LE (1971) Synthesis of hydroxyhydrapachol and lapachol. *J Chem Soc C* 3:509–511
- Pires SMG, de Paula R, Simoes MMQ et al (2011) Novel biomimetic oxidation of lapachol with H<sub>2</sub>O<sub>2</sub> catalysed by a manganese(III) porphyrin complex. *RSC Advances* 1:1195–1199
- Prakash L, Garg MG (1981) Chemical constituents of the roots of *Millingtonia hortensis* Linn. and *Acacia nilotica* (Linn.) Del. *J Ind Chem Soc* 58:96–97
- Prakash L, Singh R (1980a) Phytochemical screening of *Bignonia gracilis* Lodd. (Bignoniaceae). *Pharmazie* 35:649–650
- Prakash L, Singh R (1980b) Chemical constituents of the stem bark and stem heartwood of *Dolichandrone crispera* Seem. *Pharmazie* 35:122–123
- Prakash L, Singh R (1980c) Chemical constituents of stem bark and root heartwood of *Tabebuia pentaphylla* Hemsl. (Bignoniaceae). *Pharmazie* 35:813
- Prakash L, Singh R (1981) Chemical examination of the leaves and stem heartwood of *Tabebuia pentaphylla* Hemsl. (Bignoniaceae). *J Ind Chem Soc* 58:1122–1123
- Purushothaman KK, Natarajan RK (1974) Chemical examination of Patala (*Stereospermum tetragonum* DC). *J Res Ind Med* 9:107–108
- Rennie EH (1895) A colouring matter from *Lomatia illicifolia* and *Lomatia longifolia*. *J Chem Soc* 67:784–793
- Rodrigues SV, Viana LM, Baumann W (2006) UV/Vis spectra and solubility of some naphthoquinones, and the extraction behaviour of plumbagin from *Plumbago scandens* roots in supercritical CO<sub>2</sub>. *Anal Bioanal Chem* 385:895–900
- Rodriguez C, Erika DC, Torrenegra R (1999) Antifungal and antibacterial furanonaphthoquinones from *Tabebuia coralibe*. *Rev Lat Quim* 27:89–93
- Rohatgi BK, Gupta RB, Roy D et al (1983) Quinones from *Tecoma pentaphylla*: constitution of tecomaquinones I and II. *Ind J Chem B* 22:886–889
- Sagrero-Nieves L (1986) Isolation of lapachol from *Diphysa robinoides*. *J Nat Prod* 49:547
- Salustiano EJS, Netto CD, Fernandes RF et al (2010) Comparison of the cytotoxic effect of lapachol, alpha-lapachone and pentacyclic 1,4-naphthoquinones on human leukemic cells. *Invest New Drugs* 28:139–144
- Sanderman W, Simatupang MH, Wenderborn W (1968) New quinones from the heartwood of *Paratecoma peroba*. *Naturwiss* 55:38
- Sandermann W, Simatupang MH (1964) Constituents from teak (*Tectona grandis*) II. Constitution and synthesis of tectol and dehydrotectol. *Chem Ber* 97:588–597
- Sandermann W, Simatupang MH (1965) New quinones from *Tectona grandis*. *Naturwiss* 52:262–263
- Santan DP, Fonseca SGC, Bedor DCG et al (2008) Application of thermal analysis in the development and characterization of PLGA microparticles containing lapachol. *Rev Ciencias Farm Bas Aplic* 29:261–266
- Sawhney SS, Matta SD, Jain R et al (1983) Investigation on the interaction of 2-hydroxy-3-(3-methyl-2-butenyl)-1,4-naphthoquinone (lapachol) with copper(II) and iron(II). *Thermochim Acta* 70:367–371
- Schmeda-Hirschmann G, Papastergiou F (2003) Naphthoquinone derivatives and lignans from the Paraguayan crude drug *Tayá pyta* (Bignoniaceae). *Zeitsch Naturforsch C* 58:495–501
- Shetgiri NP, Kokitkar SV, Sawant SN (2001) *Radermachera xylocarpa*: the highly efficient source of lapachol and synthesis of its derivatives. *Acta Pol Pharm* 58:133–135
- Singh P, Singh A (1980) Quinonoid constituents of the bark of *Markhamia tipulate* Wall. *Pharmazie* 35:701–702
- Singh P, Prakash L, Joshi KC (1972) Lapachol and other constituents from the Bignoniaceae. *Phytochemistry* 11:1498
- Singh P, Jain S, Bhargava S (1989) A 1,4-anthraquinone derivative from *Tectona grandis*. *Phytochemistry* 28:1258–1259
- Singh P, Natani K, Jain S et al (2006) Microwave-assisted rapid cyclization of lapachol, a main constituents of *Heterophragma adenophyllum*. *Nat Prod Res* 20:207–212
- Singh P, Khandelwal P, Hara N et al (2008a) Radermachol and naphthoquinone derivatives from *Tecomella tipular*: complete <sup>1</sup>H and <sup>13</sup>C NMR assignments of radermachol with the aid of computational <sup>13</sup>C shift prediction. *Ind J Chem B* 47:1865–1870
- Singh P, Pandey D, Mathur J et al (2008b) Barleriaquinone-I from the heartwood of *Tectona grandis* Linn. *J Ind Chem Soc* 85:1060–1063
- Singh P, Khandelwal P, Sharma K et al (2010) Cetyl triacetoate and other constituents from *Acacia jacquemontii* and *Kigelia pinnata*. *J Ind Chem Soc* 87:1403–1407
- Steinert J, Khalaf H, Rimpler M (1996) High performance liquid chromatography separation of some naturally occurring

- naphthoquinones and anthraquinones. J Chrom A 723: 206–209
- Sun JS, Geiser AH, Frydman B (1998) A preparative synthesis of lapachol and related naphthoquinones. Tetrahedron Lett 39:8221–8224
- Velasquez J, Rojas LB, Usubillaga A (2004) Antifungal activity of naphthoquinone from *Tabebuia serratifolia* Vahl. Nicholson Ciencia 12:64–69
- Vessecchi R, Emery FS, Galembeck SE et al (2010) Fragmentation studies and electrospray ionization mass spectrometry of lapachol: protonated, deprotonated and cationized species. Rapid Commun Mass Spectr 24:2101–2108
- Viana EP, Santa Rosa RS, Almeida SSMS et al (1999) Constituents of the stem bark of *Bauhinia guianensis*. Fitoterapia 70:111–112
- Viana LM, Freitas MR, Rodrigues SV et al (2003) Extraction of lapachol from *Tabebuia avellanedae* wood with supercritical CO<sub>2</sub>: an alternative to soxhlet extraction? Braz J Chem Eng 20:317–325
- Vidal-Tessier AM, Delaveau P, Champion B et al (1988) Lipophilic quinones of the trunk wood of *Tabebuia serratifolia*. Ann Pharm Franc 46:55–57
- Villegas JR, Amato S, Castro I et al (1995) 4-Aryltetraline tipul and furanonaphthoquinone from *Tabebuia palmeri* wood. Fitoterapia 66:281–282
- Wagner H, Kreher B, Lotter H et al (1989) Structure determination of new isomeric naphtha[2,3-b]furan-4,9-diones from *Tabebuia avellanedae* by the selective INEPT technique. Helv Chim Acta 72:659–667
- Windeisen E, Klassen A, Wegener G (2003) On the chemical characterization of plantation teakwood from Panama. Holz Roh Werkst 61:416–418
- Zani C, de Oliveira AB, de Oliveira GG (1991) Furanonaphthoquinones from *Tabebuia ochracea*. Phytochemistry 30:2379–2381
- Zhou J, Duan L, Chen H et al (2009) Atovaquone derivatives as potent cytotoxic and apoptosis inducing agents. Bioorg Med Chem Lett 19:5091–5094
- Zoghbi MGB, Oliveira J, Guilhon GMSP (2009) The genus *Mansoa* (Bignoniaceae): a source of organosulfur compounds. Rev Bras Farmac 19:795–804