

# Repetitive Magnetic Stimulation Promotes Neural Stem Cells Proliferation by Upregulating MiR-106b *In Vitro*\*

Hua LIU (柳 华)<sup>1</sup>, Xiao-hua HAN (韩肖华)<sup>1#</sup>, Hong CHEN (陈 红)<sup>1</sup>, Cai-xia ZHENG (郑彩霞)<sup>1</sup>, Yi YANG (杨 翼)<sup>2</sup>, Xiao-lin HUANG (黄晓琳)<sup>1#</sup>

<sup>1</sup>Department of Rehabilitation Medicine, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, China

<sup>2</sup>College of Health Science, Wuhan Institute of Physical Education, Wuhan 430079, China

© Huazhong University of Science and Technology and Springer-Verlag Berlin Heidelberg 2015

**Summary:** Neural stem cells (NSCs) proliferation can be influenced by repetitive transcranial magnetic stimulation (rTMS) *in vivo* via microRNA-106b-25 cluster, but the underlying mechanisms are poorly understood. This study investigated the involvement of microRNA-106b-25 cluster in the proliferation of NSCs after repetitive magnetic stimulation (rMS) *in vitro*. NSCs were stimulated by rMS (200/400/600/800/1000 pulses per day, with 10 Hz frequency and 50% maximum machine output) over a 3-day period. NSCs proliferation was detected by using ki-67 and EdU staining. Ki-67, p21, p57, cyclinD1, cyclinE, cyclinA, cdk2, cdk4 proteins and miR-106b, miR-93, miR-25 mRNAs were detected by Western blotting and qRT-PCR, respectively. The results showed that rMS could promote NSCs proliferation in a dose-dependent manner. The proportions of ki-67+ and Edu+ cells in 1000 pulses group were 20.65% and 4.00%, respectively, significantly higher than those in control group (9.25%, 2.05%). The expression levels of miR-106b and miR-93 were significantly upregulated in 600–1000 pulses groups compared with control group ( $P < 0.05$  or  $0.01$  for all). The expression levels of p21 protein were decreased significantly in 800/1000 pulses groups, and those of cyclinD1, cyclinA, cyclinE, cdk2 and cdk4 were obviously increased after rMS as compared with control group ( $P < 0.05$  or  $0.01$  for all). In conclusion, our findings suggested that rMS enhances the NSCs proliferation *in vitro* in a dose-dependent manner and miR-106b/p21/cdks/cyclins pathway was involved in the process.

**Key words:** repetitive magnetic stimulation; neural stem cells; EdU; ki67; microRNA-106b; cyclin-dependent kinase; cyclin-dependent kinase inhibitor

As a non-invasive brain stimulation (NIBS) technique, repetitive transcranial magnetic stimulation (rTMS) is considered to be safe and effective<sup>[1]</sup>. rTMS induces electric currents in the brain repeatedly so as to influence the excitability of the cortex, which has been proved by positron emission tomography (PET)<sup>[2]</sup> and functional magnetic resonance imaging (fMRI)<sup>[3]</sup> in many clinic tests. The therapeutic effect of rTMS was found to involve neural plasticity<sup>[4]</sup> and hippocampal neurogenesis in rats<sup>[5]</sup>. Ten Hz rTMS was demonstrated to promote the proliferation of neural stem cells (NSCs) in rat subventricular zone in our previous study<sup>[6]</sup>. However, the underlying mechanism is unclear.

By recognizing the 3' untranslated region (3'UTR)<sup>[7–9]</sup>, microRNAs (miRNAs), short (approximately 22 nucleotides) non-coding RNAs, can silence the target mRNAs. A number of miRNAs can determine the cell fate, especially neurogenesis, neural differentiation and neural plasticity in the center nerve system<sup>[10–12]</sup>. MiR106b-25 cluster, which includes miR-106b, miR-25 and miR-93 with the similar sequence and targets, is strongly linked with NSCs functions<sup>[13–15]</sup>. Guo *et al*

showed that miR-106b-25 cluster in the brain of rat with focal cerebral ischemia could increase EdU-positive NSCs after high frequency rTMS<sup>[6]</sup>.

In contrast to numerous studies *in vivo*, there were only few studies on rTMS *in vitro*. In the present study, we used NSCs from neonatal rats to confirm that repetitive magnetic stimulation (rMS) could promote NSCs proliferation by regulating miR106b-25 cluster. We measured the expression of miR-106b, miR-25, miR-93 mRNAs and p21, p57, cyclinD1, cyclinA, cyclinE, cdk2 and cdk4 proteins after rMS, and found that miR-106b was the primary factor that affected the proliferation of NSCs. Because the stimulus intensity is an important parameter related to the rTMS effect<sup>[1]</sup>, we also studied the relationship between the pulses and NSCs proliferation. Our findings indicated that rMS enhances the NSCs proliferation *in vitro* in a pulse dose-dependent manner. All of these results suggest that rMS at 10 Hz or higher can promote the proliferation of NSCs *in vitro*, and miR-106b/p21/cdks/cyclins pathway is involved in the process.

## 1 MATERIALS AND METHODS

### 1.1 NSCs Culture

NSCs were harvested from the bilateral hippocampus of neonatal 3-day-old SD rats provided by the Experimental Animal Center, Huazhong University of

Hua LIU, E-mail: hualiu2005@163.com

#Corresponding authors, Xiao-lin HUANG, E-mail: xiaolinh2-006@126.com; Xiao-hua HAN, E-mail: 491695008@qq.com

\*This project was supported by grants from the National Natural Science Foundation of China (No. 81071601, and No. 81171858).

Science and Technology (Wuhan, China). After mechanical isolation, NSCs were cultured in DMEM/F12 (Gibco, USA) supplemented with 1×B27 supplement minus vitamin A (Invitrogen, USA), 20 ng/mL recombinant human bFGF (Pepro Tech, USA) and 20 ng/mL recombinant human EGF (Pepro Tech, USA) at 5% CO<sub>2</sub> in a 37°C incubator at a density of 1×10<sup>5</sup>/mL. The media were replaced every 2–3 days and the cells were passaged every 5 days when the neurospheres' diameter reached 100–200 μm. To obtain the single-cell suspension, we used 1×TrypLE™ express enzyme (Gibco, USA) to dissociate cells at 5% CO<sub>2</sub> in a 37°C incubator for 3 min, then pipetted cells mechanically with a 1-mL pipet for almost 30 times. Cells from original tissue were regarded as passage 1 cells.

### 1.2 rMS Exposure


rMS was administered by a customized magnetic stimulator with a 90-mm figure-of-eight coil (Yirui De, CCY-I, Wuhan, China). Passage 2 NSCs in dishes and flasks were put on the cross of the “8” coil, where the magnetic field was strongest (fig. 1). The rMS parameters were as follows: the frequency, 10 Hz; the intensity, 50% of the device's maximum power (peak value 3.5T); the interval time of the stimulation, 10 s. A series of rMS (200/400/600/800/1000 pulses per day) were administered once a day for 3 consecutive days. For control group, the NSCs in dishes and flasks were also put on the cross of the “8” coil but subjected to no stimulation.

### 1.3 Immunofluorescence Assay

Nestin and SOX<sub>2</sub> staining was used to identify the hippocampus-derived NSCs. Besides the microscopy analysis, the proliferation of cells was assessed by ki-67 and 5-ethynyl-2-deoxyuridine (EdU) staining. After 1-h exposure to rTMS, neurospheres were plated on 48-well plate coverslips coated with poly-L-lysine (0.1 mg/mL) (Sigma, USA) at 5% CO<sub>2</sub> in a 37°C incubator for at least 30 min. Then NSCs were fixed with absolute methyl alcohol for 20 min for ki-67/nestin/SOX<sub>2</sub> immunofluorescence before 200 μL solution A (1:1000) of EdU (Ruibo Company, China) was added into each well for incubation at 5% CO<sub>2</sub> in a 37°C incubator for 4 h. Briefly, each coverslip was covered with 20 μL mouse anti-nestin (1:100; BD, USA) and goat anti-SOX<sub>2</sub> (1:100; Santa Cruz Biotechnology, USA) antibody at 4°C overnight for NSCs immunostaining. FITC-anti-ki67 (1:50; eBioscience, USA) was added to cells for staining directly. The next day, nestin- and SOX<sub>2</sub>-staining cells were incubated with secondary FITC-labeled rabbit anti-mouse IgG (1:400; Jackson ImmunoResearch, USA) and Cy3-labeled rabbit anti-goat IgG (1:100; EarthOx, USA) for 2 h in the dark. DAPI was added for nuclei staining for 15 min.

Wells were rinsed by PBS (5 min×3), 0.5% TritonX-100 PBS (10 min) and PBS (5 min×1) for preparing for EdU staining. 1× Apollo® (500 μL) containing solution B 25 μL, solution C 5 μL, solution D 1.5 μL and solution E 5 mg was applied and incubated for 30 min at room temperature in the dark. Then cells were washed by 0.5% TritonX-100 PBS (5 min×3) and absolute methyl alcohol (5 min×1) respectively. 1× Hoechst 33342 staining was used for nuclei staining for 30 min. Apollo® solution excitation wavelength was 550 nm, emission wavelength 565 nm, Hoechst 33342 solution excitation wavelength 350 nm, and emission wavelength 461 nm.

The time of exposure was below 30 ms<sup>[16]</sup>.



**Fig. 1** The position of the NSCs culture flask and the stimulation parameters

A: the flask put on the cross of “8” coil; B: rMS stimulator. The cambered line means the magnetic field.

Immunofluorescence images were observed and recorded under the Olympus B×51 microscope. Five microscopic fields were randomly selected and the cells counted using image processing software (Image J, USA) for quantification.

### 1.4 Quantitative Real-time PCR (qRT-PCR)

According to the manufacturer's instructions, total RNA was extracted from stimulated passage 2 neurospheres. RNA was quantified by using a spectrophotometer. To quantify the expression of miR-106b, miR-25 and miR-93, reverse transcribed RNA was carried out by using TaqMan MicroRNA Reverse Transcription Kit (Applied Biosystems, Foster City, CA) at 70°C for 5 min, 42°C for 60 min, and 95°C for 5 min. Each 20 μL qPCR system contained 100 μmol/L forward (F) primer, 100 μmol/L reverse (R) primer, 10 μL SYBR Green Supermix/Flourescein qPCR Master Mix (2×) (TaKaRa, Japan) and 4 μL cDNA (10×). The program was 1 cycle of 50°C for 2 min; 95°C for 10 min; 40 cycles of 95°C for 30 s, and 60°C for 30 s. Samples were run in triplicate and miRNA expressions were analyzed by 2<sup>-ΔΔCT</sup> method<sup>[17]</sup>. The primers are listed in the table 1.

### 1.5 Western Blotting

The crude lysates of whole cells were extracted by using RIPA buffer, and then they were transferred to prechilled eppendorf tubes and centrifuged at 12 000 g for 10 min at 4°C. The protein concentrations were quantified by using the Protein Assay Kit for bicinchoninic acid (Pierce, UK) according to the manufacturer's protocol. Proteins (50 μg) were resolved on a 15% SDS PAGE, transferred onto PVDF membranes and probed sequentially with antibodies. Antibodies were listed as follows: ki-67 (1:500, eBioscience, USA), p21 (1:200, Santa Cruz, USA), p57 (1:200, Santa Cruz, USA), cyclinA (1:400, Abcam, USA), cyclinD1 (1:1000, Cell Signaling Technology, USA), cyclinE (1:200, Santa Cruz, USA), cdk2 (1:200, Santa Cruz, USA) and cdk4 (1:200, Santa Cruz, USA). The expression levels of protein were analyzed by Odyssey IR imaging system (LI-COR Biosciences, USA).

**Table 1 Primer sequences of miR-106b, miR-93 and miR-25**

Primers	Loop	Forward	Reverse
U6		5'-CGCTTCGGCAGCACAT-ATAC-3'	5'-AAATATGGAACGCT-TCACGA-3'
rno-miR-106b	5'-GTCGTATCCAGTGCAGGGTCCGAGGT-ATTCGCACTGGAACGACATCTG-3'	5'-TGCGCTAAAGTGCTGACAGTG-3'	5'-CTCAAGTGTCGTGG-AGTCGGCAA-3'
rno-miR-25-3p	5'-GTCGTATCCAGTGCAGGGTCCGAGGT-ATTCGCACTGGATACGACTCAGA-3'	5'-TGCGCCATTGCACTTG-TCTCGG-3'	5'-CTCAAGTGTCGTGG-AGTCGGCAA-3'
rno-miR-93-5p	5'-GTCGTATCCAGTGCAGGGTCCGAGGT-ATTCGCACTGGATACGACCTACC-3'	5'-TGCGCCAAAGTGCTGT-TCGTGCA-3'	5'-CTCAAGTGTCGTGG-AGTCGGCAA-3'

### 1.6 Data Analysis

Results are expressed as  $\bar{x} \pm s_{\bar{x}}$  and the data were analyzed by one-way ANOVA followed by a subsequent sidark post-hoc test via the statistical software SPSS (version 17.0; SPSS, USA). Statistical significance was set at  $P < 0.05$ .

## 2 RESULTS


### 2.1 Identification of NSCs

Light microscopy revealed smooth and shiny passage 2 neurospheres, indicating a good activity of the

neurospheres (fig. 2A). Nestin and SOX<sub>2</sub>, the common markers of fetal and adult NSCs<sup>[18]</sup>, were found to be highly expressed in neurospheres (fig. 2B–2E).

### 2.2 Proliferation of NSCs *In Vitro*

The percentage of ki-67+ cells was significantly increased in 800 and 1000 pulses groups ( $P < 0.01$  for both; fig. 3, and the expression levels of ki-67 protein were also significantly increased in 600, 800 and 1000 pulses groups as compared with control group ( $P < 0.01$  for all; fig. 4).

**Fig. 2** Immunocharacterization of NSCs

NSCs are cultured and identified by nestin and SOX<sub>2</sub> fluorescent staining. A: Neurospheres were observed under the light microscope. Bar=50 μm. B: DAPI expressing nuclei (blue). Bar=20 μm. C: nestin (green) in neurosphere. Bar=20 μm. D: SOX<sub>2</sub> (red) in neurosphere. Bar=20 μm. E: merged picture. Bar=20 μm.

We evaluated NSCs proliferation by EdU staining, which has recently been considered to be an alternative of BrdU staining<sup>[19]</sup>. The results showed that EdU+ cells were significantly increased in 600, 800 and 1000 pulses groups as compared with control group ( $P < 0.01$  for all; fig. 5). The percentage of ki-67+ and Edu+ cells in 1000 pulses group was 20.65% and 4.00%, respectively, significantly higher than that in control group (9.25%, 2.05%).


### 2.3 Expression of miRNA-106b Cluster in NSCs Stimulated by rMS

The expression levels of miR-106b and miR-93 were significantly increased in 600 pulses stimulation

group ( $P < 0.05$  for all; fig. 6) and in 800/1000 pulses groups (all  $P < 0.01$ ; fig. 6) as compared with control group. However, the expression levels of miR-25 were significantly reduced in 400/600/800/1000 pulses groups (all  $P < 0.01$ , fig. 6).


### 2.4 Expression of p21 and p57 in NSCs Treated by rMS

The expression levels of p21 protein were decreased significantly in 800/1000 pulses groups ( $P < 0.01$  for all; fig. 7), while p57 protein expression levels were increased significantly in 600 pulses group ( $P < 0.05$ ; fig. 7) and in 800/1000 pulses groups as compared with control group ( $P < 0.01$  for all; fig. 7).




**Fig. 3** Ki-67 staining showing the effect of rMS on the proliferation of NSCs *in vitro*

A: control group; B-F: 200, 400, 600, 800 and 1000 pulses groups, respectively. Ki-67 fluorescence (green) was expressed in nuclei, and all nuclei were counterstained using DAPI (blue). The arrows indicate ki-67+ cells. Bar=50 μm. G: the statistical graph. \*\* $P<0.01$  vs. control group



**Fig. 4** Ki-67 protein expression in NSCs after rMS

A: expression changes of ki-67 in different groups; B: the statistical graph. \*\* $P<0.01$  vs. control group



**Fig. 5** EdU staining showing the effect of rMS on the proliferation of NSCs *in vitro*

A: control group; B-F: 200, 400, 600, 800 and 1000 pulses groups, respectively. Proliferative nuclei were counterstained by EdU (red), and nuclei were counterstained using Hoechst 33342 (blue). The arrows indicate EdU+ cells. Bar=50 μm. G: the statistical graph. \* $P<0.05$ , \*\* $P<0.01$  vs. control group


## 2.5 Regulatory Effects of Cyclin-dependent Kinase Inhibitor p21 on Cyclin-dependent Kinases

The expression levels of cyclinA protein were significantly enhanced in 600/800/1000 pulses groups ( $P<0.05$  or  $P<0.01$ ; fig. 7) as compared with control group. Those of cyclinD1 protein were significantly in-

creased in 800/1000 pulses groups ( $P<0.01$  for all); cyclinE protein expression levels were much higher in 200/400/600/800/1000 pulses groups ( $P<0.05$  or  $P<0.01$ ); cdk2 protein expression levels were obviously increased in 400/600/800/1000 pulses groups ( $P<0.05$  or  $P<0.01$ ), and cdk4 protein expression levels were also markedly




increased in 1000 pulses group ( $P < 0.05$ ).



**Fig. 6** Effect of rMS on the expression of the miR-106b family members

A: The expression of miR-106b cluster members were assessed by qRT-PCR in different groups; 1: DL2000, 2: control group, 3–7: 200–1000 pulses groups; B: the statistical graph. \* $P < 0.05$ , \*\* $P < 0.01$



**Fig. 7** p21-regulated potential signaling pathways

A: cyclin-dependent kinase protein expressions in different groups; 1: control group; 2–6: 200–1000 pulses groups; B: p21 and p57 protein expressions in different groups; C: the statistical graph. \* $P < 0.05$ , \*\* $P < 0.01$  vs. control group

### 3 DISCUSSION

#### 3.1 Proliferation of NSCs and rMS Pluses

Ki-67 is a marker of cells at each stage of cell cycle except  $G_0$  phase<sup>[20]</sup>. As an alternative to BrdU, EdU is recently used to examine the proliferation of cells in S phase<sup>[16]</sup>. In our present study, ki-67 and EdU immunofluorescence showed that the percentage of ki-67+ and EdU+ cells was increased by 2.23 and 1.95 folds, respectively, in 1000 pluses group as compared with control group, suggesting that rMS promoted NSCs proliferation *in vitro*.

It has been reported that rTMS has therapeutic effects on depression, because it can reverse the decrease of NSCs<sup>[21]</sup> and enhance neurogenesis<sup>[22]</sup>. Neurogenesis is believed to be a potential therapeutic method for center nerve system diseases based on regulation of endogenous and exogenous factors<sup>[23–25]</sup>. A variety of cells are reported to be influenced by rTMS, including neurons<sup>[26]</sup>, gails<sup>[27]</sup>, NSCs<sup>[28]</sup>, p12 and HL-60 cells<sup>[29]</sup>. However, the mechanism of neurogenesis by rMS remains elusive.

Our data demonstrated that rMS could promote the proliferation of NSCs in a dose-dependent manner. rMS protocol consists of stimulation frequency, amplitude duration and coil shape, which have been reported to be important in the rMS treatment<sup>[1]</sup>. As the frequency of 10 Hz was shown to have an exciting effect on rat brain<sup>[30]</sup>, we chose 10 Hz in the present study. Additionally, the

effect of rTMS depends on the number of stimuli *in vivo*<sup>[31]</sup>. We investigated the relationship between the number of pluses and proliferation of NSCs *in vitro* in the current study and observed that the highest pulses had the best proliferation effect as *in vivo*.

#### 3.2 Relationship between MicroRNA Cluster miR-106b-25 and NSCs Proliferation after rMS

The miR-106b cluster members were found to be correlated with cancer<sup>[32–33]</sup>, NSCs proliferation<sup>[13]</sup> and neovascularization<sup>[34]</sup>. They are located in the 13th intronic region of the Mcm7 oncogene and they can control DNA synthesis and cells entry into S phase<sup>[35]</sup>. In the present study, the result showed that the expression levels of miR-106b and miR-93 were significantly increased and those of miR-25 were reduced with the number of pluses increasing. This indicates that rMS can drive the proliferation of NSCs by regulating miR-106b cluster.

Our study showed that miR-25 has an opposite trend to miR-106b and miR-93 after rMS *in vitro*. It is known that miR106b-25 belongs to intronic microRNAs<sup>[36]</sup>. Given that intronic microRNAs can regulate host genes by synergistic or antagonistic effects<sup>[37]</sup>, miR-25 may silence the promoter of host gene to keep the equilibrium of miR106b-25 cluster. The proliferative potential of NSCs is finite *in vivo*<sup>[38]</sup>. The miR-25 down-expression after rMS may be a contributor to this limitation. miR-25 may play a complicated role in the proliferation of NSCs after rMS, and further work will be

needed to find out how miR-25 is influenced by rMS in rat NSCs.

### 3.3 Potential Signaling Pathways Regulated by p21

MicroRNAs affect the proliferation of cells through four common mechanisms<sup>[39]</sup>. Targeting cell cycle regulators is a very important one<sup>[40]</sup>. To predict all conserved targets of rat miR-106 cluster members, TargetScan (www.targetscan.org) and KEGG database were used in this study. We found that the cell cycle progression regulator p21 was a direct target of miR-106b, and p57 was a target of miR-25<sup>[40-42]</sup>.

Overexpressing the entire miR-106b-25 clusters in adult NSCs could promote cells proliferation<sup>[13]</sup>. This is related to the expression of cyclin-dependent kinase inhibitors such as p21, which can control G<sub>1</sub>-to-S cell cycle transition<sup>[40]</sup>. P21 that represses the downstream protein of cyclinD1-cdk4/6, cyclin A-cdk2 and cyclin E-cdk2 complexes regulates cells maintenance by negatively prohibiting cell progression through G<sub>1</sub> phase<sup>[43-45]</sup>. Marques-Torreson *et al* showed that p21 could also repress the expression of Sox2 gene<sup>[46]</sup>, which is highly expressed in NSCs<sup>[47]</sup>. Lange *et al* reported that overexpression of cdk4/cyclinD1 could shorten the G<sub>1</sub> phase of progenitors and increase the expansion of progenitor<sup>[48]</sup>, while loss of cdk2 and cdk4 could reduce the length of S-phase and be prone to neuronal differentiation<sup>[49]</sup>. Moreover, as a repressor, p21 can bind to E2F1, c-Myc or STAT3 to control cells fate<sup>[50]</sup>. Our data indicated that rMS suppressed the expression of p21 in NSCs and up-regulated the expressions of its downstream protein (cyclinD1, cyclinA, cyclinE, cdk2 and cdk4) to promote the proliferation of NSCs.

In conclusion, we demonstrated that rMS can promote the proliferation of NSCs in a dose-dependent manner *in vitro*. miR-106b/p21/cdks/cyclins pathway is involved in the proliferation of NSCs after rMS by modulating cell cycle progression.

### Conflict of Interest Statement

The authors declare no conflict of interest in this study.

### REFERENCES

- Pell GS, Roth Y, Zangen A. Modulation of cortical excitability induced by repetitive transcranial magnetic stimulation: influence of timing and geometrical parameters and underlying mechanisms. *Prog Neurobiol*, 2011,93(1):59-98
- Touge T, Gerschlagel W, Brown P, *et al*. Are the after-effects of low-frequency rTMS on motor cortex excitability due to changes in the efficacy of cortical synapses? *Clin Neurophysiol*, 2001,112(11):2138-2145
- Bilek E, Schafer A, Ochs E, *et al*. Application of high-frequency repetitive transcranial magnetic stimulation to the DLPFC alters human prefrontal-hippocampal functional interaction. *J Neurosci*, 2013,33(16):7050-7056
- Lisanby SH, Datto CJ, Szuba MP. ECT and TMS: past, present, and future. *Depress Anxiety*, 2000,12(3):115-117
- Rothkegel H, Sommer M, Paulus W. Breaks during 5Hz rTMS are essential for facilitatory after effects. *Clin Neurophysiol*, 2009,121(3):426-430
- Guo F, Han X, Zhang J, *et al*. Repetitive transcranial magnetic stimulation promotes neural stem cell proliferation via the regulation of MiR-25 in a rat model of focal cerebral ischemia. *PLoS One*, 2014,9(10):e109267
- Zeng Y, Yi R, Cullen BR. Recognition and cleavage of primary microRNA precursors by the nuclear processing enzyme Drosha. *EMBO J*, 2005,24(1):138-148
- Saito K, Ishizuka A, Siomi H, *et al*. Processing of pre-microRNAs by the Dicer-1-Loquacious complex in *Drosophila* cells. *PLoS Biol*, 2005,3(7):e235
- Kim VN, Han J, Siomi MC. Biogenesis of small RNAs in animals. *Nat Rev Mol Cell Biol*, 2009,10(2):126-139
- Cremisi F. MicroRNAs and cell fate in cortical and retinal development. *Front Cell Neurosci*, 2013,7:141
- Perruisseau-Carrier C, Jurga M, Forraz N, *et al*. miRNAs stem cell reprogramming for neuronal induction and differentiation. *Mol Neurobiol*, 2011,43(3):215-227
- Zhao C, Sun G, Li S, *et al*. A feedback regulatory loop involving microRNA-9 and nuclear receptor TLX in neural stem cell fate determination. *Nat Struct Mol Biol*, 2009,16(4):365-371
- Brett JO, Renault VM, Rafalski VA, *et al*. The microRNA cluster miR-106b~25 regulates adult neural stem/progenitor cell proliferation and neuronal differentiation. *Aging*, 2011,3(2):108-124
- Peck B, Schulze A. A role for the cancer-associated miR-106b~25 cluster in neuronal stem cells. *Aging*, 2011,3(4):329-331
- Landgraf P, Rusu M, Sheridan R, *et al*. A mammalian microRNA expression atlas based on small RNA library sequencing. *Cell*, 2007,129(7):1401-1414
- Morte MI, Carreira BP, Machado V, *et al*. Evaluation of proliferation of neural stem cells *in vitro* and *in vivo*. *Curr Protoc Stem Cell Biol*, 2013, Chapter 2: Unit 2D.14 doi: 10.1002/9780470151808.sc02d14s24.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*, 2001,25(4):402-408
- Conti L, Cattaneo E. Neural stem cell systems: physiological players or *in vitro* entities? *Nat Rev Neurosci*, 2010,11(3):176-187
- Salic A, Mitchison TJ. A chemical method for fast and sensitive detection of DNA synthesis *in vivo*. *Proc Natl Acad Sci U S A*, 2008,105(7):2415-2420
- Hayashi Y, Takei H, Kurosumi M. Ki67 immunohistochemical staining: the present situation of diagnostic criteria. *Nihon Rinsho*, 2013,70(Suppl 7):428-432
- Reif A, Fritzen S, Finger M, *et al*. Neural stem cell proliferation is decreased in schizophrenia, but not in depression. *Mol Psychiatry*, 2006,11(5):514-522
- Arias-Carrion O, Verdugo-Diaz L, Feria-Velasco A, *et al*. Neurogenesis in the subventricular zone following transcranial magnetic field stimulation and nigrostriatal lesions. *J Neurosci Res*, 2004,78(1):16-28
- Werner S, Unsicker K, von Bohlen und Halbach O. Fibroblast growth factor-2 deficiency causes defects in adult hippocampal neurogenesis, which are not rescued by exogenous fibroblast growth factor-2. *J Neurosci Res*, 2011,89(10):1605-1617
- Jeong CH, Kim SM, Lim JY, *et al*. Mesenchymal stem cells expressing brain-derived neurotrophic factor enhance endogenous neurogenesis in an ischemic stroke model. *Biomed Res Int*, 2014,2014:129145
- Emsley JG, Hagg T. Endogenous and exogenous ciliary neurotrophic factor enhances forebrain neurogenesis in adult mice. *Exp Neurol*, 2003,183(2):298-310

- 26 Rotem A, Moses E. Magnetic stimulation of one-dimensional neuronal cultures. *Biophys J*, 2008, 94(12):5065-5078
- 27 Kim JY, Choi GS, Cho YW, *et al.* Attenuation of spinal cord injury-induced astroglial and microglial activation by repetitive transcranial magnetic stimulation in rats. *J Korean Med Sci*, 2013,28(2):295-299
- 28 Ueyama E, Ukai S, Ogawa A, *et al.* Chronic repetitive transcranial magnetic stimulation increases hippocampal neurogenesis in rats. *Psychiatry Clin Neurosci*, 2011,65(1):77-81
- 29 Sontag W, Kalka D. No effect of pulsed electromagnetic fields on PC12 and HL-60 cells. *Radiat Environ Biophys*, 2006,45(1):63-71
- 30 Vlachos A, Muller-Dahlhaus F, Roskopp J, *et al.* Repetitive magnetic stimulation induces functional and structural plasticity of excitatory postsynapses in mouse organotypic hippocampal slice cultures. *J Neurosci*, 2012,32(48):17514-17523
- 31 Gilio F, Conte A, Vanacore N, *et al.* Excitatory and inhibitory after-effects after repetitive magnetic transcranial stimulation (rTMS) in normal subjects. *Exp Brain Res*, 2007,176(4):588-593
- 32 Yang TS, Yang XH, Chen X, *et al.* MicroRNA-106b in cancer-associated fibroblasts from gastric cancer promotes cell migration and invasion by targeting PTEN. *FEBS Lett*, 2014,588(13):2162-2169
- 33 Tan W, Li Y, Lim SG, *et al.* miR-106b-25/miR-17-92 clusters: polycistrons with oncogenic roles in hepatocellular carcinoma. *World J Gastroenterol*, 2014,20(20):5962-5972
- 34 Semo J, Sharir R, Afek A, *et al.* The 106b~25 microRNA cluster is essential for neovascularization after hindlimb ischaemia in mice. *Eur Heart J*, 2013,35(45):3212-3223
- 35 Zhang XY, Tang LZ, Ren BG, *et al.* Interaction of MCM7 and RACK1 for activation of MCM7 and cell growth. *Am J Pathol*, 2013,182(3):796-805
- 36 Ying SY, Chang CP, Lin SL. Intron-mediated RNA interference, intronic microRNAs, and applications. *Methods Mol Biol*, 2010,629:205-237
- 37 Lutter D, Marr C, Krumsiek J, *et al.* Intronic microRNAs support their host genes by mediating synergistic and antagonistic regulatory effects. *BMC Genomics*, 2010, 11:224
- 38 Kippin TE, Martens DJ, van der Kooy D. p21 loss compromises the relative quiescence of forebrain stem cell proliferation leading to exhaustion of their proliferation capacity. *Genes Dev*, 2005,19(6):756-767
- 39 Ivey KN, Srivastava D. MicroRNAs as regulators of differentiation and cell fate decisions. *Cell Stem Cell*, 2010,7(1):36-41
- 40 Ivanovska I, Ball AS, Diaz RL, *et al.* MicroRNAs in the miR-106b family regulate p21/CDKN1A and promote cell cycle progression. *Mol Cell Biol*, 2008,28(7):2167-2174
- 41 Joaquin M, Gubern A, Posas F. A novel G1 checkpoint mediated by the p57 CDK inhibitor and p38 SAPK promotes cell survival upon stress. *Cell Cycle*, 2012,11(18):3339-3340
- 42 Kan T, Sato F, Ito T, *et al.* The miR-106b-25 polycistron, activated by genomic amplification, functions as an oncogene by suppressing p21 and Bim. *Gastroenterology*, 2009,136(5):1689-1700
- 43 Nishioka S, Nakano D, Kitada K, *et al.* The cyclin-dependent kinase inhibitor p21 is essential for the beneficial effects of renal ischemic preconditioning on renal ischemia/reperfusion injury in mice. *Kidney Int*, 2013,85(4):871-879
- 44 von Harsdorf R, Hauck L, Mehrhof F, *et al.* E2F-1 overexpression in cardiomyocytes induces downregulation of p21CIP1 and p27KIP1 and release of active cyclin-dependent kinases in the presence of insulin-like growth factor I. *Circ Res*, 1999,85(2):128-136
- 45 Puri PL, Balsano C, Burgio VL, *et al.* MyoD prevents cyclinA/cdk2 containing E2F complexes formation in terminally differentiated myocytes. *Oncogene*, 1997,14(10):1171-1184
- 46 Marques-Torrejon MA, Porlan E, Banito A, *et al.* Cyclin-dependent kinase inhibitor p21 controls adult neural stem cell expansion by regulating Sox2 gene expression. *Cell Stem Cell*, 2012,12(1):88-100
- 47 Miyagi S, Nishimoto M, Saito T, *et al.* The Sox2 regulatory region 2 functions as a neural stem cell-specific enhancer in the telencephalon. *J Biol Chem*, 2006,281(19):13374-13381
- 48 Lange C, Huttner WB, Calegari F. Cdk4/cyclinD1 overexpression in neural stem cells shortens G1, delays neurogenesis, and promotes the generation and expansion of basal progenitors. *Cell Stem Cell*, 2009,5(3):320-331
- 49 Lim S, Kaldis P. Loss of Cdk2 and Cdk4 induces a switch from proliferation to differentiation in neural stem cells. *Stem Cells*, 2012,30(7):1509-1520
- 50 Devgan V, Mammucari C, Millar SE, *et al.* p21WAF1/Cip1 is a negative transcriptional regulator of Wnt4 expression downstream of Notch1 activation. *Genes Dev*, 2005,19(12):1485-1495

(Received Dec. 3, 2014; revised Mar. 10, 2015)