

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/47809779>

Trophic Transfer of Lead Through a Model Marine Four-Level Food Chain: *Tetraselmis suecica*, *Artemia franciscana*, *Litopenaeus vannamei*, and *Haemulon scudder*

ARTICLE in ARCHIVES OF ENVIRONMENTAL CONTAMINATION AND TOXICOLOGY · NOVEMBER 2010

Impact Factor: 1.9 · DOI: 10.1007/s00244-010-9620-4 · Source: PubMed

CITATIONS

10

READS

93

6 AUTHORS, INCLUDING:



[Martin F Soto-Jiménez](#)

Instituto de Ciencias del Mar y Limnología, ...

50 PUBLICATIONS 825 CITATIONS

SEE PROFILE



[Claudia arellano fiore](#)

University of Milan

2 PUBLICATIONS 15 CITATIONS

SEE PROFILE



[Jorge Ruelas-Inzunza](#)

Instituto Tecnológico de Mazatlán

52 PUBLICATIONS 424 CITATIONS

SEE PROFILE

Environmental Toxicology

BIOLOGICAL RESPONSES OF A SIMULATED MARINE FOOD CHAIN TO LEAD ADDITION

MARTÍN F. SOTO-JIMÉNEZ,*† CLAUDIA ARELLANO-FIORE,‡ RUTH ROCHA-VELARDE,§ MARTÍN E. JARA-MARINI,||
JORGE RUELAS-INZUNZA,‡ DOMENICO VOLTOLINA,# MARTÍN G. FRÍAS-ESPERICUETA,†† JESÚS M. QUINTERO-ALVAREZ,†
and FEDERICO PÁEZ-OSUNA†

†Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Mazatlán, Sinaloa, México

‡Technological Institute of Mazatlán, Mazatlán, Sinaloa, México

§Mazatlán Aquarium, Mazatlán, Sinaloa, México

||Centro de Investigación en Alimentación y Desarrollo, Hermosillo, Sonora, México

#Centro de Investigaciones Biológicas del Noroeste, Mazatlán, Sinaloa, México

††Laboratorio de Estudios Ambientales, Universidad Autónoma de Sinaloa, Mazatlán, Sinaloa, México

(Submitted 29 September 2010; Returned for Revision 22 November 2010; Accepted 11 March 2011)

Abstract—This investigation sought to assess the biological responses to Pb along a simplified four-level food chain, from the primary producer, the microalgae *Tetraselmis suecica*, grown in a control medium with < 1 µg/L of Pb and exposed to a sublethal dose (20 µg/L of Pb) and used as the base of a simulated food chain, through the primary-, secondary-, and tertiary-level consumers, namely, the brine shrimp, *Artemia franciscana*; the white-leg shrimp, *Litopenaeus vannamei*; and the grunt fish, *Haemulon scudder*, respectively. Growth of Pb-exposed *T. suecica* was 40% lower than that of the control cultures, and survival of *A. franciscana* fed this diet was 25 to 30% lower than the control. No differences in the growth rates of Pb-exposed and control shrimp and fish and no gross morphological changes were evident in the exposed specimens. However, the exposed shrimp and fish had 20 and 15% higher mortalities than their controls, respectively. In addition, behavioral alterations were observed in exposed shrimp and fish, including reduction in food consumption or cessation of feeding, breathing air out of the water, reduction of motility, and erratic swimming. The negative correlation between Pb concentration in whole body of shrimp and fish and Fulton's condition factor suggested also that the exposed organisms were stressed because of Pb accumulation. Environ. Toxicol. Chem. 2011;9999:1–7. © 2011 SETAC

Keywords—Lead effects Trophic transfer Bioassay experiment Marine food chain Lead pollution

INTRODUCTION

Lead (Pb) is known to affect growth, development, and reproduction in humans and animals; therefore, Pb contamination of the marine ecosystems has become an environmental problem worldwide, especially in coastal ecosystems, in which it may reach levels of risk for aquatic organisms and for their consumers, including humans [1–3]. The metal content of the diet has been recognized as the main contribution to metal body burden of aquatic organisms and as a major route for transfer through marine food webs [4–7]. Because the metals ingested may become toxic when accumulation in the tissues of aquatic animals reaches a critical level [8], evaluation of the uptake, biotransference, and biomagnification of metals from the low trophic levels to top consumers is critical for determining their effects on marine food chains and their influence on ecosystems and on human life [9–11].

However, the study of transfer processes of metals along food webs and the evaluation of their effects is difficult in the natural environment, so knowledge of their impact on organisms, populations, communities, and ecosystems is still limited. In particular, little information is available on the trophic transfer of Pb in food chains in subtropical ecosystems [12,13], and almost none is available on the effects of the chronic toxicity of Pb for aquatic organisms.

Many bioassays reproducing the natural conditions of an ecosystem have been developed over the past decades, in an effort to understand the relationships between metal transfer and its effects on aquatic organisms [13]. However, experimental studies with aquatic food chains exposed to environmentally realistic concentrations of metal through the consumption of whole, living prey organisms are surprisingly rare [5,8,14,15]. Most previous studies have focused on a single species or on no more than two levels of an aquatic food chain and did not use natural chain links, because metals were added by injection into prey items or into ground fish or through direct exposure to waterborne metals in concentrations far in excess of those occurring in most natural waters [14–16].

The present study determined the biological responses to dietary exposure to Pb of organisms of different trophic levels in a simulated marine food chain, chosen in part because of their availability and ease of culture as well as because they are the normal links of some common food chains used in aquaculture, such as microalgae–*Artemia*–shrimp and fish larvae or post-larvae, or because they are links of natural food chains (e.g., shrimp–carnivorous demersal fish).

MATERIALS AND METHODS

The experimental design consisted of feeding the brine shrimp *Artemia franciscana* with the microalgae *Tetraselmis suecica* grown in f/2 medium (control diet) or in the same medium with 20 µg/L Pb. This concentration, which is of the same order of magnitude observed in contaminated coastal environments [17] and between four and five times higher than

* To whom correspondence may be addressed
(martin@ola.icmyl.unam.mx).

Published online xx Month 2011 in Wiley Online Library
(wileyonlinelibrary.com).

concentrations found in Mazatlán Harbor, Sinaloa, México [13], is equivalent to 2.5 times the continuous concentration criterion for Pb (CCC = 8.1 $\mu\text{g/L}$) ([18]; <http://www.epa.gov/ost/criteria/wqtable/>). This is an estimate of the level of Pb to which an aquatic community can be exposed indefinitely without an undesired effect and, according to the European Community Dangerous Substances Directive, is considered dangerous for the aquatic environment [19].

The *Artemia* biomass obtained with the control cultures was fed to juveniles of the white-leg shrimp, which in turn served as diet for the carnivorous fish *Haemulon scudderi*. The same food chain, starting with *Artemia* fed with Pb-grown *T. suecica*, served for studying the transference of this metal and assessing its effect through the food chain. Because metal bioaccumulated through dietary exposure can be deposited in different tissues and affect different physiological processes [8,20], we also determined the concentration of Pb in different organs and tissues of shrimp and fish.

All experiments were carried out with natural seawater from Mazatlán Bay, filtered through 2- μm cartridge filters, with an ultraviolet and ozone treatment system. Mean water temperature, salinity, and pH were $28.2 \pm 2.0^\circ\text{C}$, 34.6 ± 1.2 ppt, and 8.1 ± 0.1 , and the respective ranges were 28 to 32°C , 34 to 36 ppt, and 7.9 to 8.2. With the exception of microalgae cultures, continuous aeration maintained dissolved oxygen saturation close to 90 to 95% ($> 7 \text{ mg/L}$). Light was provided by the natural 14:10 h light:dark photoperiod, and irradiance during daylight ranged from 250 to 1,500 $\mu\text{mol/m}^2/\text{s}$.

Trophic level 1: Microalgae

The control diet was the prasinophyte *T. suecica* (strain TES-1 of the Centro de Investigación Científica y de Educación Superior de Ensenada Culture Collection) [21] grown with the multistep procedure (1, 15, 120 L, with 72 h for each step) in three 160-L fiberglass cylindrical tanks with 120 L of f/2 medium prepared with natural seawater (mean background dissolved Pb concentration $0.29 \pm 0.12 \mu\text{g/L}$, range 0.18–0.50 $\mu\text{g/L}$). The experimental diet was grown in three similar tanks with f/2 medium and a suitable volume (depending on the initial Pb concentration) of a 1 mg/L Pb (NO_3)₂ stock solution, to obtain a concentration of 20 $\mu\text{g/L}$ Pb.

Pb(NO_3)₂ was chosen because Pb(II) is the most stable ionic species and is thought to be the form in which Pb is mostly bioaccumulated by aquatic organisms [22]. This procedure was repeated 10 times to obtain a sufficient amount of *Artemia* biomass for shrimp feeding.

Trophic level 2: Artemia

Nauplii of *A. franciscana* hatched from commercial cysts (INVE Aquaculture) as described by Van Stappen [23] were harvested and transferred to three fiberglass tanks (3,000 L) with approximately 2,500 L of filtered seawater and an initial concentration of 350 to 400 nauplii/L. The Pb-exposed and control microalgae cultures had different cell concentrations. For this reason, daily feeding was provided in two rations of approximately 10 L of the control cultures and approximately 20 L of those exposed to Pb. The resulting concentrations in the *Artemia* cultures ranged between 2.5 and 4×10^6 cells/L. The final harvest was after 19 to 23 d in three (control diet) and six (experimental diet) separate cultures, which were necessary to obtain the amount of biomass for feeding the next links of the control and experimental food chains, respectively.

Trophic level 3: Shrimp

Litopenaeus vannamei juveniles (total wet wt 1.0–1.5 g and total length 5.3–5.7 cm) were obtained from a commercial farm, acclimated during 3 d in three 3,000-L fiberglass tanks with filtered seawater, and fed with nonexposed *Artemia* biomass. After acclimation, groups of 36 specimens were distributed in 16 100-L tanks. The diet of 12 groups, chosen at random, was adult *Artemia* fed with Pb-exposed microalgae. The remaining four received the control diet. In both cases, feeding was ad libitum, twice daily. Excess food was removed after 1 h and, to avoid ingestion of nonexperimental food through cannibalism, molting specimens were removed from the tank.

Trophic level 4: Fish

After a preventive bath [24] (0.025 ml/L formalin and 0.1 mg/L malachite green), in total 120 *H. scudderi* of similar size and weight (7.15 ± 1.2 cm total length and 4.63 ± 1.7 g) captured in Mazatlán Bay were acclimated for 3 d in three 3,000-L fiberglass tanks filled with 5- μm -filtered seawater. After acclimation, groups of 15 fish were separated into each of the eight 300-L tanks and fed daily to satiation with nonexposed shrimp for 6 d. After this period, the fish in five tanks were fed the Pb-exposed shrimp, and the remaining three were fed the control diet.

Determination of sublethal effects

The yields of the microalgae cultures were assessed as cell concentration, determined with a hemocytometer, and growth rates were calculated using the initial and final cell numbers (N). The dry biomass yields were determined using triplicate 20-ml samples, concentrated on precalibrated Whatman GF-C glass fiber filters, rinsed with 3 to 5 ml of a 3% ammonium formate solution, and dried to constant weight at 65°C in a convection oven.

The concentration of the *A. franciscana* cultures was determined at approximately weekly intervals by direct counts of triplicate samples of known volume, from nauplii to the day of the final harvest. In each case, 10 specimens taken at random were measured from the oral to the aboral region under a stereomicroscope with a calibrated eyepiece, and the specimens contained in 1-L sample were concentrated on a 100- μm mesh sieve, counted, and dried as for microalgae to estimate the individual dry weight and the biomass yield.

During the feeding experiments of all trophic levels, dead shrimp and exuviae, as well as dead fish, were removed immediately before adding each of the half-daily food rations. The wet weights and total lengths of five active and apparently healthy specimens of shrimp and fish were obtained weekly, and their individual total weights (W) and lengths (TL) were used to calculate their Fulton's condition factors (K) with the equation $K = 100 \times W/\text{TL}^3$. A decrease in this factor may be interpreted as depletion of body reserves, which could be a response to external stressors such as heavy metals [25].

Sampling for Pb analysis

Sample collection and analysis were performed following clean technique protocols [3]. The seawater samples used to determine background Pb concentrations were collected in 1-L polyethylene (LDPE) bottles, previously acid cleaned, filled with 6 N HCl Optimum Grade (Fisherbrand), and stored for at least one month.

The samples (100–300 ml) of *T. suecica* cultures obtained at the time of harvest were concentrated on precombusted (500°C ,

4 h), acid-cleaned (2 M HCl) Whatman GF/F glass fiber filters; dried at 55°C to constant weight; and analyzed as for the rest of the samples to obtain the mean Pb contents. Composite samples of *Artemia* (4–5 g wet wt per batch) were obtained with a 200- μ m mesh sieve at least 6 h after the addition of microalgae, at approximately weekly intervals and at the end of the growth experiment. By that time, no microalgae were available, *Artemia* had ceased feeding, and their guts were practically empty. For this reason, no further depuration was deemed necessary.

To avoid depuration procedures, the weekly shrimp (four to 12 specimens) and fish (three to five organisms) samples were obtained several hours (6–12 h) after feeding. Shrimp specimens were dissected to separate the muscle, exoskeleton (antennules, carapace, pleopods, and pereopods), hepatopancreas, and the remaining organs contained in the cephalothorax; in the case of fish, muscle, skeleton (including skull and dorsal and anal fin-rays), skin and scales, gills, liver, viscera, and remaining tissues were separated [13,26]. These samples were frozen and freeze-dried for 72 h. The dried samples were ground, homogenized, and stored in sealed polypropylene containers until analysis.

Analysis

All samples were processed and analyzed in high-efficiency particulate-filtered air (HEPA class 1000) and a trace-metal-clean laboratory using high-purity reagents and water. The powdered samples (100–200 mg) were digested in 30-ml Teflon[®] vessels with 8 to 10 ml of trace-metal-grade HNO₃:HCl (3:1 v/v), heated overnight at 130°C on a Mod Block unit, transferred to clean polypropylene vials, and diluted to 25 ml with MilliQ water.

The dissolved Pb concentrations of the acidified water and of the biological samples were determined using graphite furnace atomic absorption spectroscopy (GFAAS; Varian SpectrAA 220). Blanks and certified reference material (SLEW-2 estuarine water [$n=5$], DORM-2 dogfish muscle [$n=6$], and National Institute of Standards and Technology [NIST] 1566b oyster tissue [$n=4$]) were analyzed to verify the accuracy of the extraction method. The certified reference material gave the following results: 0.008 ± 0.001 μ g/L for SLEW-2, 0.0611 ± 0.012 for DORM-2, and 0.32 ± 0.02 for NIST 1566b (certified values 0.009 ± 0.001 μ g/L, 0.065 ± 0.007 μ g/g, and 0.31 ± 0.01 μ g/g, respectively).

Data processing

The mean cell concentrations, growth rates, dry biomass yields, and Pb contents of the Pb-exposed and control *T. suecica* cultures were compared using Student's *t* tests. The biological responses over time of the Pb-exposed and control cultures of the higher trophic levels were compared using two-way analysis of variance repeated-measures tests, followed by paired *t* tests

Table 1. Mean cell concentrations (N , in 10^6 cells/ml), growth rates (μ in cell divisions/day), dry biomass yields (mg/L), and Pb content (μ g/g dry wt) of *Tetraselmis suecica* cultures in f/2 medium (control) or in f/2 medium with 20 μ g/L Pb^a

	Control	Exposed
Cell count (N)	$0.87 \pm 0.05^*$	0.47 ± 0.06
Growth rates (μ)	$1.29 \pm 0.11^*$	1.02 ± 0.16
Biomass	$98 \pm 10^*$	48 ± 20
Pb	$3.5 \pm 1.1^*$	18.4 ± 3.4

^a Results are means \pm standard deviation; $n=10$.

* Significant differences (analysis of variance, $\alpha=0.05$) were observed between control and exposed microalgae.

with the Bonferroni correction. Because the response data collected at different times from the same groups of organisms are not independent, Mauchly's test statistic was used to evaluate whether the relationships between pairs of experimental conditions were similar (assumption of sphericity). The level of significance in all statistical tests was 0.05. Statistica 7.0 (Statistica for Windows; Statsoft) and Office Excel 2007 (Microsoft) were used for all the statistical analyses.

RESULTS

The mean final cell concentrations, growth rates, and total dry biomass yields of the control cultures of *T. suecica* were significantly higher than those obtained with the Pb-exposed cultures (Table 1). The absorption efficiencies (AE) of these cultures were estimated with the equation $AE = 100 (P_{bi} - P_{bf})/P_{bw}$, where P_{bi} and P_{bf} are the initial and final Pb concentrations of the total microalgal biomass and P_{bw} is the initial amount of Pb present in the whole volume of the culture medium. The mean Pb content of *T. suecica* grown for 72 h in the Pb-enriched growth medium was 18.4 ± 3.4 μ g/g, with an absorption efficiency of 10 to 20% of the added Pb, whereas that of the control cultures averaged 3.5 ± 1.1 μ g/g, with an assimilation efficiency $>70\%$.

Artemia franciscana (level 2)

A tendency to lower survival and lower biomass yields was noted in the exposed cultures, but the differences were significant ($p < 0.05$) only on the day of final harvest (Table 2). A similar tendency existed in the case of total length, but no significant differences were observed throughout the experiment (Table 2). The mean Pb contents of the exposed *A. franciscana* used as food for the next food chain link were 12.0 ± 0.9 to 15.0 ± 1.7 μ g/g, which was significantly higher than that of the control *Artemia* cultures (1.3 ± 0.4 to 1.8 ± 0.3 μ g/g; Student's *t* tests for paired observations, $\alpha=0.05$).

Table 2. Mean cell count (ind/L), total length (mm), and dry biomass (mg/L) of *Artemia franciscana* cultures fed *T. suecica* grown in f medium with the background Pb concentration (control) or with 20 μ g/L of Pb (exposed)^a

Day	Cell count (ind/L)		Length (mm; $n=10$)		Dry biomass (mg/L)	
	Control	Exposed	Control	Exposed	Control	Exposed
1	$390 \pm 15D$	$395 \pm 17D$	$0.7 \pm 0.1A$	$0.6 \pm 0.2A$	$12.2 \pm 2.8A$	$11.5 \pm 3.0A$
7	$379 \pm 10D$	$368 \pm 14CD$	$3.5 \pm 0.7B$	$2.9 \pm 1.4B$	$65.0 \pm 2.3B$	$60.0 \pm 2.5B$
13	$353 \pm 12C$	$342 \pm 22CB$	$6.0 \pm 0.8C$	$5.0 \pm 1.2C$	$102 \pm 4.0C$	$94.3 \pm 7.6C$
19–23	$320 \pm 20B$	$255 \pm 30A$	$9.2 \pm 1.4D$	$8.2 \pm 1.8D$	$195 \pm 12E$	$153 \pm 18D$

^a Results are means \pm standard deviation. Equal letters means no significant differences: $A < B < C < D < E$.

Litopenaeus vannamei (level 3)

A general tendency toward lower survival and total length and weight was observed in the Pb-exposed shrimp cultures, but the differences were significant only on the final day for survival and size, whereas the controls had higher wet weight from day 28 until the day of harvest. Although no Pb-related differences were observed between the intermediate and the final mean values of the Fulton's condition factor calculated for the two treatments (Table 3), a significant negative correlation existed between Pb concentrations (whole body) and this factor ($r = -0.63$, $p < 0.05$).

After 30 d of exposure, we observed abnormal behavior of the exposed organisms, such as surfacing (presumably for air breathing as a response to stress), reduction of motility, and erratic swimming. No behavioral alterations were observed throughout the experiment in the control cultures.

The Pb contents of the different organs and tissues showed a clear trend to increase as a function of time, and the final concentrations in all tissues of the exposed organisms were significantly ($p < 0.05$) higher than the initial values (Table 4). Lead levels in control shrimp also increased, indicating a natural bioaccumulation with regard to age of organism. In

addition, the Pb levels in organs and tissues were consistently higher in the Pb-exposed organisms than in controls ($p < 0.05$).

The weight-normalized Pb concentration showed the following order in the final destination of transferred Pb: exoskeleton (28–38% of total Pb) > muscle (23–31%) > remaining tissues of cephalothorax (27–35%) > hepatopancreas (7–11%). Muscle and exoskeleton, which made up approximately 90% of the body mass, contained only between 51 and 69% of the total Pb burden, whereas the remaining organs and tissues contained more than one-third of total Pb, although they made up < 5% of the total body mass.

Haemulon scudderii (level 4)

No significant differences were observed in intermediate and final length, mean wet weight, or Fulton's condition factor throughout the experiment, but final survival was significantly lower in the exposed organisms (Table 5). The behavior of the Pb-exposed and control fish was similar during most of the experiment, but during the last week the exposed fish were more lethargic during feeding, and their food consumption was lower than that of the control fish. In addition, several exposed specimens were observed surfacing, presumably for air breathing.

Table 3. Initial, intermediate, and final mean values of survival, total length (mm), wet weight (g), and Fulton's condition index (K) of *Litopenaeus vannamei* fed the control and the experimental diets (*Artemia franciscana* fed *Tetraselmis suecica* grown in f medium with the background Pb concentration or with $20 \mu\text{g L}^{-1}$ of Pb)^a

Day	Survival (%)		Length (mm)		Wet weight (g)		K	
	Control	Exposed	Control	Exposed	Control	Exposed	Control	Exposed
1	100C	100C	$6.3 \pm 1.6\text{A}$	$6.4 \pm 1.1\text{A}$	$1.2 \pm 0.3\text{A}$	$1.3 \pm 0.2\text{A}$	0.50 ± 0.1	0.51 ± 0.13
15	94–96C	92–96C	$10.8 \pm 0.6\text{B}$	$10.5 \pm 1.6\text{B}$	$5.1 \pm 0.3\text{B}$	$4.8 \pm 0.7\text{B}$	0.48 ± 0.09	0.45 ± 0.17
28	90–95C	88–92BC	$13.3 \pm 0.6\text{C}$	$12.1 \pm 0.6\text{C}$	$7.2 \pm 0.4\text{D}$	$6.2 \pm 0.3\text{C}$	0.41 ± 0.13	0.40 ± 0.2
42	84–90B	67–78A	$14.8 \pm 1.0\text{D}$	$13.1 \pm 0.6\text{C}$	$8.5 \pm 0.6\text{E}$	$7.1 \pm 0.3\text{D}$	0.38 ± 0.11	0.31 ± 0.14

^a Results are means \pm standard deviation; $n = 12$. Equal letters means no significant differences: $\text{A} < \text{B} < \text{C} < \text{D} < \text{E}$.

Table 4. Weighted normalized concentration of Pb ($\mu\text{g g}^{-1}$) in different organs and tissues of white shrimp *Litopenaeus vannamei* fed with *Artemia franciscana* exposed to Pb via diet^a

Day	Exoskeleton		Muscle		Hepatopancreas		Remaining tissues of cephalo-thorax	
	Control	Exposed	Control	Exposed	Control	Exposed	Control	Exposed
0	$0.12 \pm 0.06\text{A}$	$0.11 \pm 0.09\text{A}$	$0.06 \pm 0.06\text{A}$	$0.06 \pm 0.09\text{A}$	$0.03 \pm 0.03\text{A}$	$0.05 \pm 0.04\text{A}$	$0.10 \pm 0.02\text{A}$	$0.09 \pm 0.05\text{A}$
15	$0.46 \pm 0.17\text{B}$	$0.67 \pm 0.11\text{C}$	$0.21 \pm 0.08\text{B}$	$0.33 \pm 0.05\text{C}$	$0.14 \pm 0.01\text{B}$	$0.19 \pm 0.02\text{C}$	$0.12 \pm 0.04\text{A}$	$0.48 \pm 0.08\text{B}$
28	$0.51 \pm 0.09\text{C}$	$0.99 \pm 0.14\text{D}$	$0.23 \pm 0.11\text{B}$	$0.52 \pm 0.12\text{D}$	$0.15 \pm 0.09\text{C}$	$0.33 \pm 0.05\text{D}$	$0.11 \pm 0.04\text{A}$	$0.53 \pm 0.11\text{B}$
42	$0.53 \pm 0.21\text{C}$	$1.18 \pm 0.18\text{E}$	$0.21 \pm 0.05\text{B}$	$0.98 \pm 0.14\text{E}$	$0.18 \pm 0.09\text{C}$	$0.34 \pm 0.09\text{D}$	$0.13 \pm 0.05\text{A}$	$1.13 \pm 0.14\text{C}$

^a All results are means \pm standard deviation. One-way analysis of variance followed by Tukey's honestly significant difference test was performed for comparing Pb concentrations among groups and treatments. Equal letters means no significant differences: $\text{A} < \text{B} < \text{C} < \text{D} < \text{E}$.

Table 5. Initial, intermediate, and final mean values of survival, total length (mm), wet weight (g), and Fulton's condition index (K) of *Haemulon scudderii* fed the control and the experimental diets (*Litopenaeus vannamei* fed *Artemia franciscana* fed *Tetraselmis suecica* grown in f medium with the background Pb concentration or with 20mg L^{-1} of Pb)^a

Day	Survival (%)		Length (mm)		Wet weight (g)		K	
	Control	Exposed	Control	Exposed	Control	Exposed	Control	Exposed
1	100D	100D	$6.9 \pm 1.2\text{A}$	$7.0 \pm 0.8\text{A}$	$4.6 \pm 1.8\text{A}$	$4.9 \pm 1.2\text{A}$	1.9 ± 0.4	1.9 ± 0.4
15	> 95D	> 90C	$7.7 \pm 0.2\text{B}$	$7.3 \pm 0.2\text{B}$	$7.9 \pm 0.2\text{B}$	$6.9 \pm 0.8\text{B}$	1.8 ± 0.3	1.8 ± 0.2
28	88–96C	72–88BC	$9.0 \pm 1.0\text{C}$	$8.6 \pm 0.8\text{C}$	$10.9 \pm 2.9\text{C}$	$9.8 \pm 1.0\text{C}$	1.7 ± 0.4	1.7 ± 0.4
42	88–91B	65–75A	$10.2 \pm 1.0\text{C}$	$9.3 \pm 1.0\text{C}$	$15.5 \pm 1.0\text{D}$	$14.5 \pm 0.8\text{D}$	1.7 ± 0.4	1.6 ± 0.3

^a Results are means \pm standard deviation; $n = 12$. Equal letters means no significant differences: $\text{A} < \text{B} < \text{C} < \text{D} < \text{E}$.

During the experiment, the mean total Pb body burden increased from 0.61 ± 0.08 to $1.24 \pm 0.27 \mu\text{g/g}$, and from 0.55 ± 0.29 to $3.32 \pm 0.41 \mu\text{g/g}$ in the control and the exposed fish, respectively. The weight-normalized concentration of Pb in different organs and tissues of *H. scudleri* showed a differential accumulation during the exposure period (Table 6). Skeleton, skin, scales, and muscle represented between 82 and 90% of the body mass and accumulated between 70 and 73% of the total metal transferred to this level. Liver and other internal organs and tissues had higher relative contents, indicating a preferential accumulation of Pb in these tissues. The internal organs of exposed fish showed from two to four times more Pb than the controls, with a net contribution close to 30% of total accumulated Pb but with a body dry weight contribution to the total biomass lower than 10%. Small amounts of Pb were accumulated in the gills (~3% of accumulated Pb).

DISCUSSION

Ecotoxicological effects

The effects on the growth rate and cell division observed for *T. suecica* resulting from Pb exposure have been reported previously for different microalgae species, including *Tetraselmis* sp., and for different metals such as Cd [27], Cu [28,29], and Pb [30–33]. The reduced growth rate of *Tetraselmis* and other microalgae exposed to Pb has been explained by these authors as resulting from alterations of metal chelation and phytochelatin production or of enzymatic systems. In particular, Pb in microalgae induces the activity of superoxide dismutase (SOD) isoenzymes, lowers the ratio of reduced to oxidized glutathione (GSH:GSSG ratio), and causes changes in cellular chlorophyll contents generated under stress conditions, which inhibit mitotic spindle formation and consequently inhibit cell division and reduce the growth rate.

In crustaceans, Pb affects hemolymph circulation because of cytological damage such as cytoplasmic vacuolation and reduction of spaces for circulation of the hemolymph [34,35]. This causes breathing difficulties and low internal oxygen concentrations related to impaired hemolymph circulation. In addition, Pb accumulated in organs and tissues may interact with the mitochondrial membranes, decreasing the metabolic rate and consequently O_2 consumption, ammonia excretion, and activity of the enzyme that catalyzes the decomposition of adenosine triphosphate (ATP) [36–38].

We did not observe external morphological anomalies or lesions in the Pb-exposed *H. scudleri*, but some showed difficulty in breathing, lethargy, and reduced consumption of food. The first effect might be due to a Pb-related inhibition of Ca^{2+} , Na^+ , and Cl^- uptake or to cytological damage of the epithelium of the gill tissue [10,11,39], although specific analyses (e.g., number of red blood cell and the activity of δ -aminolevulinic acid dehydratase) and histological evaluations for injuries to this organ and other internal organs, such as liver and kidney, will be required to assess the impact to fish of dietary exposure to Pb.

Comparison with previous studies

As mentioned above, the effects on growth observed for *T. suecica* resulting from Pb exposure have been reported previously for different microalgae species and for different metals [26–29]. However, the grade of growth inhibition varies greatly among different metals, among phytoplankton species, and among experimental conditions (e.g., concentration of metal, initial cell densities, light illumination, temperature,

Table 6. Weight-normalized concentration of Pb ($\mu\text{g g}^{-1}$) in different organs and tissues of grunt fish *Haemulon scudleri* fed with *Litopenaneus vannamei* exposed to Pb via diet^a

Day	Skeleton			Skin/scales			Muscle			Gills			Liver			Viscera (+remain tissues)		
	Control	Exposed	Exposed	Control	Exposed	Exposed	Control	Exposed	Exposed	Control	Exposed	Exposed	Control	Exposed	Exposed	Control	Exposed	Exposed
0	0.14 ± 0.08A	0.12 ± 0.06A	0.13 ± 0.05A	0.15 ± 0.03A	0.13 ± 0.04A	0.13 ± 0.05A	0.13 ± 0.04A	0.13 ± 0.05A	< LD	< LD	< LD	< LD	0.11 ± 0.03A	0.10 ± 0.03A	0.10 ± 0.03A	0.07 ± 0.06A	0.06 ± 0.03A	0.06 ± 0.03A
15	0.31 ± 0.09B	0.27 ± 0.05B	0.33 ± 0.11B	0.28 ± 0.08B	0.20 ± 0.02B	0.29 ± 0.10B	0.20 ± 0.02B	0.29 ± 0.10B	0.03 ± 0.02A	0.01 ± 0.02A	0.03 ± 0.02A	0.03 ± 0.02A	0.12 ± 0.05A	0.24 ± 0.02B	0.24 ± 0.02B	0.09 ± 0.05A	0.25 ± 0.08B	0.25 ± 0.08B
28	0.37 ± 0.05B	0.40 ± 0.04B	0.56 ± 0.11C	0.33 ± 0.06B	0.29 ± 0.06B	0.68 ± 0.16C	0.29 ± 0.06B	0.68 ± 0.16C	0.09 ± 0.01B	< LD	0.09 ± 0.01B	0.09 ± 0.01B	0.22 ± 0.04B	0.42 ± 0.06C	0.42 ± 0.06C	0.10 ± 0.05A	0.27 ± 0.03B	0.27 ± 0.03B
42	0.30 ± 0.09B	0.79 ± 0.21C	0.82 ± 0.19C	0.31 ± 0.1B	0.29 ± 0.05B	0.74 ± 0.14C	0.29 ± 0.05B	0.74 ± 0.14C	0.09 ± 0.02B	0.01 ± 0.03A	0.09 ± 0.02B	0.09 ± 0.02B	0.19 ± 0.06B	0.52 ± 0.16C	0.52 ± 0.16C	0.14 ± 0.04A	0.41 ± 0.05C	0.41 ± 0.05C

^a All results are means ± standard deviation. One-way analysis of variance followed by Tukey's honestly significant difference test was performed for comparing Pb concentrations among groups and treatments. LD = detection limit. Equal letters means no significant differences; A < B < C < D < E.

composition of culture media, and exposure time). Previous experiments with *Tetraselmis* sp. have shown that this genus is relatively more tolerant to metals compared with other micro-algae genera. The tolerance of the *Tetraselmis* genus has been related to detoxification mechanisms for sequestering and expelling metal, such as metal chelating complexes, most probably as class III metallothioneins and insoluble salts in the cytoplasm; elevated vacuolization of the cytoplasm; formation of multilayered cell walls; and increase in the excretion of organic matter [27–29,32–34].

Multiple authors have reported negative effects on motility, fecundity and reproduction, and morphological alterations in crustacean species associated with exposure to high levels of Pb (> 100 ppb) [8,35–39]. However, previous bioassays with fish exposed to long-term high Pb concentrations (e.g., > 100 d to 235 ppb Pb, ~200 d to 120 ppb) have reported a wide range of effects, including neurological degeneration and destruction, lordoscoliosis, paralysis, muscular atrophy, and such morphological alterations as spinal deformities, black tails, and degeneration of the caudal fin [40,41]. Chronic exposure to sublethal Pb levels also causes alteration in physiological functions of fish, such as reduction in food consumption or cessation of feeding, disturbed ionic regulation, reduced swimming speed, growth inhibition, interference with reproduction, and mortality [42,43]. In our short-term exposure with Pb doses lower than those used in most reported studies, none of these effects was verified.

Although the exposed shrimp and fish did not show morphological changes and appeared healthy and active during most of the exposure period, this study concerned only short-term, subtle effects of moderate doses of Pb. Health implications of long-term exposure to moderate doses of Pb are beyond the scope of the present study.

Lead trophic transference

This experiment did not follow the traditional linear responses of aquatic species exposed to unrealistic concentrations of a toxicant for time periods that are short relative to the organism's life span. Rather, our experiment was designed to study the trophic transfer and any biological effects of Pb through a multispecies system covering a four-trophic-level food chain. We exposed the primary producer of our food chain to environmentally realistic concentrations of dissolved Pb; examined its effect on the next trophic levels caused by the consumption of whole, living prey organisms; and quantified the Pb content in different tissues of shrimp and fish.

The trophic transfer of Pb and the biological responses depend on the structure and composition of the aquatic food chain, so the selection of species is a determining factor when studying the effects of a pollutant, because different trophic relations can yield different scenarios for metal transference and variability of responses. This study included some organisms that are common residents of the coastal waters of the south-eastern Gulf of California and, for practical reasons, some species that are commercially important. For example, *T. suecica* is widely distributed and is commonly utilized in aquaculture as food for fishes, crustaceans, and mollusks. *Litopenaeus vannamei* is the most widely cultivated shrimp species in the world and grunt fish is a commercially important resource in the region; this issue is of particular relevance for the safety of marine products. *Artemia franciscana* was the only species used that is not a resident in the regional coastal waters, although it is widely used in commercial hatcheries. Among marine invertebrates, *Artemia* is one of the most commonly

used species as a test organism to assess toxic effects of metals because of several characteristics that guarantee reliability and feasibility in ecotoxicology studies.

Because *T. suecica* is the basis of our simulated marine food chain, the effects of Pb on the growth and development of phytoplankton are transferred to the zooplankton and subsequently to the upper trophic levels. However, the concentration of Pb did not increase as the metal moved up through the food chain [13]. This suggests mechanisms of self-regulation and elimination of Pb in the species used in our simulated food chain.

Harmful effects in our simulated marine food chain exposed to chronic levels of Pb were not always obvious in the organisms exposed, presumably because they might take more time to appear. However, sustaining a laboratory-based food chain during the entire-life stage is complicated and expensive, which makes entire-life experiments difficult. To the best of our knowledge, no studies following this strategy are available in the literature. In fact, few studies in the literature have related the whole-body Pb concentrations in an organism and concentrations in their tissues and organs to sublethal effects of Pb via diet, which makes it difficult to reach definitive conclusions. Even so, the results revealed that added Pb affected physiological functions and behavior of organisms without killing them.

Ecological implications

A major challenge in this study was to translate the biological response data in terms of ecological significance and possible application under natural circumstances. Our design took into account some of the environmental conditions and interactions in subtropical coastal ecosystems. For this reason, the adverse effects on individual fitness and survival observed in our organisms exposed to Pb via diet probably allow inference of the direct consequences of Pb contamination to the local populations and whole communities.

Eutrophication has been increasing in coastal waters from Gulf of California as a result of inputs of nutrients (C, N, and P) associated with the increase of population living in coastal areas and the rise in demand for foods that require a greater use of fertilizers and animal feed. In addition, the Gulf of California coastal zone is also being influenced by metals inputs from agricultural, mining, industrial, and domestic activities. Although it is known that alteration in the cycling of nutrients in coastal waters provokes perturbations in the biogeochemical cycles of trace metals, knowledge about the influence of nutrient enrichments on the metal uptake in natural ecosystems and, consequently, the toxicity and trophic transfers in aquatic food webs is very limited. For this reason, it is necessary to perform additional studies on the metal transfer processes and on the short- and long-term effects on local species to assess their ecological implications, including their possible consequences for human life.

Acknowledgement—J.F. Ontiveros-Cuadras and S. Rendón-Rodríguez helped during the bioassays, H. Bojórquez-Leyva performed the laboratory analyses, and V. Montes and Y. Montaña-Ley provided support in manuscript preparation. We especially thank the personnel of Mazatlán Aquarium for their support. Financial support was provided by the grant UNAM-PAPIIT IN206409 and SEP-CONACYT 60215.

REFERENCES

1. Bryan GW, Langston WJ. 1992. Bioavailability, accumulation and effects of heavy metals in sediments with special reference to United Kingdom estuaries: A review. *Environ Pollut* 76:89–131.

2. Stamatis N, Kamidis N, Sylaios G. 2006. Sediment and suspended matter lead contamination in the Gulf of Kavala, Greece. *Environ Monit Assess* 115:433–449.
3. Soto-Jiménez MF, Páez-Osuna F, Scelfo G, Hibdon S, Franks R, Aggarawl J, Flegal AR. 2008. Lead pollution in subtropical ecosystems on the SE Gulf of California coast: A study of concentrations and isotopic composition. *Mar Environ Res* 66:451–458.
4. Schlegel CE, Lee BG, Luoma SN. 2002. Dietary metals exposure and toxicity to aquatic organisms: Implications for ecological risk assessment. In Newman MC, Roberts MH Jr, Hale RC, eds *Coastal and Estuarine Risk Assessment*. Lewis, Boca Raton, FL, USA, pp 151–188.
5. Wang WX. 2002. Interactions of trace metals and different marine food chains. *Mar Ecol Prog Ser* 243:295–309.
6. Barwick M, Maher W. 2003. Biotransference and biomagnification of selenium, copper, cadmium, zinc, arsenic and lead in a temperate seagrass ecosystem from Lake Macquarie Estuary, NSW, Australia. *Mar Environ Res* 56:471–502.
7. Boyle D, Amlund H, Lundbye AK, Hogstrand C, Buryy NR. 2010. Bioavailability of a natural lead-contaminated invertebrate diet to zebrafish. *Environ Toxicol Chem* 29:708–714.
8. Fisher NS, Hook SE. 2002. Toxicology tests with aquatic animals need to consider the trophic transfer of metals. *Toxicology* 181/182:531–536.
9. Amiard JC, Amiard-Triquet C, Metayer C, Marchand J, Ferre R. 1980. Study on the transfer of Cd, Pb, Cu and Zn in neritic and estuarine trophic chains. I. The inner estuary of the Loire (France) in the summer of 1978. *ater Res* 14:665–673.
10. Mount DR, Barth AK, Garrison TD, Barten KA, Hockett JR. 1994. Dietary and waterborne exposure of rainbow trout (*Oncorhynchus mykiss*) to copper, cadmium, lead and zinc using a live diet. *Environ Toxicol Chem* 13:2031–2041.
11. Rainbow PS. 1996. Biomonitoring of heavy metal availability in the marine environment. *Environ Pollut* 94:244.
12. Ruelas-Inzunza J, Páez-Osuna F. 2008. Trophic distribution of Cd, Pb, and Zn in a food web from Altata-Ensenada del Pabellón subtropical lagoon, SE Gulf of California. *Arch Environ Contam Toxicol* 54:584–596.
13. Soto-Jiménez MF, Arellano-Fiore C, Rocha-Velarde R, Jara-Marini ME, Ruelas-Inzunza J, Páez-Osuna F. 2011. Trophic transfer of lead through a model marine 1 four-level food chain: *Tetraselmis suecica*, *Artemia franciscana*, *Litopenaeus vannamei* and *Haemulon scudderii*. *Arch Environ Contam Toxicol* DOI: 10.007/s00244.0109620.4.
14. Mathews T, Fisher NS. 2008. Evaluating the trophic transfer of cadmium, polonium, and methylmercury in an estuarine food chain. *Environ Toxicol Chem* 27:1093–1101.
15. Mathews T, Fisher NS. 2008. Trophic transfer of seven trace metals in a four-step marine food chain. *Mar Ecol Prog Ser* 367:23–33.
16. Luoma SN. 1996. The developing framework of marine ecotoxicology: Pollutants as a variable in marine ecosystems? *J Exp Mar Biol Ecol* 200:29–55.
17. Cuong DT, Karupiah S, Obbard JP. 2008. Distribution of heavy metals in the dissolved and suspended phase of the sea-surface microlayer, seawater column and in sediments of Singapore's coastal environment. *Environ Monit Assess* 138:255–272.
18. U.S. Environmental Protection Agency. 2009. National recommended water quality criteria. EPA-822-4-02-047. Washington, DC.
19. Council of European Communities. 1976. Dangerous Substances Directive: Council Directive 76/464/EEC of 4 May 1976 on pollution caused by certain dangerous substances discharged into the aquatic environment of the Community. *Official J Eur Commun* 129:23–29.
20. Wang WX, Fisher NS. 1999. Assimilation efficiencies of chemical contaminants in aquatic invertebrates: A synthesis. *Environ Toxicol Chem* 18:2034–2045.
21. Trujillo-Valle ML. 1993. La colección de microalgas del CICESE. Comunicaciones Académicas, Serie Acuicultura. Reporte Técnico CIACT9301. Centro de Investigaciones Científicas y de Educación Superior de Ensenada. Ensenada, Baja California Sur, México, p 38.
22. Nussey G, Van Vuren JHJ, Du Preez HH. 2000. Bioaccumulation of chromium, manganese, nickel and lead in the tissues of the moggel, *Labeo umbratus* (Cyprinidae), from Witbank Dam, Mpumalanga. *Water SA* 26:269–284.
23. Van Stappen G. 1996. Artemia: Use of cysts. In Lavens P, Sorgeloos P, eds, *Manual on the Production and Use of Live Food for Aquaculture Fisheries Technical Paper 361*. Food and Agriculture Organization of the United Nations, Viale delle Terme di Caracalla, Rome, Italy, pp 107–123.
24. Leteux F, Meyer FP. 1972. Mixtures of malachite green and formalin for controlling *Ichthyophthirius multifiliis* and other protozoan parasites of fish. *Prog Fish Culturist* 34:21–26.
25. Farkas A, Salánki J, Specziár A, Varanka I. 2001. Metal pollution as health indicator of lake ecosystems. *Int J Occup Med Environ Health* 14:163–170.
26. Schmitt CJ, Finger SE. 1987. The effects of sample preparation on measured concentrations of eight elements in edible tissues of fish from streams contaminated by lead mining. *Arch Environ Contam Toxicol* 16:185–207.
27. Pérez-Rama M, Alonso JA, Lopez CH, Vaamonde ET. 2002. Cadmium removal by living cells of the marine microalga *Tetraselmis suecica*. *Bioresource Technol* 84:265–270.
28. Levy JL, Stauber JL, Jolley DF. 2007. Sensitivity of marine microalgae to copper: The effect of biotic factors on copper adsorption and toxicity. *Sci Total Environ* 387:141–154.
29. Levy JL, Angel BM, Stauber JL, Poon WL, Simpson SL, Cheng SH, Jolley DF. 2008. Uptake and internalisation of copper by three marine microalgae: Comparison of copper-sensitive and copper-tolerant species. *Aquat Toxicol* 89:82–93.
30. Debelius B, Forja JM, DelValls A, Lubián LM. 2009. Toxicity and bioaccumulation of copper and lead in five marine microalgae. *Ecotoxicol Environ Saf* 72:1503–1513.
31. Muhaemin M. 2004. Toxicity and bioaccumulation of lead in *Chlorella* and *Dunaliella*. *J Coast Dev* 8:27–33.
32. Cordero J, Guevara M, Morales E, Lodeiros C. 2005. Efecto de metales pesados en el crecimiento de la microalga tropical *Tetraselmis chuii* (Prasinophyceae). *Rev Biol Trop* 53:325–330.
33. Satoh A, Vudikaria LQ, Kurano N, Miyachi S. 2005. Evaluation of the sensitivity of marine microalgal strains to the heavy metals, Cu, As, Sb, Pb, and Cd. *Environ Int* 31:713–722.
34. Pérez-Rama M, Torres E, Herrero Cy, Abalde J. 2001. Class III metallothioneins in response to cadmium toxicity in the marine microalga *Tetraselmis suecica* (Kyllin) Butcher. *Environ Toxicol Chem* 20:2061–2066.
35. Rainbow PS. 1997. Ecophysiology of trace metal uptake in crustaceans. *Estuar Coast Shelf Sci* 44:169–175.
36. Spicer JJ, Weber RE. 1991. Respiratory impairment in crustaceans and molluscs due to exposure to heavy metals. *Comp Biochem Physiol C* 100:339–342.
37. Chinni S, Khan RN, Yallapragada PR. 2000. Oxygen consumption, ammonia-N excretion, and metal accumulation in *Penaeus indicus* postlarvae exposed to lead. *Bull Environ Contam Toxicol* 64:144–151.
38. Chinni S, Khan RN, Yallapragada PR. 2002. Acute toxicity of lead on tolerance, oxygen consumption, ammonia-N excretion, and metal accumulation in *Penaeus indicus* postlarvae. *Ecotoxicol Environ Saf* 51:79–84.
39. Fingerhann M, Devi M, Reddy PS, Katayani R. 1996. Impact of heavy metal exposure on the nervous system and endocrine mediated processes in crustaceans. *Zool Stud* 35:1–8.
40. Sorenson E. 1991. *Metal Poisoning in Fish*. CRC, Boca Raton, FL, USA.
41. Rogers JT, Patel M, Gilmour KM, Wood CM. 2005. Mechanisms behind Pb-induced disruptions of Na⁺ and Cl⁻ balance in rainbow trout (*Oncorhynchus mykiss*). *Am J Physiol Reg Integr Comp Physiol* 289:463–472.
42. Alves LC, Wood CM. 2006. The chronic effects of dietary lead in freshwater juvenile rainbow trout (*Oncorhynchus mykiss*) fed elevated calcium diets. *Aquat Toxicol* 78:217–232.
43. Roach AC, Maher W, Krikowa F. 2008. Assessment of metals in fish from Lake Macquarie, New South Wales, Australia. *Arch Environ Contam Toxicol* 54:292–308.



111 RIVER STREET, HOBOKEN, NJ 07030

*****IMMEDIATE RESPONSE REQUIRED*****

Your article will be published online via Wiley's EarlyView® service (www.interscience.wiley.com) shortly after receipt of corrections. EarlyView® is Wiley's online publication of individual articles in full text HTML and/or pdf format before release of the compiled print issue of the journal. Articles posted online in EarlyView® are peer-reviewed, copyedited, author corrected, and fully citable via the article DOI (for further information, visit www.doi.org). EarlyView® means you benefit from the best of two worlds--fast online availability as well as traditional, issue-based archiving.

Please follow these instructions to avoid delay of publication.

☐ **READ PROOFS CAREFULLY**

- This will be your only chance to review these proofs. **Please note that once your corrected article is posted online, it is considered legally published, and cannot be removed from the Web site for further corrections.**
- Please note that the volume and page numbers shown on the proofs are for position only.

☐ **ANSWER ALL QUERIES ON PROOFS** (Queries for you to answer are attached as the last page of your proof.)

- Mark all corrections directly on the proofs. Note that excessive author alterations may ultimately result in delay of publication and extra costs may be charged to you.

☐ **CHECK FIGURES AND TABLES CAREFULLY**

- Check size, numbering, and orientation of figures.
- All images in the PDF are downsampled (reduced to lower resolution and file size) to facilitate Internet delivery. These images will appear at higher resolution and sharpness in the printed article.
- Review figure legends to ensure that they are complete.
- Check all tables. Review layout, title, and footnotes.

RETURN

- ☐ **PROOFS**
☐ **PAGE CHARGE FORM**
☐ **CTA (If you have not already signed one)**

RETURN IMMEDIATELY AS YOUR ARTICLE WILL BE POSTED ONLINE SHORTLY AFTER RECEIPT

QUESTIONS?

Production Editor
E-mail: etcprod@wiley.com

Refer to journal acronym and article production number
(i.e., ETC 00-001 for ETC ms 00-001).

SOCIETY OF ENVIRONMENTAL TOXICOLOGY AND CHEMISTRY – COPYRIGHT TRANSFER AGREEMENT

Please sign and return immediately to the Editorial Office

Environmental Toxicology and Chemistry

C. H. Ward, Editor-in-Chief

Civil and Environmental Engineering, Rice University, MS 316, 6100 Main Street, Houston TX 77005 USA

Email: etcj@rice.edu • Fax: +001 713 348 5948

Manuscript No.: _____ Issue (Editorial Office only): _____

Corresponding Author ("Contributor"): _____

Co-Contributors: _____

Manuscript Title ("Contribution"): _____

for publication in *Environmental Toxicology and Chemistry* (the "Journal") published by Wiley-Blackwell or any successor publisher ("Wiley-Blackwell") on behalf of the Society of Environmental Toxicology and Chemistry ("the Society").

If the Contribution is not accepted for publication, or if the Contribution is subsequently rejected, this Agreement shall be null and void. **Publication cannot proceed without a signed copy of this Agreement.**

- A. Copyright Assignment** The Contributor assigns to the Society, during the full term of copyright and any extensions or renewals, all copyright in and to the Contribution, and all rights therein, including but not limited to the right to publish, republish, transmit, sell, distribute, and otherwise use the Contribution in whole or in part in electronic and print editions of the Journal and in derivative works throughout the world, in all languages and in all media of expression now known or later developed, and to license or permit others to do so.
- B. Citation and Credit** Reproduction, posting, transmission, or other distribution or use of the final Contribution in whole or in part in any medium by the Contributor as permitted by this Agreement requires a citation to the Journal and an appropriate credit to the Society and Wiley-Blackwell as Publisher, suitable in form and content as follows: [Title of Article], [Contributor], *Environmental Toxicology and Chemistry* [Volume/Issue], Copyright © [year] Society of Environmental Toxicology and Chemistry, Wiley-Blackwell Publisher. Links to the final article on Wiley-Blackwell's website are encouraged where appropriate.
- C. Retained Rights by Contributor or Contributor Employer** Notwithstanding the above, the Contributor or, if applicable, the Contributor's Employer, retains all proprietary rights other than copyright, such as patent rights, in any process, procedure, or article of manufacture described in the Contribution.
- D. Permitted Uses of Contribution by Contributor**
- 1. Submitted Version.** The Society licenses back the following rights to the Contributor in the version of the Contribution as originally submitted for publication:
 - a. After the final version is published, the right to self-archive on the Contributor's personal website or in the Contributor's employer's institutional repository or archive on both intranets and the Internet. The Contributor may not update the submitted version or replace it with the published Contribution. The version posted must contain a legend as follows: This is the pre-peer-reviewed version of the following article: FULL CITE, which has been published in final form at [Link to final article].
 - b. The right to transmit, print, and share copies with colleagues.
 - 2. Accepted Version.** Re-use of the accepted and peer-reviewed (but not final) version of the Contribution shall be by separate agreement with Wiley-Blackwell. Requests for permission should be addressed to the permissions department at journalsrights@wiley.com. Wiley-Blackwell has agreements with certain funding agencies governing re-use of the accepted version. For details of those agreements, and other offerings allowing open web use, see <http://www.wiley.com/go/funderstatement>. **NOTE: NIH grantees** should check the box at the end of this document. Pursuant to NIH mandate, Wiley-Blackwell will post the accepted version of Contributions authored by NIH grant-holders to PubMed Central upon acceptance. The accepted version will be made publicly available 12 months after publication. For more information, see www.wiley.com/go/nihmandate.
 - 3. Final Published Version.** The Society hereby licenses back to the Contributor the following rights with respect to

the final published version of the Contribution:

- a. Copies for colleagues. The personal right of the Contributor only to send or transmit individual copies of the final published version in any format to colleagues upon their specific request provided that no fee is charged, and further provided that there is no systematic distribution of the Contribution, e.g., posting on a listserv, website, or automated delivery.
- b. Re-use in other publications. The right to re-use the final Contribution or parts thereof for any publication authored or edited by the Contributor (excluding journal articles) where such re-used material constitutes less than half of the total material in such publication. In such case, any modifications should be accurately noted.
- c. Teaching duties. The right to include the Contribution in teaching or training duties at the Contributor's institution or place of employment including in course packs, e-reserves, presentation at professional conferences, in-house training, or distance learning. The Contribution may not be used in seminars outside of normal teaching obligations (e.g., commercial seminars). Electronic posting of the final published version in connection with teaching or training at the Contributor's institution or place of employment is permitted subject to the implementation of reasonable access control mechanisms, such as user name and password. Posting the final published version on the open Internet is not permitted.
- d. Oral presentations. The right to make oral presentations based on the Contribution.

4. Article Abstracts, Figures, Tables, Data Sets, Artwork, and Selected Text (up to 250 words).

- a. Contributors may re-use unmodified abstracts for any non-commercial purpose. For on-line uses of the abstracts, the Society encourages but does not require linking back to the final published versions.
- b. Contributors may re-use figures, tables, data sets, artwork, and selected text up to 250 words from their Contributions, provided the following conditions are met:
 - i. Full and accurate credit must be given to the Contribution.
 - ii. Modifications to figures, tables, and data must be noted. Otherwise, no changes may be made.
 - iii. Re-use may not be made for direct commercial purposes, or for financial consideration to the Contributor.
 - iv. Nothing herein shall permit dual publication in violation of journal ethical practices.

E. Contributions Owned by Employer

1. If the Contribution was written by the Contributor in the course of the Contributor's employment (as a "work-made-for-hire" in the course of employment), the Contribution is owned by the company or employer, which must sign this Agreement (in addition to the Contributor's signature). In such case, the company or employer hereby assigns to the Society, during the full term of copyright, all copyright in and to the Contribution for the full term of copyright throughout the world as specified in paragraph A above.
2. In addition to the rights specified as retained in paragraph B above and the rights granted back to the Contributor pursuant to paragraph C above, the Society hereby grants back, without charge, to such company or employer, its subsidiaries and divisions, the right to make copies of and distribute the final published Contribution internally in print format or electronically on the Company's intranet. Copies so used may not be resold or distributed externally. However, the company or employer may include information and text from the Contribution as part of an information package included with software or other products offered for sale or license or included in patent applications. Posting of the final published Contribution by the company or employer on a public-access website may be done only with Wiley-Blackwell's written permission and payment of any applicable fees. Also, upon payment of Wiley-Blackwell's reprint fee, the institution may distribute print copies of the published Contribution externally.

F. Government Contracts In the case of a Contribution prepared under U.S. Government contract or grant, the U.S. Government may reproduce, without charge, all or portions of the Contribution and may authorize others to do so, for official U.S. Government purposes only, if the U.S. Government contract or grant so requires.

G. Government Employees

1. U.S. Government Employees: A contribution prepared by a U.S. federal government employee as part of the employee's official duties, or which is an official U.S. Government publication, is called a "U.S. Government work," and is in the public domain in the United States. In such case, the employee may cross out Paragraph A.1 but must sign (in the Contributor's signature line) and return this Agreement. If the Contribution was not

prepared as part of the employee's duties or is not an official U.S. Government publication, it is not a U.S. Government work.

2. U.K. Government Employees: The rights in a Contribution prepared by an employee of a U.K. government department, agency, or other Crown body as part of his or her official duties, or which is an official government publication, belong to the Crown. U.K. government authors should submit a signed declaration form together with this Agreement. The form can be obtained via <http://www.opsi.gov.uk/advice/crown-copyright/copyright-guidance/publication-of-articles-written-by-ministers-and-civil-servants.htm>
3. Non-U.S., Non-U.K. Government Employees: If your status as a government employee legally prevents you from signing this Agreement, please contact the Editorial Office.

- H. Copyright Notice** The Contributor and the company or employer agree that any and all copies of the final published version of the Contribution or any part thereof distributed or posted by them in print or electronic format as permitted herein will include the notice of copyright as stipulated in the Journal and a full citation to the Journal as published by Wiley-Blackwell.
- I. Contributor's Representations** The Contributor represents that the Contribution is the Contributor's original work, all individuals identified as Contributors actually contributed to the Contribution, and all individuals who contributed are included. If the Contribution was prepared jointly, the Contributor agrees to inform the co-Contributors of the terms of this Agreement and to obtain their signature to this Agreement or their written permission to sign on their behalf. The Contribution is submitted only to this Journal and has not been published before. (If excerpts from copyrighted works owned by third parties are included, the Contributor will obtain written permission from the copyright owners for all uses as set forth in Wiley-Blackwell's permissions form or in the Journal's Instructions for Contributors, and show credit to the sources in the Contribution.) The Contributor also warrants that the Contribution contains no libelous or unlawful statements, does not infringe upon the rights (including without limitation the copyright, patent, or trademark rights) or the privacy of others, or contain material or instructions that might cause harm or injury.
- J. Signature** All Contributors must sign below and check the box or boxes that apply. If your Contribution was written during the course of employment, your employer must also sign where indicated. **NOTE:** NIH grantees must also check the NIH grantee box.

<input type="checkbox"/> Contributor-owned work	_____	_____
<input type="checkbox"/> Contributing author authorized to sign for all	Contributor's signature	Date
<input type="checkbox"/> U.S. Government work	_____	_____
<input type="checkbox"/> U.K. Government work (Crown Copyright)	Type or print contributing author name and title	
<input type="checkbox"/> Other Government work	_____	
<input type="checkbox"/> NIH Grantee	Co-contributor's signature	Date
Type or print co-contributor's name and title		

<input type="checkbox"/> Company- or Institution-owned work (made-for-hire in course of employment)	_____	_____
	Company or Institution (Employer-for-Hire)	Date
	Authorized signature of Employer	Date

Environmental Toxicology and Chemistry
Page Charge Form

PLEASE RETURN WITH YOUR PAGE PROOFS TO: John Wiley & Sons, 111 River Street, Hoboken, NJ 07030.
ATTENTION: Jeffrey Collins (etcp@wiley.com). Telephone: (201) 748-8864

Article Number: _____

Authors: _____

Please calculate your page charge based on the information in the table below.

Select one	Author Category	Pages 1-6	Pages 7-12	Pages 13 and beyond
<input type="checkbox"/>	Critical Review Author	FREE	\$50 per page	\$150 per page
<input type="checkbox"/>	SETAC member in good standing	FREE	\$50 per page	\$150 per page
<input type="checkbox"/>	Other	\$50 per page	\$150 per page	\$150 per page

SETAC member in good standing: Membership is current at the time of submission and has been continuous for 2 years prior to submission

Other: Non-member, or membership is not current at the time of submission or has not been continuous for 2 years prior to submission

Total article length _____ pages

Cost for pages 1-6 _____

Additional cost for pages 7-12 _____

Additional cost for pages 13 and beyond _____

TOTAL page charges _____

Form of payment (in \$US drawn from a US Bank; payable to Wiley-Blackwell):

• Visa • MasterCard • Check • Purchase Order No. _____

Card # _____ Expiration Date _____

Signed _____ Date _____

Institution _____

Name _____

Address _____

BILL TO (if your payment does not accompany this form):

Name: _____ Institution _____

Address: _____



Additional reprint and journal issue purchases

Should you wish to purchase additional copies of your article, please click on the link and follow the instructions provided:
<https://caesar.sheridan.com/reprints/redir.php?pub=10089&acro=ETC>

Corresponding authors are invited to inform their co-authors of the reprint options available.

Please note that regardless of the form in which they are acquired, reprints should not be resold, nor further disseminated in electronic form, nor deployed in part or in whole in any marketing, promotional or educational contexts without authorization from Wiley. Permissions requests should be directed to mailto: permissionsus@wiley.com

For information about 'Pay-Per-View and Article Select' click on the following link: <http://wileyonlinelibrary.com/ppv>