

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/224826746>

Photodissociation of heme distal methionine in ferrous cytochrome c revealed by subpicosecond time-resolved resonance Raman spectroscopy

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · NOVEMBER 2004

Impact Factor: 12.11 · DOI: 10.1021/ja046422i

CITATIONS

18

READS

13

5 AUTHORS, INCLUDING:



Simona Cianetti

Novartis

14 PUBLICATIONS 234 CITATIONS

SEE PROFILE



Marten H Vos

École Polytechnique

142 PUBLICATIONS 3,658 CITATIONS

SEE PROFILE



Jean-Louis Martin

Ecole Polytechnique & French Institute of He...

236 PUBLICATIONS 7,292 CITATIONS

SEE PROFILE



Sergei G Kruglik

Pierre and Marie Curie University - Paris 6

99 PUBLICATIONS 919 CITATIONS

SEE PROFILE

Photodissociation of Heme Distal Methionine in Ferrous Cytochrome *c* Revealed by Subpicosecond Time-Resolved Resonance Raman Spectroscopy

Simona Cianetti,^{†,‡} Michel Négrerie,[†] Marten H. Vos,[†] Jean-Louis Martin,[†] and Sergei G. Kruglik^{*,†,§}

Laboratory for Optics and Biosciences, INSERM U451, CNRS UMR 7645, Ecole Polytechnique, 91128 Palaiseau Cedex, France, LENS, University of Florence, Sesto Fiorentino, Italy, and B. I. Stepanov Institute of Physics, National Academy of Sciences, Minsk, Belarus

Received June 16, 2004; E-mail: sergei.kruglik@polytechnique.fr

In heme proteins, binding and release of diatomic gaseous ligands (O_2 , CO, NO) are controlled by the coordination and the redox state of the heme iron.¹ A number of proteins have been identified in which the displacement of a heme-ligated internal amino acid by an external diatomic gaseous ligand plays a physiological role. These include, among others, EcDOS (bacterial oxygen sensor),² CooA (bacterial carbon monoxide sensor),³ and mammalian cytochrome *c* (cyt *c*). Indeed, mitochondrial cyt *c* has recently been shown to bind nitric oxide in apoptotic cells.⁴ Here, since the heme is six-coordinated in the resting state, binding of a diatomic ligand requires the cleavage of one axial bond. The first step in understanding the activation mechanism of these enzymes is to structurally demonstrate the possibility of such a bond breaking and to identify its nature.

Generally, diatomic axial ligands can be photodissociated from the heme iron with high yield, providing a unique possibility to study the ultrafast heme structural dynamics after axial ligand bond breaking by femtosecond spectroscopy.¹ For cyt *c* in the resting state, the heme iron is coordinated by the two *endogenous* ligands, histidine (His-18) and methionine (Met-80). Previous photophysical studies on ferrous cyt *c* were based on femtosecond transient absorption (TA) spectroscopy with high-energy excitation in the Soret band of the heme.^{5,6} These studies suggested the photolysis of either the Fe–His proximal bond⁵ or the Fe–Met distal bond.⁶ The latter conclusion was based on a similarity of the reconstructed photoproduct absorption spectrum with the stationary absorption spectrum of a model five-coordinated histidine-ligated complex, microperoxidase.⁶ Also, femtosecond coherence spectroscopy study of cyt *c*,⁶ performed along with TA, allowed to reconstruct the vibrational modes at ~ 40 , ~ 80 , and ~ 220 cm^{-1} , which were interpreted as Met-80 detachment.

However, TA spectroscopy monitors changes in the electronic system of the heme upon photoexcitation. For cyt *c*, the observed TA kinetics is extremely fast, with the longest time constant of ~ 6 ps interpreted as ligand geminate rebinding.^{5,6} This is on the same time scale as electronic relaxation without ligand detachment, the effect that may complicate the photoproduct TA spectra. Time-resolved resonance Raman (TR³) is known as the most direct structure-sensitive tool in studying the processes of axial ligand(s) binding and release.^{3,7–9} In this study, we applied subpicosecond TR³ (i) to directly assess whether the event of photodissociation of an internal axial ligand in ferrous cyt *c* takes place, (ii) to identify unequivocally which side chain is involved in bond cleavage, and (iii) to monitor heme structural changes during this process.

For this study, we have developed a novel subpicosecond Raman spectrometer, which will be described in detail elsewhere. Briefly,

this device operates at 1 kHz repetition rate and consists of ~ 50 -fs Ti:Sapphire oscillator–amplifier laser source and a spectrometer part based on optical parametric generator–amplifier systems in both pump and probe channels. Sample photolysis was performed by pump pulses with an energy of 2.0–2.5 μJ , centered at 550 nm, the maximum of the Q_{0-0} band, the lowest-lying electronic transition in cyt *c*. Thus, we minimized the effect of heme vibrational heating. Raman scattering was excited by 20–30 nJ probe pulses at 435 nm. This wavelength is near the photoproduct absorption maximum⁶ and is considerably red-shifted with respect to the Soret band of the resting ferrous cyt *c* ($\lambda_{max} = 416$ nm). To achieve appropriate spectral resolution, the probe pulses were tailored by a specially designed narrow-band interference filter. The instrumental time response function was Gaussian with a half-width $T_G = 630 \pm 40$ fs, and the spectral resolution was 30 cm^{-1} (fwhm). Raman spectra were obtained in a 90° light-collection geometry using a 1-m single spectrograph with 1200 grooves/mm grating and a liquid nitrogen-cooled CCD.

Horse heart cyt *c* was purchased from Sigma. The ferrous sample, with concentration of 0.3 mM, was prepared by reducing the protein in 0.1 M Tris/HCl buffer (pH 7.4) using 1 mM sodium dithionite. A cyt *c* N-fragment (residues 1–56)^{10,11} was obtained by limited proteolysis of the parental protein. Samples were placed under anaerobic conditions in a standard 1-cm quartz cuvette with a stirring bar. Ammonium sulfate (0.5 M) was added to the samples for Raman intensity calibration using the 980 cm^{-1} line of SO_4^{2-} . Raman spectra normalization and reconstruction of TR³ difference spectra representing pure photoproduct contribution have been described elsewhere.⁹

Figure 1 shows the results for ferrous cyt *c*. The high-resolution ground-state spectrum (Figure 1A) recorded with cw excitation at 441.6 nm is in agreement with literature data.¹² Apart from the spectral resolution, which illustrates the chosen spectral-temporal compromise, the ground-state spectrum (Figure 1B) recorded with subpicosecond excitation and negative time delay between pump and probe pulses ($\Delta t = -5$ ps) is identical to that in Figure 1A. This finding implies that (i) the probe pulse is weak enough not to induce noticeable photolysis and (ii) no long-lived (millisecond or longer) pump-induced spectral changes take place.

The distinctive feature in the 1-ps spectrum of the cyt *c* photoproduct (Figure 1D) is the appearance of a strong 216 cm^{-1} band, which is assigned to the Fe–His stretching (ν_{Fe-His}).¹³ This band is not Raman-active in six-coordinated cyt *c*,¹² but is specifically enhanced in the case of five-coordinated ferrous high-spin domed heme structures.¹³ Hence, instant rising of ν_{Fe-His} (Figure 1D) readily indicates the photodissociation of the internal axial ligand, namely Met-80. Indeed, although the Fe–Met stretch has not been identified for cyt *c*, it is expected¹² at much higher frequency, around 350 cm^{-1} . The observed Fe–His frequency is

[†] Ecole Polytechnique.

[‡] University of Florence.

[§] B. I. Stepanov Institute of Physics.




Figure 1. TR³ spectra of ferrous cyt *c*. Accumulation time was 50 min for each spectrum. Spectra A and B are ground-state spectra recorded with cw excitation at 441.6 nm and subpicosecond excitation at 435 nm, respectively; spectrum B was recorded at $\Delta t = -5$ ps. Spectra C–F are difference spectra at various Δt : -1 (C), 1 (D), 2 (E), and 15 ps (F).

lower than that of the *b*-type hemes in deoxy Mb^{3,13} or R-state HbA (220 cm⁻¹).⁷ As it is equal to that of T-state HbA (216 cm⁻¹)⁷ and higher than that of photodissociated CoxA (211 cm⁻¹)³, this frequency value is presumably indicative of the distortion, due to strain exerted by the protein, of the proximal His-18 side chain coupled to the domed cyt *c* heme. This effect, however, is less pronounced than in the sensor protein CoxA³.

Other prominent spectral changes in TR³ spectra (Figure 1B versus Figure 1D,E) are (i) the transformation of the unresolved contour in ~ 365 – 430 cm⁻¹ range into two broad bands and (ii) disappearance of the band at 569 cm⁻¹. The former contour contains $\delta(C_\beta-C_\alpha-C_\beta)$ and $\delta(C_\beta-C_\alpha-S)$ bending modes of thioether substituents¹² participating in cyt *c*-specific interaction of the heme with cysteine side chains. The latter band is the out-of-plane pyrrole folding mode γ_{21} , which is activated because of protein-induced heme distortion; this mode is known to disappear when the heme geometry is relaxed.¹² Thus, the observed spectral changes indicate, for the cyt *c* five-coordinated transient heme structure, both an alteration in cysteine coupling and a relaxation of the protein-induced deformations of the heme macrocycle.

Concerning temporal evolution of the photoinduced changes, our TR³ data are in a general accord with the reported TA kinetic measurements.⁶ The instrument-limited rise of the $\nu_{\text{Fe-His}}$ (Figure 1C,D) implies axial ligand photolysis in the femtosecond time scale. The photoproduct spectra change modestly between $\Delta t = 1$ ps and $\Delta t = 2$ ps. The main difference is the red shift of the porphyrin ν_8^* band (ground-state ν_8 -mode¹² frequency is 345 cm⁻¹ (Figure 1A)), from 329 to 333 cm⁻¹ (Figure 1D,E). This shift can be associated with the heme cooling during the first few picoseconds, as deduced from TR³ spectra in the high-frequency region of porphyrin marker bands (data not shown), where the effect is most pronounced. The reported recombination time of ~ 6 ps^{5,6} is fully supported here as the photoproduct TR³ spectrum is almost




Figure 2. Same as in Figure 1 but for ferrous cyt *c* N-fragment (residues 1–56), in buffer pH 9. Spectrum C is a difference spectrum at $\Delta t = 2$ ps.

completely vanished at $\Delta t = 15$ ps (Figure 1F). Unfortunately, a quantitative Raman kinetic analysis is complicated by the changes in Raman excitation profiles due to spectral evolution of the photoproduct absorption bands during the first few picoseconds.¹⁴

Finally, to identify unequivocally the side chain that is photo-dissociated in native cyt *c*, a specially prepared¹⁰ cyt *c* N-fragment has been studied (Figure 2). The ground-state spectrum (Figure 2A) recorded with cw excitation at 441.6 nm is in accord with the results of a recent study,¹¹ which demonstrates that, at pH 9, cyt *c* N-fragment exists predominantly as a six-coordinated bis-histidine complex, but with a small proportion of a five-coordinated mono-histidine form. Thus, the ground-state N-fragment heme structure differs from that of the native protein, as reflected in the marked steady-state spectral differences (Figure 2A,B versus Figure 1A,B). By contrast, the photoproduct spectra, originated exclusively from the five-coordinated hemes, are almost identical (Figure 2C and Figure 1E). Since, for N-fragment, the five-coordinated species can only be a heme–His complex, the same structural state is obtained for native cyt *c*, thus confirming the photodissociation of Met-80 distal side chain.

Acknowledgment. This work was supported by a Marie Curie Fellowship to S.C. We thank J.-C. Lambry for technical assistance with the setup and S. Santucci for providing the cyt *c* fragment.

References

- (1) Martin, J.-L.; Vos, M. H. *Methods Enzymol.* **1994**, *232*, 416–429.
- (2) Delgado-Nixon, V. M.; Gonzalez, G.; Gilles-Gonzalez, M. A. *Biochemistry* **2000**, *39*, 2685–2691.
- (3) Uchida, T.; Ishikawa, H.; Ishimori, K.; Morishima, I.; Nakajima, H.; Aono; Mizutani, Y.; Kitagawa, T. *Biochemistry* **2000**, *39*, 12747–12752.
- (4) Schonhoff, C. M.; Gaston, B.; Mannick, J. J. *Biol. Chem.* **2003**, *278*, 18265–18270.
- (5) Jongeward, K. A.; Magde, D.; Taube, D. J.; Traylor, T. G. *J. Biol. Chem.* **1988**, *263*, 6027–6030.
- (6) Wang, W.; Ye, X.; Demidov, A. A.; Rosca, F.; Sjödin, T.; Cao, W.; Sheeran, M.; Champion, P. M. *J. Phys. Chem B* **2000**, *104*, 10789–10801.
- (7) Friedman, J. M. *Science* **1985**, *228*, 1273–1280.
- (8) Franzen, S.; Bohn, B.; Poyart, C.; Martin, J.-L. *Biochemistry* **1995**, *34*, 1224–1237.
- (9) Kruglik, S. G.; Mojzes, P.; Mizutani, Y.; Kitagawa, T.; Turpin P.-Y. *J. Phys. Chem. B* **2001**, *105*, 5018–5031.
- (10) Santucci, R.; Fiorucci, L.; Sinibaldi, F.; Polizio, F.; Desideri, A.; Ascoli, F. *Arch. Biochem. Biophys.* **2000**, *379*, 331–336.
- (11) Santoni, E.; Scatragli, S.; Sinibaldi, F.; Fiorucci, L.; Santucci, R.; Smulevich, G. *J. Inorg. Biochem.* **2004**, *98*, 1067–1077.
- (12) Hu, S.; Morris, I. K.; Singh, J. P.; Smith, K. M.; Spiro, T. G. *J. Am. Chem. Soc.* **1993**, *115*, 12446–12458.
- (13) Kitagawa, T. In *Biological Applications of Raman Spectroscopy*; Spiro, T. G., Ed.; Wiley: New York, 1988; Volume 3, Chapter 3.
- (14) Ye, X.; Demidov, A.; Rosca, F.; Wang, W.; Kumar, A.; Ionascu, D.; Zhu, L.; Barrick, D.; Wharton, D.; Champion, P. *J. Phys. Chem A* **2003**, *107*, 8156–8165.

JA046442I