

See discussions, stats, and author profiles for this publication at:  
<https://www.researchgate.net/publication/265143173>

# Phytochemical profile of *Paulownia tomentosa* (Thunb). Steud.

ARTICLE *in* PHYTOCHEMISTRY REVIEWS · AUGUST 2014

Impact Factor: 2.41 · DOI: 10.1007/s11101-014-9376-y

---

READS

68

2 AUTHORS, INCLUDING:



Šmejkal Karel

University of Veterinary and Pharmace...

46 PUBLICATIONS 375 CITATIONS

SEE PROFILE

# Phytochemical profile of *Paulownia tomentosa* (Thunb.) Steud.

Kristýna Schneiderová · Karel Šmejkal



Received: 25 April 2014 / Accepted: 2 August 2014  
© Springer Science+Business Media Dordrecht 2014

**Abstract** *Paulownia tomentosa*, a member of the plant family Paulowniaceae and a rich source of biologically active secondary metabolites, is traditionally used in Chinese herbal medicine. Flavonoids, lignans, phenolic glycosides, quinones, terpenoids, glycerides, phenolic acids, and miscellaneous other compounds have been isolated from different parts of *P. tomentosa* plant. Recent interest in this species has focused on isolating and identifying of prenylated flavonoids, that exhibit potent antioxidant, antibacterial, and antiphlogistic activities and inhibit severe acute respiratory syndrome coronavirus papain-like protease. They show cytotoxic activity against various human cancer cell lines and inhibit the effects of human cholinesterase, butyrylcholinesterase, and bacterial neuraminidases. Most of the compounds considered here have never been isolated from any other species of plant. This review summarizes the information about the isolated compounds that are active, their bioactivities, and the structure–activity relationships that have been worked out for them.

**Keywords** *Bignonia tomentosa* · Flavonoid · Lignan · *Paulownia tomentosa* · Paulowniaceae · Phenolic glycosides

## Introduction

The plant *Paulownia tomentosa* (Thunb.) Siebold & Zucc. ex Steud. is a very adaptable and extremely fast-growing timber tree native to central and western China traditionally used in Chinese Medicine. This deciduous tree is now grown in many areas worldwide, mostly as a decorative ornamental tree (Erbar and Gülden 2011; Zhu et al. 1986).

Two related varieties of *P. tomentosa* have been described—var. *tsinlingensis* has a round to shallowly cordate leaf base and a glabrous or sparsely hairy lower leaf surface, whereas var. *tomentosa* is characterized by a cordate leaf blade base and an abaxial surface that is densely hairy when mature (Hong et al. 1998).

*Paulownia* was named *paulownia* in honour of Anna Paulowna (1795–1865), queen consort of The Netherlands and a daughter of Tsar Paul I of Russia. Nowadays, it is commonly known under its synonym *Bignonia tomentosa* (Zhu et al. 1986) or as the princess-tree, empress-tree, foxglove tree, royal paulownia, kiri, or mao pao tong (yuan bian zhong) (Erbar and Gülden 2011; Hong et al. 1998).

The genus *Paulownia* was first assigned to family Bignoniaceae by Swiss botanist Thunberg (1781). It was then transferred to the Scrophulariaceae family by the Dutch scholars Zuccarini and Siebold (1835) (Zhu et al. 1986). At last, Paulownia has been categorized as a family of its own, Paulowniaceae, based on data from the latest molecular phylogenetic studies. This

K. Schneiderová · K. Šmejkal (✉)  
Department of Natural Drugs, Faculty of Pharmacy,  
University of Veterinary and Pharmaceutical Sciences  
Brno, Palackého 1/3, 612 42 Brno, Czech Republic  
e-mail: karel.mejkal@post.cz

**Table 1** Non-prenylated flavonoid aglycones isolated from *P. tomentosa*




Name of compound and ID	Isolation source	Biological activity	Chemical structure				
Matteucinol (syn. 4'-O-methylfarrerol) ( <b>1</b> ) (2S)	Leaves (Zhu et al. 1986)	Weak cytotoxic activity in vitro against human leukaemia (HL-60) and human hepatoma (SMMC-7721) cell lines (Zhao et al. 2012); no $\alpha$ -glucosidase inhibitory effect (Zhao et al. 2009); weak aldose-reductase inhibitory activity (Kadota et al. 1994); moderate inhibitory effect on human immunodeficiency virus-1 protease (Lee et al. 2008); no activity against lipopolysaccharide (LPS)-induced NO production in RAW 264.7 macrophages (Li et al. 2011)					
Apigenin ( <b>2</b> )	Leaves (Zhao et al. 2012), flowers (Chen et al. 2009; Jiang et al. 2004)	Antioxidant (Prince Vijaya Singh et al. 2004); antibacterial, anti-inflammatory, antispasmodic, antidiarrhoic, antiproliferative, vasorelaxant (Jiang et al. 2004); neuroprotective (Losi et al. 2004); cardioprotective (Psotová et al. 2004); chemopreventive activity against skin cancer (Chen et al. 2009); activity reviewed by Shukla and Gupta (2009) and Patel et al. (2007)					
Kaempferol ( <b>3</b> )	Leaves (Si et al. 2008a)	Cytotoxic, pro-apoptotic (suppresses cell metastasis via inhibition of the ERK-p38-JNK and AP-1 signaling pathways in U-2 OS human osteosarcoma cells) (Chen et al. 2013); antioxidant (e.g., attenuates bladder hyperactivity caused by potassium chloride after protamine sulphate-induced bladder injury) (Huang et al. 2014); impact on human health and cancer chemoprevention reviewed by Chen and Chen (2013), and Calderon-Montano et al. (2011)					

Table 1 continued

		R1	R2	R3	R4	R5
Luteolin (4)	Leaves (Si et al. 2008a; Zhang and Li 2011)	H	OH	OH	H	OH
Quercetin (5)	Bark (Si et al. 2011b), leaves (Si et al. 2008a)	OH	H	OH	OH	OH
7,3'- Dimethylquercetin (syn. rhamnazin) (6)	Leaves, green immature fruit (Wollenweber et al. 2008)	OH	OMe	OH	H	OMe

Memory-improving (Yoo et al. 2013); inhibition of  $\alpha$ -amylase activity (Funke and Melzig 2005); cytotoxic (BGC-823 gastric carcinoma xenografts in nude mice) (Lu et al. 2013); vascular protective (Si et al. 2013); biological activity reviewed by Lopez-Lazaro (2009)

Antifibrotic (Yoon et al. 2012); antioxidant, antiproliferative, anti-inflammatory, cardioprotective (Chen et al. 2013); neuroprotective (Ghosh et al. 2013); anti-diabetic [dose-dependent inhibition of both  $\text{Na}^+/\text{K}^+$  ATPase and sodium hydrogen exchanger in type 2 diabetic erythrocytes (Mishra and Rizvi 2012)]; inhibition of PI3K (Koch et al. 2013); antiviral [anti HCV (Khachatoorian et al. 2012); HCMV (Cotin et al. 2012)]; no anti-HIV activity in vitro tested on H9 cells in the absence of toxicity (Tang et al. 1994); review of bioactivities by Russo et al. (2012)

Antimicrobial activity, poor antifungal effect (Omosa et al. 2014); low antimicrobial activity, low toxicity against human lymphocytes and monocytes, antioxidant/anti-inflammatory activity (Martini et al. 2004; Pelzer et al. 1998); low affinity to acetylcholinesterase (Remya et al. 2012); low inhibitory effect on NO production in RAW264.7 cells (Sudalai et al. 2013); no activity against HSV-1, low toxicity against Vero cells (Tian et al. 2009); no antioxidant activity (Takamatsu et al. 2003); no trypanocidal activity (Grael et al. 2000); activity against lipid peroxidation in rat liver microsomes (Yun et al. 2000); cytotoxicity against TK-10, MCF-7 and UACC-62 cells (Lopez-Lazaro et al. 2000); no inhibition of glycolysis in various tumor cells (Suolinna et al. 1975)

Table 1 continued


	R1	R2	R3	R4	R5
7,3',4'-Trimethylquercetin (7)	OH	OMe	OMe	H	OMe
Thrombin inhibition (Shi et al. 2012); inhibition of IL-4 synthesis in basophils (Hirano et al. 2006); weak trypanocidal activity (Jordao et al. 2004); weak inhibition of NO production in LPS-activated mouse peritoneal macrophages (Matsuda et al. 2003); weak inhibition of degranulation of RBL-2H3 cells (Mastuda et al. 2002); inhibition of Pgp activity (Scambia et al. 1994); no inhibition of glycolysis in a variety of tumor cells (Szuolinná et al. 1975)					
7,3',4'-Trimethylmyricetin (8)	OH	OMe	OMe	OH	OMe
Weak inhibition of NO production in LPS-activated mouse peritoneal macrophages (Matsuda et al. 2003); weak inhibition of degranulation of RBL-2H3 cells (Mastuda et al. 2002)					
7,3',4',5'-Tetramethylmyricetin (9)	OH	OMe	OMe	OMe	OMe
					
	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
(+)-Catechin (10) (2R, 3S)	OH	H	OH	OH	OH
Leaves (Si et al. 2008a) Antiviral (HCV) (Khachatoorian et al. 2012); antimicrobial (Mankovskaia et al. 2013); antioxidant, e.g., adjunctive therapy for the treatment of Parkinson's disease (Teixeira et al. 2013); osteogenic (Wei et al. 2011); anti-proliferative, anti-inflammatory, antifibrotic (GRX cells) (Bragança de Moraes et al. 2012); activity reviewed, for example, by Bansal et al. (2013)					
Dihydrotricin (11) (2S)	H	OMe	OH	OMe	OH
Fruits (Šmejkal et al. 2007a) Antioxidant, vasorelaxant, low anti-platelet aggregation activity (Chang et al. 2010)					

Table 1 continued

		R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
(–)-Epicatechin (12) (2R, 3R)	Leaves (Si et al. 2008a)	OH	H	OH	OH	OH
Homoeriodictyol (syn. hesperetin) (13) (2S)	Leaves (Zhang and Li 2011)	H	H	OH	OMe	OH
Naringenin (14) (2S)	Bark (Si et al. 2011b), leaves (Si et al. 2008a)	H	H	OH	H	OH
Taxifolin (15) (2R, 3R)	Leaves (Si et al. 2008a)	OH	OH	OH	H	OH


Antifungal (Betts et al. 2013); improves muscle mitochondrial structure (Taub et al. 2012); antioxidant, cardioprotective (Paneerselvam et al. 2010); cytotoxic (HT29 human colon adenocarcinoma cells) (Sánchez-Tena et al. 2013); anti-inflammatory (Fu et al. 2013); activity reviewed, for example, by Fraga and Oteiza (2011)


Major flavonoid of citrus, activity reviewed by Khan and Zill-E-Huma (2014)

Anti-diabetic (inhibition of PI3K) (Koch et al. 2013); antiviral [anti HCV (Khachatoorian et al. 2012), HCMV (Cotin et al. 2012)]; antimicrobial (de Aguiar et al. 2013); anti-inflammatory (Jayaraman et al. 2012); antioxidant (Wang et al. 2012); cytotoxic (ER $\alpha$ -transfected HeLA cells, ER $\alpha$ -containing HepG2 cells) (Bulzomi et al. 2010); no antiparasitic activity against *Leishmania* spp. (*major*, *chagasi*, *braziliensis*, *amazonensis*), *Trypanosoma cruzi* (Grecco Sdos et al. 2012); major flavonoid of citrus, activity reviewed by Khan and Zill-E-Huma (2014)

Inhibition of  $\beta$ -amyloid aggregation (Sato et al. 2013); potential for atopic dermatitis treatment (blocks hERG K<sup>+</sup> channels) (Yun et al. 2013); cytotoxic, pro-apoptotic (DU145 human prostate carcinoma cells) (Zhang and Coultas 2013); hypotensive (ACEI), anti-inflammatory, antioxidant (diminished 5-lipoxygenase and NADPH oxidase activity) (Arutyunyan et al. 2012); activity reviewed by Weidmann (2012)

Table 1 continued

	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
5,4'-Dihydroxy-7,3'-dimethoxyflavanone (16) (2S)	H	H	OH	OMe	OMe
Flowers (Chen et al. 2009; Jiang et al. 2004; Kim et al. 2010a, b)	Inactive against glutamate-induced neurotoxicity studied in primary-cultured rat cortical cells (Kim et al. 2010a, b); no trypanocidal activity (Grael et al. 2000)				
5-Hydroxy-7,3',4'-trimethoxyflavanone (17) (2S)	H	H	OMe	OMe	OMe
Flowers (Kim et al. 2010a, b)	Inactive against glutamate-induced neurotoxicity studied in primary-cultured rat cortical cells (Kim et al. 2010a, b); inhibition of inflammation by induction of synovial apoptosis of fibroblast-like synoviocytes through caspase 3 activation in rats with adjuvant arthritis (Li et al. 2010, 2013); inhibition of JAK2-STAT3 signal pathway in rats (Li et al. 2012); inhibition of phosphodiesterase 4 (Yang et al. 2011); low toxicity on B16F10 and SK-MEL-1 melanoma cells (Rodriguez et al. 2002); antimutagenic activity (Miyazawa et al. 2000)				
7-Caffeoyl-acacetin (syn. 7-caffeoyl-4'-methoxyapigenin) (18)					
Stem bark (Si et al. 2009)					



**Fig. 1** Schematic of the pathways of the biosynthesis of geranylated flavonoids

family includes only one genus and between six and ten species. It was originally introduced in 1949 by the research of Nakai (Erbar and Glden 2011).

### Biological activity and traditional uses of *P. tomentosa* extracts


According to both legends and records, people were already using *Paulownia* for various purposes about 2,600 years ago. Chinese herbal medicine has used *P. tomentosa* traditionally to relieve bronchitis, especially by reducing coughing, asthma, and phlegm (Zhu et al. 1986). It has also been used to treat conjunctivitis, dysentery, enteritis, erysipelas, gonorrhea, hemorrhoids, parotitis, traumatic bleeding, and tonsillitis (Jiang et al. 2004; Si et al. 2009). Solutions prepared from the leaves and fruit extracted in water have been used in daily applications to promote the healthy growth of hair and turn grey hair dark. An extract prepared from the wood and leaves may relieve swollen feet. Pharmacological experiments have shown that extracts from the fruit can reduce blood pressure. Nowadays, injections and tablets prepared from *Paulownia* flowers and fruit are used for the

herbal treatment of chronic bronchitis and other kinds of inflammation (Zhu et al. 1986).

More than 130 physiologically active constituents have been isolated from different parts of the *Paulownia* plant. Their biological activity has been tested using both the isolated compounds and different types of extracts. For example, *n*-butanol, EtOAc, and MeOH extracts obtained from the fruit have displayed antiradical activity in anti-DPPH and peroxynitrite assays, due to mainly the presence of flavonoids and phenolic glycosides, but not of all compounds present in these extracts have been identified) (Šmejkal et al. 2007b). An MeOH extract obtained from the fruit inhibited hAChE ( $IC_{50} = 1.44$  mg/ml) and BChE ( $IC_{50} = 0.97$  mg/ml) significantly more strongly than  $CHCl_3$  and water extracts (possibly due to the content of phenolic glycosides and C-prenylated dihydroflavonols and flavanones) (Cho et al. 2012). Significant concentration-dependent anti-inflammatory properties of EtOH extracts of the bark of the tree have also been observed recently using a lipopolysaccharide (LPS)-induced nitric oxide production inhibition model in the murine macrophages cell line RAW264.7 (Si et al. 2011a). Kim et al. (2010a, b) showed potential of aqueous extract of *P. tomentosa* to suppress glutamate



**Table 2** C-prenylated flavonoids isolated from *P. tomentosa*

Name of compound and ID	Isolation	Biological activity	Chemical structure
			
			R
6-Isopentenyl-3'-O-methyltaxifolin ( <b>19</b> ) (2 <i>R</i> , 3 <i>R</i> )	Fruit (Šmejkal et al. 2007a)		H
3,4',5,5',7-Pentahydroxy-3'-methoxy-6-(3-methyl-2-butenyl) flavanone ( <b>20</b> ) (2 <i>R</i> , 3 <i>R</i> )	Fruit (Asai et al. 2008)	Antioxidant (Asai et al. 2008); cytotoxic (A2780 human ovarian cancer cell line) (Murphy et al. 2005, 2006)	OH

induced toxicity in primary cultured rat cortical cells (with possible sesquiterpene lactone as active substance). The bio-activities of individual compounds and, where possible, the structure–activity relationships are in Tables 1, 2, 3 and 4 and discussed in separate chapters.

### *P. tomentosa* flavonoids and their biological activity

Flavonoids represent the most numerous group of secondary metabolites isolated from *P. tomentosa*. They can be divided into simple non-prenylated flavonoids **1–18** (Table 1), C-prenylated and C-geranylated flavonoids **19–59** (Tables 2, 3, respectively), and flavonoid glycosides **60–65**. Compounds **16**, **17**, **19**, **20**, **25–31**, **33–36**, **38–59** have never been isolated from any other species. Most of the isolated flavanones are characterized by a 2*S* configuration in contrast to the dihydroflavonols, for which 2*R*, 3*R* isomer is often observed.

Some of the flavonoid compounds found in *P. tomentosa* have been categorized as dietary flavonoids. Consuming these compounds is believed to deliver health benefits. The activities of these flavonoids are frequently been reviewed, for example in the papers of Romano et al. (2013), Marzocchella et al. (2011), Ross and Kasum (2002), and Havsteen (2002). Some simple flavonoid substances isolated from fruit of *P. tomentosa* are commonly observed as the lipophilic components of different plant exudates. Flavonoids **1–18** show broad-spectrum pharmacological activities

(Table 1). Many papers report their potential role as anticancer compounds for use against human cervical and breast (Bulzomi et al. 2010, 2012), hepatoma, leukemic (Zhao et al. 2012), osteosarcoma (Chen et al. 2013), gastric carcinoma (Lu et al. 2013), colon adenocarcinoma (Sánchez-Tena et al. 2013), or prostatic (Zhang and Coultas 2013) cancer cell lines. Their antioxidant properties could explain their promising cardioprotective (Paneerselvam et al. 2010; Psotová et al. 2004) and neuroprotective effects (Losi et al. 2004). Antimicrobial (Betts et al. 2013; Jiang et al. 2004; Mankovskaia et al. 2013) and antiviral (Khachatoorian et al. 2012) bio-activities against different pathogens have also been discovered. In some cases, no specific biological activity has been observed (Kim et al. 2010a, 2010b; Tang et al. 1994).

*Paulownia tomentosa* is also a rich source of prenylated and geranylated flavonoids (**19–59**), whose occurrence is limited to relatively a few plant families (Yazaki et al. 2009). There are sometimes misunderstandings in the naming of these compounds. A compound containing both a phenolic skeleton and a terpenoid side chain is often designated as a “prenylated” phenolic substance, even though it contains not a prenyl, but rather a geranyl or other type of terpenoid side-chain. Prenylated flavonoids are biosynthesized by a combination of several pathways: the acetate, shikimate, and mevalonate (Fig. 1). It is now known that the prenyl and geranyl moieties are biosynthesized via mevalonate or deoxyxylulose pathway. The connection of terpenoid and flavonoid biosynthetic

**Table 3** C-geranylated flavonoids isolated from *P. tomentosa*


Name of compound and ID	Isolation	Biological activity	Chemical structure				
Mimulone (syn. 6-geranylnaringenin, bonannione A) ( <b>21</b> ) (2S)	Flowers (Chen et al. 2009; Jiang et al. 2004; Kim et al. 2010a, b), fruit (Asai et al. 2008; Cho et al. 2012, 2013; Lee et al. 2014; Šmejkal et al. 2007a, 2008b)	Antioxidant (Šmejkal et al. 2007b; Zima et al. 2010), antioxidant and anti-inflammatory properties in vitro in S100B-induced human monocytes (Lin et al. 2011), anti-inflammatory activity in LPS-activated THP-1 cells (Peluso et al. 2010); antimicrobial against seven Gram-positive bacteria, no antimicrobial activity against three types of Gram-negative bacteria or yeast (Šmejkal et al. 2008b), antimicrobial against <i>S. aureus</i> and various methicillin resistant strains of <i>S. aureus</i> (MRSA 287, MRSA 4211, MRSA, 6975, MRSA 630, MRSA 62059) (Navrátilová et al. 2013); cytotoxic (epithelioid cell line WB344 (Šmejkal et al. 2007b), human erythro-leukaemia cell line K562 (Šmejkal et al. 2008a), human ovarian cancer cell line A2780 (Murphy et al. 2005, 2006; Yoder et al. 2007), KB, KB-VIN, A549, DU145 (Rosselli et al. 2011), low activity against T-lymphoblastic leukaemia CEM, multiple myeloma RPMI 8226 and U266 cell lines, BJ human fibroblast cell line (Šmejkal et al. 2010)]; SARS-CoV PLpro activity (Cho et al. 2013); hAChE, BChE inhibitor (Cho et al. 2012); significant inhibitory effects against <i>Cp</i> -NanI (a sialidase from <i>C. perfringens</i> ) (Lee et al. 2014); weak estrogenic, no progestogenic and androgenic activity (Milligan et al. 2000); inactive against glutamate-induced neurotoxicity studied in primary-cultured rat cortical cells (Kim et al. 2010a, b)		R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>
			H	H	OH	H	

Table 3 continued

		R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>
Diplacone (syn. propolin C, nymphaeol A) (22) (25)	Flowers (Chen et al. 2009; Jiang et al. 2004; Kim et al. 2010a, b), fruit (Asai et al. 2008; Cho et al. 2012, 2013; Lee et al. 2014; Šmejkal et al. 2007a, 2008b)	H	OH	OH	H
	<p>Antioxidant (Kumazawa et al. 2007; Asai et al. 2008; Šmejkal et al. 2007b; Zima et al. 2010; Trusheva et al. 2011), prooxidant and anti-inflammatory (Hošek et al. 2013); antihelvetic (Kobayashi et al. 2008); antimicrobial against seven Gram-positive bacteria, no antimicrobial activity against three types of Gram-negative bacteria or yeast (Šmejkal et al. 2008b), antimicrobial against <i>S. aureus</i>, various methicillin resistant strains of <i>S. aureus</i> (MRSA 287, MRSA 4211, MRSA, 6975, MRSA 630, MRSA 62059)—most potent together with 31 (Navrátilová et al. 2013) and methicillin-susceptible <i>S. aureus</i> strain. Anti-MRSA, inactive against <i>P. aeruginosa</i> (Raghukumar et al. 2010); antiproliferative (TNF-<math>\alpha</math>, similar to indomethacin) (Hošek et al. 2010), but no anti-inflammatory effect was observed in COX-2 inhibition assay (Phommart et al. 2005); cytotoxic (epithelioid cell line WB344 (Šmejkal et al. 2007a), T-lymphoblastic leukaemia CEM, multiple myeloma RPMI 8226 and U266 cell lines, BJ human fibroblast cell line (Šmejkal et al. 2010), human erythro-leukaemia cell line K562 (Šmejkal et al. 2008a), KB human oral carcinoma, human breast cancer, human small cell lung cancer NCI-H187 (Phommart et al. 2005), induction of apoptosis via caspase activation, bid and cytochrome c release in human melanoma cells (Chen et al. 2004), human ovarian cancer cell line A2780 (Murphy et al. 2005, 2006; Yoder et al. 2007), PC-3, DU-145, Hep-3B cell line (Chen et al. 2012); hAChE, BChE inhibitor (Cho et al. 2012), SARS-CoV PL<sub>pro</sub> activity (Cho et al. 2013); significant inhibitory effects against <i>Cp-NanI</i> (a sialidase from <i>C. perfringens</i>) (Lee et al. 2014); selective docking to <i>Leishmania major</i> N-myristoyltransferase (Ogunbe et al. 2014); antileishmanial activity (<i>L. donovani</i> amastigotes) (Salem et al. 2011); inactive against glutamate-induced neurotoxicity studied in primary-cultured rat cortical cells (Kim et al. 2010a, b)</p>				

Table 3 continued

		R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>
Diplacol ( <b>23</b> ) (2R, 3R)	Fruit (Asai et al. 2008), flowers, leaves (Kobayashi et al. 2008)	OH	OH	OH	H
3'-O-methyldiplacone ( <b>24</b> )	Fruit (Asai et al. 2008; Cho et al. 2012, 2013; Lee et al. 2014; Šmejkal et al. 2007a, 2008b), flowers and leaves (Kobayashi et al. 2008)	H	OMe	OH	H
3'-O-methyldiplacol (syn. diplacol 3'-O-methylether) ( <b>25</b> ) (2R, 3R)	Flowers (Kobayashi et al. 2008), fruit (Asai et al. 2008; Cho et al. 2012, 2013; Šmejkal et al. 2008b)	OH	OMe	OH	H

Antioxidant (Asai et al. 2008); cytotoxic (human ovarian cancer cell line A2780) (Murphy et al. 2006; Yoder et al. 2007), low activity against T-lymphoblastic leukaemia CEM, multiple myeloma RPMI 8226 and U266 cell lines, BJ human fibroblast cell line (Šmejkal et al. 2010)

Antioxidant (Šmejkal et al. 2007a; Yazaki et al. 2009); SARS-CoV PLpro activity (Cho et al. 2013); weak hAChE, BChE inhibitor (Cho et al. 2012); significant inhibitory effects against *Cp-NanI* (a sialidase from *C. perfringens*) (Lee et al. 2014); strong docking energy to *L. major* pteridine reductase 1, selective docking to glycerol-3-phosphate dehydrogenase (Ogungbe et al. 2014), antileishmanial activity (*L. donovani* amastigotes) (Salem et al. 2011); activity against lymphoblastic leukaemia CEM, multiple myeloma RPMI 8226 and U266 cell lines, BJ human fibroblast cell line (Šmejkal et al. 2010), human erythro-leukaemia cell line K562 (Šmejkal et al. 2008a); weak antimicrobial against seven Gram-positive bacteria, no antimicrobial activity against three types of Gram-negative bacteria or yeast (Šmejkal et al. 2008b), weak antimicrobial activity against *S. aureus*, various methicillin resistant strains of *S. aureus* (MRSA 287, MRSA 4211, MRSA, 6975, MRSA 630, MRSA 62059) (Navrátilová et al. 2013)

Antioxidant (Asai et al. 2008; Yazaki et al. 2009), antimicrobial against seven Gram-positive bacteria, no antimicrobial activity against three types of Gram-negative bacteria or yeast (Šmejkal et al. 2008b), antimicrobial against *S. aureus* and various methicillin resistant strains of *S. aureus* (MRSA 287, MRSA 4211, MRSA, 6975, MRSA 630, MRSA 62059)—most potent together with 27 (Navrátilová et al. 2013); cytotoxic (human erythro-leukaemia cell line K562) (Šmejkal et al. 2008a), low activity against T-lymphoblastic leukaemia CEM, multiple myeloma RPMI 8226 and U266 cell lines, BJ human fibroblast cell line (Šmejkal et al. 2010); SARS-CoV PLpro activity (Cho et al. 2013); hAChE, BChE inhibitor (Cho et al. 2012); antileishmanial activity (*L. donovani* amastigotes) (Salem et al. 2011), selective docking to *L. major* methionyl tRNA synthetase (Ogungbe et al. 2014)

Table 3 continued

			R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>
3'-O-methyl-5'-O-methyldiplacone (syn. 6-geranyl-4',5,7-trihydroxy-3',5'-dimethoxyflavanone) ( <b>26</b> ) (2S)	Fruit (Asai et al. 2008; Cho et al. 2012, 2013; Šmejkal et al. 2008b)	Antioxidant (Yazaki et al. 2009), antimicrobial against seven Gram-positive bacteria, no antimicrobial activity against three types of Gram-negative bacteria or yeast (Šmejkal et al. 2008b), antimicrobial against <i>S. aureus</i> and various methicillin resistant strains of <i>S. aureus</i> (MRSA 287, MRSA 4211, MRSA, 6975, MRSA 630, MRSA 62059) (Navrátilová et al. 2013); cytotoxic (multiple myeloma RPMI 8226 and U266 cell lines, human cervical cancer cells HeLa, BJ human fibroblast cell line (Šmejkal et al. 2010), human erythro-leukaemia cell line K562) (Šmejkal et al. 2008a); SARS-CoV PLpro activity (Cho et al. 2013); hAChE, BChE inhibitor (Cho et al. 2012)	H	OMe	OH	OMe
3'-O-methyl-5'-hydroxydiplocone (syn. 6-geranyl-4',5,5',7-tetrahydroxy-3'-methoxyflavanone) ( <b>27</b> ) (2S)	Immature fruit (Asai et al. 2008), fruit (Šmejkal et al. 2008b), leaves (Kobayashi et al. 2008)	Antioxidant (Asai et al. 2008; Yazaki et al. 2009; Zima et al. 2010); antimicrobial activity against seven Gram-positive bacteria, no antimicrobial activity against three types of Gram-negative bacteria or yeast (Šmejkal et al. 2008b), antimicrobial against <i>S. aureus</i> and various methicillin resistant strains of <i>S. aureus</i> (MRSA 287, MRSA 4211, MRSA, 6975, MRSA 630, MRSA 62059) (Navrátilová et al. 2013); cytotoxic (human erythro-leukemia cell line K562) (Šmejkal et al. 2008a), activity against T-lymphoblastic leukaemia CEM, multiple myeloma RPMI 8226 and U266 cell lines, BJ human fibroblast cell line (Šmejkal et al. 2010)	H	OMe	OH	OH
3'-O-methyl-5'-methoxydiploacol ( <b>28</b> ) (racemic mixture of 2R, 3R and 2S, 3S)	Fruit (Šmejkal et al. 2007a)		OH	OMe	OH	OMe
6-Geranyl-5,7-dihydroxy-3',4'-dimethoxyflavanone ( <b>29</b> ) (2S)	Immature fruit (Asai et al. 2008)	Antioxidant (Asai et al. 2008)	H	H	OMe	OMe
6-Geranyl-3',5,7-tri-hydroxy-4'-methoxyflavanone (syn. 4'-O-methyldiplacone) ( <b>30</b> ) (2S)	Immature fruit (Asai et al. 2008), fruits (Cho et al. 2012, 2013; Lee et al. 2014)	Antioxidant (Asai et al. 2008); SARS-CoV PLpro activity (Cho et al. 2013); hAChE, BChE inhibitor (Cho et al. 2012); significant inhibitory effects against <i>Cp-NanI</i> (a sialidase from <i>C. perfringens</i> ) (Lee et al. 2014); antileishmanial activity ( <i>L. donovani</i> amastigotes) (Salem et al. 2011)	H	H	OMe	OH
6-Geranyl-3,3',5,7-tetrahydroxy-4'-methoxyflavanone (syn. 4'-O-methyldiploacol) ( <b>31</b> ) (2R, 3R)	Immature fruit (Asai et al. 2008), fruit (Cho et al. 2012, 2013)	Antioxidant (Asai et al. 2008), SARS-CoV PLpro activity (Cho et al. 2013); hAChE, BChE inhibitor (Cho et al. 2012)	OH	H	OMe	OH
Schizolaenone C ( <b>32</b> ) (2S)	Fruit (Šmejkal et al. 2010)	Cytotoxic (T-lymphoblastic leukaemia CEM, multiple myeloma RPMI 8226 and U266 cell lines, human cervical cancer cells HeLa, BJ human fibroblast cell line) (Šmejkal et al. 2010), cytotoxic (human ovarian cancer cell line A2780) (Murphy et al. 2006; antioxidant (Zima et al. 2010)	H	OH	H	OH

Table 3 continued




	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>
6-Geranyl-3,3',5,5',7-pentahydroxy-4'-methoxyflavanone ( <b>33</b> ) (2 <i>R</i> , 3 <i>R</i> )	OH	OH	OMe	OH
6-Geranyl-3',5,5',7-tetrahydroxy-4'-methoxyflavanone ( <b>34</b> ) (2 <i>S</i> )	H	OH	OMe	OH
Tomentodiplacone B ( <b>35</b> ) (2 <i>S</i> )	<p>Fruit (Cho et al. 2012)</p> <p>hAChE, BChE inhibitor (Cho et al. 2012)</p> <p>Antioxidant (Yazaki et al. 2009; Zima et al. 2010); low direct cytotoxicity (human erythro-leukemia cell line K562) (Šmejkal et al. 2008a); low direct activity against T-lymphoblastic leukaemia CEM, multiple myeloma RPMI 8226 and U266 cell lines, BJ human fibroblast cell line (Šmejkal et al. 2010), effect on cell cycle (THP-1 human monocytic leukaemia cells) (Kollár et al. 2011); antimicrobial against <i>S. aureus</i> and various methicillin resistant strains of <i>S. aureus</i> (MRSA 287, MRSA 4211, MRSA, 6975, MRSA 630, MRSA 62059) (Navrátilová et al. 2013), no antimicrobial activity on Gram-negative bacteria or yeast and on seven Gram-positive bacteria (Navrátilová et al. 2013; Šmejkal et al. 2008b)</p> 			
	<p>Fruits (Šmejkal et al. 2008b)</p> 			
	R			
3,3',4',5,7-Pentahydroxy-6-[7-hydroxy-3,7-dimethyl-2( <i>E</i> )octenyl] flavanone ( <b>36</b> ) (2 <i>R</i> , 3 <i>R</i> )	OH			
Prokinawan ( <b>37</b> ) (2 <i>S</i> )				
	<p>Immature fruit (Asai et al. 2008)</p> <p>Antioxidant (Asai et al. 2008)</p>			
	<p>Flowers (Kobayashi et al. 2008), immature fruit (Asai et al. 2008)</p> <p>Antioxidant (Asai et al. 2008)</p>			
				

Table 3 continued

	R1	R2	R3
Tomentodiplacone ( <b>38</b> ) (2 <i>S</i> )	H	OMe	H
Fruit (Šmejkal et al. 2008b)			
Antioxidant (Yazaki et al. 2009); antimicrobial against seven Gram-positive bacteria, no antimicrobial activity against three types of Gram-negative bacteria or yeast (Šmejkal et al. 2008b), weak erythroid differentiation activity (Šmejkal et al. 2008a)			
Tomentodiplacol ( <b>39</b> ) (2 <i>R</i> , 3 <i>R</i> )	OH	OMe	H
Fruit (Šmejkal et al. 2007a)			
Antioxidant (Šmejkal et al. 2007a)			
4',5,5',7-Tetrahydroxy-6-[6-hydroxy-3,7-dimethyl-2( <i>E</i> ),7-octadienyl]-3'-methoxyflavanone ( <b>40</b> ) (2 <i>S</i> )	H	OMe	OH
Immature fruit (Asai et al. 2008)			
Antioxidant (Asai et al. 2008)			
3,3',4',5,7-Pentahydroxy-6-[6-hydroxy-3,7-dimethyl-2( <i>E</i> ),7-octadienyl]flavanone ( <b>41</b> ) (2 <i>R</i> , 3 <i>R</i> )	OH	OH	H
Immature fruit (Asai et al. 2008)			
Antioxidant (Asai et al. 2008)			
Tanariflavanone D ( <b>42</b> ) (2 <i>S</i> )	H	H	OH
Immature fruit (Asai et al. 2008), fruit (Schneiderová et al. 2013)			
Antioxidant (Asai et al. 2008; Phommart et al. 2005); cytotoxic (human breast cancer cell line BC) (Phommart et al. 2005); no anti-inflammatory effect in COX-2 inhibition assay (Phommart et al. 2005)			
Tomentomimulol ( <b>43</b> ) (2 <i>R</i> , 3 <i>R</i> )	OH	H	H
Fruit (Schneiderová et al. 2013)			
Mimulone B ( <b>44</b> ) (2 <i>S</i> )	H	H	H
Fruit (Schneiderová et al. 2013)			




Table 3 continued





	R	
Tomtentodiplacone C (45) (racemic mixture of 2 <i>R</i> , 2 <i>S</i> )	Fruit (Navrátilová et al. 2013)	Antimicrobial against <i>S. aureus</i> and various methicillin resistant strains of <i>S. aureus</i> (MRSA 287, MRSA 4211, MRSA, 6975, MRSA 630, MRSA 62059), no antimicrobial activity on Gram-negative bacteria, minor ability to affect the initiation of eukaryotic translation via dual-luciferase reported assay (firefly and renilla) in comparison with anisomycin (Navrátilová et al. 2013)
Mimulone E (46) (2 <i>S</i> )	Fruit (Navrátilová et al. 2013)	 H
Tomtentodiplacone D (47) (2 <i>S</i> )	Fruit (Navrátilová et al. 2013)	Antimicrobial against <i>S. aureus</i> and various methicillin resistant strains of <i>S. aureus</i> (MRSA 287, MRSA 4211, MRSA, 6975, MRSA 630, MRSA 62059), no antimicrobial activity on Gram-negative bacteria, no ability to affect the initiation of eukaryotic translation via dual-luciferase reported assay (firefly and renilla) (Navrátilová et al. 2013)
Tomtentodiplacone E (48) (2 <i>S</i> )	Fruit (Navrátilová et al. 2013)	 H
Tomtentodiplacone F (49) (2 <i>S</i> )	Fruit (Navrátilová et al. 2013)	Antimicrobial against <i>S. aureus</i> and various methicillin resistant strains of <i>S. aureus</i> (MRSA 287, MRSA 4211, MRSA, 6975, MRSA 630, MRSA 62059), no antimicrobial activity on Gram-negative bacteria, no ability to affect the initiation of eukaryotic translation via dual-luciferase reported assay (firefly and renilla) (Navrátilová et al. 2013)
Tomtentodiplacone G (50) (2 <i>S</i> )	Fruit (Navrátilová et al. 2013)	 H
Tomtentodiplacone H (51) (racemic mixture 2 <i>R</i> , 2 <i>S</i> )	Fruit (Navrátilová et al. 2013)	Antimicrobial against <i>S. aureus</i> and various methicillin resistant strains of <i>S. aureus</i> (MRSA 287, MRSA 4211, MRSA, 6975, MRSA 630, MRSA 62059), no antimicrobial activity on Gram-negative bacteria, no ability to affect the initiation of eukaryotic translation via dual-luciferase reported assay (firefly and renilla) (Navrátilová et al. 2013)
Tomtentodiplacone I (52) (racemic mixture of 2 <i>R</i> , 2 <i>S</i> )	Fruit (Navrátilová et al. 2013)	 H



Table 3 continued

		R				
Mimulone C ( <b>53</b> ) (2 <i>S</i> )	Fruit (Navrátilová et al. <a href="#">2013</a> )	Antimicrobial against <i>S. aureus</i> and various methicillin resistant strains of <i>S. aureus</i> (MRSA 287, MRSA 4211, MRSA, 6975, MRSA 630, MRSA 62059), no antimicrobial activity on Gram-negative bacteria, <b>53</b> has no ability to affect the initiation of eukaryotic translation via dual-luciferase reported assay (firefly and renilla) (Navrátilová et al. <a href="#">2013</a> )				
Mimulone D ( <b>54</b> ) (2 <i>S</i> )	Fruit (Navrátilová et al. <a href="#">2013</a> )					
			R1	R2	R3	R4
Tomentin A ( <b>55</b> ) (2 <i>S</i> )	Fruit (Cho et al. <a href="#">2013</a> )	Activity against Severe acute respiratory syndrome corona virus (SARS-CoV PLpro, a reversible inhibitor <b>63</b> was the most potent compound tested (Cho et al. <a href="#">2013</a> )	H	H	OH	OH
Tomentin B ( <b>56</b> ) (2 <i>S</i> )			H	H	OMe	OH
Tomentin C ( <b>57</b> ) (2 <i>S</i> )			H	OH	OMe	OMe
Tomentin D ( <b>58</b> ) (2 <i>S</i> )			H	OMe	OH	OMe
Tomentin E ( <b>59</b> ) (2 <i>R</i> , 3 <i>S</i> )			OH	H	OH	OMe

**Table 4** Phenolic glycosides isolated from *P. tomentosa*



Name of compound and ID	Isolation	Biological activity	Chemical structure
Coniferin (syn. abietin, coniferosid, laricin) ( <b>69</b> )	Bark (Sticher and Lahloub 1982; Zhu et al. 1986)	Antiphlogistic (Choi et al. 2004), low COX-2/1 and 5-LOX inhibition (Wang et al. 2013; Diaz Lanza et al. 2001); low antioxidant activity in vitro (DPPH) (She et al. 2013); stimulation osteoblastic bone formation in vitro (Ding et al. 2010); concentration-dependent contractions in rat aortic rings (Deliorman et al. 2000); inhibition of ADP-induced platelet aggregation (Panossian et al. 1998); hypotensive in rats (Matsubara et al. 1991)	
Syringin (syn. eleutherosid B) ( <b>70</b> )	Bark (Sticher and Lahloub 1982; Zhu et al. 1986)	Antidepressant (Kurkin et al. 2006), antiphlogistic, antinociceptive (Choi et al. 2004); peroxyl radical scavenging capacity (Kim et al. 2010a, b), inhibition of NO production in LPS-induced RAW 264.7 cells (Lee et al. 2009); enhances memory in aged rats (Huang et al. 2013a, b); protective effects against Aβ(25–35)-induced atrophies of axons and dendrites (Bai et al. 2011; Tohda et al. 2008); immunomodulatory activity in PMN phagocytic function test (Sharma et al. 2012); hepatoprotective in mice (Gong et al. 2014); adiponectin receptor 2 agonist (Sun et al. 2013); low COX and 5-LOX inhibition (Diaz Lanza et al. 2001); concentration-dependent contractions in rat aortic rings (Deliorman et al. 2000); inhibition of ADP-induced platelet aggregation (Panossian et al. 1998; Iizuka et al. 2005); moderate in vitro cytotoxic activities against A549 and HL-60 (Yang et al. 2012); antitumour activity in sarcoma S180 transplanted mice (Zhang et al. 2007); weak inhibition of snake PDE I (Chai et al. 2009); hypoglycemic effect (Liu et al. 2008; Niu et al. 2008a, b; Niu et al. 2007); adaptogenic activity reviewed by Panossian and Wagner (2005)	

Table 4 continued

	R1	R2	R3
Acteoside (syn. verbascoside) ( <b>71</b> )	OH	OH	H
Bark (Si et al. 2011b; Sticher and Lahloub 1982; Zhu et al. 1986), wood (Si et al. 2008d), sapwood (Ota et al. 1993), fruit (Šmejkal et al. 2007b), leaves (Schilling et al. 1982), young plant (Damtoft and Jensen 1993)			
$\beta$ -Oxoacteoside (syn. tomentoside A) ( <b>72</b> )			
Martynoside ( <b>73</b> )	OH	OH	=O
Antioxidant (Tozuka et al. 2005)			
	OMe	OMe	H
Antioxidant (Jiménez and Riguera 1994); antihypertensive (Kang et al. 2003); cytotoxic (antiestrogen in breast cancer cells MCF7 via the ER-pathway, osteoblasts KS483, endometrial cancer cells Ishikawa) (Papoutsi et al. 2006)			
Campneoside I ( <b>74</b> )	OH	OH	OMe
Antibacterial against <i>Staphylococcus</i> and <i>Streptococcus</i> species (MIC 150 $\mu$ g/ml) (Kang et al. 1994)			
Campneoside II (syn. $\beta$ -hydroxy-acteoside) ( <b>75</b> )	OH	OH	OH
Antibacterial, anti-inflammatory (Jiménez and Riguera 1994), anticomplement (Si et al. 2008b)			
Ilicifolioside A ( <b>76</b> )	OH	OH	OEt
Anticomplement (Si et al. 2008b)			






Table 4 continued

			R
Isocampneoside (syn. isoverbascoside) (77)	Bark (Si et al. 2011b), wood (Si et al. 2008d), sapwood (Ota et al. 1993), fruit (Šmejkal et al. 2007b), leaves (Schilling et al. 1982), young plant (Damtoft and Jensen 1993)	Antioxidant (Šmejkal et al. 2007b), xanthine oxidase inhibition (Kong et al. 1999); suppression of glutamate induced neurotoxicity (Koo et al. 2005); antioxidant, immunosuppressive (Jiménez and Riguera 1994); cytotoxic (e.g., human erythro-leukemia cell line K562) (Šmejkal et al. 2008a)	H
Isocampneoside I (78)	Wood (mixture of <i>R</i> - and <i>S</i> -epimers) (Si et al. 2008d)	Inhibition of D-galactosamine induced cytotoxicity in primary cultured mouse hepatocytes (Pan et al. 2010)	OMe
Isocampneoside II (79)	Bark (Si et al. 2011b), wood (Si et al. 2008b)	Aldose reductase inhibitor (potential against diabetic complications) (Kim et al. 2011); anticomplement (Si et al. 2008b); antioxidant, neuro-protective on hydrogen peroxide-induced oxidative injury in PC12 cells (Si et al. 2013).	OH
Isollicifolioside A (80)	Wood (Si et al. 2008b)	Anticomplement (Si et al. 2008b)	OEt
Cistanoside F (81)	Bark (Si et al. 2011b)	Antioxidant (Xiong et al. 1996); vasorelaxant (Yoshikawa et al. 2006); interaction with bovine serum albumin (Wu et al. 2012)	



**Fig. 2** Flavonoid glycosides

pathways is provided by prenyltransferases (Šmejkal 2014; Andersen and Markham 2006; Kuzuyama et al. 2005). The side chain can later be converted into different moieties by several reactions. It is unclear, whether the described modifications of the prenyl side-chain are natural—sunlight, the presence of oxygen and an elevated temperature can all affect the terpenoid metabolism—or are artefacts of isolation. Some of the changes in the prenyl moiety have been observed after treatment of an extract, or during the separation of a mixture in the acidic environment of silica gel (Navrátilová et al. 2013; Šmejkal 2014) (Fig. 2).

Most of the prenylated flavonoids isolated from *P. tomentosa* belong to C-geranylated group of flavonoids (21–52). Some of them have their side-chain further modified by hydroxylation (35–44, 50, 51, 53–59), methoxylation (45–47, 49, 50), oxidation (47, 48, 52), cyclization (51, 53–59), or reduction (19, 20, 47–52). Interestingly, these compounds have been isolated from the leaves, flowers and fruit, but they are most commonly isolated from the roots, root bark, or bark of different plants (Botta et al. 2005). Compounds 23, 24, and 28 have been found in the yellow dendritic trichomes on the adaxial side of the *P. tomentosa* leaves. On the other hand, no significant detected concentration of secondary metabolites has been detected in the white dendritic trichomes on the adaxial side of the leaves or the brown dendritic trichomes on the flower buds (Kobayashi et al. 2008). Glandular hairs found on the young reproductive

organs of *P. tomentosa* are rich in flavonoids, with concentrations over 1,000 times greater than those on the surfaces of the young leaves (Kobayashi et al. 2008). Some seasonal variations in the appropriate time for harvesting the fruit have been discovered. Autumn is the best time to obtain high concentrations of flavonoids whereas early summer is better for phenylpropanoid glycosides (Holubová and Šmejkal 2011).

The antioxidant (Asai et al. 2008; Šmejkal et al. 2007a, b; Zima et al. 2010), antibacterial (Navrátilová et al. 2013; Šmejkal et al. 2008b), antiphlogistic (Hošek et al. 2010), cytotoxic (Kollár et al. 2011; Šmejkal et al. 2007a, 2008a, 2010), and severe acute respiratory syndrome coronavirus papain-like protease (SARS-CoV PLpro) activities (Cho et al. 2013) of isolated geranylated flavonoids have been described recently, together with their inhibitory effect on both human acetylcholinesterase and butyrylcholinesterase (Cho et al. 2012). The great ability of several geranylated flavanones to interact with bacterial sialidase isolated from *Clostridium perfringens* (Cp-NanI), a bacterium causing various gastrointestinal diseases, has recently been reported (Lee et al. 2014).

Possible relationships between structure and activity have been proposed for each of these biological activities. Nevertheless, it is difficult to evaluate real structure–activity relationships only by comparing the published studies, because the study conditions and the assays employed are often different. Generally, the

addition of an isoprenoid chain renders the derivative molecule more pharmacologically effective than the parent compound, probably because the prenyl group increases the lipophilicity and confers a strong affinity for biological membranes (Botta et al. 2005; Epifano et al. 2007; Chen et al. 2014).

Interestingly, neither the geranyl side-chain nor its substitution affects the antioxidant activity of flavonoids. The spatial arrangement of the substitution of the flavanone skeleton is more important. For example, the antiradical activity is increased by 2'-hydroxyl substitution of the ring B, whereas 4'-methoxyl substitution diminishes it. The general rules postulated for the antioxidant activity of flavonoids in vitro are applicable (Havsteen 2002; Chen et al. 2002), there are many review publications that touch on this topic, and it is not aim of this paper to delve deeply into this (Plaza et al. 2014). The type of antioxidant assays used for the experiment could also be a factor that significantly affects this activity (Zima et al. 2010).

Numerous reports about structure related antimicrobial activity have been published, but comparison shows the results to be widely conflicting (Cushnie and Lamb 2005). Based on several studies, it has been postulated that hydroxylation at position C-5 or C-7 of ring A and positions C-2' or C-4' of ring B increases the antibacterial activity (Šmejkal et al. 2008b; Navrátilová et al. 2013). However, contrary to this assumption, **45** and **50** had no significant antibacterial activity (Navrátilová et al. 2013). Interestingly, some C-geranylated flavonoids do not able inhibit the growth of Gram-negative bacteria (**21**, **22**, **25–27**, **35**, **38**, **45**, **47–48**, **50**, **53**, and **54**). Resistance to these compounds is probably due to the more complex structure and hydrophilic nature of G-cell walls (Navrátilová et al. 2013; Šmejkal et al. 2008b). Furthermore, substitution of the geranyl side-chain with carbonyl, hydroxyl or methoxyl groups diminishes the antibacterial activity in a manner similar to what is seen when the geranyl substituent at C-6 forms a ring by reacting with the hydroxyl group at C-7. Compounds **45**, **47**, **48**, **50**, **53**, and **54** exhibit some degree of activity in the range of the concentrations tested on the Gram-positive bacteria *Staphylococcus aureus* and various methicillin resistant strains of *S. aureus*. The level of activity varied in depending on both the structure and the bacterial strain used in the assay. Flavonoid structures like **24** are more effective in protecting plants from water loss because of their

reduced polarity (Kobayashi et al. 2008; Navrátilová et al. 2013). Compounds **45**, **47**, **48**, **50**, and **53** were also tested for their ability to affect the initiation of the eukaryotic translation via dual-luciferase reported assay (firefly and renilla), but only **45** showed a modest activity in comparison with anisomycin (Navrátilová et al. 2013).

The cytotoxicity of the prenylated flavonoids obtained from *P. tomentosa* has been tested in more than 20 different cell lines. The type of prenylation strongly affects the cytotoxicity of a flavonoid in the traditional P-388 murine leukemia model. The unmodified 3-prenyl group and the presence of corresponding *ortho*-dihydroxy or trihydroxy substitution of the flavonoid ring B are crucial to its activity against P-388 as compared with other prenyl substituents (Hakim et al. 1999, 2002, 2006). Similar findings have been observed for modified C-6 geranyl groups and the substitution of the ring B for other cell lines tested (Šmejkal et al. 2010). However, replacing the *para*-hydroxy group of the ring B of a C-8 prenylated flavanone with several different acyl substituents resulted in greater cytotoxicity (Aniol et al. 2012; Šmejkal 2014). It has also been found that cytotoxic activity is diminished by the presence of a C-3 hydroxyl substituent on the ring C, 4'-methoxy substitution of the ring B or a *para*-hydroxy substituted ring B (Šmejkal et al. 2008a). For this reason, it is important to emphasize that the relative importance of the substitution of ring B can differ according to the cell line used. Modification of a prenyl or geranyl side-chain can not only change the direct cytotoxicity, but it may also affect the proliferation cells or trigger apoptosis (Kollár et al. 2011; Šmejkal et al. 2007a, 2008a, 2010; Šmejkal 2014).

Prenyated flavonoids **55–59**, modified with an unusual 3,4-dihydro-2*H*-pyran moiety, have been found to inhibit the severe acute respiratory syndrome corona virus PLpro enzyme more effectively than their parent compounds, the precursors from which were they derived (Cho et al. 2013).

The presence of a geranyl group at the C-6 position (**21**, **22**, **24–26**, **30**, **31**, **33**, and **34**) seems to be crucial for the hAChE and BChE inhibitory effects. The most effective inhibitor was **22**. All of the geranylated flavonoids, apart from **26**, inhibited hAChE dose-dependently. It appears that greater inhibition is observed when ring B of the flavanone bears free hydroxyl groups (Cho et al. 2012).


The crystal structure of the bacterial sialidase *Cp*-NanI catalytic domain in a complex with the geranylated flavonoid diplacone (**22**) provides structural insights into the binding mode of natural flavonoid-based inhibitors at atomic resolution. It shows how the geranyl and C-3' hydroxyl groups of **22** contribute to the stability of the enzyme-inhibitor complex. Time-dependent competitive inhibition patterns have been observed. Structural comparison of the human sialidases Neu1–Neu4 with the *Cp*-NanI–diplacone (**22**) complex suggests that the interaction between human sialidases and **22** is likely to be unfavourable because of polar or ionic repulsion (Lee et al. 2014).

Six flavonoid glycosides have been extracted from the stem bark **60–62** (Si et al. 2009) and flowers **63–66** (Scogin 1980) of *P. tomentosa*. Apigenin-7-*O*- $\beta$ -D-glucopyranoside (synonym: cosmosiin) (Kurkina et al. 2011) (**60**) shows anti-HIV activity in vitro on H9 cells (Tang et al. 1994) and enhances the secretion of adiponectin, the phosphorylation of the tyrosine residue of insulin receptor- $\beta$ , and the translocation of GLUT5 (Rao et al. 2011). Compound **60** shows no significant hepatic Glc-6-phosphatase inhibitory activity in vitro (Kumar et al. 2010). Apigenin-7-*O*- $\beta$ -D-glucuronopyranoside (**61**) was more potent than apigenin (**2**) and omeprazole in an assay analysing the inhibition of reflux esophagitis and gastritis promoted surgically and by application of indomethacine in rats (Min et al. 2005). No information about the biological activity of apigenin-7-*O*-[ $\beta$ -D-glucuronopyranosyl (1  $\rightarrow$  2)-*O*- $\beta$ -D-glucuronopyranoside] (**62**) has been found in the literature. Delphinidin-3-*O*-glucoside (synonym:

myrtillin) (**63**) protects microglia from inflammation-induced stress signaling (Carey et al. 2013), is cytotoxic against the MCF-7 (breast) cancer cell line (Vareed et al. 2006), and has an affinity for the estrogenic ER $\beta$  receptor (Hidalgo et al. 2012). Delphinidin-3,5-di-*O*-glucoside (**64**) and cyanidin-3,5-di-*O*-glucoside (**65**) are potent antioxidants (Lee et al. 2011; Tanaka et al. 2013). Compounds **63** and **65** are potent ACE inhibitors in vitro (Hidalgo et al. 2012; Persson et al. 2009).

### *P. tomentosa* lignans and their biological activity

Only three lignans: (+)-paulownin (**66**), (+)-sesamin (**67**), (+)-piperitol (**68**) have been isolated from the wood of *P. tomentosa* (Ina et al. 1987; Takahashi and Nakagawa 1966; Zhu et al. 1986) (Fig. 3). Several reports have described their pharmacological effects (Zhang et al. 2014; Pan et al. 2009). Compound **66** is a promising antifungal agent that acts against the basidiomycetes *Fomitopsis palustris* and *Trametes versicolor* (Kawamura et al. 2004). Sesamin (**67**) has been previously isolated from the wood of *P. tomentosa*, and *P. kawakamii* and the hybrid of *P. elongata*  $\times$  *P. fortunei* (Takahashi and Nakagawa 1966; Zhu et al. 1986). Compound **67** is the major lignan in sesame, and has various biological activities, such as antioxidant, angiogenic (Chung et al. 2010), antiphlogistic (Chatrattanakunchai et al. 2000), antihypertensive (Nakano et al. 2006), insecticidal (Nascimento et al. 2004), neuroprotective (Khan et al. 2010), and anticarcinogenic (Yasuda and Sakaki



**Fig. 3** Lignans


2012). However, this compound showed no significant ability to reduce the formation of atherosclerotic lesions in the ApoE (−/−) gene-knockout mouse, whereas specific dietary polyphenols, especially quercetin and theaflavin were more active (Loke et al. 2010). It has been suggested that some of the biological effects attributed to sesamine (67) may, in fact, be caused by its metabolites and that 67 may be acting as a proactive substance in the body (Yasuda and Sakaki 2012). The metabolism of sesamin (67) by cytochrome P450 and UDP-glucuronosyltransferase has been found remarkably different in humans as compared to other animals. Further investigation into the safety of taking sesamin (67) with therapeutic drugs that are metabolised by CYP2C9 is also needed (Yasuda and Sakaki 2012).

#### *P. tomentosa* phenolic glycosides and their biological activity

A total of thirteen phenylpropanoid glycosides (69–81) with multifarious bioactivities, natural compounds derived biosynthetically from the amino acid phenylalanine, have been isolated from different parts of the *P. tomentosa* plant (Table 4) (Damtoft and Jensen 1993; Kang et al. 1994; Ota et al. 1993; Schilling et al. 1982; Si et al. 2008b, d, Si et al. 2011b; Sticher and Lahloub 1982; Šmejkal et al. 2007b; Zhu et al. 1986). Phenylpropanoids are of great interest for fabricating effective tonic, anticancer, hepatoprotective, immunostimulating, antimicrobial, and anti-inflammatory phytopreparations (Kurkin 2003; Galvez et al. 2006; Panossian and Wagner 2005; Fu et al. 2008; Pan et al. 2003).

#### *P. tomentosa* quinones and their biological activity

A new furanoquinone, methyl-5-hydroxy-dinaphthol [1,2–2',3']furan-7,12-dione-6-carboxylate (MHDDC) (82) (Fig. 4) identified in *P. tomentosa* stem bark (MeOH extract) has been shown to possess antiviral activity against poliovirus types 1 and 3, using HeLa cells in vitro (Kang et al. 1999). Its cathepsin K and L inhibitory activities in vitro have recently been discovered. The 5-OH functional group may have a favourable effect on this reduction potential which prevents the degradation of bone the matrix carried out by osteoclasts (Park et al. 2009).



**Fig. 4** Quinones

The naphthoquinone plumbagin (83) has been detected in the leaves and fruit of *P. tomentosa* (Babula et al. 2006). It has been used in traditional systems of medicine since ancient times (Pile et al. 2013). It has exhibited promising antimalarial activity in vitro with IC<sub>50</sub> of 580 (270–640) nM and 370 (270–490) nM, respectively, against 3D7 chloroquine-sensitive *P. falciparum* and K1 chloroquine-resistant *P. falciparum* clones, using an assay based on SYBR Green I. Toxicity testing indicated relatively low toxicity at dose levels up to 100 mg/kg body weight (for a single oral dose) and 25 mg/kg (for daily dose given for consecutive 14 days) for acute and subacute toxicity, respectively. Based on the results of in vivo antimalarial testing, plumbagin administered at a the dose of 25 mg/kg of body weight for 4 consecutive days exhibited moderate to weak antimalarial activity with regard to its ability to reduce parasitaemia and prolong survival time (Sumsakul et al. 2014). Other published studies of plumbagin (83) described effects such as antifungal, antibacterial, cytotoxic (Krishnaswamy and Purushothaman 1980), anticoagulant (Santhakumari et al. 1978), anthelmintic (antischistosomal) (Zhang and Coultas 2013) and an anti-inflammatory effect on the amelioration of experimental ulcerative colitis in mice at a dose of 8–10 mg/kg (Pile et al. 2013).

#### *P. tomentosa* terpenoids and their biological activity

Six iridoids: 7-β-hydroxyharpagide (84), paulownioside (85), catalpol (86), aucubin (87), tomentoside (88) and 7-hydroxytomentoside (89) have been isolated from the leaves of *P. tomentosa* (84–87) (Adriani et al. 1981; Franzyk et al. 1999), the bark of the trunk and



roots (**86**) (Plouvier 1971) and parts of the young plant (**85**, **86**, **88**, and **89**) (Damtoft and Jensen 1993) (Fig. 5). Compound **86** shows neuroprotective activity (Li et al. 2004), a cardioprotective effect against myocardial infarction—specifically reperfusion damage (Huang et al. 2013a, b), and radioprotective effects (Chen et al. 2013). It also increases glucose utilization by increasing the secretion of  $\beta$ -endorphin from the adrenal gland (Shieh et al. 2011). Compounds **86** and **87** possess antispasmodic activity (Deurbina et al. 1994). Aucubin (**87**) had antioxidant and pancreas-protective effects on streptozotocin-induced diabetes in rats (Jin et al. 2008) and it may improve obesity-induced atherosclerosis by attenuating the TNF- $\alpha$ -induced inflammatory response (Park 2013). A weak antibacterial effect (against *Streptococcus pneumoniae* and MG-hemolytic streptococcus, MIC 28.946 mg/ml) (Zheng et al. 2012) and neuroprotective effects of **87** on diabetes and diabetic encephalopathy (Xue et al. 2012) have also been observed.


The sesquiterpenic lacton isoatriplicolide tiglate (**90**) has been isolated from *P. tomentosa* flowers. It has neuroprotective effects against glutamate-induced neurotoxicity (Kim et al. 2010a, 2010b) and cytotoxic activity against several cancer cell lines: A549 lung carcinoma; SK-OV-3 adenocarcinoma; SK-Mel-2 malignant melanoma; XF498 central nervous system tumors; HCT15 colon adenocarcinoma, against which its effect is comparable to that of reference substance adriamycin (Moon and Zee 2001); human breast cancers MDA-MB-231, MCF7, HS578T, and T47D; and the HeLa, SiHa and C33A cervical cancer cell lines (Jung et al. 2012).

Seven phytosterols have been isolated from *P. tomentosa* leaves: ursolic acid (**91**) (Zhu et al. 1986; Zhang and Li 2011), 3-epiursolic acid (**92**), pomolic acid (**93**), corosolic acid (**94**), maslinic acid (**95**),  $\beta$ -sitosterol (**96**), and daucosterol (**97**) (Zhang and Li 2011). Most of these show various biological activities, e.g., **91** is a potentially useful for treating Alzheimer's disease (because of its ability to block the interactions of the amyloid  $\beta$ -CD36) (Wilkinson et al. 2011). It can prevent the recruitment of the monocytes that accelerate atherosclerosis, a major complication of diabetes, in mice (Ullevig et al. 2011), and it also possesses antibacterial (Wong et al. 2012), anti-trypanosomal, and anti-leishmanial properties (Bero et al. 2011). Compound **91** has also shown cytotoxic effects

against K562 and K562/ADR human chronic myelogenous leukaemia, HL60 and HL60/ADR human acute myelocytic leukemia cancer cells, and the human colon cancer cell lines SW480 and SW620 (Shan et al. 2011). Compounds **93–96** also show promising cytotoxic potential, e.g., **93** is effective against the human chronic myelogenous leukaemia cell line 562 and also against cells derived from chronic myeloid leukaemia patients (Vasconcelos et al. 2007). Compound **94** shows cytotoxic effects against osteosarcoma MG-63 cells (Cai et al. 2011) and immunosuppressive activity on myeloid-derived suppressor cells in the murine sarcoma model (Holland et al. 2013). Compound **95** is effective against HT29 human colon cancer cells (Reyes-Zurita et al. 2011) and **96** promotes apoptosis in various cancer cells (Hac-Wydro 2013). Anti-inflammatory activities have been revealed for compounds **91**, **92**, and **96** (Deepak and Handa 2000), and **93** (Schinella et al. 2008). An immunomodulatory effect of **97** in protecting mice against candidiasis disseminated by the CD4+ Th1 immune response has been seen (Lee et al. 2007). Other recently discovered pharmacological effects include antimalarial (**95**) (Moneriz et al. 2011), hypotensive and endothelium-dependent vasorelaxant effects (**93**) (Estrada et al. 2011). Compound **95** is potentially useful for treating cerebral ischemic injuries (Guan et al. 2011), and **94** has promising antidiabetic effects (Sivakumar et al. 2009).


### *P. tomentosa* glycerides and their biological activity


Thirty acylglycerols (**98–127**) belonging to 1,3-di-*O*-acetyl-2-*O*-(acyl)-glycerols, 1-*O*-acetyl-2-*O*-(acyl)-*sn*-glycerols and 2-*O*-(acyl)-glycerols, including only five known compounds (**99**, **106**, **108**, **114**, and **119**), have been identified in the exudates secreted from the glandular trichomes on the leaves: 1,3-di-*O*-acetyl-2-*O*-[(3*R*,6*S*)-3-(acetyloxy)-6-hydroxyeicosanoyl]-glycerol (**98**), 1-*O*-acetyl-2-*O*-[(3*R*)-3-(acetyloxy)octadecanoyl]-*sn*-glycerol (**99**), 1-*O*-acetyl-2-*O*-[(3*R*,7*R*)-3,7-*bis*(acetyloxy)octadecanoyl]-*sn*-glycerol (**100**), 1-*O*-acetyl-2-*O*-[(3*R*,6*S*)-3-(acetyloxy)-6-hydroxyoctadecanoyl]-*sn*-glycerol (**101**), 2-*O*-[(3*R*,7*R*)-3,7-dihydroxy-eicosanoyl]-glycerol (**102**), 2-*O*-[(3*R*,6*S*)-3,6-dihydroxy-eicosanoyl]-glycerol (**103**), 2-*O*-[(3*R*,8*R*)-3,8-dihydroxy-eicosanoyl]-glycerol (**104**), 2-*O*-[(3*R*,

**Fig. 5** Terpenoids

9*R*)-3,9-dihydroxy-eicosanoyl]-glycerol (**105**), 1-*O*-acetyl-2-*O*-[(3*R*)-3-(acetyloxy) eicosanoyl]-*sn*-glycerol (**106**), 1-*O*-acetyl-2-*O*-[(3*R*,6*S*)-3,6-*bis*(acetyloxy)eicosanoyl]-*sn*-glycerol (**107**), 1-*O*-acetyl-2-*O*-

[(3*R*,7*R*)-3,7-*bis*(acetyloxy)eicosanoyl]-*sn*-glycerol (**108**), 1-*O*-acetyl-2-*O*-[(3*R*,8*R*)-3,8-*bis*(acetyloxy)eicosanoyl]-*sn*-glycerol (**109**), 1-*O*-acetyl-2-*O*-[(3*R*,9*R*)-3,9-*bis*(acetyloxy)eicosanoyl]-*sn*-glycerol (**110**), 1-*O*-

**Fig. 6** Glycerides



**Fig. 7** Miscellaneous compounds

acetyl-2-*O*-[(3*R*,6*S*)-3-(acetyloxy)-6-hydroxyeicosanoyl]-*sn*-glycerol (**111**), 1-*O*-acetyl-2-*O*-[(3*R*,8*R*)-3-(acetyloxy)-8-hydroxyeicosanoyl]-*sn*-glycerol (**112**), 1-*O*-acetyl-2-*O*-[(3*R*,9*R*)-3-(acetyloxy)-9-hydroxyeicosanoyl]-*sn*-glycerol (**113**), 2-*O*-[(3*R*)-3-(acetyloxy) octadecanoyl]-glycerol (**114**), 2-*O*-[(3*R*,6*S*)-3-(acetyloxy)-6-hydroxy-octadecanoyl]-glycerol (**115**), 2-*O*-[(3*R*,7*R*)-3-(acetyloxy)-7-hydroxy-octadecanoyl]-glycerol (**116**), 2-*O*-[(3*R*,8*R*)-3-(acetyloxy)-8-hydroxy-octadecanoyl]-glycerol (**117**), 2-*O*-[(3*R*,9*R*)-3-(acetyloxy)-9-hydroxy-octadecanoyl]-glycerol (**118**), 2-*O*-[(3*R*,7*R*)-3,7-*bis*(acetyloxy) octadecanoyl]-glycerol (**119**), 2-*O*-[(3*R*,8*R*)-3,8-*bis*(acetyloxy)octadecanoyl]-glycerol (**120**), 2-*O*-[(3*R*)-3-(acetyloxy)eicosanoyl]-glycerol (**121**), 2-*O*-[(3*R*,6*S*)-3-(acetyloxy)-6-hydroxy-eicosanoyl]-glycerol (**122**), 2-*O*-[(3*R*,7*R*)-3-(acetyloxy)-7-hydroxy-eicosanoyl]-glycerol (**123**), 2-*O*-[(3*R*,8*R*)-3-(acetyloxy)-8-hydroxy-eicosanoyl]-glycerol (**124**), 2-*O*-[(3*R*,9*R*)-3-(acetyloxy)-9-hydroxy-eicosanoyl]-glycerol (**125**), 2-*O*-[(3*R*,8*R*)-3,8-*bis*(acetyloxy) eicosanoyl]-glycerol (**126**), 2-*O*-[(3*R*,9*R*)-3,9-*bis*(acetyloxy)eicosanoyl]-glycerol (**127**). The relatively most abundant constituents were **111** (20 % of the total glycerides), **126** (14 %), and **120** and **127** (12 %) (Asai et al. 2009). Another analysis of the secretions of the glandular hairs on the leaves of both bud flushes and adult trees as well as the flowers of adult trees showed that these secretions contain ten glycerides (**106**, **108–110**, **118**, **122**, **124–127**). These compounds showed sticky character, but they were not toxic to several insects. Therefore, the glandular hairs on the leaves and flowers may serve only to physically deter herbivores (Kobayashi et al. 2008) (Fig. 6).

### Miscellaneous *P. tomentosa* compounds and their biological activity

Six known phenolic acids, namely *p*-hydroxybenzoic acid (**128**), vanillic acid (**129**), gallic acid (**130**), cinnamic acid (**131**), *p*-coumaric acid (**132**), and caffeic acid (**133**) have been identified in the leaves of *P. tomentosa* (**128–133**), bark (**130** and **131**) and wood (**133**) (Ota et al. 1993; Si et al. 2008c, 2011b), and *p*-ethoxybenzaldehyde (**134**) is present in flowers (Yuan et al. 2009). The 5,7-dihydroxy-6-geranylchromone (**135**) isolated from fruits shows only moderate cytotoxic activity against a suspension culture of *Nicotiana tabacum* cv. Bright Yellow (BY-2), which confirms that ring B of the flavonoid skeleton is important for this activity (Šmejkal et al. 2008a). A large group of essential oil substances found in the flowers of *P. tomentosa* have also been identified (Oprea et al. 2004; Ibrahim et al. 2013) (Fig. 7).

### Conclusions

Plants are still highly esteemed all over the world as rich sources of therapeutic agents. In this context, traditional Chinese herbal medicines, such as *Paulownia* continue to influence a modern healthcare. It has been estimated that approximately 420,000 plant species exist on Earth, but little is known about the phytochemical or therapeutic qualities of most of them. Thanks to the development of the spectral and other analytical methods used in modern phytochemistry, many of the principal physiologically active

secondary metabolites have been identified and researched in detail. The accumulated knowledge of traditional medicine is therefore, playing an important role in enhancing the success of drug discovery in herbal medicine. Approximately 80 % of the currently known antimicrobial, cardiovascular, immunosuppressive, and anticancer drugs are of plant origin (Pan et al. 2013).

As mentioned in this review, *P. tomentosa* is a rich source of multifarious secondary metabolites, mainly prenylated flavonoids. As of today 135 compounds, including flavonoids, lignans, phenolic glycosides, quinones, terpenoids, glycerides, phenolic acids, and other miscellaneous compounds have been isolated from various extracts of this plant.

Of increasing interest are the isolation and identification of *P. tomentosa* prenylated flavonoids, as they have shown promising pharmacological effects. In the first experiments, their antioxidant, antibacterial, antiphlogistic, cytotoxic, and SARS-CoV PL activities have been discovered along with inhibitory effects on human acetylcholinesterase, butyrylcholinesterase, and bacterial neuraminidases. More than 40 compounds with modified prenyl or geranyl side-chains attached at C-6 of the flavonoid skeleton have been isolated from the flowers, fruit, and leaves of *P. tomentosa*. Only two of these compounds have shown the presence of a five-carbon side-chain; for the others a ten-carbon side-chain is typical. Further, only a few compounds have shown a geranyl moiety modified by hydroxylation at C-6'' or C-7''. Some compounds have a geranyl group modified by forming a heterocyclic moiety, which is also unusual. Most of them have never been isolated from any other plant species. However, further in vivo pharmacology studies are needed to precisely elucidate biological mechanism of action, efficacy, and toxicity of these promising therapeutic agents. Elucidation of the structure–activity relationships is also crucial for their further total syntheses and introduction into medical practice. For these reasons, the study of *P. tomentosa* as a source of biologically active metabolites is significant and future interest in this plant is ensured.

## References

- Adriani C, Bonini C, Iavarone C et al (1981) Isolation and characterization of paulownioside, a new highly oxygenated iridoid glucoside from *Paulownia tomentosa*. J Nat Prod 44:739–744
- Ahmad M, Rizwani GH (1995) Acteoside: a new antihypertensive drug. Phytother Res 9:525–527
- Akbay P, Calis I, Ünder U et al (2002) *In vitro* immunomodulatory activity of verbascoside from *Nepeta ucrainica* L. Phytother Res 16:593–595
- Andersen OM, Markham KR (2006) Flavonoids: chemistry, biochemistry, applications. CRC Press, Boca Raton
- Aniol M, Swiderska A, Stompor M et al (2012) Antiproliferative activity and synthesis of 8-prenylnaringenin derivatives by demethylation of 7-*O*- and 4'-*O*-substituted isoxanthohumols. Med Chem Res 21:4230–4238
- Arutyunyan TV, Korystova AF, Kublik LN et al (2012) Effects of taxifolin on the activity of angiotensin-converting enzyme and reactive oxygen and nitrogen species in the aorta of aging rats and rats treated with nitric oxide synthase inhibitor and dexamethasone. Age 35:2089–2097
- Asai T, Hara N, Kobayashi S et al (2008) Geranylated flavanones from the secretion on the surface of the immature fruits of *Paulownia tomentosa*. Phytochemistry 69: 1234–1241
- Asai T, Hara N, Kobayashi S et al (2009) Acylglycerols (=glycerides) from the glandular trichome exudate on the leaves of *Paulownia tomentosa*. Helv Chim Acta 92:1473–1494
- Babula P, Mikelová R, Adam V et al (2006) Chromatografické stanovení naftochinonů v rostlinách (Chromatographic evaluation of naphthochinones in plants). Chem Listy 100:271–276
- Bai Y, Tohda C, Zhu S et al (2011) Active components from Siberian ginseng (*Eleutherococcus senticosus*) for protection of amyloid  $\beta$ (25–35)-induced neuritic atrophy in cultured rat cortical neurons. J Nat Med 65:417–423
- Bansal S, Vyas S, Bhattacharya S et al (2013) Catechin prodrugs and analogs: a new array of chemical entities with improved pharmacological and pharmacokinetic properties. Nat Prod Rep 30(11):1438–1454
- Bero J, Hannaert V, Chataigné G et al (2011) *In vitro* antitrypanosomal and antileishmanial activity of plants used in Benin in traditional medicine and bio-guided fractionation of the most active extract. J Ethnopharmacol 137:998–1002
- Betts JW, Wareham DW, Haswell SJ et al (2013) Antifungal synergy of theaflavin and epicatechin combinations against *Candida albicans*. J Microbiol Biotechnol 23:1322–1326
- Botta B, Vitali A, Menendez P (2005) Prenylated flavonoids: pharmacology and biotechnology. Curr Med Chem 12:713–739
- Bragança de Moraes CM, Melo DA, Santos RC et al (2012) Antiproliferative effect of catechin in GRX cells. Biochem Cell Biol 90:575–584
- Bulzomi P, Bolli A, Galluzzo P et al (2010) Naringenin and 17 $\beta$ -estradiol coadministration prevents hormone-induced human cancer cell growth. IUBMB Life 62:51–60
- Bulzomi P, Bolli A, Galluzzo P et al (2012) The naringenin-induced proapoptotic effect in breast cancer cell lines holds out against a high bisphenol A background. IUBMB Life 64:690–696
- Cai X, Zhang H, Tong D et al (2011) Corosolic acid triggers mitochondria and caspase-dependent apoptotic cell death in osteosarcoma MG-63 cells. Phytother Res 25:1354–1361

- Calderon-Montano JM, Burgos-Moron E et al (2011) A review on the dietary flavonoid kaempferol. *Mini-Rev Med Chem* 11(4):298–344
- Carey AN, Fisher DR, Rimando AM et al (2013) Stilbenes and anthocyanins reduce stress signaling in BV-2 mouse microglia. *J Agric Food Chem* 61:5979–5986
- Chai X-Y, Ren H-Y, Xu Z-R et al (2009) Investigation of two Flacourtiaceae plants: *Bennettiodendron leprosipes* and *Flacourtia ramontchi*. *Planta Med* 75:1246–1252
- Chang CL, Wang GJ, Zhang LJ et al (2010) Cardiovascular protective flavonolignans and flavonoids from *Calamus quiquesei*. *Phytochemistry* 71:271–279
- Chatrattanakunchai S, Fraser T, Stobart K (2000) Sesamin inhibits lysophosphatidylcholine acyltransferase in *Mortierella alpine*. *Biochem Soc Trans* 28:718–721
- Chen AY, Chen YC (2013) A review of the dietary flavonoid, kaempferol on human health and cancer chemoprevention. *Food Chem* 138(4):2099–2107
- Chen J-W, Zhu Z-Q, Hu T-X et al (2002) Structure–activity relationship of natural flavonoids in hydroxyl radical-scavenging effects. *Acta Pharmacol Sin* 23(7):667–672
- Chen CN, Wu CL, Lin JK (2004) Propolin C from propolis induces apoptosis through activating caspases, Bid and cytochrome *c* release in human melanoma cells. *Biochem Pharmacol* 67:53–66
- Chen J, Liu Y, Shi YP et al (2009) Determination of flavonoids in the flowers of *Paulownia tomentosa* by high-performance liquid chromatography. *J Anal Chem* 64:282–288
- Chen C-N, Hsiao C-J, Lee S-S et al (2012) Chemical modification and anticancer effect of prenylated flavanones from Taiwanese propolis. *Nat Prod Res* 26(2):116–124
- Chen C, Chen Z, Xu F et al (2013) Radio-protective effect of catalpol in cultured cells and mice. *J Radiat Res* 54:76–82
- Chen X, Mukwaya E, Wong M-S et al (2014) A systematic review on biological activities of prenylated flavonoids. *Pharm Biol (Lond, UK)* 52(5):655–660
- Cho JK, Ryu YB, Curtis-Long MJ et al (2012) Cholinesterase inhibitory effects of geranylated flavonoids from *Paulownia tomentosa* fruits. *Bioorg Med Chem* 20:2595–2602
- Cho JK, Curtis-Long MJ, Lee KH et al (2013) Geranylated flavonoids displaying SARS-CoV papain-like protease inhibition from the fruits of *Paulownia tomentosa*. *Bioorg Med Chem* 21:3051–3057
- Choi J, Shin KM, Park HJ et al (2004) Anti-inflammatory and antinociceptive effects of sinapyl alcohol and its glucoside syringin. *Planta Med* 70:1027–1032
- Chung BH, Lee JJ, Kim JD et al (2010) Angiogenic activity of sesamin through the activation of multiple signal pathways. *Biochem Biophys Res Commun* 391:254–260
- Cotin S, Calliste CA, Mazon MC et al (2012) Eight flavonoids and their potential as inhibitors of human cytomegalovirus replication. *Antivir Res* 96:181–186
- Cushnie TP, Lamb AJ (2005) Antimicrobial activity of flavonoids. *Int J Antimicrob Agents* 26:343–356
- Damtoft S, Jensen SR (1993) Tomentoside and 7-hydroxytomentoside, two new iridoid glucosides from *Paulownia tomentosa*. *Phytochemistry* 34:1636–1638
- de Aguiar SC, Zeoula LM, Franco SL et al (2013) Antimicrobial activity of Brazilian propolis extracts against rumen bacteria *in vitro*. *World J Microbiol Biotechnol* 29:1951–1959
- Deepak M, Handa SS (2000) Antiinflammatory activity and chemical composition of extracts of *Verbena officinalis*. *Phytother Res* 14:463–465
- Delormann D, Calis I, Ergun F et al (2000) Studies on the vascular effects of the fractions and phenolic compounds isolated from *Viscum album* ssp. *album*. *J Ethnopharmacol* 72:323–329
- Deurbina AVO, Martini ML, Fernandez B et al (1994) *In vitro* antispasmodic activity of peracetylated penstemoniside, aucubin and catalpol. *Planta Med* 60:512–515
- Diaz Lanza AM, Abad Martinez MJA, Matellano L et al (2001) Lignan and phenylpropanoid glycosides from *Phillyrea latifolia* and their *in vitro* anti-inflammatory activity. *Planta Med* 67:219–223
- Ding Y, Liang C, Yang SY et al (2010) Phenolic compounds from *Artemisia iwayomogi* and their effects on osteoblastic MC3T3-E1 cells. *Biol Pharm Bull* 33:1448–1453
- Epifano F, Genovese S, Menghini L et al (2007) Chemistry and pharmacology of oxyprenylated secondary plant metabolites. *Phytochemistry* 68:939–953
- Erbar C, Gülden C (2011) Ontogeny of the flowers in *Paulownia tomentosa*—a contribution to the recognition of the resurrected monogeneric family Paulowniaceae. *Flora* 206:205–218
- Estrada O, González-Guzmán JM, Salazar-Bookaman M et al (2011) Pomolic acid of *Licania pittieri* elicits endothelium-dependent relaxation in rat aortic rings. *Phytomedicine* 18:464–469
- Fraga CG, Oteiza PI (2011) Dietary flavonoids: role of (–)-epicatechin and related procyanidins in cell signaling. *Free Radic Biol Med* 51(4):813–823
- Franzyk H, Jensen SR, Thale Z et al (1999) Halohydrins of antirrhinoside—the correct structures of muralioside and epimuralioside. *J Nat Prod* 62:275–278
- Fu G, Pang H, Wong YH (2008) Naturally occurring phenyl-ethanoid glycosides: potential leads for new therapeutics. *Curr Med Chem* 15:2592–2613
- Fu Z, Yuskavage J, Liu D (2013) Dietary flavonol epicatechin prevents the onset of type 1 diabetes in nonobese diabetic mice. *J Agric Food Chem* 61:4303–4309
- Funke I, Melzig MF (2005) Effect of different phenolic compounds on  $\alpha$ -amylase activity: Screening by microplate-reader based kinetic assay. *Pharmazie* 60:796–797
- Galvez M, Martin-Cordero C, Ayuso MJ (2006) Pharmacological activities of phenylpropanoids glycosides. *Stud Nat Prod Chem* 33:675–718
- Ghosh A, Sarkar S, Mandal AK, Das N (2013) Neuroprotective role of nanoencapsulated quercetin in combating ischemia-reperfusion induced neuronal damage in young and aged rats. *PLoS One* 8(4):1–12
- Gong X, Zhang L, Jiang R et al (2014) Hepatoprotective effects of syringin on fulminant hepatic failure induced by D-galactosamine and lipopolysaccharide in mice. *J Appl Toxicol* 34:265–271
- Grael CFF, Vichnewski W, De Souza GEP et al (2000) A study of the trypanocidal and analgesic properties [of substances] from *Lychnophora grammongolense* (Duarte) Semir & Leitao Filho. *Phytother Res* 14(3):203–206
- Grecco Sdos S, Reimão JQ, Tempone AG et al (2012) *In vitro* antileishmanial and antitrypanosomal activities of

- flavanones from *Baccharis retusa* DC. (Asteraceae). *Exp Parasitol* 130:141–145
- Guan T, Qian Q, Tang X et al (2011) Maslinic acid, a natural inhibitor of glycogen phosphorylase, reduces cerebral ischemic injury in hyperglycaemic rats by GLT-1 up-regulation. *J Neurosci Res* 89:1829–1839
- Hac-Wydro K (2013) Studies on  $\beta$ -sitosterol and ceramide-induced alterations in the properties of cholesterol/sphingomyelin/ganglioside monolayers. *Biochim Biophys Acta* 1828:2460–2469
- Hakim EH, Fahriyati A, Kau MS et al (1999) Artoindonesianins A and B, two new prenylated flavones from the root of *Artocarpus champeden*. *J Nat Prod* 62:613–615
- Hakim EH, Asnizar Y, Aimi N et al (2002) Artoindonesianin P, a new prenylated flavone with cytotoxic activity from *Artocarpus lanceifolius*. *Fitoterapia* 73:668–673
- Hakim EH, Achmad SA, Juliawaty LD et al (2006) Prenylated flavonoids and related compounds of the Indonesian *Artocarpus* (Moraceae). *J Nat Med* 60:161–184
- Havsteen BH (2002) The biochemistry and medical significance of the flavonoids. *Pharmacol Ther* 96:67–202
- He J, Hu XP, Zeng Y et al (2011) Advanced research on acetoside for chemistry and bioactivities. *J Asian Nat Prod Res* 13:449–464
- Hidalgo M, Martin-Santamaria S, Recio I et al (2012) Potential anti-inflammatory, anti-adhesive, anti/estrogenic, and angiotensin-converting enzyme inhibitory activities of anthocyanins and their gut metabolites. *Genes Nutr* 7:295–306
- Hirano T, Higa S, Arimitsu J et al (2006) Luteolin, a flavonoid, inhibits AP-1 activation by basophils. *Biochem Biophys Res Commun* 340(1):1–7
- Holubová P, Šmejkal K (2011) Changes in the level of bioactive compounds in *Paulownia tomentosa* fruits. *J Liq Chromatogr Relat Technol* 34:276–288
- Hong D, Yang H, Jin C et al (1998) Scrophulariaceae through Gesneriaceae. *Flora China* 18:8–10
- Horland H, Fujiwara Y, Takemura K et al (2013) Corosolic acid impairs tumor development and lung metastasis by inhibiting the immunosuppressive activity of myeloid-derived suppressor cells. *Mol Nutr Food Res* 57(6):1046–1054
- Hošek J, Závalová V, Šmejkal K et al (2010) Effect of diplacene on LPS-induced inflammatory gene expression in macrophages. *Folia Biol (Praha)* 56:124–130
- Hošek J, Toniolo A, Neuwirth O et al (2013) Prenylated and geranylated flavonoids increase production of reactive oxygen species in mouse macrophages but inhibit the inflammatory response. *J Nat Prod* 76(9):1586–1591
- Huang C, Cui Y, Ji L et al (2013a) Catalpol decreases peroxynitrite formation and consequently exerts cardioprotective effects against ischemia/reperfusion insult. *Pharm Biol* 51:463–473
- Huang D, Hu Z, Yu Z (2013b) Eleutheroside B or E enhances learning and memory in experimentally aged rats. *Neural Regen Res* 8:1103–1112
- Huang YB, Lin MW, Chao Y et al (2014) Anti-oxidant activity and attenuation of bladder hyperactivity by the flavonoid compound kaempferol. *Int J Urol* 21(1):94–98
- Ibrahim NA, El-Hawary SS, Mohammed MMD et al (2013) Chemical composition, antimicrobial activity of the essential oil of the flowers of *Paulownia tomentosa* (Thunb.) Steud. growing in Egypt. *J Appl Sci Res* 9(4):3228–3232
- Iizuka T, Nagai M, Moriyama H et al (2005) Antiplatelet aggregatory effects of the constituents isolated from the flower of *Carthamus tinctorius*. *Nat Med (Tokyo, Jpn)* 59:241–244
- Ina H, Ono M, Sashida Y et al (1987) (+)-Piperitol from *Paulownia tomentosa*. *Planta Med* 53:504
- Jayaraman J, Jesudoss VA, Menon VP et al (2012) Anti-inflammatory role of naringenin in rats with ethanol induced liver injury. *Toxicol Mech Methods* 22:568–576
- Jiang TF, Du X, Shi YP et al (2004) Determination of flavonoids from *Paulownia tomentosa* (Thunb.) Steud. by micellar electrokinetic capillary electrophoresis. *Chromatographia* 59:255–258
- Jiménez C, Riguera R (1994) Phenylethanoid glycosides in plants: structure and biological activity. *Nat Prod Rep* 11:591–606
- Jin L, Xue HY, Jin LJ et al (2008) Antioxidant and pancreas-protective effect of aucubin on rats with streptozotocin-induced diabetes. *Eur J Pharmacol* 582:162–167
- Jordao CO, Vichnewski W, Petto de Souza GE et al (2004) Trypanocidal activity of chemical constituents from *Lychophora salicifolia* Mart. *Phytother Res* 18(4):332–334
- Jung S, Moon HI, Ohk J et al (2012) Inhibitory effect and mechanism on antiproliferation of isoaiplicolide tiglate (PCAC) from *Paulownia coreana*. *Molecules* 17:5945–5951
- Kadota S, Basnet P, Hase K et al (1994) Matteuorientate A and B, two new and potent aldose reductase inhibitors from *Matteuccia orientalis* (Hook.) Trev. *Chem Pharm Bull* 42(8):1712–1714
- Kang KH, Jang SJ, Kim BK et al (1994) Antibacterial phenylpropanoid glycosides from *Paulownia tomentosa* Steud. *Arch Pharm Res* 17:470–475
- Kang KH, Huh H, Kim BK et al (1999) An antiviral furanoquinone from *Paulownia tomentosa* Steud. *Phytother Res* 13:624–626
- Kang DG, Lee YS, Kim HJ et al (2003) Angiotensin converting enzyme inhibitory phenylpropanoid glycosides from *Clerodendron trichotomum*. *J Ethnopharmacol* 89:151–154
- Kawamura F, Ohara S, Nishida A (2004) Antifungal activity of constituents from the heartwood of *Gmelina arborea*: part 1. Sensitive antifungal assay against Basidiomycetes. *Holzforschung* 58:189–192
- Khachatoorian R, Arumugaswami V, Raychaudhuri S et al (2012) Divergent antiviral effects of bioflavonoids on the hepatitis C virus life cycle. *Virology* 433:346–355
- Khan MK, Zill-E-Huma DO (2014) A comprehensive review on flavanones, the major citrus polyphenols. *J Food Compos Anal* 33(1):85–104
- Khan MM, Ishrat T, Ahmad A et al (2010) Sesamin, attenuates behavioural, biochemical and histological alterations induced by reversible middle cerebral artery occlusion in the rats. *Chem Biol Interact* 183:255–263
- Kim HJ, Woo ER, Shin CG et al (2001) HIV-1 integrase inhibitory phenylpropanoid glycosides from *Clerodendron trichotomum*. *Arch Pharm Res* 24:286–291
- Kim SJ, Kwon DY, Kim YS et al (2010a) Peroxyl radical scavenging capacity of extracts and isolated components



- from selected medicinal plants. Arch Pharm Res 33:867–873
- Kim SK, Cho SB, Moon HI et al (2010b) Neuroprotective effects of a sesquiterpene lactone and flavanones from *Paulownia tomentosa* Steud against glutamate-induced neurotoxicity in primary cultured rat cortical cells. Phytother Res 24:1898–1900
- Kim JK, Lee YS, Kim SH et al (2011) Inhibition of aldose reductase by phenylethanoid glycoside isolated from the seeds of *Paulownia coreana*. Biol Pharm Bull 34:160–163
- Kobayashi S, Asai T, Fujimoto Y et al (2008) Anti-herbivore structures of *Paulownia tomentosa*: morphology, distribution, chemical constituents and changes during shoot and leaf development. Ann Bot 101:1035–1047
- Koch CE, Ganjam GK, Steger J et al (2013) The dietary flavonoids naringenin and quercetin acutely impair glucose metabolism in rodents possibly via inhibition of hypothalamic insulin signalling. Br J Nutr 109:1040–1051
- Kollár P, Bárta T, Závalová V et al (2011) Geranylated flavanone tomentodiplacone B inhibits proliferation of human monocytic leukaemia (THP-1) cells. Br J Pharmacol 162(7):1534–1541
- Kong LD, Wolfender JL, Cheng CH et al (1999) Xanthine oxidase inhibitors from *Brandisia hancei*. Planta Med 65:744–746
- Koo KA, Sung SH, Park JH et al (2005) *In vitro* neuroprotective activities of phenylethanoid glycosides from *Callicarpa dichotoma*. Planta Med 71:778–780
- Krishnaswamy M, Purushothaman KK (1980) Plumbagin: a study of its anticancer, antibacterial and antifungal properties. Indian J Exp Biol 18:876–877
- Kumar M, Rawat P, Khan MF et al (2010) Phenolic glycosides from *Dodecadenia grandiflora* and their glucose-6-phosphatase inhibitory activity. Fitoterapia 81:475–479
- Kumazawa S, Ueda R, Hamasaka T et al (2007) Antioxidant prenylated flavonoids from poplars collected in Okinawa, Japan. J Agric Food Chem 55(19):7722–7725
- Kurkin VA (2003) Phenylpropanoids from medicinal plants: distribution, classification, structural analysis, and biological activity. Chem Nat Compd 39:123–153
- Kurkin VA, Dubishchev AV, Ezhkov VN et al (2006) Antidepressant activity of some phytopharmaceuticals and phenylpropanoids. Pharm Chem J 40:614–619
- Kurkina AV, Khusainova AI, Daeva ED et al (2011) Flavonoids from *Tanacetum vulgare* flowers. Chem Nat Compd 47:284–285
- Kuzuyama T, Noel JP, Richard SB (2005) Structural basis for the promiscuous biosynthetic prenylation of aromatic natural products. Nature 435:983–987
- Lee JH, Lee JY, Park JH et al (2007) Immunoregulatory activity of daucosterol, a beta-sitosterol glycoside, induces protective Th1 immune response against disseminated Candidiasis in mice. Vaccine 25:3834–3840
- Lee JS, Miyashiro H, Nakamura N et al (2008) Two new triterpenes from the rhizome of *Dryopteris crassirhizoma*, and inhibitory activities of its constituents on human immunodeficiency virus-1 protease. Chem Pharm Bull 56:711–714
- Lee S-H, Jung MJ, Heo S-I et al (2009) Anti-inflammatory effect and HPLC analysis of extract from edible *Cirsium setidens*. J Korean Soc Appl Biol Chem 52:437–442
- Lee JH, Lee HJ, Choung MG (2011) Anthocyanin compositions and biological activities from the red petals of Korean edible rose (*Rosa* hybrid cv. Noblered). Food Chem 129:272–278
- Lee Y, Ryu YB, Youn H-S et al (2014) Structural basis of sialidase in complex with geranylated flavonoids as potent natural inhibitors. Acta Cryst D70:1357–1365
- Li DQ, Duan YL, Bao YM et al (2004) Neuroprotection of catalpol in transient global ischemia in gerbils. Neurosci Res 50:169–177
- Li R, Cai L, Xie X-f et al (2010) 7,3'-Dimethoxy hesperetin induces apoptosis of fibroblast-like synoviocytes in rats with adjuvant arthritis through caspase 3 activation. Phytother Res 24(12):1850–1856
- Li Y-L, Wu L, Ouyang D-W et al (2011) Phenolic Compounds of *Abies nephrolepis* and their NO production inhibitory activities. Chem Biodivers 8(12):2299–2309
- Li R, Cai L, Ren D-y et al (2012) Therapeutic effect of 7,3'-dimethoxy hesperetin on adjuvant arthritis in rats through inhibiting JAK2-STAT3 signal pathway. Int Immunopharmacol 14(2):157–163
- Li R, Cai L, Xie X-f et al (2013) 7,3'-dimethoxy hesperetin inhibits inflammation by inducing synovial apoptosis in rats with adjuvant-induced arthritis. Immunopharmacol Immunotoxicol 35(1):139–146
- Lin LC, Wang YH, Hou YC et al (2006) The inhibitory effect of phenylpropanoid glycosides and iridoid glucosides on free radical production and  $\beta$ 2-integrin expression in human leucocytes. J Pharm Pharmacol 58:129–135
- Lin J-A, Fang S-C, Wu C-H et al (2011) Anti-inflammatory effect of the 5,7,4'-Trihydroxy-6-geranylfavanone isolated from the fruit of *Artocarpus communis* in S100B-induced human monocytes. J Agric Food Chem 59(1):105–111
- Liu KY, Wu Y-C, Liu I-M et al (2008) Release of acetylcholine by syringin, an active principle of *Eleutherococcus senticosus*, to raise insulin secretion in Wistar rats. Neurosci Lett 434:195–199
- Loke WM, Proudfoot JM, Hodgson JM et al (2010) Specific dietary polyphenols attenuate atherosclerosis in apolipoprotein E-knockout mice by alleviating inflammation and endothelial dysfunction. Arterioscler Thromb Vasc Biol 30:749–757
- Lopez-Lazaro M (2009) Distribution and biological activities of the flavonoid luteolin. Mini Rev Med Chem 9(1):31–59
- Lopez-Lazaro M, Martin-Cordero C, Cortes F et al (2000) Cytotoxic activity of flavonoids and extracts from *Retama sphaerocarpa* Boissier. Z Naturforsch C (J Biosci) 55(1/2):40–43
- Losi G, Puia G, Garzon G et al (2004) Apigenin modulates GABAergic and glutamatergic transmission in cultured cortical neurons. Eur J Pharmacol 502:41–46
- Lu XY, Li YH, Xiao XW, Li XB (2013) Inhibitory effects of luteolin on human gastric carcinoma xenografts in nude mice and its mechanism. Zhonghua Yi Xue Za Zhi 93:142–146
- Mankovskaia A, Lévesque CM, Prakki A (2013) Catechin-incorporated dental copolymers inhibit growth of *Streptococcus mutans*. J Appl Oral Sci 21:203–207
- Martini ND, Katerere DRP, Eloff JN (2004) Biological activity of five antibacterial flavonoids from *Combretum*



- erythrophyllum* (Combretaceae). J Ethnopharmacol 93(2–3):207–212
- Marzocchella L, Fantini M et al (2011) Dietary flavonoids: molecular mechanisms of action as anti-inflammatory agents. Recent Pat Inflamm Allergy Drug Discov 5(3):200–220
- Mastuda H, Morikawa T, Ueda K et al (2002) Structural requirements of flavonoids for inhibition of antigen-induced degranulation, TNF- $\alpha$  and IL-4 production from RBL-2H3 cells. Bioorg Med Chem 10(10):3123–3128
- Matsubara Y, Yusa T, Sawab A et al (1991) Studies on physiologically active substances in citrus fruit peel. Part XX. Structure and physiological activity of phenyl propanoid glycosides in lemon (*Citrus limon* Burm. f.) peel. Agric Biol Chem 55(3):647–650
- Matsuda H, Morikawa T, Ando S et al (2003) Structural requirements of flavonoids for nitric oxide production inhibitory activity and mechanism of action. Bioorg Med Chem 11(9):1995–2000
- Milligan SR, Kalita JC, Pocock V et al (2000) The endocrine activities of 8-prenylnaringenin and related hop (*Humulus lupulus* L.) flavonoids. J Clin Endocrinol Metab 85(12):4912–4915
- Min YS, Yim SH, Bail KL et al (2005) The effects of apigenin-7-O- $\beta$ -D-glucuronopyranoside on reflux oesophagitis and gastritis in rats. Auton Autacoid Pharmacol 25:85–91
- Mishra N, Rizvi SI (2012) Quercetin modulates Na<sup>+</sup>/K<sup>+</sup> ATPase and sodium hydrogen exchanger in type 2 diabetic erythrocytes. Cell Mol Biol 58(1):148–152
- Miyazawa M, Okuno Y, Nakamura S-I et al (2000) Antimutagenic activity of flavonoids from *Pogostemon cablin*. J Agric Food Chem 48(3):642–647
- Moneriz G, Mestres J, Bautista JM et al (2011) Multi-targeted activity of maslinic acid as an antimalarial natural compound. FEBS J 278:2951–2961
- Moon HI, Zee OP (2001) Anticancer compound of *Paulownia tomentosa*. Nat Prod Sci 7:21–22
- Murphy BT, Cao S, Norris A et al (2005) Cytotoxic flavanones of *Schizolaena hystrix* from the Madagascar rainforest. J Nat Prod 68:417–419
- Murphy BT, Cao S, Norris A et al (2006) Cytotoxic compounds of *Schizolaena hystrix* from the Madagascar rainforest. Planta Med 72(13):1235–1238
- Nakano D, Kwak CJ, Fujii K et al (2006) Sesamin metabolites induce an endothelial nitric oxide-dependent vasorelaxation through their antioxidative property-independent mechanisms: possible involvement of the metabolites in the antihypertensive effect of sesamin. J Pharmacol Exp Ther 318:328–335
- Nascimento IR, Murata AT, Bortoli SA et al (2004) Insecticidal activity of chemical constituents from *Aristolochia pubescens* against *Anticarsia gemmatilis* larvae. Pest Manag Sci 60:413–416
- Navrátilová A, Schneiderová K, Veselá D et al (2013) Minor C-glycosylated flavanones from *Paulownia tomentosa* fruits with MRSA antibacterial activity. Phytochemistry 89:104–113
- Niu H-S, Hsu F-L, Liu I-M et al (2007) Increase of  $\beta$ -endorphin secretion by syringin, an active principle of *Eleutherococcus senticosus*, to produce antihyperglycemic action in type 1-like diabetic rats. Hormone Metab Res 39:894–898
- Niu H-S, Hsu F-L, Liu I-M (2008a) Role of sympathetic tone in the loss of syringin-induced plasma glucose lowering action in conscious Wistar rats. Neurosci Lett 445:113–116
- Niu H-S, Liu I-M, Cheng J-T et al (2008b) Hypoglycemic effect of syringin from *Eleutherococcus senticosus* in streptozotocin-induced diabetic rats. Planta Med 74:109–113
- Ogungbe IV, Erwin WR, Setzer WN (2014) Antileishmanial phytochemical phenolics: molecular docking to potential protein targets. J Mol Graph Model 48:105–117
- Omosa L, Amugune B, Ndunda B et al (2014) Antimicrobial flavonoids and diterpenoids from *Dodonaea angustifolia*. S Afr J Bot 91:58–62
- Oprea E, Radulescu V, Chiliment S (2004) The analysis of the volatile and semi-volatile compounds of the *Paulownia tomentosa* flowers by gas chromatography coupled with mass spectrometry. Revista de chimie 55:410–412
- Ota M, Azuma T, Onodera S et al (1993) The chemistry of color changes in kiri wood (*Paulownia tomentosa* Steud.) III. A new caffeic acid sugar ester from Kiri wood. Mozukai Gakkaishi 39:479–485
- Pan J, Yuan C, Lin C et al (2003) Pharmacological activities and mechanisms of natural phenylpropanoid glycosides. Pharmazie 58:767–775
- Pan J-Y, Chen S-L, Yang M-H et al (2009) An update on lignans: natural products and synthesis. Nat Prod Rep 26(10):1251–1292
- Pan Y, Morikawa T, Ninomiya K et al (2010) Bioactive constituents from Chinese natural medicines. XXXVI. Four new acylated phenylethanoid oligoglycosides, Kankanosides J1, J2, K1, and K2, from stems of *Cistanche tubulosa*. Chem Pharm Bull 58:575–578
- Pan SY, Zhou SF, Gao SH et al (2013) New perspectives on how to discover drugs from herbal medicines: CAM's outstanding contribution to modern therapeutics. Evid Based Complement Altern Med 2013:1–25, doi:10.1155/2013/627375
- Paneerselvam M, Kawaraguchi Y, Horikawa YT et al (2010) Effect of epicatechin and naloxone on cardioprotective phenotype. FASEB J 24 (Meeting Abstract Supplement) 1029.8
- Panossian A, Wagner H (2005) Stimulating effect of adaptogens: an overview with particular reference to their efficacy following single dose administration. Phytother Res 19:819–838
- Panossian A, Kocharian A, Matinian K et al (1998) Pharmacological activity of phenylpropanoids of the mistletoe, *Viscum album*, host. *Pyrus caucasica*. Phytomedicine 5:11–17
- Papoutsis Z, Kassi E, Mitakou S et al (2006) Acteoside and martynoside exhibit estrogenic/antiestrogenic properties. J Steroid Biochem Mol Biol 98:63–71
- Park KS (2013) Aucubin, a naturally occurring iridoid glycoside inhibits TNF- $\alpha$ -induced inflammatory responses through suppression of NF- $\kappa$ B activation in 3T3-L1 adipocytes. Cytokine 62:407–412
- Park Y, Kong JY, Cho H (2009) A Furanquinone from *Paulownia tomentosa* stem for a new cathepsin K inhibitor. Phytother Res 23:1485–1488
- Patel D, Shukla S et al (2007) Apigenin and cancer chemoprevention: progress, potential and promise (review). Int J Oncol 30(1):233–245

- Peluso MR, Miranda CL, Hobbs DJ et al (2010) Xanthohumol and related prenylated flavonoids inhibit inflammatory cytokine production in LPS-activated THP-1 monocytes: structure-activity relationships and in Silico binding to myeloid differentiation protein-2 (MD-2). *Planta Med* 76(14):1536–1543
- Pelzer LE, Guardia T, Juárez AO et al (1998) Acute and chronic antiinflammatory effects of plant flavonoids. *Farmacol* 53(6):421–424
- Persson IA, Persson K, Andersson RG (2009) Effect of *Vaccinium myrtillus* and its polyphenols on angiotensin-converting enzyme activity in human endothelial cells. *J Agric Food Chem* 57:4626–4629
- Phommart S, Sutthivaiyakit P, Chimnoi N et al (2005) Constituents of the leaves of *Macaranga tanarius*. *J Nat Prod* 68:927–930
- Pile JE, Navalta JW, Davis CD et al (2013) Interventional effects of plumbagin on experimental ulcerative colitis in mice. *J Nat Prod* 76:1001–1006
- Plaza M, Pozzo T, Liu J et al (2014) Substituent effects on in vitro antioxidizing properties, stability, and solubility in flavonoids. *J Agric Food Chem* 62(15):3321–3333
- Plouvier V (1971) The heterosides of *Catalpa bignonioides* Walt. (Bignoniaceae). *Comp Rend Acad Sci* 272(D):1443–1446
- Prince Vijaya Singh J, Selvendiran K, Mumtaz Banu S et al (2004) Protective role of apigenin on the status of lipid peroxidation and antioxidant defense against hepatocarcinogenesis in Wistar albino rats. *Phytomedicine* 11:309–314
- Psotová J, Chlopčíková S, Miketová P et al (2004) Chemoprotective effect of plant phenolics against anthracycline-induced toxicity on rat cardiomyocytes. Part III. Apigenin, baicalein, kaempferol, luteolin and quercetin. *Phytother Res* 18:516–521
- Raghukumar R, Vali L, Watson D et al (2010) Antimethicillin-resistant *Staphylococcus aureus* (MRSA) activity of “pacific propolis” and isolated prenylflavanones. *Phytother Res* 24:1181–1187
- Rao YK, Lee MJ, Chen K et al (2011) Insulin-mimetic action of rhoifolin and cosmosiin isolated from *Citrus grandis* (L.) osbeck leaves: enhanced adiponectin secretion and insulin receptor phosphorylation in 3T3-L1 cells. *Evid Based Complement Altern Med* 2011:1–9, doi:10.1093/ecam/nep204
- Remya C, Dileep KV, Tintu I et al (2012) Design of potent inhibitors of acetylcholinesterase using morin as the starting compound. *Front Life Sci* 6(3–4):107–117
- Reyes-Zurita FJ, Pachón-Peña G, Lizárraga D et al (2011) The natural triterpene maslinic acid induces apoptosis in HT29 colon cancer cells by a JNK-p53-dependent mechanism. *BMC Cancer* 11:154
- Rodríguez J, Yanez J, Vicente V et al (2002) Effects of several flavonoids on the growth of B16F10 and SK-MEL-1 melanoma cell lines: relationship between structure and activity. *Melanoma Res* 12(2):99–107
- Romano B, Pagano E et al (2013) Novel insights into the pharmacology of flavonoids. *Phytother Res* 27(11):1588–1596
- Ross JA, Kasum CM (2002) Dietary flavonoids: bioavailability, metabolic effects, and safety. *Annu Rev Nutr* 22:19–34
- Rosselli S, Bruno M, Maggio A et al (2011) Cytotoxic geranylflavonoids from *Bonannia graeca*. *Phytochemistry* 72:942–945
- Russo M, Spagnuolo C et al (2012) The flavonoid quercetin in disease prevention and therapy: facts and fancies. *Biochem Pharmacol* 83(1):6–15
- Salem MM, Capers J, Rito S et al (2011) Antiparasitic activity of C-geranyl flavonoids from *Mimulus bigelovii*. *Phytother Res* 25(8):1246–1249
- Sánchez-Tena S, Alcarraz-Vizán G, Marín S et al (2013) Epicatechin gallate impairs colon cancer cell metabolic productivity. *J Agric Food Chem* 61:4310–4317
- Santhakumari G, Rathinam K, Seshadri C (1978) Angicoagulant activity of plumbagin. *Indian J Exp Biol* 16:485–487
- Sato M, Murakami K, Uno M et al (2013) Site-specific inhibitory mechanism for amyloid  $\beta$ 42 aggregation by catechol-type flavonoids targeting the Lys residues. *J Biol Chem* 288(32):23212–23224
- Scambia G, Ranelletti FO, Panici P et al (1994) Quercetin potentiates the effect of Adriamycin in a multidrug-resistant MCF-7 human breast-cancer cell line: P-glycoprotein as a possible target. *Cancer Chemother Pharmacol* 34(6):459–464
- Schilling G, Hügel M, Mayer W (1982) Verbascoside and isoverbascoside from *Paulownia tomentosa* Steud. *Zeit Naturforsch* 37(B):1633–1635
- Schinella G, Aquila S, Dade M et al (2008) Anti-inflammatory and apoptotic activities of pomolic acid isolated from *Cecropia pachystachya*. *Planta Med* 74:215–220
- Schneiderová K, Šlapetová T, Hrabal R et al (2013) Tomentomimulol and mimulone B: two new C-geranylated flavonoids from *Paulownia tomentosa* fruits. *Nat Prod Res* 27:613–618
- Scogin R (1980) Anthocyanins of the Bignoniaceae. *Biochem Syst Ecol* 8:273–276
- Shan JZ, Xuan YY, Ruan SQ et al (2011) Proliferation-inhibiting and apoptosis-inducing effects of ursolic acid and oleanolic acid on multi-drug resistance cancer cells *in vitro*. *Chin J Integr Med* 17:607–611
- Sharma U, Bala M, Kumar N et al (2012) Immunomodulatory active compounds from *Tinospora cordifolia*. *J Ethnopharmacol* 141:918–926
- She G-M, Zhang Y-J, Yang C-R (2013) A new phenolic constituent and a cyanogenic glycoside from *Balanophora involucreata* (Balanophoraceae). *Chem Biodivers* 10:1081–1087
- Shi Z-H, Li N-G, Tang Y-P et al (2012) Metabolism-based synthesis, biologic evaluation and SARs analysis of O-methylated analogs of quercetin as thrombin inhibitors. *Eur J Med Chem* 54:210–222
- Shieh JP, Cheng KC, Chung HH et al (2011) Plasma glucose lowering mechanisms of catalpol, an active principle from roots of *Rehmannia glutinosa*, in streptozotocin-induced diabetic rats. *J Agric Food Chem* 59:3747–3753
- Shukla S, Gupta S (2009) Role of apigenin in human health and disease. In: Preedy VR (ed) *Beer in health and disease prevention*, Academic Press, San Diego, pp e202–e216
- Si C, Deng X, Liu Z (2008a) Structure and activity relationship of antioxidant flavonoids from leaves of *Paulownia tomentosa* var. *tomentosa*. In: 2nd international papermaking and environment conference, Tianjin University of Science and Technology, Tianjin, pp 263–266

- Si CL, Deng XJ, Liu Z et al (2008b) Studies on the phenylethanoid glycosides with anti-complement activity from *Paulownia tomentosa* var. *tomentosa* wood. *J Asian Nat Prod Res* 10:1003–1008
- Si C, Deng X, Xu Q et al (2008c) Characterization of phenolic acids and antioxidant activities of *Paulownia tomentosa* var. *tomentosa* leaves. In: Proceedings of the international conference on pulping, papermaking and biotechnology, pp 31–33
- Si CL, Liu Z, Kim JK et al (2008d) Structure elucidation of phenylethanoid glycosides from *Paulownia tomentosa* Steud. var. *tomentosa* wood. *Holzforschung* 62:197–200
- Si CL, Wu L, Zhu ZY et al (2009) Apigenin derivatives from *Paulownia tomentosa* Steud. var. *tomentosa* stem barks. *Holzforschung* 63:440–442
- Si CL, Lu YY, Hu HY et al (2011a) Evaluation of total phenolics, flavonoids and anti-inflammatory property of ethanolic extracts of *Paulownia tomentosa* var. *tomentosa* bark. *Planta Med* 77:SL53
- Si CL, Lu YY, Qin PP et al (2011b) Phenolic extractives with chemotaxonomic significance from the bark of *Paulownia tomentosa* var. *tomentosa*. *BioResources* 6:5086–5098
- Si CL, Shen T, Jiang YY et al (2013) Antioxidant properties and neuroprotective effects of isocampneoside II on hydrogen peroxide-induced oxidative injury in PC12 cells. *Food Chem Toxicol* 59:145–152
- Sivakumar G, Vail DR, Nair V et al (2009) Plant-based corosolic acid: future anti-diabetic drug? *Biotechnol J* 4:1704–1711
- Šmejkal K (2014) Cytotoxic potential of *C*-prenylated flavonoids. *Phytochem Rev* 13:245–275
- Šmejkal K, Grycová L, Marek R et al (2007a) *C*-geranyl compounds from *Paulownia tomentosa* Fruits. *J Nat Prod* 70:1244–1248
- Šmejkal K, Holubová P, Zima A et al (2007b) Antiradical activity of *Paulownia tomentosa* (Scrophulariaceae) extracts. *Molecules* 12:1210–1219
- Šmejkal K, Babula P, Šlapetová T et al (2008a) Cytotoxic activity of *C*-geranyl compounds from *Paulownia tomentosa* fruits. *Planta Med* 74:1488–1491
- Šmejkal K, Chudík S, Klouček P et al (2008b) Antibacterial *C*-geranylflavonoids from *Paulownia tomentosa* (Scrophulariaceae) fruits. *J Nat Prod* 71:706–709
- Šmejkal K, Svačinová J, Šlapetová T et al (2010) Cytotoxic activities of several geranyl-substituted flavanones. *J Nat Prod* 73:568–572
- Sticher O, Lahloub MF (1982) Phenolic glycosides of *Paulownia tomentosa* bark. *Planta Med* 46:145–148
- Sudsai T, Wattanapiromsakul C, Tewtrakul S (2013) Inhibition of nitric oxide production by compounds from *Boesenbergia longiflora* using lipopolysaccharide-stimulated RAW264.7 macrophage cells. *Songklanakarin J Sci Technol* 35(3):317–323
- Sumsakul W, Plensuriyakarn T, Chaijaroenkul W et al (2014) Antimalarial activity of plumbagin in vitro and in animal models. *BMC Complement Altern Med* 14:15
- Sun Y, Zang Z, Zhong L et al (2013) Identification of adiponectin receptor agonist utilizing a fluorescence polarization based high throughput assay. *PLoS One* 8(5):e63354
- Suolinna EM, Buchsbaum RN, Racker E (1975) Effect of flavonoids on aerobic glycolysis and growth of tumor cells. *Cancer Res* 35(7):1865–1872
- Takahashi K, Nakagawa T (1966) Studies on constituents of medicinal plants. VII. The stereochemistry of paulownin and isopaulownin. *Chem Pharm Bull* 14:641–647
- Takamatsu S, Galal AM, Ross SA et al (2003) Antioxidant effect of flavonoids on DCF production in HL-60 cells. *Phytother Res* 17(8):963–966
- Tanaka J, Kadekaru T, Ogawa K et al (2013) Maqui berry (*Aristotelia chilensis*) and the constituent delphinidin glycoside inhibit photoreceptor cell death induced by visible light. *Food Chem* 139:129–137
- Tang R, Chen K, Cosentino M et al (1994) Apigenin-7-*O*- $\beta$ -D-glucopyranoside, an anti-HIV principle from *Kummerowia striata*. *Bioorg Med Chem Lett* 4:455–458
- Taub PR, Ramirez-Sanchez I, Ciaraldi TP et al (2012) Alterations in skeletal muscle indicators of mitochondrial structure and biogenesis in patients with type 2 diabetes and heart failure: effects of epicatechin rich cocoa. *Clin Transl Sci* 5(1):43–47
- Teixeira MD, Souza CM, Menezes AP et al (2013) Catechin attenuates behavioral neurotoxicity induced by 6-OHDA in rats. *Pharmacol Biochem Behav* 110:1–7
- Tian L-W, Pei Y, Zhang Y-J et al (2009) 7-*O*-methylkaempferol and -quercetin glycosides from the whole plant of *Nervilia fordii*. *J Nat Prod* 72(6):1057–1060
- Tohda C, Ichimura M, Bai Y et al (2008) Inhibitory effects of *Eleutherococcus senticosus* extracts on amyloid  $\beta$ (25–35)-induced neuritic atrophy and synaptic loss. *J Pharmacol Sci* 107:329–339
- Tozuka H, Ota M, Kofujita H et al (2005) Synthesis of dihydroxyphenacyl glycosides for biological and medicinal study:  $\beta$ -oxoacetoside from *Paulownia tomentosa*. *J Wood Sci* 51:48–59
- Trusheva B, Popova M, Koendhori EB et al (2011) Indonesian propolis: chemical composition, biological activity and botanical origin. *Nat Prod Res* 25:606–613
- Ullevig SL, Zhao Q, Zamora D et al (2011) Ursolic acid protects diabetic mice against monocyte dysfunction and accelerated atherosclerosis. *Atherosclerosis* 219:409–416
- Vareed SK, Reddy MK, Schutzki RE (2006) Anthocyanins in *Cornus alternifolia*, *Cornus controversa*, *Cornus kousa* and *Cornus florida* fruits with health benefits. *Life Sci* 78:777–784
- Vasconcelos FC, Gattass CR, Rumjanek VM et al (2007) Pomolic acid-induced apoptosis in cells from patients with chronic myeloid leukemia exhibiting different drug resistance profile. *Invest New Drugs* 25:525–533
- Wang J, Yang Z, Lin L et al (2012) Protective effect of naringenin against lead-induced oxidative stress in rats. *Biol Trace Elem Res* 146:354–359
- Wang Y-M, Xu M, Wang D et al (2013) Anti-inflammatory compounds of “Qin-Jiao”, the roots of *Gentiana dahurica* (Gentianaceae). *J Ethnopharmacol* 147(2):341–348
- Wei YJ, Tsai KS, Lin LC et al (2011) Catechin stimulates osteogenesis by enhancing PP2A activity in human mesenchymal stem cells. *Osteoporos Int* 22(5):1469–1479
- Weidmann AE (2012) Dihydroquercetin: more than just an impurity? *Eur J Pharmacol* 684(1–3):19–26

- Wilkinson K, Boyd JD, Glicksman M et al (2011) A high-content drug screen identifies ursolic acid as an inhibitor of amyloid- $\beta$  interactions with its receptor CD36. *J Biol Chem* 286:34914–34922
- Wollenweber E, Wehde R, Christ M et al (2008) Surface flavonoids in *Catalpa ovata*, *Greyia sutherlandii* and *Paulownia tomentosa*. *Nat Prod Commun* 3:1285–1287
- Wong KC, Haq Ali DM, Boey PL (2012) Chemical constituents and antibacterial activity of *Melastoma malabathricum* L. *Nat Prod Res* 26:609–618
- Wu A, Lin C, Zhao X et al (2012) Spectroscopic study on interaction between cistanoside F and bovine serum albumin. *Zhongguo Zhong Yao Za Zhi* 37:1392–1398
- Xiong Q, Kadota S, Tani T et al (1996) Antioxidative effects of phenylethanoids from *Cistanche deserticola*. *Biol Pharm Bull* 19:1580–1585
- Xiong QB, Hase K, Tezuka Y et al (1998) Hepatoprotective activity of phenylethanoids from *Cistanche deserticola*. *Planta Med* 64:120–125
- Xue HY, Lu YN, Fang XM et al (2012) Neuroprotective properties of aucubin in diabetic rats and diabetic encephalopathy rats. *Mol Biol Rep* 39:9311–9318
- Yang Y-L, Hsu H-T, Wang K-H et al (2011) Hesperetin-7,3'-O-dimethylether selectively inhibits phosphodiesterase 4 and effectively suppresses ovalbumin-induced airway hyper-responsiveness with a high therapeutic ratio. *J Biomed Sci (Lond, UK)* 18:84
- Yang X, Yuan J, Wan J (2012) Cytotoxic phenolic glycosides from *Boschniakia himalaica*. *Chem Nat Comp* 48(4):555–558
- Yasuda K, Sakaki T (2012) How is sesamin metabolised in the human liver to show its biological effects? *Expert Opin Drug Metab Toxicol* 8:93–102
- Yazaki K, Sasaki K, Tsurumaru Y (2009) Prenylation of aromatic compounds, a key diversification of plant secondary metabolites. *Phytochemistry* 70:1739–1745
- Yoder BJ, Cao S, Norris A et al (2007) Antiproliferative prenylated stilbenes and flavonoids from *Macaranga alnifolia* from the Madagascar rainforest. *J Nat Prod* 70:342–346
- Yoo DY, Choi JH, Kim W, Nam SM, Jung HY, Kim JH, Won M-H, Yoon YS, Hwang IK (2013) Effects of luteolin on spatial memory, cell proliferation, and neuroblast differentiation in the hippocampal dentate gyrus in a scopolamine-induced amnesia model. *Neurol Res* 35:813–820
- Yoon JS, Chae MK, Jang SY, Lee SY, Lee EJ (2012) Antifibrotic effects of quercetin in primary orbital fibroblasts and orbital fat tissue cultures of graves' orbitopathy. *Invest Ophthalm Vis Sci* 53:5921–5929
- Yoshikawa M, Matsuda H, Morikawa T et al (2006) Phenylethanoid oligoglycosides and acylated oligosugars with vasorelaxant activity from *Cistanche tubulosa*. *Bioorg Med Chem* 14:7468–7475
- Yuan ZL, Luo L, Zang AM et al (2009) Isolation and bioassay of herbicidal active ingredient from *Paulownia tomentosa*. *Chin J Pestic Sci* 2:239–243
- Yun B-S, Lee I-K, Kim J-P et al (2000) Lipid peroxidation inhibitory activity of some constituents isolated from the stem bark of *Eucalyptus globulus*. *Arch Pharm Res* 23(2):147–150
- Yun J, Bae H, Choi SE et al (2013) Taxifolin glycoside blocks human ether-a-go-go related gene K(+) channels. *Korean J Physiol Pharmacol* 17(1):37–42
- Zhang SM, Coultas KA (2013) Identification of plumbagin and sanguinarine as effective chemotherapeutic agents for treatment of schistosomiasis. *Int J Parasitol Drugs Drug Resist* 3:28–34
- Zhang DL, Li XQ (2011) Studies on the chemical constituents from the leave of *Paulownia tomentosa*. *Zhong Yao Cai* 34:232–234
- Zhang W, Zhang W-D, Zhang C et al (2007) Antitumor activities of extracts and compounds from the roots of *Daphne tangutica* Maxim. *Phytother Res* 21(11):1113–1115
- Zhang J, Chen J, Liang Z et al (2014) New lignans and their biological activities. *Chem Biodivers* 11(1):1–54
- Zhao J, Zhou X-W, Chen X-B et al (2009)  $\alpha$ -Glucosidase inhibitory constituents from *Toona sinensis*. *Chem Nat Compd* 45:244–246
- Zhao J, Ding HX, Wang CM (2012) Isolation, modification and cytotoxic evaluation of flavonoids from *Rhododendron hainanense*. *J Pharm Pharmacol* 64:1785–1792
- Zheng J, Liu D, Zhao SQ et al (2012) Enzymatic extraction and antibacterial activity from *Eucommia ulmoides* leaves. *Zhong Yao Cai* 35:304–306
- Zhu ZH, Chao CJ, Lu XY et al (1986) Paulownia in China: cultivation and utilization. Asian Network for Biological Science and International Development Research Centre, Chinese Academy of Forestry, Beijing. <http://hdl-bnc.idrc.ca/dspace/bitstream/10625/8226/1/71235.pdf>. (Cited 13 Mar 2013)
- Zima A, Hošek J, Trembl J et al (2010) Antiradical and cytoprotective activities of several C-geranyl-substituted flavanones from *Paulownia tomentosa* Fruit. *Molecules* 15:6035–6049