

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/251405815>

ChemInform Abstract: Microwave-Assisted Synthesis of Symmetrical and Unsymmetrical N,N'-Disubstituted Thioureas and Ureas over MgO in dry Media.

ARTICLE in MONATSHFTE FUER CHEMIE/CHEMICAL MONTHLY · MAY 2012

Impact Factor: 1.22 · DOI: 10.1007/s00706-011-0583-6

READS

33

2 AUTHORS:



Hassan Valizadeh

Azarbaijan Shahid Madani University

43 PUBLICATIONS 173 CITATIONS

SEE PROFILE



Leila Dinparast

Azarbaijan Shahid Madani University

16 PUBLICATIONS 24 CITATIONS

SEE PROFILE

Microwave-assisted synthesis of symmetrical and unsymmetrical N,N#-disubstituted thioureas and ureas over MgO in dry media

Hassan Valizadeh & Leila Dinparast

Monatshefte für Chemie - Chemical Monthly

An International Journal of Chemistry

ISSN 0026-9247


Monatsh Chem

DOI 10.1007/s00706-011-0583-6

**Monatshefte
für Chemie
Chemical Monthly**

An International Journal of Chemistry

Vol. 139 No. 3 2008



 SpringerWienNewYork

 **Springer**

Your article is protected by copyright and all rights are held exclusively by Springer-Verlag. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your work, please use the accepted author's version for posting to your own website or your institution's repository. You may further deposit the accepted author's version on a funder's repository at a funder's request, provided it is not made publicly available until 12 months after publication.

Microwave-assisted synthesis of symmetrical and unsymmetrical *N,N'*-disubstituted thioureas and ureas over MgO in dry media

Hassan Valizadeh · Leila Dinparast

Received: 10 May 2010 / Accepted: 12 July 2011
© Springer-Verlag 2011

Abstract Under mild microwave irradiation conditions a variety of symmetrical and unsymmetrical *N,N'*-disubstituted thioureas and ureas were prepared via the reaction of *N*-monosubstituted hydroxylamines with isocyanate and isothiocyanate derivatives over MgO under solvent-free conditions. This new method afforded satisfactory results with good yields, short reaction time, and simplicity in the experimental procedure.

Keywords *N*-Monosubstituted hydroxylamines · Isocyanate · Isothiocyanate · Microwave irradiation · Solvent-free

Introduction

Green chemistry has been attracting great interest from chemists because of the potential environmental benefits of using such methods. A number of procedures involving new eco-friendly reagents and catalysts, selected media such as water, supercritical fluids, ionic liquids, solvent-free reactions, non-classical modes of activation such as ultrasound or microwaves are now recommended for green chemistry. There are several advantages of performing organic reactions in a solventless system such as short reaction time, increased safety, and economic advantages from the absence of solvent [1]. Recently, microwave irradiation (MWI) has

become an established tool in organic synthesis [2–4], because of the rate enhancements, higher yields, and often improved selectivity with respect to conventional reaction conditions. In addition, solvent-free MWI processes are also clean and efficient; moreover, the use of organic or inorganic solid supports has received increased attention [5, 6].

Thiourea and urea derivatives are important synthetic intermediates for amidines or guanidines [7]. Thioureas have also been used as neutral receptors for various anions [8] and natural product mimics. There are many methods for the preparation of disubstituted ureas [9–14] and thioureas [15–19] including condensation of amine hydrochlorides with potassium thiocyanate [20]. The most direct approach involves interaction of carbon disulfide with primary amines [9]. However, this reaction is often associated with several by-products depending on the reaction conditions employed [9]. To the best of our knowledge, there are no reports on the use of *N*-substituted hydroxylamines in the reaction with isothiocyanate and isocyanate derivatives for the synthesis of the related *N,N'*-disubstituted thioureas and ureas.

In continuation of our work in studying organic reactions in ionic liquids, water, or solventless systems as a green reaction medium [21–27], we wish to report here the synthesis of symmetrical and unsymmetrical *N,N'*-disubstituted thioureas and ureas from the reaction of *N*-substituted hydroxylamines with isothiocyanate and isocyanate derivatives over MgO as basic solid support in a solventless system under MWI (Scheme 1).

Results and discussions

In a pilot experiment, *N*-methylhydroxylamine (**2a**) and phenylisothiocyanate (**1a**) were mixed with excess of MgO

Electronic supplementary material The online version of this article (doi:10.1007/s00706-011-0583-6) contains supplementary material, which is available to authorized users.

H. Valizadeh (✉) · L. Dinparast
Department of Chemistry, Faculty of Sciences,
Azarbaijan University of Tarbiat Moallem, Tabriz, Iran
e-mail: hvalizadeh2@yahoo.com

Scheme 1





Table 1 Microwave-assisted synthesis of thioureas and ureas over MgO in solventless system

Entry	Reactant 1	Reactant 2	MWI or thermal heating	Time/min	Product 3, yield ^a /%	Product 4, yield ^a /%
1	1a	2a	MWI	3	3a (76)	4a (9)
2	1a	2b	MWI	4.5	–	4a (81)
3	1b	2a	MWI	3.5	3b (75)	4b (8)
4	1b	2b	MWI	3	–	4b (78)
5	1c	2a	MWI	3.5	3c (71)	4c (13)
6	1c	2b	MWI	4	3d (76)	4c (12)
7	1d	2a	MWI	3.5	3e (74)	4d (10)
8	1d	2b	MWI	4	3f (72)	4d (9)
9	1a	2a	MWI (neat conditions)	60	3a (22)	–
10	1c	2a	MWI (neat conditions)	60	3c (23)	–
11	1a	2a	Thermal heating	300	3a (25)	–
12	1c	2a	Thermal heating	300	3c (21)	–

^a Isolated yields after recrystallization

and irradiated by microwave to obtain *N*-methyl-*N'*-phenylthiourea (**3a**) in 76% yield and *N,N'*-diphenylthiourea (**4a**) as a minor product in less than 10% yield (Table 1). By this method, other isocyanates and isothiocyanates with *N*-(phenyl and methyl)hydroxylamines gave satisfactory results (Table 1, entries 2–8). In order to exhibit the catalytic effect of MgO, the condensation of *N*-methylhydroxylamine with two isothiocyanates (**1a** and **1c**) was studied in the absence of MgO under MWI conditions. It was found that MgO was more effective, and very poor yield of the corresponding thioureas were isolated in these conditions (Table 1, entries 9–10). In the absence of MWI, the reactions proceeded in longer reaction times at 80 °C in lower yields in comparison with MWI conditions (Table 1, entries 11–12). So MWI can shorten the reaction time and also assist the reaction process effectively. We conducted these reactions on a 25-mmol scale and found that they underwent a smooth transformation to the thiourea and urea derivatives in good yields. Thus, the present procedure is amenable to scaling up. Optimization of the reaction conditions was studied with different molar ratios of MgO and under different microwave powers. The best ratio was found to be 200 mol% of MgO under 200-W microwave power. Increasing the power of microwave decreases the yields of the products.

A plausible mechanism for the formation of the selected products **3a** and **4a** over MgO is outlined in Scheme 2. Intermediate **A** apparently results from the initial nucleophilic attack of *N*-methylhydroxylamine on phenylisothiocyanate in the presence of MgO as base catalyst. Subsequent nucleophilic addition of intermediate **A** to another phenylisothiocyanate yields the intermediate **B**, which fragments to CSO, aniline, and **3a** as a major product. The fragmentation of **B** perhaps accrued in the



Scheme 2

presence of water or oxygen present in the open reaction vessel under MWI. The compound *N,N'*-diphenylthiourea (**4a**) was obtained from the nucleophilic attack of aniline on unreacted phenylisothiocyanate in very low yield.

Conclusion

Under solvent-free conditions, microwave-assisted synthesis of symmetrical and unsymmetrical *N,N'*-disubstituted thioureas and ureas over MgO can be achieved from *N*-monosubstituted hydroxylamines and isocyanates and isothiocyanates efficiently. The advantages of this protocol are mild reaction conditions, short reaction times, simple work-up procedure, absence of organic solvent, and good yields. Therefore, we believe that the new synthetic method reported here constitutes an environmentally greener and safer process than existing approaches.

Experimental

All reagents were purchased from Merck and used without further purification. NMR spectra of samples in CDCl₃ solution were measured using a Bruker Avance AC-400 instrument (¹H at 400 MHz and ¹³C at 100 MHz). Mass spectra were recorded on a Platform II (Micromass, Manchester, UK) quadrupole mass spectrometer fitted with an electrospray interface. Elemental analyses were carried out on a Perkin-Elmer 240C elemental analyzer. Melting points were measured in open glass capillaries using a Stuart melting point apparatus. Microwave experiments were conducted in a Milestone MicroSYNTH apparatus.

General procedure for the preparation of thioureas and ureas over MgO under MWI

Isocyanate and isothiocyanate derivatives (20 mmol) and *N*-monosubstituted hydroxylamine (20 mmol) were added

to 10 cm³ acetone containing 1.61 g MgO (40 mmol) and stirred for 10 min at room temperature. After evaporation of acetone the mixture was subjected to MWI at 200 W for a few minutes (depending on the reactants, see Table 1). The completion of reaction was monitored by TLC using EtOAc/petroleum 1:2 as eluent. After completion of the reaction, the mixture was cooled to room temperature and extracted with ether or ethyl acetate. The extracts were concentrated on a rotary evaporator and the crude mixture was purified by silica gel (Merck 230–240 mesh) column chromatography using a 1:2 ethyl acetate/*n*-hexane mixture as eluent to give pure products **3a–3f** and **4a–4d**, which were characterized by their spectroscopic data (¹H NMR, ¹³C NMR, and IR), elemental analysis, and by comparison of their melting points with those reported in the literature (Table 2).

Acknowledgments The partial financial assistance from the Research Vice Chancellor of Azarbaijan University of Tarbiat Moallem is gratefully acknowledged.

References

1. Tanaka K, Toda F (2000) Chem Rev 100:1025
2. Lu YJ, Sun N, Huang ZS, Gu LQ (2008) Chin Chem Lett 19:518
3. Zarei A, Hajipour AR, Khazdooz L (2010) Dyes Pigm 85:133
4. Castanheiro RAP, Pinto MMM, Cravo SMM, Pinto DCGA, Silva AMS, Kijjoa A (2009) Tetrahedron 65:3848
5. Rostamizadeh S, Ghaieni HR, Aryan R, Amani AM (2010) Tetrahedron 66:494
6. Schroeder DC (1955) Chem Rev 55:181
7. Esteves-Souza A, Echevarría A, Vencato J, Jimeno ML, Elguero J (2001) Tetrahedron 57:6147
8. Tobe Y, Sasaki S, Mizuno M, Hirose K, Naemura K (1998) J Org Chem 63:7481 (and references cited therein)
9. Becker A, Heizler W (1983) Helv Chim Acta 66:1011
10. Criton M, Le VM-H (2008) Bioorg Med Chem Lett 18:3607
11. Ludwig BJ, Dürsch F, Auerbach M, Tomeczek K, Berger FM (1967) J Med Chem 10:556
12. Luk'yanov OA, Gordeev PB (1999) Russian Chem Bull 48:2302
13. Grigat E, Pütter R, König C (1965) Chem Ber 98:144
14. Douglas GH, Diamond J, Studt WL, Mir GN, Alioto RL, Auyang K, Burns BJ, Cias J, Darkes PR, Dodson SA, O'Connor S, Santora NJ, Tsuei CT, Zalipsky JJ, Zimmerman HK (1978) Arzneimittelforschung 28:1435
15. Vogel AI (1954) A textbook of organic chemistry. Longmans, London, p 614
16. Ballini R, Bosica G, Fiorini D, Maggi R, Righi P, Sartori G, Sartorio R (2002) Tetrahedron Lett 43:8445
17. Aravindakshan PA, Bhramaramba A, Nair GV, Nambury CNV (1963) Indian J Chem 1:395
18. Blanco LJ, Barría CS, Benito JM, Mellet CO, Fuentes J, Santoyo-González F, Fernández JMG (1999) Synthesis 1907
19. Moutou J-L, Schmitt M, Collot V, Bourguignon J-J (1981) Tetrahedron 37:1781
20. Herr RJ, Kuhler JL, Meckler H, Opalka CJ (2000) Synthesis 1569 (and references cited therein)
21. Valizadeh H, Vaghefi S (2009) Synth Commun 39:1666
22. Valizadeh H, Heravi MM, Amiri M (2010) Mol Divers 14:575
23. Valizadeh H (2010) Heteroatom Chem 21:78

Table 2 Melting points of products

Product	M.p./°C	Lit. m.p./°C
3a	95.5–99	92 [28]
3b	146–147	148–149 [29]
3c	87–89	54 [30]
3d	91–93	98 [31]
3e	99–102	54–55 [29]
3f	95.7–97	94–95 [32]
4a	152–154	153–154 [11]
4b	229–233	232 [11]
4c	77–78.5	78 [12]
4d	111–113	112 [33]

24. Valizadeh H, Shockravi A (2005) Tetrahedron Lett 46:3501
25. Valizadeh H, Mamaghani M, Badrian A (2005) Synth Commun 35:785
26. Valizadeh H, Fakhari A (2010) Molecules 15:2972
27. Valizadeh H, Gholipour H (2010) Synth Commun 40:1477
28. Ambati NB, Anand V, Hanumanthu P (1997) Synth Commun 27:1487
29. Snyder JK, Stock LM (1980) J Org Chem 45:886
30. Hofmann AW (1868) Chem Ber 1:27
31. Nair GV (1965) J Indian Chem Soc 42:359
32. Izdebski J, Pawlak D (1989) Synthesis 423
33. Zywucki B, Jadzyn J, Stockhausen M, Steffen A (1989) J Mol Struct 193:51