

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/273382892>

Design, synthesis, molecular modeling and anti-breast cancer activity of novel quinazolin-4-one derivatives linked to thiazolidinone, oxadiazole or pyrazole moieties

ARTICLE in MEDICINAL CHEMISTRY RESEARCH · MARCH 2015

Impact Factor: 1.4 · DOI: 10.1007/s00044-015-1357-1

CITATION

1

READS

47

3 AUTHORS:



Marwa Farag

9 PUBLICATIONS 61 CITATIONS

SEE PROFILE



Amany Belal

Taif University

12 PUBLICATIONS 19 CITATIONS

SEE PROFILE



Mahmoud Youns

Helwan University

24 PUBLICATIONS 463 CITATIONS

SEE PROFILE

Design, synthesis, molecular modeling and anti-breast cancer activity of novel quinazolin-4-one derivatives linked to thiazolidinone, oxadiazole or pyrazole moieties

Marwa F. Ahmed · Amany Belal · Mahmoud Youns

Received: 28 May 2014 / Accepted: 2 March 2015 / Published online: 8 March 2015
© Springer Science+Business Media New York 2015

Abstract 6,8-Dibromo-2-(4-chlorophenyl)-quinazolin-4-one linked directly to oxadiazole **5**, pyrazole **6** or through amide linkage to thiazolidinone **2a–d** and **3a–h** were synthesized; their chemical structures were confirmed by spectral and elemental analyses. Their anti-breast cancer activity was evaluated against human breast cancer cell line (MCF-7) using resazurin reduction method and doxorubicin as a reference drug. Linking quinazolin-4-one scaffold to oxadiazole or pyrazole gave compounds **5** and **6** with a closely similar activity as doxorubicin; their IC₅₀ was 23, 22 and 22 nmol/ml, respectively; however, the hybridization of quinazolin-4-one with thiazolidinone gave much better activity than doxorubicin. The most active compounds of the hybrid molecules between quinazolin-4-one and thiazolidinone are **2c,d** and **3a,f**. Their IC₅₀ range was (3–9 nmol/ml). In an attempt to explore the mode of action of the best active compounds, docking on the ATP binding site of EGFR was performed. In vitro screening of

these four compounds against EGFR tyrosine kinases showed inhibitory activity range 54–77.2 %.

Keywords Quinazolin-4-one · Thiazolidinone · Oxadiazole · Pyrazole · MCF-7

Introduction

Breast cancer occurs in humans and other mammals and originates from the tissues of the breast, either from milk ducts known as ductal carcinomas or from the lobules that supply milk for the ducts called lobular carcinomas (Santiago, 2010). The majority of human breast cancer cases are in women; however, it can also occur in men (Weiss *et al.*, 2005). Breast cancer is the most common tumor among women worldwide. Its incidence is increasing around the world, and it is believed to be the leading cause of cancer mortality, according to American Cancer Society. The estimated new breast cancer cases in USA (2014) are 235,030 cases, 232,670 of them estimated in women and the other 2360 in men. In addition, the estimated breast cancer deaths are 40,430 for the same year (Siegel *et al.*, 2014).

Quinazolines play a major role in the field of medicinal chemistry; they are of particular interest and considered as promising scaffolds in the search for new bioactive agents due to their diverse pharmacological activities such as anti-inflammatory (Alagarsamy *et al.*, 2009), antimicrobial (Mohamed *et al.*, 2010), antihypertensive (Ismail *et al.*, 2006), anticonvulsant (Georgey *et al.*, 2008), cholinesterase inhibitors (Decker, 2005), anticancer (Chandrika *et al.*, 2008) and anti-diabetic (Dongamanti *et al.*, 2012). Moreover, quinazolin-4-one derivative **I** (Fig. 1) bearing phenyl moiety at position 2 and 3 was synthesized and showed to inhibit about twenty percent of MCF-7 cells at 10 μM


M. F. Ahmed (✉)
Department of Pharmaceutical Chemistry, Faculty of Pharmacy,
Helwan University, Cairo, Egypt
e-mail: marwafarag80@yahoo.com

A. Belal
Department of Medicinal Chemistry, Faculty of Pharmacy,
Beni-Suef University, Beni Suef 62514, Egypt

A. Belal (✉)
Department of Pharmaceutical Chemistry, Faculty of Pharmacy,
Taif University, Taif 21974, Kingdom of Saudi Arabia
e-mail: abilalmoh1@yahoo.com

M. Youns
Department of Biochemistry and Molecular Biology, Faculty of
Pharmacy, Helwan University, Cairo, Egypt

Fig. 1 Active compounds against MCF-7 cell line



concentration (Al-Omary *et al.*, 2013). Another quinazolin-4-one having a phenyl moiety at the same positions 2 and 3 compound **II** (Fig. 1) showed a 49 % growth of MCF-7 inhibition at the same concentration 10 μ M; replacing the NH **II** with O **III** (Fig. 1) affected the activity and caused a 34 % growth inhibition, so it was observed that a minor change in the side chain has a great effect on the activity

against MCF-7 cell line (Al-Suwaidan *et al.*, 2013). In addition, thiazolidinone **IV** (Havrylyuk *et al.*, 2010), oxadiazole **V** (Galal *et al.*, 2010) and pyrazole **VI** (Fig. 1) containing compounds proved to have a great activity against breast cancer cells. Based on these findings, quinazolin-4-one scaffold substituted with phenyl at position 2,3 linked directly to oxadiazole **5**, pyrazole **6** or

through amide linkage to thiazolidinone **2a** was synthesized, screened against MCF-7 cell line using doxorubicin as a reference drug; **2a** was the best active one and showed better activity than both compounds **5,6** and doxorubicin itself. Furthermore, synthesis for a series **2a–d**, **3a–h** of quinazolin-4-one hybrid molecules with thiazolidinone was performed aiming to obtain potent and active compounds helpful in breast cancer treatment.

Epidermal growth factor receptors are family of receptor tyrosine kinases that play a crucial rule in cell growth regulation and survival (Olayioye *et al.*, 2000); they are highly expressed in a number of human tumors as breast cancer, colon, prostate and ovarian cancer, thus considered as attractive targets for the design and development of new anticancer active agents. It was also reported that quinazoline derivatives have a potent inhibitory activity against EGFR (Barlesi *et al.*, 2005; Burris *et al.*, 2005; Hennequin *et al.*, 2002; Kopper, 2008; Wood *et al.*, 2004); hence, docking the best active compounds **2c,d** and **3a,f** into ATP binding site of EGFR was performed to explore the possible interactions and amino acid binding for these compounds with EGFR.

Result and discussion

Chemistry

Compounds **1a–d** and **4** were synthesized according to the reported methods (Ahmed *et al.*, 2016); cyclocondensation of the Schiff bases **1a–d** with mercapto acetic acid in glacial acetic acid afforded the corresponding quinazolin-4-one linked to thiazolidinone nucleus through amide linkage **2a–d**, respectively. Their IR spectrum revealed the appearance of an additional absorption band corresponding to the carbonyl of thiazolidinone nucleus, and ¹H-NMR spectra showed the appearance of two signals at 3.8–3.9 and 5.9 which are characteristic for CH₂ and CH–Ar in thiazolidinone moiety. Moreover, compounds **2b** and **2c** showed signals at 3.8 and 2.3 ppm corresponding to OCH₃ and CH₃, respectively. To make substitution at position 5 of the thiazolidinone nucleus aiming to obtain another new derivatives of quinazolinone–thiazolidinone hybrid molecules, reacting **2a–d** with paraformaldehyde and 2-ry amines as piperidine or morpholine to give the corresponding Mannich bases **3a–h** (Scheme 1), ¹H-NMR showed the appearance of morpholine and piperidine protons and a signal for CH₂–N protons at 2.9 ppm in addition to the other signals characterizing the structures.

To prepare a hybrid molecule of quinazolin-4-one with oxadiazole, heating the acid hydrazide **4** with acetic anhydride was done and the corresponding methyl oxadiazole derivative **5** was obtained. Furthermore, reaction of **4** with


diketonic esters such as ethyl acetoacetate afforded the corresponding pyrazolyl-quinazolinone molecule **6** (Scheme 2), microanalysis and mass spectrum confirmed both structures **5,6**, and in addition the ¹H-NMR of compound **5** showed a characteristic signal for the CH₃ group at 2.6 ppm.

Biological screening

All the synthesized compounds were screened for their in vitro cytotoxic and growth inhibitory activities against MCF-7 cell line using resazurin cell growth inhibition assay method in comparison with the activity of the known anticancer drug doxorubicin. The cytotoxic activities of our tested compounds were expressed as IC₅₀ values (the dose that reduces survival to 50 %). The tested compounds showed anti-breast cancer activity with IC₅₀ values ranging from 3 to 23 nmol/ml (Table 1). The two series **2a–d** and **3a–h** showed better activity than doxorubicin except **3d** that showed to be closely similar to the reference drug. Compounds **5** and **6** showed less to similar activity as doxorubicin. The best active compounds are **2c,d** and **3a,f**; their IC₅₀ values range from 3 to 9 nmol/ml. In vitro screening of these four compounds against EGFR-TKs was also performed, they showed inhibitory activity range from 54 to 77.2 % (Table 3), and the best activity was assigned for compound **3f** that showed inhibitory activity at 77.2 %.

Molecular docking studies


The epidermal growth factor receptor (EGFR) is a trans-membrane receptor tyrosine kinase that transmits signals for the cell causing proliferation, differentiation or motility. Overexpression of the EGFR is noticed in numerous human malignancies; hence, targeting the ATP binding site of the EGFR by small molecules that act as tyrosine kinase inhibitors is considered a strategy for cancer treatment (Yun *et al.*, 2007). Moreover, several 4-anilinoquinazolines act as tyrosine kinase inhibitors (TKIs) which are currently in clinical investigation as gefitinib (Iressa) (Wakeling *et al.*, 2002), erlotinib (Tarceva) (Pollack *et al.*, 1999) and lapatinib (Rusnak *et al.*, 2001). Molecular docking studies for the most active compounds **2c,d** and **3a,f** that showed to be more potent than doxorubicin against MCF-7 were performed on the ATP binding site of EGFR to explore possible interactions and amino acid binding with these receptors. To perform accurate validation of the docking protocol, docking of IRE should be carried out to study the root mean standard deviation (RMSD), scoring energy (S) and amino acid interactions. Docking was performed using London dG force, and the results were refined using force field energy. IRE was fitted in the pocket of EGFR



Scheme 1 Synthesis of compounds **2a–d** and **3a–h**. Reagents and conditions: *a* thioglycolic acid, glacial acetic acid, reflux 6 h. *b* Paraformaldehyde, secondary amine, DMF, reflux 8 h

with $S = -22.28$ kcal/mol, $RSMD = 1.52$ and showed interactions with two amino acids Asp 855 and Lys 745 by two hydrogen bonds (Fig. 2). The aforementioned docking protocol was used for all the docked compounds **2c,d** and **3a,f** on the active site of EGFR. Docking scores, amino acids interactions and interacting groups are shown in Table 2. All the docked compounds exhibited proper fitting on the active site with docking score energy range from -13.15 to -19.16 kcal/mol (Figs. 3, 4, 5, 6). Compound **3f** fit with the active site with the best fitting score energy

(-19.16 kcal/mol) and amino acid interaction with Cys 797, Lys 728 and Lys 716 (Fig. 6). Compound **2c** showed amino acid interactions with Asp 855 and Lys 754 with score energy -13.37 kcal/mol (Fig. 3). Docking score energy of compounds **2d** and **3a** are -13.15 and -13.49 kcal/mol, respectively (Figs. 4, 5). These results support the postulation of the ability of these compounds to act as EGFR-TKIs and give a promising venue for further investigation of the quinazolin-4-one hybrid molecules with thiazolidinone (Table 3).



Scheme 2 Synthesis of compounds **5** and **6**. Reagents and conditions: *a* acetic anhydride, reflux 6 h, *b* ethyl acetoacetate, absolute ethanol, reflux 6 h

Table 1 IC₅₀ values expressed in $\mu\text{mol/ml}$ (nmol/ml) of the new compounds and doxorubicin against MCF-7 cell line

Compound number	IC ₅₀	Compound number	IC ₅₀
2a	0.011 (11)	3d	0.023 (23)
2b	0.016 (16)	3e	0.014 (14)
2c	0.006 (6)	3f	0.009 (9)
2d	0.009 (9)	3g	0.011 (11)
3a	0.003 (3)	3h	0.018 (18)
3b	0.012 (12)	5	0.023 (23)
3c	0.012 (12)	6	0.022 (22)
Doxorubicin			0.022 (22)

Conclusion

We have synthesized three hybridized molecules of quinazolin-4-one with thiazolidinone **2a**, oxadiazole **5** or pyrazole **6**, and we have evaluated their anti-breast cancer activity against MCF-7 cell line. The best activity was assigned for compound **2a**, so more derivatives of this hybrid molecule were synthesized **2a–d**, **3a–h** and screened also against the same cell line. From the observed results, we can conclude that:

1. Linking quinazolin-4-one with thiazolidinone in one molecule **2a** gave better activity against MCF-7 cell line than linking it with oxadiazole **5** or pyrazole **6**; the obtained hybrid molecule **2a** showed better activity than doxorubicin itself.
2. All hybrid molecules that have no substitution at position 5 of the thiazolidinone nucleus **2a–d** showed better activity than doxorubicin.
3. Substituting position 5 of thiazolidinone moiety **3a–h** gave better active compounds than doxorubicin except compound **3d**; moreover, substituting this position with morpholine decreased the activity than substituting it with piperidine except compound **3f**.
4. The best active compounds **2c,d** and **3a,f** that showed IC₅₀ range 3–9 nmol/ml are quinazolin-4-one and thiazolidinone hybrid molecules. In vitro screening of these compounds against EGFR tyrosine kinases revealed inhibitory activity range 54–77.2 %, and the best inhibitory activity 77.2 % was shown by compound **3f** followed by **2c** (70 %).
5. Molecular docking studies for the best active compounds **2c,d** and **3a,f** against ATP binding site of EGFR proved their ability to quick fit this active site; docking results supported the in vitro EGFR tyrosine kinase inhibitory activities.

Fig. 2 IRE in the ATP binding site of EGFR (RMDS = 1.52, $S = -22.28$ kcal/mol)




Table 2 Amino acid interactions and the binding scores of the docked compounds on the active site EGFR

Compound number	S (kcal/mol)	Amino acid interactions	Interacting groups
2c	−13.37	Asp 855 Lys 745	NH of the amide spacer CO of thiazolidinone
2d	−13.15		
3a	−13.49		
3f	−19.16	Lys 728 Lys 716 Cys 797	O of morpholine O of morpholine CO of the amide spacer
IRE	−22.28	Asp 855 Lys 745	NH of morpholine O of OCH ₃

Experimental

Chemistry

Melting points were detected using Electrothermal Stuart SMP3 digital melting point apparatus and were uncorrected. Elemental analyses were carried out in the micro-analytical units of National Research Centre and Cairo

University, Egypt. IR spectra were recorded on FT-IR spectrophotometer (Nexus 670-Nicolet, USA) and PerkinElmer-9712 spectrophotometer. ¹H-NMR spectra were determined on a Varian-Gemmi-300 MHz and Joel-Ex270 MHz NMR spectrometer using TMS as an internal standard. ¹³C NMR (DMSO-d₆) spectra were recorded at 100.62 MHz at the aforementioned research center in Cairo University. Mass spectra were recorded on Finnigan Mat SSQ 7000 mode EI 70 eV (Thermo Inst. Sys. Inc., USA). Thin-layer chromatography was carried out on silica gel 60 F254 (Merck) thin-layer chromatography plates using a chloroform, petroleum ether, methanol mixture (7:4:1 v/v) as the mobile phase. Compounds **1a–d** and **4** were synthesized with the same previously reported methods (Ahmed *et al.*, 2016).

4-[6,8-Dibromo-2-(4-chlorophenyl)-4-oxo-4H-quinazolin-3-yl]-benzoic acid benzylidene-hydrazide (1a) Yellow crystals (methanol); yield 70 %; mp 150–152 °C; IR (KBr) ν_{max} 3370 (NH), 1710, 1685 (2CO) and 1596 (C=N) cm^{-1} ; ¹H NMR (300 MHz, DMSO-d₆): $\delta = 7.35\text{--}7.39$ (4H, m, H-2', H-6', H-2'', H-6''), 7.5–7.64 (3H, m, H-3''', H-4''', H-5'''), 7.65–7.79 (6H, m, H-3', H-5', H-3'', H-5'', H-2''', H-6'''), 8.1 (1H, s, H-5), 8.2 (1H, s, H-7), 8.36 (1H, s, CH=N), and at 11.80 (1H, s, NH, exchangeable with D₂O); ¹³C NMR (100 MHz, DMSO-d₆): 164.4 (C=N), 163.3 (C=O), 160.5 (C=O), 154.3 (Ar/olefinic

Fig. 3 Compound **2c** in the ATP binding site of EGFR ($S = -13.37$ kcal/mol)




Fig. 4 Compound **2d** in the ATP binding site of EGFR ($S = -13.15$ kcal/mol)





Fig. 5 Compound **3a** in the ATP binding site of EGFR ($S = -13.49$ kcal/mol)



carbon), 146.8 (CH), 139.2 (Ar/olefinic carbon), 136 (Ar/olefinic carbon), 135.4 (Ar/olefinic carbon), 133.6 (Ar/olefinic carbon), 131.2 (Ar/olefinic carbon), 131 (Ar/olefinic carbon), 129.6 (Ar/olefinic carbon), 129.2 (Ar/olefinic carbon), 129.1 (Ar/olefinic carbon), 128.9 (Ar/olefinic carbon), 128.8 (Ar/olefinic carbon), 128.4 (Ar/olefinic carbon), 126.7 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 124.7 (Ar/olefinic carbon), 122 (Ar/olefinic carbon), 112.9 (Ar/olefinic carbon); MS: $m/z = 635.94$; Anal. Calcd for: $C_{28}H_{17}Br_2ClN_4O_2$: C, 52.82; H, 2.69; N, 8.80 %. Found C, 55.70; H, 2.65; N, 8.6 %.

4-[6,8-Dibromo-2-(4-chlorophenyl)-4-oxo-4H-quinazolin-3-yl]-benzoic acid (4-methoxy-benzylidene)-hydrazide (1b) Yellow crystals (acetic acid); yield 80 %; mp 190–192 °C; IR (KBr) ν_{max} 3355 (NH), 1715, 1690 (2CO) and 1598 (C=N) cm^{-1} ; 1H NMR (300 MHz, DMSO- d_6): $\delta = 3.83$ (3H, s, OCH₃), 7.0–7.3 (4H, m, H-2', H-6', H-2'', H-6''), 7.4–7.6 (2H, m, H-3''', H-5'''), 7.7–7.79 (6H, m, H-3', H-5', H-3'', H-5'', H-2''', H-6'''), 8.1 (1H, s, H-5), 8.25 (1H, s, H-7), 8.36 (1H, s, CH=N), and at 11.87 (1H, s, NH, exchangeable with D₂O); ^{13}C NMR (100 MHz, DMSO- d_6): 164 (C=N), 163.2 (C=O), 162.9 (Ar/olefinic carbon), 160.6 (C=O), 154.3 (Ar/olefinic carbon), 146.8 (CH), 139.4 (Ar/olefinic carbon), 136.1 (Ar/olefinic carbon), 135.7 (Ar/

olefinic carbon), 131.3 (Ar/olefinic carbon), 130.2 (Ar/olefinic carbon), 129.6 (Ar/olefinic carbon), 129.1 (Ar/olefinic carbon), 128.9 (Ar/olefinic carbon), 128.4 (Ar/olefinic carbon), 126.7 (Ar/olefinic carbon), 126 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 124.5 (Ar/olefinic carbon), 122 (Ar/olefinic carbon), 114.4 (Ar/olefinic carbon), 113.2 (Ar/olefinic carbon), 55.7 (CH₃); MS: $m/z = 665.95$; Anal. Calcd for: $C_{29}H_{19}Br_2ClN_4O_3$: C, 52.24; H, 2.87; N, 8.40 %. Found C, 52.20; H, 2.90; N, 8.56 %.

4-[6,8-Dibromo-2-(4-chlorophenyl)-4-oxo-4H-quinazolin-3-yl]-benzoic acid (4-methyl-benzylidene)-hydrazide (1c) White crystals (ethanol); yield 80 %; mp 200–202 °C; IR (KBr) ν_{max} 3315 (NH), 1720, 1690 (2CO) and 1600 (C=N) cm^{-1} ; 1H NMR (300 MHz, DMSO- d_6): $\delta = 2.35$ (3H, s, CH₃), 7.06–7.25 (4H, m, H-2', H-6', H-2'', H-6''), 7.35–7.59 (2H, m, H-3''', H-5'''), 7.7–7.79 (6H, m, H-3', H-5', H-3'', H-5'', H-2''', H-6'''), 8.2 (1H, s, H-5), 8.28 (1H, s, H-7), 8.36 (1H, s, CH=N), and at 11.8 (1H, s, NH, exchangeable with D₂O); ^{13}C NMR (100 MHz, DMSO- d_6): 163.9 (C=N), 163 (C=O), 160.5 (C=O), 154.7 (Ar/olefinic carbon), 146.4 (CH), 140.7 (Ar/olefinic carbon), 139.9 (Ar/olefinic carbon), 136 (Ar/olefinic carbon), 135.7 (Ar/olefinic carbon), 131.1 (Ar/olefinic carbon), 130.5 (Ar/olefinic carbon), 129.6 (Ar/olefinic carbon), 129.1 (Ar/

Fig. 6 Compound **3f** in the ATP binding site of EGFR ($S = -19.16$ kcal/mol)




Table 3 EGFR tyrosine kinase assay of compounds **2c,d** and **3a,f** at single dose (10 μ M)

Compound number	% Inhibition of EGFR tyrosine kinase
2c	70
2d	54
3a	54.3
3f	77.2
Gefitinib	100

olefinic carbon), 128.9 (Ar/olefinic carbon), 128.4 (Ar/olefinic carbon), 126.7 (Ar/olefinic carbon), 126.1 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 124.4 (Ar/olefinic carbon), 122.1 (Ar/olefinic carbon), 113.5 (Ar/olefinic carbon), 21.3 (CH_3); MS: $m/z = 649.95$; Anal. Calcd for: $\text{C}_{29}\text{H}_{19}\text{Br}_2\text{ClN}_4\text{O}_2$: C, 53.52; H, 2.94; N, 8.61 %. Found C, 53.49; H, 3.00; N, 8.67 %.

4-[6,8-Dibromo-2-(4-chlorophenyl)-4-oxo-4H-quinazolin-3-yl]-benzoic acid (2-hydroxy-benzylidene)-hydrazide (1d) Yellowish white crystals (methanol); yield 80 %; mp 154–156 $^{\circ}\text{C}$; IR (KBr) ν_{max} 3315 (NH), 1710, 1690

(2CO) and 1600 ($\text{C}=\text{N}$) cm^{-1} ; ^1H NMR (300 MHz, $\text{DMSO}-d_6$): $\delta = 7.2\text{--}7.45$ (4H, m, H-2', H-6', H-2'', H-6''), 7.5–7.7 (2H, m, H-3''', H-5'''), 7.73–8.00 (6H, m, H-3', H-5', H-3'', H-5'', H-4''', H-6'''), 8.24 (1H, s, H-5), 8.3 (1H, s, H-7), 8.54 (1H, s, $\text{CH}=\text{N}$), 10.1 (1H, s, NH, exchangeable with D_2O) and at 11.2 (1H, s, OH); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): 163.5 ($\text{C}=\text{N}$), 163.2 ($\text{C}=\text{O}$), 160.1 ($\text{C}=\text{O}$), 157.2 (Ar/olefinic carbon), 154.3 (Ar/olefinic carbon), 146 (CH), 139.2 (Ar/olefinic carbon), 136.1 (Ar/olefinic carbon), 135.7 (Ar/olefinic carbon), 132.4 (Ar/olefinic carbon), 131.3 (Ar/olefinic carbon), 129.6 (Ar/olefinic carbon), 129.1 (Ar/olefinic carbon), 128.9 (Ar/olefinic carbon), 128.4 (Ar/olefinic carbon), 127.4 (Ar/olefinic carbon), 126.7 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 124.5 (Ar/olefinic carbon), 122.6 (Ar/olefinic carbon), 121.4 (Ar/olefinic carbon), 118.5 (Ar/olefinic carbon), 113.2 (Ar/olefinic carbon); MS: $m/z = 651.93$; Anal. Calcd for: $\text{C}_{28}\text{H}_{17}\text{Br}_2\text{ClN}_4\text{O}_3$: C, 51.52; H, 2.63; N, 8.58 %. Found C, 51.39; H, 2.60; N, 8.65 %.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(4-oxo-2-(aryl)thiazolidin-3-yl)benzamide (2a–d) General method: A mixture of compounds

1a–d (10 mmol) and thioglycolic acid (0.35 ml; 10 mmol) in glacial acetic acid (20 ml) was refluxed for 6 h. The excess solvent was evaporated under reduced pressure, and the obtained residue was poured on crushed ice. The solid product was filtered off and washed with water to obtain the desired products **2a–d**, respectively.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(4-oxo-2-phenyl thiazolidin-3-yl)benzamide (2a) Orange crystals (methanol); yield 75 %; mp 210–212 °C; IR (KBr) ν_{max} 3270 (NH), 3067 (CH, aromatic), 2940 (CH, thiazolidinone), 1720 (CO, cyclic amide thiazolidinone), 1690 (CO, cyclic amide quinazoline), 1660 (CONH) and 1595 (C=N) cm^{-1} ; ^1H NMR (300 MHz, DMSO- d_6): δ = 3.90 (2H, s, CH_2 , thiazolidinone ring), 5.92 (1H, s, CH of thiazolidinone ring), 7.35–7.45 (4H, m, H-2', H-6', H-2'', H-6''), 7.5–7.59 (3H, m, H-3''', H-4''', H-5'''), 7.65–7.79 (6H, m, H-3', H-5', H-3'', H-5'', H-2''', H-6'''), 8.1 (1H, s, H-5), 8.3 (1H, s, H-7), 10.2 (1H, s, NH, exchangeable with D_2O); ^{13}C NMR (100 MHz, DMSO- d_6): 168.8 (C=O), 164 (C=N), 163.7 (C=O), 160.6 (C=O), 154.3 (Ar/olefinic carbon), 139.2 (Ar/olefinic carbon), 136.1 (Ar/olefinic carbon), 135.7 (Ar/olefinic carbon), 131.3 (Ar/olefinic carbon), 130.1 (Ar/olefinic carbon), 129.6 (Ar/olefinic carbon), 129.2 (Ar/olefinic carbon), 128.8 (Ar/olefinic carbon), 128.6 (Ar/olefinic carbon), 128.2 (Ar/olefinic carbon), 127.6 (Ar/olefinic carbon), 127.1 (Ar/olefinic carbon), 126.9 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 124.5 (Ar/olefinic carbon), 122 (Ar/olefinic carbon), 113.2 (Ar/olefinic carbon), 57.3 (CH), 35.6 (CH_2); MS: m/z = 675.96; Anal. Calcd for: $\text{C}_{30}\text{H}_{19}\text{Br}_2\text{ClN}_4\text{O}_3\text{S}$: C, 50.69; H, 2.69; N, 7.88 %. Found C, 50.60; H, 2.72; N, 7.76 %.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(2-(4-methoxy phenyl)-4-oxothiazolidin-3-yl)benzamide (2b) White crystals (acetic acid); yield 75 %; mp 155–157 °C; IR (KBr) ν_{max} 3365 (NH), 3070 (CH, aromatic), 2950 (CH, thiazolidinone), 1715 (CO, cyclic amide thiazolidinone), 1690 (CO, cyclic amide quinazoline), 1665 (CONH) and 1600 (C=N) cm^{-1} ; ^1H NMR (300 MHz, DMSO- d_6): δ = 3.9 (2H, s, CH_2 , thiazolidinone ring), 3.8 (3H, s, OCH_3), 5.90 (1H, s, CH of thiazolidinone ring), 7.3–7.5 (4H, m, H-2', H-6', H-2'', H-6''), 7.55–7.64 (2H, m, H-3''', H-5'''), 7.7–8.1 (6H, m, H-3', H-5', H-3'', H-5'', H-2''', H-6'''), 8.2 (1H, s, H-5), 8.45 (1H, s, H-7), 9.27 (1H, s, NH, exchangeable with D_2O); ^{13}C NMR (100 MHz, DMSO- d_6): 170 (C=O), 165.2 (C=N), 164.5 (C=O), 161.6 (C=O), 159 (Ar/olefinic carbon), 155.2 (Ar/olefinic carbon), 140.2 (Ar/olefinic carbon), 136.4 (Ar/olefinic carbon), 135.9 (Ar/olefinic carbon), 131.5 (Ar/olefinic carbon), 131.3 (Ar/olefinic carbon), 130.1 (Ar/olefinic carbon), 129.6 (Ar/olefinic carbon), 129 (Ar/olefinic carbon), 128.6 (Ar/olefinic carbon), 128.2 (Ar/olefinic

carbon), 127.9 (Ar/olefinic carbon), 125.4 (Ar/olefinic carbon), 124.6 (Ar/olefinic carbon), 122 (Ar/olefinic carbon), 114.2 (Ar/olefinic carbon), 113.4 (Ar/olefinic carbon), 58.4 (CH), 55.8 (CH_3), 36.4 (CH_2); MS: m/z = 739.93; Anal. Calcd for: $\text{C}_{31}\text{H}_{21}\text{Br}_2\text{ClN}_4\text{O}_4\text{S}$: C, 50.26; H, 2.86; N, 7.56 %. Found C, 52.30; H, 2.90; N, 7.69 %.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(4-oxo-2-p-tolyl thiazolidin-3-yl)benzamide (2c) Yellowish white crystals (ethanol); yield 70 %; mp 179–181 °C; IR (KBr) ν_{max} 3365 (NH), 3070 (CH, aromatic), 2930 (CH, thiazolidinone), 1700 (CO, cyclic amide thiazolidinone), 1685 (CO, cyclic amide quinazoline), 1665 (CONH) and 1600 (C=N) cm^{-1} ; ^1H NMR (300 MHz, DMSO- d_6): δ = 2.34 (3H, s, CH_3), 3.9 (2H, s, CH_2 , thiazolidinone ring), 5.90 (1H, s, CH of thiazolidinone ring), 7.25–7.45 (4H, m, H-2', H-6', H-2'', H-6''), 7.5–7.65 (2H, m, H-3''', H-5'''), 7.7–8.1 (6H, m, H-3', H-5', H-3'', H-5'', H-2''', H-6'''), 8.2 (1H, s, H-5), 8.3 (1H, s, H-7), 9.25 (1H, s, NH, exchangeable with D_2O); ^{13}C NMR (100 MHz, DMSO- d_6): 166.9 (C=O), 164.2 (C=N), 163.5 (C=O), 161.2 (C=O), 154.3 (Ar/olefinic carbon), 139.2 (Ar/olefinic carbon), 136.8 (Ar/olefinic carbon), 136.2 (Ar/olefinic carbon), 136.1 (Ar/olefinic carbon), 135.7 (Ar/olefinic carbon), 131.3 (Ar/olefinic carbon), 129.6 (Ar/olefinic carbon), 129.1 (Ar/olefinic carbon), 128.9 (Ar/olefinic carbon), 128.6 (Ar/olefinic carbon), 127.6 (Ar/olefinic carbon), 126.7 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 124.5 (Ar/olefinic carbon), 122.1 (Ar/olefinic carbon), 113.2 (Ar/olefinic carbon), 57.4 (CH), 35.4 (CH_2), 21.3 (CH_3); MS: m/z = 739.93; Anal. Calcd for: $\text{C}_{31}\text{H}_{21}\text{Br}_2\text{ClN}_4\text{O}_3\text{S}$: C, 51.73; H, 2.92; N, 7.73 %. Found C, 51.88; H, 2.70; N, 7.67 %.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(2-(2-hydroxy phenyl)-4-oxothiazolidin-3-yl)benzamide (2d) White crystals (methanol); yield 65 %; mp 140–142 °C; IR (KBr) ν_{max} 3360 (NH), 3060 (CH, aromatic), 2930 (CH, thiazolidinone), 1720 (CO, cyclic amide thiazolidinone), 1685 (CO, cyclic amide quinazoline), 1665 (CONH) and 1600 (C=N) cm^{-1} ; ^1H NMR (300 MHz, DMSO- d_6): δ = 4.1 (2H, s, CH_2 , thiazolidinone ring), 6.00 (1H, s, CH of thiazolidinone ring), 7.5–7.59 (4H, m, H-2', H-6', H-2'', H-6''), 7.6–7.7 (2H, m, H-3''', H-5'''), 7.73–8.00 (6H, m, H-3', H-5', H-3'', H-5'', H-4''', H-6'''), 8.25 (1H, s, H-5), 8.4 (1H, s, H-7), 9.50 (1H, s, NH, exchangeable with D_2O); ^{13}C NMR (100 MHz, DMSO- d_6): 170.2 (C=O), 164.6 (C=N), 163.9 (C=O), 161.5 (C=O), 155.5 (Ar/olefinic carbon), 140.2 (Ar/olefinic carbon), 137.3 (Ar/olefinic carbon), 136.1 (Ar/olefinic carbon), 135.2 (Ar/olefinic carbon), 132.7 (Ar/olefinic carbon), 131.3 (Ar/olefinic carbon), 130.2 (Ar/olefinic carbon), 129.6 (Ar/olefinic carbon), 129.1 (Ar/olefinic carbon), 128.7 (Ar/olefinic carbon), 128.5 (Ar/olefinic carbon), 127.6 (Ar/olefinic

carbon), 126.5 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 124.7 (Ar/olefinic carbon), 123.2 (Ar/olefinic carbon), 115.4 (Ar/olefinic carbon), 57.9 (CH), 36.4 (CH₂); MS: m/z = 743.88; Anal. Calcd for: C₃₀H₁₈Br₂Cl₂N₄O₃S: C, 48.35; H, 2.43; N, 7.52 %. Found C, 48.15; H, 2.32; N, 7.49 %.

4-(2-Phenyl-6,8-dibromo-2-(4-chlorophenyl)-4-oxo-(4H)quinazolin-3-yl)-N-[5-piperidino methyl and/or morpholinomethyl]-4-oxo-2-arylthiazolidin-3-yl]benzamides (3a–h): (Mannich bases) General method: A mixture of compounds (2a–d), paraformaldehyde (0.045 g; 5 mol) and the appropriate secondary amine (5 mmol), namely piperidine and/or morpholine in DMF (20 ml), was refluxed for 8 h. The excess solvent was evaporated under vacuum, and the obtained residue was poured on crushed ice. The solid product was filtered off and washed with water to obtain the desired products 3a–h, respectively.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(4-oxo-2-phenyl-5-(piperidin-1-ylmethyl)thiazolidin-3-yl)benzamide (3a) Yellow crystals (methanol); yield 60 %; mp 130–132 °C; IR (KBr) ν_{\max} 3365 (NH), 3090 (CH, aromatic), 1710 (CO, cyclic amide thiazolidinone), 1685 (CO, cyclic amide quinazoline), 1665 (CONH) and 1610 (C=N) cm⁻¹; ¹H NMR (300 MHz, DMSO-d₆): δ = 1.53 (6H, m, 3CH₂ of piperidine), 2.45 (4H, m, CH₂NCH₂ of piperidine), 2.9 (2H, s, –CH₂–N), 3.73 (1H, s, CH, thiazolidinone ring), 5.90 (1H, s, CH of thiazolidinone ring), 7.5–7.62 (4H, m, H-2', H-6', H-2'', H-6''), 7.65–7.79 (3H, m, H-3''', H-4''', H-5'''), 7.8–8.0 (6H, m, H-3', H-5', H-3'', H-5'', H-2''', H-6'''), 8.15 (1H, s, H-5), 8.4 (1H, s, H-7), 10.3 (1H, s, NH, exchangeable with D₂O); ¹³C NMR (100 MHz, DMSO-d₆): 173.4 (C=O), 165.4 (C=N), 164.7 (C=O), 160.5 (C=O), 154.9 (Ar/olefinic carbon), 139.4 (Ar/olefinic carbon), 139.2 (Ar/olefinic carbon), 136.3 (Ar/olefinic carbon), 135.7 (Ar/olefinic carbon), 131.3 (Ar/olefinic carbon), 129.9 (Ar/olefinic carbon), 129.4 (Ar/olefinic carbon), 128.9 (Ar/olefinic carbon), 128.6 (Ar/olefinic carbon), 127.4 (Ar/olefinic carbon), 127.1 (Ar/olefinic carbon), 126.9 (Ar/olefinic carbon), 126.7 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 124.5 (Ar/olefinic carbon), 124 (Ar/olefinic carbon), 114.5 (Ar/olefinic carbon), 56.9 (CH₂), 54.9 (CH), 53.9 (CH₂), 52.3 (CH), 25.9 (CH₂), 24.5 (CH₂); MS: m/z = 807.01; Anal. Calcd for: C₃₆H₃₀Br₂ClN₅O₃S: C, 53.51; H, 3.74 N, 8.67 %. Found C, 53.48; H, 3.70; N, 8.53 %.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(5-(morpholinomethyl)-4-oxo-2-phenylthiazolidin-3-yl)benzamide (3b) Brown crystals (ethanol); yield 70 %; mp 222–224 °C; IR (KBr) ν_{\max} 3360 (NH), 3086 (CH, aromatic), 1695 (CO, cyclic amide thiazolidinone), 1685 (CO, cyclic amide quinazoline), 1660 (CONH) and 1601

(C=N) cm⁻¹; ¹H NMR (300 MHz, DMSO-d₆): δ = 2.6 (4H, t, 2CH₂ of morpholine), 2.8 (2H, s, –CH₂–N), 3.4 (4H, t, 2CH₂ of morpholine) 3.65 (1H, s, CH, thiazolidinone ring), 5.90 (1H, s, CH of thiazolidinone ring), 7.4–7.50 (4H, m, H-2', H-6', H-2'', H-6''), 7.6–7.69 (3H, m, H-3''', H-4''', H-5'''), 7.71–7.79 (6H, m, H-3', H-5', H-3'', H-5'', H-2''', H-6'''), 8.00 (1H, s, H-5), 8.1 (1H, s, H-7), 10.1 (1H, s, NH, exchangeable with D₂O); ¹³C NMR (100 MHz, DMSO-d₆): 174.5 (C=O), 165.7 (C=N), 165.2 (C=O), 161.3 (C=O), 155.9 (Ar/olefinic carbon), 139.7 (Ar/olefinic carbon), 139.5 (Ar/olefinic carbon), 137.3 (Ar/olefinic carbon), 136.7 (Ar/olefinic carbon), 131.5 (Ar/olefinic carbon), 129.7 (Ar/olefinic carbon), 129.2 (Ar/olefinic carbon), 128.7 (Ar/olefinic carbon), 128.4 (Ar/olefinic carbon), 127.9 (Ar/olefinic carbon), 127.4 (Ar/olefinic carbon), 126.4 (Ar/olefinic carbon), 126.1 (Ar/olefinic carbon), 125.6 (Ar/olefinic carbon), 124.7 (Ar/olefinic carbon), 124 (Ar/olefinic carbon), 117.5 (Ar/olefinic carbon), 66.7 (CH₂), 56.6 (CH₂), 54.9 (CH), 52.9 (CH₂), 52.4 (CH); MS: m/z = 808.99; Anal. Calcd for: C₃₅H₂₈Br₂ClN₅O₄S: C, 51.90; H, 3.48; N, 8.65 %. Found C, 51.80; H, 3.50; N, 8.77 %.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(2-(4-methoxy phenyl)-4-oxo-5-(piperidin-1-ylmethyl)thiazolidin-3-yl)benzamide (3c) White crystals (methanol); yield 60 %; mp 196–197 °C; IR (KBr) ν_{\max} 3360(NH), 3070(CH, aromatic), 1695(CO, cyclic amide thiazolidinone), 1685 (CO, cyclic amide quinazoline), 1665 (CONH) and 1601 (C=N) cm⁻¹; ¹H NMR (300 MHz, DMSO-d₆): δ = 1.5 (6H, m, 3CH₂ of piperidine), 2.6 (4H, m, CH₂NCH₂ of piperidine), 2.9 (2H, s, –CH₂–N), 3.71 (1H, s, CH, thiazolidinone ring), 3.83(3H, s, OCH₃), 5.72 (1H, s, CH of thiazolidinone ring), 7.5–7.7 (4H, m, H-2', H-6', H-2'', H-6''), 7.74–7.82 (2H, m, H-3''', H-5'''), 7.85–8.2 (6H, m, H-3', H-5', H-3'', H-5'', H-2''', H-6'''), 8.3 (1H, s, H-5), 8.5 (1H, s, H-7), 11.3 (1H, s, NH, exchangeable with D₂O); ¹³C NMR (100 MHz, DMSO-d₆): 175.2 (C=O), 166.5 (C=N), 165.9(C=O), 163.5 (C=O), 157.4 (Ar/olefinic carbon), 140.4 (Ar/olefinic carbon), 140.2 (Ar/olefinic carbon), 137.3 (Ar/olefinic carbon), 136.7 (Ar/olefinic carbon), 133.3 (Ar/olefinic carbon), 130.9 (Ar/olefinic carbon), 129.4 (Ar/olefinic carbon), 129.2 (Ar/olefinic carbon), 128.6 (Ar/olefinic carbon), 127.7 (Ar/olefinic carbon), 127.4 (Ar/olefinic carbon), 127.1 (Ar/olefinic carbon), 126.9 (Ar/olefinic carbon), 125.7 (Ar/olefinic carbon), 125.5 (Ar/olefinic carbon), 125 (Ar/olefinic carbon), 116.2 (Ar/olefinic carbon), 57.9 (CH₂), 56.3 (CH₃), 55.4 (CH), 54.2 (CH₂), 53.4 (CH), 26.7 (CH₂), 25.5 (CH₂); MS: m/z = 837.02; Anal. Calcd for: C₃₇H₃₂Br₂ClN₅O₄S: C, 53.03; H, 3.85; N, 8.36 %. Found C, 53.07; H, 3.90; N, 8.43 %.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(2-(4-methoxy phenyl)-5-(morpholinomethyl)-4-oxothiazolidin-3-yl)benzamide (3d) Yellowish white crystals (acetic acid); yield 65 %; mp 175–177 °C; IR (KBr) ν_{max} 3350 (NH), 3065 (CH, aromatic), 1695 (CO, cyclic amide thiazolidinone), 1685 (CO, cyclic amide quinazoline), 1660 (CONH) and 1601 (C=N) cm^{-1} ; ^1H NMR (300 MHz, DMSO- d_6): δ = 2.8 (4H, t, 2CH_2 of morpholine), 2.9 (2H, s, $-\text{CH}_2-\text{N}$), 3.4 (3H, s, OCH_3), 3.7 (4H, t, 2CH_2 of morpholine) 3.8 (1H, s, CH, thiazolidinone ring), 5.95 (1H, s, CH of thiazolidinone ring), 7.6–7.8 (4H, m, H-2', H-6', H-2'', H-6''), 7.82–7.95 (2H, m, H-3''', H-5'''), 8.0–8.3 (6H, m, H-3', H-5', H-3'', H-5'', H-2''', H-6'''), 8.4 (1H, s, H-5), 8.6 (1H, s, H-7), 10.5 (1H, s, NH, exchangeable with D_2O); ^{13}C NMR (100 MHz, DMSO- d_6): 173.5 (C=O), 166.2 (C=O), 164.6 (C=N), 163.4 (C=O), 155.9 (Ar/olefinic carbon), 139.9 (Ar/olefinic carbon), 139.6 (Ar/olefinic carbon), 138.2 (Ar/olefinic carbon), 137.2 (Ar/olefinic carbon), 132.5 (Ar/olefinic carbon), 130.6 (Ar/olefinic carbon), 130.2 (Ar/olefinic carbon), 128.9 (Ar/olefinic carbon), 128.3 (Ar/olefinic carbon), 128.0 (Ar/olefinic carbon), 127.6 (Ar/olefinic carbon), 126.7 (Ar/olefinic carbon), 126.4 (Ar/olefinic carbon), 125.4 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 125 (Ar/olefinic carbon), 118.5 (Ar/olefinic carbon), 65.6 (CH_2), 57.6 (CH_2), 55.8 (CH_3), 55.1 (CH), 53.2 (CH_2), 52.6 (CH); MS: m/z = 839.00; Anal. Calcd for: $\text{C}_{36}\text{H}_{30}\text{Br}_2\text{ClN}_5\text{O}_5\text{S}$: C, 51.48; H, 3.60; N, 8.34 %. Found C, 51.50; H, 3.70; N, 8.40 %.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(4-oxo-5-(piperidin-1-ylmethyl)-2-p-tolylthiazolidin-3-yl)benzamide (3e) Yellow crystals (ethanol); yield 65 %; mp 212–214 °C; IR (KBr) ν_{max} 3360 (NH), 3090 (CH, aromatic), 1720 (CO, cyclic amide thiazolidinone), 1690 (CO, cyclic amide quinazoline), 1665 (CONH) and 1610 (C=N) cm^{-1} ; ^1H NMR (300 MHz, DMSO- d_6): δ = 1.53 (6H, m, 3CH_2 of piperidine), 2.34 (3H, s, CH_3), 2.5 (4H, m, CH_2NCH_2 of piperidine), 2.9 (2H, s, $-\text{CH}_2-\text{N}$), 3.75 (1H, s, CH, thiazolidinone ring), 5.90 (1H, s, CH of thiazolidinone ring), 7.5–7.65 (4H, m, H-2', H-6', H-2'', H-6''), 7.69–7.74 (2H, m, H-3''', H-5'''), 7.79–8.1 (6H, m, H-3', H-5', H-3'', H-5'', H-2''', H-6'''), 8.2 (1H, s, H-5), 8.3 (1H, s, H-7), 9.3 (1H, s, NH, exchangeable with D_2O); ^{13}C NMR (100 MHz, DMSO- d_6): 172.3 (C=O), 163.2 (C=N), 162.4 (C=O), 160.5 (C=O), 154.2 (Ar/olefinic carbon), 139.2 (Ar/olefinic carbon), 139 (Ar/olefinic carbon), 136.3 (Ar/olefinic carbon), 136.5 (Ar/olefinic carbon), 132.1 (Ar/olefinic carbon), 129.7 (Ar/olefinic carbon), 129.2 (Ar/olefinic carbon), 129 (Ar/olefinic carbon), 127.6 (Ar/olefinic carbon), 126.7 (Ar/olefinic carbon), 126.3 (Ar/olefinic carbon), 126.1 (Ar/olefinic carbon), 124.9 (Ar/olefinic carbon), 123.3 (Ar/olefinic carbon), 122.5 (Ar/olefinic

carbon), 120 (Ar/olefinic carbon), 114.1 (Ar/olefinic carbon), 56.7 (CH_2), 53.4 (CH), 52.2 (CH_2), 50.4 (CH), 24.7 (CH_2), 23.5 (CH_2), 22.4 (CH_3); MS: m/z = 821.03; Anal. Calcd for: $\text{C}_{37}\text{H}_{32}\text{Br}_2\text{ClN}_5\text{O}_5\text{S}$: C, 54.06; H, 3.92 N, 8.52 %. Found C, 54.12; H, 3.79; N, 8.53 %.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(5-(morpholinomethyl)-4-oxo-2-p-tolylthiazolidin-3-yl)benzamide (3f) White crystals (ethanol); yield 60 %; mp 165–167 °C; IR (KBr) ν_{max} 3355 (NH), 3060 (CH, aromatic), 1695 (CO, cyclic amide thiazolidinone), 1680 (CO, cyclic amide quinazoline), 1660 (CONH) and 1600 (C=N) cm^{-1} ; ^1H NMR (300 MHz, DMSO- d_6): δ = 2.34 (3H, s, CH_3), 2.7 (4H, t, 2CH_2 of morpholine), 2.9 (2H, s, $-\text{CH}_2-\text{N}$), 3.6 (4H, t, 2CH_2 of morpholine) 3.75 (1H, s, CH, thiazolidinone ring), 5.90 (1H, s, CH of thiazolidinone ring), 7.53–7.69 (4H, m, H-2', H-6', H-2'', H-6''), 7.71–7.77 (2H, m, H-3''', H-5'''), 7.79–8.1 (6H, m, H-3', H-5', H-3'', H-5'', H-2''', H-6'''), 8.15 (1H, s, H-5), 8.25 (1H, s, H-7), 10.3 (1H, s, NH, exchangeable with D_2O); ^{13}C NMR (100 MHz, DMSO- d_6): 171.2 (C=O), 162.5 (C=N), 161.2 (C=O), 160.5 (C=O), 153.2 (Ar/olefinic carbon), 138.7 (Ar/olefinic carbon), 138.4 (Ar/olefinic carbon), 137.2 (Ar/olefinic carbon), 137.0 (Ar/olefinic carbon), 130.9 (Ar/olefinic carbon), 130.6 (Ar/olefinic carbon), 130.0 (Ar/olefinic carbon), 127.9 (Ar/olefinic carbon), 127.3 (Ar/olefinic carbon), 126.2 (Ar/olefinic carbon), 125.5 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 124.3 (Ar/olefinic carbon), 124.1 (Ar/olefinic carbon), 122.2 (Ar/olefinic carbon), 120.4 (Ar/olefinic carbon), 114.9 (Ar/olefinic carbon), 63.7 (CH_2), 55.4 (CH_2), 53.1 (CH), 50.6 (CH_2), 49.9 (CH), 23.8 (CH_3); MS: m/z = 823.01; Anal. Calcd for: $\text{C}_{36}\text{H}_{30}\text{Br}_2\text{ClN}_5\text{O}_4\text{S}$: C, 52.48; H, 3.67; N, 8.50 %. Found C, 52.38; H, 3.70; N, 8.48 %.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(2-(2-hydroxy phenyl)-4-oxo-5-(piperidin-1-ylmethyl)thiazolidin-3-yl)benzamide (3g) Brown crystals (methanol); yield 60 %; mp 145–147 °C; IR (KBr) ν_{max} 3365 (NH), 3100 (CH, aromatic), 1725 (CO, cyclic amide thiazolidinone), 1710 (CO, cyclic amide quinazoline), 1670 (CONH) and 1615 (C=N) cm^{-1} ; ^1H NMR (300 MHz, DMSO- d_6): δ = 1.53 (6H, m, 3CH_2 of piperidine), 2.5 (4H, m, CH_2NCH_2 of piperidine), 2.9 (2H, s, $-\text{CH}_2-\text{N}$), 3.75 (1H, s, CH, thiazolidinone ring), 5.90 (1H, s, CH of thiazolidinone ring), 7.45–7.55 (4H, m, H-2', H-6', H-2'', H-6''), 7.64–7.76 (2H, m, H-3''', H-5'''), 7.79–8.00 (6H, m, H-3', H-5', H-3'', H-5'', H-4''', H-6'''), 8.3 (1H, s, H-5), 8.4 (1H, s, H-7), 9.3 (1H, s, NH, exchangeable with D_2O), 10.2 (1H, s, OH); ^{13}C NMR (100 MHz, DMSO- d_6): 173.4 (C=O), 164 (C=N), 163.7 (C=O), 160.6 (C=O), 154.3 (Ar/olefinic carbon), 153.7 (Ar/olefinic carbon), 139.4 (Ar/olefinic carbon), 136.1 (Ar/olefinic carbon), 135.7 (Ar/olefinic carbon), 131.3 (Ar/olefinic carbon), 129.6 (Ar/olefinic carbon),

129.1 (Ar/olefinic carbon), 128.9 (Ar/olefinic carbon), 128.5 (Ar/olefinic carbon), 128 (Ar/olefinic carbon), 127.6 (Ar/olefinic carbon), 126.7 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 124.5 (Ar/olefinic carbon), 122 (Ar/olefinic carbon), 121.2 (Ar/olefinic carbon), 115.1 (Ar/olefinic carbon), 113.2 (Ar/olefinic carbon), 56.6 (CH₂), 53.9 (CH₂), 52.4 (CH), 48.6 (CH), 25.9 (CH₂), 24.5 (CH₂); MS: m/z = 823.01; Anal. Calcd for: C₃₆H₃₀Br₂ClN₅O₄S: C, 52.48; H, 3.6 N, 8.50 %. Found C, 52.43; H, 3.55; N, 8.53 %.

4-(6,8-Dibromo-2-(4-chlorophenyl)-4-oxoquinazolin-3(4H)-yl)-N-(2-(2-hydroxy phenyl)-5-(morpholinomethyl)-4-oxothiazolidin-3-yl)benzamide (3h) White crystals (ethanol); yield 60 %; mp 160–162 °C; IR (KBr) ν_{\max} 3360(NH), 3065 (CH, aromatic), 1670 (CO, cyclic amide thiazolidinone), 1685 (CO, cyclic amide quinazoline), 1660 (CONH) and 1610 (C=N) cm⁻¹; ¹H NMR (300 MHz, DMSO-d₆): δ = 2.7 (4H, t, 2CH₂ of morpholine), 2.9 (2H, s, -CH₂-N), 3.6 (4H, t, 2CH₂ of morpholine) 3.75 (1H, s, CH, thiazolidinone ring), 5.90 (1H, s, CH of thiazolidinone ring), 7.5–7.65 (4H, m, H-2', H-6', H-2'', H-6''), 7.67–7.79 (2H, m, H-3''', H-5'''), 8.0–8.2 (6H, m, H-3', H-5', H-3'', H-5'', H-4''', H-6'''), 8.3 (1H, s, H-5), 8.4 (1H, s, H-7), 9.3 (1H, s, NH, exchangeable with D₂O), 10.34 (1H, s, OH); ¹³C NMR (100 MHz, DMSO-d₆): 172.4 (C=O), 163.6 (C=N), 162.4 (C=O), 161.5 (C=O), 155.9 (Ar/olefinic carbon), 140.7 (Ar/olefinic carbon), 140.4 (Ar/olefinic carbon), 139.2 (Ar/olefinic carbon), 139.0 (Ar/olefinic carbon), 132.5 (Ar/olefinic carbon), 130.6 (Ar/olefinic carbon), 130.0 (Ar/olefinic carbon), 129.5 (Ar/olefinic carbon), 129.3 (Ar/olefinic carbon), 127.2 (Ar/olefinic carbon), 126.5 (Ar/olefinic carbon), 126.2 (Ar/olefinic carbon), 125.5 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 124.4 (Ar/olefinic carbon), 122.6 (Ar/olefinic carbon), 116.7 (Ar/olefinic carbon), 66.7 (CH₂), 56.4 (CH₂), 52.9 (CH₂), 52.4 (CH), 48.6 (CH); MS: m/z = 824.98; Anal. Calcd for: C₃₅H₂₈Br₂ClN₅O₅S: C, 50.90; H, 3.42; N, 8.48 %. Found C, 50.89; H, 3.42; N, 8.48 %.

4-[6,8-Dibromo-2-(4-chlorophenyl)-4-oxo-4H-quinazolin-3-yl]-benzoic acid hydrazide (4) Orange crystals (methanol); yield 80 %; mp 170/172 °C; IR (KBr) ν_{\max} 3315, 3140 (NH, NH₂), 3060 (CH, aromatic), 1715 (CO, quinazoline ring) and 1645 (CO, amide) cm⁻¹; ¹H NMR (300 MHz, DMSO-d₆): δ = 4.5 (2H, s, NH₂, exchangeable with D₂O), 7.53–7.6 (4H, m, H-2', H-6', H-2'', H-6''), 7.72–7.8 (4H, m, H-3', H-5', H-3'', H-5''), 8.15 (1H, s, H-5), 8.25 (1H, s, H-7), and at 9.8 (1H, s, NH, exchangeable with D₂O); ¹³C NMR (100 MHz, DMSO-d₆): 167.2 (C=O), 164.1 (C=N), 160.4 (C=O), 154.1 (Ar/olefinic carbon), 139.2 (Ar/olefinic carbon), 136.3 (Ar/olefinic carbon), 131.2 (Ar/olefinic carbon), 131.1 (Ar/olefinic carbon), 130.1 (Ar/olefinic carbon), 129.4 (Ar/olefinic

carbon), 128.8 (Ar/olefinic carbon), 128.6 (Ar/olefinic carbon), 128.2 (Ar/olefinic carbon), 127.5 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 124.5 (Ar/olefinic carbon), 122.2 (Ar/olefinic carbon), 113.5 (Ar/olefinic carbon); MS: m/z = 547.91; Anal. Calcd for: C₂₁H₁₃Br₂ClN₄O₂: C, 49.97; H, 2.39; N, 10.21 %. Found C, 49.70; H, 2.59; N, 10.40 %.

6,8-Dibromo-2-(4-chlorophenyl)-3-(4-(5-methyl-1,3,4-oxadiazol-2-yl) phenyl) quinazolin-4(3H)-one (5) A mixture of the hydrazide **4** (5.48 g; 10 mmol) and acetic anhydride (20 ml) was refluxed for 6 h. The precipitated solid formed upon cooling, was filtered and recrystallized as yellowish white crystals (ethanol); yield 70 %; mp 174–176 °C; IR (KBr) ν_{\max} 3075 (CH, aromatic), 1703 (CO) and 1610 (C=N) cm⁻¹; ¹H NMR (300 MHz, DMSO-d₆): δ = 2.6 (3H, s, methyl), 7.5–7.7 (4H, m, H-2', H-6', H-2'', H-6''), 7.8–8.0 (4H, m, H-3', H-5', H-3'', H-5''), 8.15 (1H, s, H-5), 8.25 (1H, s, H-7); ¹³C NMR (100 MHz, DMSO-d₆): 164.9 (C=N), 164.5 (C=N), 164 (C=N), 160.5 (C=O), 154.9 (Ar/olefinic carbon), 139.4 (Ar/olefinic carbon), 135.4 (Ar/olefinic carbon), 132.2 (Ar/olefinic carbon), 131.3 (Ar/olefinic carbon), 129.5 (Ar/olefinic carbon), 129.1 (Ar/olefinic carbon), 129 (Ar/olefinic carbon), 127.5 (Ar/olefinic carbon), 126.3 (Ar/olefinic carbon), 125.2 (Ar/olefinic carbon), 121.5 (Ar/olefinic carbon), 122 (Ar/olefinic carbon), 113.5 (Ar/olefinic carbon), 20.5 (CH₃); MS: m/z = 571.91; Anal. Calcd for: C₂₃H₁₃Br₂ClN₄O₂: C, 48.24; H, 2.29; N, 9.78 %. Found C, 48.34; H, 2.33; N, 9.68 %.

6,8-Dibromo-2-(4-chlorophenyl)-3-(4-(3-methyl-5-oxo-4,5-dihydro-1H-pyrazol-1-yl)phenyl)quinazolin-4(3H)-one (6) A mixture of the hydrazide **4** (5.48 g; 10 mmol), and ethyl acetoacetate (1.3 g; 10 mmol) in absolute ethanol (20 ml) was refluxed for 6 h. The precipitated solid formed upon cooling, was filtered and recrystallized as white crystals (methanol); yield 70 %; mp 290–292 °C; IR (KBr) ν_{\max} 3065 (CH, aromatic), 1720 (CO, cyclic amide pyrazol), 1685 (CO, cyclic amide quinazoline) cm⁻¹; ¹H NMR (300 MHz, DMSO-d₆): δ = 1.9 (3H, s, CH₃), 2.8 (2H, s, CH₂ of pyrazol), 7.55–7.76 (4H, m, H-2', H-6', H-2'', H-6''), 7.8–8.0 (4H, m, H-3', H-5', H-3'', H-5''), 8.15 (1H, s, H-5), 8.25 (1H, s, H-7); ¹³C NMR (100 MHz, DMSO-d₆): 175.5 (C=O), 164 (C=N), 162.9 (C=N), 160.6 (C=O), 154.3 (Ar/olefinic carbon), 139.9 (Ar/olefinic carbon), 135.6 (Ar/olefinic carbon), 134.2 (Ar/olefinic carbon), 131.3 (Ar/olefinic carbon), 129.2 (Ar/olefinic carbon), 128 (Ar/olefinic carbon), 126.5 (Ar/olefinic carbon), 125.3 (Ar/olefinic carbon), 122 (Ar/olefinic carbon), 121.7 (Ar/olefinic carbon), 113.5 (Ar/olefinic carbon), 42.5 (CH₂), 16.5 (CH₃); MS: m/z = 585.92; Anal. Calcd for: C₂₄H₁₅Br₂ClN₄O₂: C, 52.20; H, 2.92; N, 10.15 %. Found C, 50.14; H, 2.58; N, 9.55 %.

Pharmacological screening

Cell culture and treatment

All reagents were handled in a sterile fume hood. DMEM medium and fetal bovine serum (FBS) were purchased from Gibco; phosphate-buffered saline pH 7.4 (PBS) and trypsin–EDTA were obtained from Sigma–Aldrich. Alamar blue or resazurin (Promega, Mannheim, Germany) reduction assay was used to assess the cytotoxicity of the studied samples. The growth medium (DMEM medium with 10 % FBS, 100 U/ml penicillin and 100 mg/l streptomycin), and alamar blue were stored at 48 °C, while trypsin–EDTA and FBS were stored frozen and thawed before use; PBS was stored at room temperature. The MCF-7 cells were obtained from the German Cancer Research Center (DKFZ). Cells were cultured in 50-cm² culture flasks (Corning) using DMEM medium supplemented with 10 % FBS, penicillin (100 IU/ml) and streptomycin (100 mg/ml). The culture was maintained at 37 °C in an atmosphere of 5 % CO₂ and 95 % relative humidity. The cells were transferred to a new flask every 2 days and treated with trypsin–EDTA to detach them from the flask. Cells were counted under a microscope using a hemacytometer (Hausser Scientific). Cell solutions were diluted with growth medium to a concentration of 1×10^5 cells/ml and transferred to a 96-well plate and treated with gradient concentrations of test compounds.

Resazurin cell growth inhibition assay

Alamar blue or resazurin (Promega, Mannheim, Germany) reduction assay was used to assess the cytotoxicity of the studied samples. The assay tests cellular viability and mitochondrial function. Briefly, adherent cells were grown in tissue culture flasks as previously described (Youns *et al.*, 2009, 2010, 2011; Youns and Fathy, 2013) and then harvested by treating the flasks with 0.025 % trypsin and 0.25 mM EDTA for 5 min. Once detached, cells were washed, counted, and an aliquot (5×10^3 cells) was placed in each well of a 96-well cell culture plate in a total volume of 100 μ l. Cells were allowed to attach overnight and then treated with samples. The final concentration of samples ranged from 0 to 100 μ M. After 48 h, 20 μ l resazurin 0.01 % w/v solution was added to each well and the plates were incubated at 37 °C for 1–2 h. Fluorescence was measured on an automated 96-well Infinite M2000 ProTM plate reader (Tecan, Crailsheim, Germany) using an excitation wavelength of 544 nm and an emission wavelength of 590 nm. After 48 h incubation, plates were treated with resazurin solution as above mentioned. Doxorubicin was used as positive control. Each assay was done at least three

times, with two replicates each. The viability was compared based on a comparison with untreated cells. IC₅₀ values (on cancer cells) were the concentration of sample required to inhibit 50 % of the cell proliferation and were calculated from a calibration curve by a linear regression using Microsoft Excel.

In vitro inhibition studies of EGFR tyrosine kinase

Compounds **2c**, **2d**, **3a**, **3f** were tested in vitro for inhibition of EGFR tyrosine kinase, using Kinase-Glo Plus luminescence kinase assay kit. In this method, tested compounds were dissolved in DMSO and tested at a single concentration of 10 μ M. They were then added to reaction plates containing the EGFR tyrosine kinase in assay buffer [20 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), pH 7.5, 10 mM MgCl₂, 1 mM ethylene glycol tetraacetic acid (EGTA), 0.02 % Brij35, 0.02 mg/ml bovine serum albumin (BSA), 0.1 mM Na₃VO₄, 2 mM dithiothreitol (DTT), 1 % DMSO]. Reactions were initiated by addition of a mixture of ATP (Sigma, St. Louis MO) and 33P ATP (PerkinElmer, Waltham MA) to a final concentration of 10 μ M. Reactions were carried out at room temperature for 120 min, followed by spotting of the reactions onto P81 ion exchange filter paper (Whatman Inc., Piscataway, NJ). Unbound phosphate was removed by extensive washing of filters in 0.75 % phosphoric acid (Ma *et al.*, 2008). Results are presented as percentage of enzyme inhibition and compared with those of gefitinib as a reference EGFR-TK inhibitor.

Molecular modeling

Molecular modeling studies were carried out on an Intel Core i5, 2.53 GHz processor, 4 GB memory with Windows XP 32-bit operating system using Molecular Operating Environment (MOE, 10.2008) software. Energy minimization was performed with MOE with RMSD gradient of 0.05 kcal/mol, MMFF94X force field and automatic calculation of the partial charges. The X-ray crystallographic structure of IRE (3-chloro-4-fluoro-*N*-[(4*Z*)-7-methoxy-6-(3-morpholin-4-ylpropoxy)quinazolin-4(1*H*)-ylidene]aniline) co-crystallized with EGFR was obtained from the protein data bank (PDB ID: 2ITO) (<http://www.rcsb.org/pdb/explore/explore.do?structureId=2ITO>). The ATP binding site of EGFR was prepared for docking studies where: (1) IRE was removed from the active site. (2) Hydrogen atoms were added to the structure with their standard geometry. (3) MOE Alpha Site Finder was used for the active sites detection. (4) The obtained model was then used in predicting interactions at the active site between the new compounds and EGFR.

References

- Ahmed MF, Youns M, Belal A (2016) Design, synthesis, molecular docking and anti-breast cancer activity of novel quinazolinones targeting ER α . *Acta Pol Pharm* 73(1)
- Alagarsamy V, Solomon VR, Sheorey RV, Jayakumar R (2009) 3-(3-Ethylphenyl)-2-substituted hydrazino-3*H*-quinazolin-4-one derivatives: new class of analgesic and anti-inflammatory agents. *Chem Biol Drug Des* 73:471–479
- Al-Omary FAM, Hassan GS, El-Messery SM, Nagi MN, Habib ESE, El-Subbagh HI (2013) Nonclassical antifolates, part 3: synthesis, biological evaluation and molecular modeling study of some new 2-heteroarylthio-quinazolin-4-ones. *Eur J Med Chem* 63:33–45
- Al-Suwaitan IA, Alanazi AM, Abdel-Aziz AA-M, Mohamed MA, El-Azab AS (2013) Design, synthesis and biological evaluation of 2-mercapto-3-phenethylquinazoline bearing anilide fragments as potential antitumor agents: molecular docking study. *Bioorg Med Chem Lett* 23:3935–3941
- Barlesi F, Tchouhadjian C, Doddoli C, Villani P, Greillier L, Kleisbauer J-P, Thomas P, Astoul P (2005) Gefitinib (ZD1839, Iressa) in non-small-cell lung cancer: a review of clinical trials from a daily practice perspective. *Fundam Clin Pharmacol* 19:385
- Burris HA, Hurwitz HI, Dees EC, Dowlati A, Blackwell KL, O'Neil B, Marcom PK, Ellis MJ, Overmoyer B, Jones SF, Harris JL, Smith DA, Koch KM, Stead A, Mangum S, Spector NL (2005) Phase I safety, pharmacokinetics, and clinical activity study of lapatinib (GW572016), a reversible dual inhibitor of epidermal growth factor receptor tyrosine kinases, in heavily pretreated patients with metastatic carcinomas. *J Clin Oncol* 23:5305–5313
- Chandrika PM, Yakaiah T, Rao ARR, Narsaiah B, Reddy NC, Sridhar V, Rao JV (2008) Synthesis of novel 4,6-disubstituted quinazolinone derivatives, their anti-inflammatory and anti-cancer activity (cytotoxic) against U937 leukemia cell lines. *Eur J Med Chem* 43:846–852
- Decker M (2005) Novel inhibitors of acetyl- and butyryl cholinesterase derived from the alkaloids dehydroevodiamine and rutaecarpine. *Eur J Med Chem* 40:305–313
- Dongamanti A, Gadiparthi R, Redamala R, Anireddy J, Burri N, Vantikommu J (2012) Synthesis, anti-bacterial, anti-asthmatic and anti-diabetic activities of novel 3-substituted quinazolin-4-ones using 1-butyl-3-methyl-imidazoliumtetrafluoro borate [BMIM+][BF $_4^-$] as a green, efficient and reusable catalyst under solvent free conditions. *J Chem Pharm Res* 4(8):3991–4000
- Galal SA, Abdelsamie AS, Rodríguez ML, Kerwin SM, El Diwani HI (2010) Synthesis and studying the antitumor activity of novel 5-(2-methyl benzimidazol-5-yl)-1,3,4-oxadiazole-2(3*H*)-thiones. *Eur J Chem* 1(2):67–72
- Georgy H, Abdel-Gawad N, Abbas S (2008) Synthesis and anticonvulsant activity of some quinazolin-4-(3*H*)-one derivatives. *Molecules* 13:2557–2569
- Havrylyuk D, Mosula L, Zimenkovsky B, Vasylenko O, Gzella A, Lesyk R (2010) Synthesis and anticancer activity evaluation of 4-thiazolidinones containing benzothiazole moiety. *Eur J Med Chem* 45:5012–5021
- Hennequin LF, Stokes ESE, Thomas AP, Johnstone C, Plé PA, Ogilvie DJ, Dukes M, Wedge SR, Kendrew J, Curwen JO (2002) Novel 4-anilinoquinazolines with C-7 basic side chains: design and structure activity relationship of a series of potent, orally active, VEGF receptor tyrosine kinase inhibitors. *J Med Chem* 45:1300
- Ismail MAH, Barker S, Abau El Ella DA, Abouzid KAM, Toubar RA, Todd MH (2006) Design and synthesis of new tetrazolyl- and carboxy-biphenylmethyl-quinazolin-4-one derivatives as angiotensin II AT $_1$ receptor antagonists. *Med Chem* 49:1526–1535
- Kopper L (2008) Lapatinib: a sword with two edges. *Pathol Oncol Res* 14:1–8
- Lv P-C, Li H-Q, Sun J, Zhou Y, Zhu H-L (2010) Synthesis and biological evaluation of pyrazole derivatives containing thiourea skeleton as anticancer agents. *Bioorg Med Chem* 18:4606–4614
- Ma H, Deacon S, Horiuchi K (2008) The challenge of selecting protein kinase assays for lead discovery optimization. *Expert Opin Drug Discov* 3:607–621
- Mohamed MS, Kamel MM, Kassem EMM, Abotaleb N, Abd El-moez SI, Ahmed MF (2010) Novel 6,8-dibromo-4(3*H*)quinazolinone derivatives of anti-bacterial and anti-fungal activities. *Eur J Med Chem* 45:3311–3319
- Olayioye MA, Neve RM, Lane HA, Hynes NE (2000) The ErbB signaling network: receptor heterodimerization in development and cancer. *EMBO J* 19:3159–3167
- Pollack VA, Savage DM, Baker DA, Tsaparikos KE, Sloan DE, Moyer JD, Barbacci EG, Pustilnik LR, Smolarek TA, Davis JA, Vaidya MP, Arnold LD, Doty JL, Iwata KK, Morin MJ (1999) Inhibition of epidermal growth factor receptor-associated tyrosine phosphorylation in human carcinomas with CP-358,774: dynamics of receptor inhibition in situ and antitumor effects in athymic mice. *J Pharmacol Exp Ther* 291:739–748
- Rusnak DW, Lackey K, Affleck K, Wood ER, Alligood KJ, Rhodes N, Keith BR, Murray DM, Knight WB, Mullin RJ, Gilmer TM (2001) The effects of the novel, reversible epidermal growth factor receptor/ErbB-2 tyrosine kinase inhibitor, GW2016, on the growth of human normal and tumor-derived cell lines in vitro and in vivo. *Mol Cancer Ther* 1:85–94
- Sariego J (2010) Breast cancer in the young patient. *Am Surg* 76(12):1397–1400
- Siegel R, Ma J, Zou Z, Jemal A (2014) Cancer statistics, 2014. *CA Cancer J Clin* 64:9–29
- Wakeling AE, Guy SP, Woodburn JR, Ashton SE, Curry BJ, Barker AJ, Gibson KH (2002) ZD1839 (Iressa): an orally active inhibitor of epidermal growth factor signaling with potential for cancer therapy. *Cancer Res* 62:5749–5754
- Weiss JR, Moysich KB, Swede H (2005) Epidemiology of male breast cancer. *Cancer Epidemiol Biomark Prev* 14:20–26
- Wood ER, Truesdale AT, McDonald OB, Yuan D, Hassell A, Dickerson SH, Ellis B, Pennisi C, Horne E, Lackey K, Alligood KJ, Rusnak DW, Gilmer TM, Shewchuk L (2004) A unique structure for epidermal growth factor receptor bound to GW572016 (Lapatinib): relationships among protein conformation, inhibitor off-rate, and receptor activity in tumor cells. *Cancer Res* 64:6652
- Youns M, Fathy G (2013) Upregulation of extrinsic apoptotic pathway in curcumin-mediated antiproliferative effect on human pancreatic carcinogenesis. *J Cell Biochem* 114:2644–2656
- Youns M, Efferth T, Reichling J, Fellenberg K, Bauer A, Hoheisel JD (2009) Gene expression profiling identifies novel key players involved in the cytotoxic effect of Artesunate on pancreatic cancer cells. *Biochem Pharmacol* 78:273–283
- Youns M, Hoheisel JD, Efferth T (2010) Toxicogenomics for the prediction of toxicity related to herbs from traditional chinese medicine. *Planta Med* 76(17):2019–2025
- Youns M, Efferth T, Hoheisel JD (2011) Transcript profiling identifies novel key players mediating the growth inhibitory effect of NS-398 on human pancreatic cancer cells. *Eur J Pharmacol* 650:170–177
- Yun C-H, Boggon TJ, Li Y, Woo MS, Greulich H, Meyerson M, Eck MJ (2007) Structures of lung cancer-derived EGFR mutants and inhibitor complexes: mechanism of activation and insights into differential inhibitor sensitivity. *Cancer Cell* 11:217–227