

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/225959266>

Gas-phase enthalpies of formation, acidities, and strain energies of the [m, n]polyprismanes ($m \geq 2$; $n = 3-8$; $m \times n \leq 16$): A CBS-Q//B3, G4MP2, and G4 theoretical study

ARTICLE in THEORETICAL CHEMISTRY ACCOUNTS · NOVEMBER 2010

Impact Factor: 2.23 · DOI: 10.1007/s00214-010-0780-0

CITATIONS

18

READS

34

2 AUTHORS:



Sierra Rayne

Chemologica Research

270 PUBLICATIONS 2,262 CITATIONS

SEE PROFILE



Kaya Forest

Saskatchewan Polytechnic, Moose Jaw, Ca...

150 PUBLICATIONS 879 CITATIONS

SEE PROFILE

Gas-phase enthalpies of formation, acidities, and strain energies of the $[m, n]$ polyprismanes ($m \geq 2$; $n = 3-8$; $m \times n \leq 16$): a CBS-Q//B3, G4MP2, and G4 theoretical study

Sierra Rayne · Kaya Forest

Received: 3 May 2010 / Accepted: 1 July 2010 / Published online: 14 July 2010
© Springer-Verlag 2010

Abstract Gas-phase standard state (298.15 K, 1 atm) enthalpies of formation ($\Delta_f H_{(g)}^\circ$) and enthalpies ($\Delta_{\text{acid}} H_{(g)}^\circ$) and free energies ($\Delta_{\text{acid}} G_{(g)}^\circ$) of acid dissociation were calculated for the $[m, n]$ polyprismanes ($m \geq 2$; $n = 3-8$; $m \times n \leq 16$) using the composite method CBS-Q//B3, G4MP2, and G4 levels of theory. Excellent agreement was obtained between the theoretical estimates and experimental values for [2, 4]polyprismane (cubane), the only member with reliable experimental data. Gas-phase acidities of the [2, 3], [2, 4], and [2, 5]polyprismanes correlate well with percent s-character of the C–H bonds as determined by nuclear magnetic resonance spectroscopy. Based on this correlation, and calculations on the [2, n] ($n = 3-8$) series, no substantial change in the gas-phase acidity or C–H bond s-character is expected with increasing [2, n] polyprismane ring size at $n = 6-8$. Where three or more stacked $[m, n]$ polyprismane rings are present ($m > 2$), geometry optimizations for the deprotonated anions converged on cage-opened non-prismatic geometries, suggesting these compounds may be structurally unstable to gas-phase deprotonation/protonation cycles and potentially preventing reliable experimental measurement or calculation of their acidities. Total strain energies (E_{str}) increase

with larger $[m, n]$ polyprismane size, and strain energies per one C–C bond ($E_{\text{str}}^{\text{CC}}$) increase with increasing stack height within a given ring size series. Within a given stack height, $E_{\text{str}}^{\text{CC}}$ reaches a minimum with a ring size of five, increasing with either decreasing or increasing ring size away from this minima. E_{str} and $E_{\text{str}}^{\text{CC}}$ do not correlate with gas-phase acidity, likely because the deprotonated anionic [2, n]polyprismane geometry retains approximately equal strain as the undissociated species despite substantial geometry changes due to carbon acid dissociation.

Keywords Polyprismanes · Enthalpy of formation · Gas-phase acidity · Strain energy · Composite methods

1 Introduction

Cage-type polyhedral hydrocarbons have attracted substantial attention because of their non-classical carbon bonding, unusual ring strain, σ -aromaticity/antiaromaticity, range of possible isomerization reactions, and unique physicochemical properties [1–12]. A subset of these general structures, the $[m, n]$ polyprismanes are of particular interest due to the presence of semiplanar carbon centers, applications in nanoengineered structures, and use as potential high energy density materials (HEDMs) and energy storage systems [4, 8, 13–16]. To help provide a common terminology for the field, Lewars [7] has developed a $[m, n]$ polyprismane numbering system whereby ‘ m ’ denotes the number of stacked rings and ‘ n ’ is the carbon size of each stacked ring. Only the [2, 3] (prismane), [2, 4] (cubane), and [2, 5] (pentaprismane) members have been synthesized [17–19]; the remainder remain high priority synthetic targets whose thermodynamic properties are of substantial significance.

Electronic supplementary material The online version of this article (doi:10.1007/s00214-010-0780-0) contains supplementary material, which is available to authorized users.

S. Rayne (✉)
Ecologica Research, Penticton, BC V2A 8J3, Canada
e-mail: rayne.sierra@gmail.com

K. Forest
Department of Chemistry, Okanagan College, Penticton,
BC V2A 8E1, Canada

In the current work, we report the results of Complete Basis Set (CBS) and Gaussian-4 (G4) method calculations on the gas-phase enthalpies of formation, acidities, and strain energies for the $[m, n]$ polyprismanes ($m \geq 2$; $n = 3-8$; $m \times n \leq 16$). Because of the interest in using these compounds as HEDMs and in nanostructures, high level theoretical estimates are needed to better constrain the thermochemical properties of widely studied members, and to also help assess the accuracy of lower cost density functional theory (DFT) approaches likely to be employed for future studies of larger $[m, n]$ polyprismanes. The physicochemical properties of the $[m, n]$ polyprismanes are also of fundamental value, given the widespread interest in studies of highly strained organic compounds that probe the limits of our understanding regarding structure and bonding.

2 Computational methods

Calculations were conducted with Gaussian 09 [20] and the CBS-Q//B3 [21, 22], G4MP2 [23], and G4 [24] composite methods using gas-phase starting geometries obtained with the PM6 semiempirical method [25] as implemented in MOPAC 2009 (<http://www.openmopac.net/>; v. 9.281). Molecular enthalpies and free energies include zero point and thermal corrections. Only the lowest energy conformation of each molecule was considered. Optimized structures were confirmed as true minima by vibrational analysis at the same level. G09 archive entries are provided in the Supplementary Materials with optimized geometries and energies at each stage of the CBS-Q//B3, G4MP2, and G4 calculation processes. All compounds have D_{nh} prismatic symmetry with no imaginary frequencies.

3 Results and discussion

Gas-phase standard state (298.15 K, 1 atm) enthalpies of formation ($\Delta_f H^\circ_{(g)}$) for the $[m, n]$ polyprismanes ($m \geq 2$; $n = 3-8$; $m \times n \leq 16$; Fig. 1) are given in Table 1. Geometrical parameters do not vary significantly with level of theory (bond lengths within $\sim \pm 0.001-0.002$ Å, bond angles within $\sim \pm 0.1-0.2^\circ$), and are in agreement with previous theoretical and experimental reports [7, 13–18, 26–44] (Fig. 1). $\Delta_f H^\circ_{(g)}$ values were calculated for all compounds using the atomization enthalpy approach [45, 46]. Selected homodesmotic and isodesmotic approaches were also used for applicable compounds. Due to computational expense, G4MP2 calculations were not completed for the [2, 8]polyprismane, and G4 calculations were only completed on the [2, 3], [2, 4], and [3, 3]polyprismanes. An experimental $\Delta_f H^\circ_{(g)}$ of 148.7 ± 1.0 kcal/mol is available for the




Fig. 1 Structures with gas-phase standard state (298.15 K, 1 atm) CBS-Q//B3 optimized bond lengths and angles for the $[m, n]$ polyprismanes ($m \geq 2$; $n = 3-8$; $m \times n \leq 16$)

[2, 4]polyprismane (cubane) [47]; our CBS-Q//B3 (148.9 kcal/mol), G4MP2 (144.2 kcal/mol), and G4 (145.2 kcal/mol) atomization-based estimates are in good agreement.

We note that the experimental $\Delta_f H^\circ_{(g)}$ for cubane has been called into question based on sublimation data [48, 49] that conflicts with the historical estimate, as well as justified critiques (i.e., concerns over subjective corrections for unburned residues in the original measurements)

Table 1 Gas-phase standard state (298.15 K, 1 atm) enthalpies of formation ($\Delta_f H^\circ_{(g)}$) for the $[m, n]$ polyprismanes ($m \geq 2$; $m \times n \leq 16$) at the CBS-Q//B3, G4MP2, and G4 levels of theory

	Current work	Literature
[2, 3]	136.1 _{CBS-Q//B3} , 132.2 _{G4MP2} , 133.2 _{G4}	136.7 _{MMP2A} [77], 121.9 _{MNDO} [77, 78], 165.0 _{AM1} [26, 77], 136.0 _{PM3} [77], 136.4/136.2 _{SCF/6-31G*(RMP2)} [26], 136.8 ± 5.8 _{(HF,MP2,B3LYP)/6-31G*} [29], 130.7/129.9 _{G2(MP2,SVP)} [79], 137.1/131.4 _{G2(MP2)} [79, 80], 133.0 _{G2(MP2)} [28], 137.8/131.9 _{G2(MP3)} [79], 136.2/130.7 _{G2} [79], 135.3/131.3 _{CBS-Q} [79], 132.3/134.3 _{CBS-APNO} [79], 135.0/134.0/135.1 _{G3(B3LYP)} [81, 82], 134.6/133.8 _{G3(MP2)} [80], 135.7/133.8 _{G3} [80]
[2, 4]	148.9 _{CBS-Q//B3} , 144.2 _{G4MP2} , 145.2 _{G4}	148.7 ± 1.0 _{expt} [47], 148.8 _{MM2} [78], 148.9 _{MM2} [37], 99.1 _{MNDO} [12 [26], 33], 151.2 _{AM1} [26], 151.1 _{AM1} [13], 113.8 _{PM3} [13], 149.9 _{HF/STO-3G} [83], 137.9 _{HF/4-31G} [83], 139.8 _{HF/6-31G*} [83], 145.1 _{SCF/3-21G} [37], 147.1 _{SCF/6-31G*} [37], 148.5/146.3 _{SCF/6-31G*(RMP2)} [84], 171.1 _{RHF/6-31G*} [14], 170.8 _{SVWN/6-31G*} [27], 124.3 _{BP/6-31G*} [27], 114.8 _{BLYP/6-31G*} [27], 122.3 _{B3LYP/6-31G*} [27], 108.5 _{BLYP/6-31+G**/BLYP/6-31G*} [27], 128.6 _{B3LYP/6-31+G**/BLYP/6-31G*} [27], 158.1 _{MP2/6-31G**/RHF/6-31G*} [14], 149.1 _{MP2/6-31G(d)**/RHF/6-31G(d)*} [16], 142.7 ± 0.9 _{(HF,MP2,B3LYP)/6-31G*} [29], 146.0 _{G3(MP2)} [85], 143.2 _{G2(MP2)} [86], 142.3 _{G2} [86], 145.1 _{G3/B3LYP} [86], 145.5 _{G3} [86], 146.5 ± 2.3 _{G2(MP2),G2,G3//B3LYP,G3[avg],homodesmic} [86], 159.0 ± 1.2^b [87], 158.5 ^c [87]
[2, 5]	124.4 _{CBS-Q//B3} , 119.5 _{G4MP2}	114.7 _{MM2} [78], 72.2 _{MNDO} [26], 117.7 _{AM1} [26], 115.4 _{SCF/3-21G} [37], 121.2/122.0 _{SCF/6-31G*} [37], 119.6/120.5 _{SCF/6-31G*(RMP2)} [37], 123.9 ± 7.2 _{(HF,MP2,B3LYP)/6-31G*} [29]
[2, 6]	158.9 _{CBS-Q//B3} , 152.9 _{G4MP2}	129.2 _{MM2} [78], 80.4 _{MNDO} [26], 139.9 _{AM1} [26], 160.3 _{SCF/3-21G} [37], 154.5/154.7 _{SCF/6-31G*} [37], 153.1/153.6 _{SCF/6-31G*(RMP2)} [37], 157.3 ± 7.9 _{(HF,MP2,B3LYP)/6-31G*} [29]
[2, 7]	214.2 _{CBS-Q//B3} , 207.1 _{G4MP2}	173.8 _{MM2} [78], 116.4 _{MNDO} [26], 191.0 _{AM1} [26], 212.0 _{SCF/6-31G*(RMP2)} [37], 215.5 ± 7.7 _{(HF,MP2,B3LYP)/6-31G*} [29]
[2, 8]	279.3 _{CBS-Q//B3}	172.7 _{MNDO} [78], 259.2 _{AM1} [26], 283.1 _{SCF/6-31G*(RMP2)} [26], 285.1 ± 6.5 _{(HF,MP2,B3LYP)/6-31G*} [29]
[3, 3]	310.7 _{CBS-Q//B3} , 303.6 _{G4MP2} , 306.1 _{G4}	n/a ^a
[3, 4]	383.2 _{CBS-Q//B3} , 373.9 _{G4MP2}	n/a
[3, 5]	385.9 _{CBS-Q//B3} , 375.4 _{G4MP2}	n/a
[4, 3]	478.1 _{CBS-Q//B3} , 467.7 _{G4MP2}	n/a
[4, 4]	598.1 _{CBS-Q//B3} , 583.9 _{G4MP2}	n/a
[5, 3]	643.1 _{CBS-Q//B3} , 629.5 _{G4MP2}	n/a

Previous theoretical and experimental values reported in the literature are also provided. Subscripts indicate levels of theory or experimental values. Values are in kcal/mol

^a Theoretical or experimental literature values not available

^b Calculated using the constant enthalpy of formation differences between the hydrocarbon and its methyl ester during studies on 1,4-dimethylcubane dicarboxylate

^c Calculated using the bicyclobutane strain energy difference between the hydrocarbon and its methyl ester during studies on 1,4-dimethylcubane dicarboxylate


regarding the quality of the solid state combustion data. Karpushenkava et al. [29] have recently proposed a revised experimental $\Delta_f H^\circ_{(g)}$ of 142.7 ± 1.0 kcal/mol for cubane based on the updated sublimation studies. Our G4 $\Delta_f H^\circ_{(g)}$ atomization-based estimate lies between these two competing $\Delta_f H^\circ_{(g)}$ values and cannot clearly discriminate as to which—if either—is correct. No other experimental $\Delta_f H^\circ_{(g)}$ values are available in the NIST database [50, 51] for these $[m, n]$ polyprismanes.

For all compounds, the G4 and G4MP2 $\Delta_f H^\circ_{(g)}$ atomization-based estimates are lower than the corresponding CBS-Q//B3 atomization-based estimates by 4–14 kcal/mol, with the difference between the CBS-Q//B3 and G4/


G4MP2 values increasing with increasing molecular size. We have previously shown that, in general, $\Delta_f H^\circ_{(g)}$ estimates using the atomization approach and CBS-Q//B3 method exhibit scaling errors with molecular size, whereas the G4/G4MP2 level methods display effective chemical accuracy without substantial scaling errors [52–54]. Thus, we expect the CBS-Q//B3 $\Delta_f H^\circ_{(g)}$ estimate of 279.3 kcal/mol for the [2, 8]polyprismane to be too high by about 14 kcal/mol; the true $\Delta_f H^\circ_{(g)}$ for the [2, 8]polyprismane is likely near 265 kcal/mol. A number of previous $\Delta_f H^\circ_{(g)}$ estimates have been reported for the [2, n]polyprismanes under consideration at various semiempirical, ab initio, density functional, and composite levels of

theory, as provided in Table 1. While there is wide variation among different methods, we note that in some cases, selected historical estimates (often at much lower levels of theory, such as SCF/3-21G) are very close to the current G4/G4MP2 values.


Various isodesmic and homodesmic approaches have been applied to estimate the $\Delta_f H_{(g)}^\circ$ of $[2, n]$ polyprismanes in the literature, particularly for the $[2, 3]$ and $[2, 4]$ members. Wang and Law [55] applied the following isodesmic reaction for $(CH)_n$ isomers (i.e., $[2, n]$ polyprismanes):



Karpushenkava et al. [29] proposed the following homodesmic reaction for $(CH)_n$ isomers:



Homodesmic reactions anchored to the experimental $\Delta_f H_{(g)}^\circ$ of cubane ($[2, 4]$ polyprismane) can also be used to estimate $\Delta_f H_{(g)}^\circ$ of other $[2, n]$ polyprismanes, yielding an equation with the following general stoichiometry:



Application of these schemes toward the $[2, n]$ polyprismanes ($n \leq 8$) using our CBS-Q//B3, G4MP2, and G4 data yields reasonable agreement with the atomization-based $\Delta_f H_{(g)}^\circ$ estimates (Table 2). However, isodesmic and homodesmic approaches have inherently lower $\Delta_f H_{(g)}^\circ$ precision than atomization methods due to the respective uncertainties of the underlying experimental data used in such calculations. The experimental $\Delta_f H_{(g)}^\circ$ for carbon (171.29 ± 0.11 kcal/mol [56]) and hydrogen (52.103 ± 0.001 kcal/mol [56]) atoms are much better constrained for the atomization method than are experimental $\Delta_f H_{(g)}^\circ$ for methane (range of $\Delta_f H_{(g)}^\circ$ from -17.5 ± 0.3 to -17.9 kcal/mol [57, 58, 59]), ethane (-20.0 ± 0.1 to -20.2 ± 0.1 kcal/mol [60, 61]), cyclohexane (-29.4 ± 0.2 to -29.8 kcal/mol [62, 63]), adamantane (-30.8 ± 1 to -33.2 kcal/mol [64, 65, 66]), and cubane (142.7 ± 1.0 to 148.7 ± 1.0 kcal/mol [29, 47]) used in the isodesmic/homodesmic approaches.

This variability in the experimental database leads to inherent variability in the calculated $\Delta_f H_{(g)}^\circ$ from isodesmic and homodesmic schemes, particularly if amplified by stoichiometries greater than unity in the respective equations (e.g., a $[2, 8]$ polyprismane $\Delta_f H_{(g)}^\circ$ isodesmic calculation employs stoichiometric coefficients of 32 methane and 24 ethane molecules per molecule of $[2, 8]$ polyprismane). Choosing the input $\Delta_f H_{(g)}^\circ$ for isodesmic/homodesmic methods based on the best $\Delta_f H_{(g)}^\circ$ agreement for a member of the target compound set with an experimental data point is not a rigorous strategy, as the output experimental data being compared to for method calibration may itself be incorrect. Cubane offers an excellent case study in this

regard, as it is the only $[m, n]$ polyprismane with experimental $\Delta_f H_{(g)}^\circ$ data, but the estimates (neither of which can be readily discounted) differ by 6 kcal/mol. In addition, the available $\Delta_f H_{(g)}^\circ$ for methane, ethane, cyclohexane, and adamantane allow us to tune the G4 predicted $\Delta_f H_{(g)}^\circ$ for cubane from 135.9 to 144.3 kcal/mol with the isodesmic scheme and from 146.3 to 151.7 kcal/mol at the G4MP2 level with the homodesmic scheme.

In contrast, a single $\Delta_f H_{(g)}^\circ$ estimate arises for hydrocarbons from the atomization method using the atomic enthalpies for which a scientific consensus has been reached. The resulting $\Delta_f H_{(g)}^\circ$ is a value all researchers using a particular level of theory and compound should be able to reproduce (ignoring the 0.003 and 0.06% experimental errors in the $\Delta_f H_{(g)}^\circ$ for the hydrogen and carbon atoms, respectively). In an atomization approach error analysis, using the lower (C: 171.18 kcal/mol; H: 52.101 kcal/mol) and upper (C: 171.40 kcal/mol; H: 52.104 kcal/mol) bounds of the atomic experimental $\Delta_f H_{(g)}^\circ$ for carbon and hydrogen atoms varies the atomization enthalpy method $\Delta_f H_{(g)}^\circ$ estimate for cubane at the G4 level by $< \pm 0.9$ kcal/mol, much less than the $\pm \sim 9$ kcal/mol $\Delta_f H_{(g)}^\circ$ estimate tuning achievable using the isodesmic scheme and range of experimental $\Delta_f H_{(g)}^\circ$ for methane and ethane. For many isodesmic and homodesmic approaches in the literature, the underlying experimental $\Delta_f H_{(g)}^\circ$ being applied are often not provided nor is an effort made to conduct a corresponding sensitivity analysis that contextualizes the likely method precision. Although isodesmic and homodesmic approaches can offer potential insights into the thermochemistry of highly strained molecules, with the high variability in the target experimental $\Delta_f H_{(g)}^\circ$ database to which these methods may be calibrated, and the imprecision of the underlying $\Delta_f H_{(g)}^\circ$ values for non-target molecules used in the calculations, these methods are susceptible to a higher level of imprecision than atomization approaches.

The Benson group additivity rules for estimating $\Delta_f H_{(g)}^\circ$ have demonstrated success for a broad range of relatively unstrained organic compounds [67–70]. Application of this approach to the $[m, n]$ polyprismanes only offers modest agreement with the CBS-Q//B3, G4MP2, and G4 atomization enthalpy results for cubane and pentaprismane ($[2, 5]$ polyprismane) (Table 3). The disagreement between the methods increases with increasing ring size (n) and stack height (m), with discrepancies approaching 200 kcal/mol for the $[5, 3]$ polyprismane. In general, the Benson additivity scheme underestimates $\Delta_f H_{(g)}^\circ$ for highly strained compounds such as the $[m, n]$ polyprismanes because appropriate strain correction factors are not available. Our Benson approach $\Delta_f H_{(g)}^\circ$ of 141.6 kcal/mol for cubane is in excellent agreement with the previous Benson estimate of 142.0 kcal/mol by Hassenruck et al.

Table 2 Gas-phase standard state (298.15 K, 1 atm) enthalpies of formation ($\Delta_f H_{(g)}^\circ$) for the $[m, n]$ polyprismanes ($m = 2$; $m \times n \leq 16$) at the CBS-Q//B3, G4MP2, and G4 levels of theory using isodesmic and homodesmic reaction schemes previously employed in the literature

	Isodesmic ^a			Homodesmic ^b			Homodesmic ^c		
	CBS-Q//B3	G4MP2	G4	CBS-Q//B3	G4MP2	G4	CBS-Q//B3	G4MP2	G4
[2, 3]	133.0 [−3.1]	132.2 [−0.7]	133.2 [−2.0]	137.1 [1.0]	135.9 [3.7]		136.0 [−0.1] ^d /131.5 ^e [−4.6]	135.6 ^d [3.4]/131.1 ^e [−1.1]	135.8 ^d [2.6]/131.3 [−1.9]
[2, 4]	144.7 [−4.2]	144.2 [−1.1]	145.2 [−2.7]	150.2 [1.3]	149.1 [4.9]		n/a ^f	n/a	n/a
[2, 5]	119.1 [−5.3]	119.5 [−1.3]	n/c ^g	126.0 [1.6]	125.7 [6.2]		124.1 ^d [−0.3]/116.6 ^e [−7.8]	125.2 ^d [5.7]/117.7 ^e [−1.8]	n/c
[2, 6]	152.6 [−6.3]	152.9 [−1.5]	n/c	160.8 [1.9]	160.3 [7.4]		158.6 ^d [−0.3]/149.6 ^e [−9.3]	159.7 ^d [6.8]/150.7 ^e [−2.2]	n/c
[2, 7]	206.8 [−7.4]	207.1 [−1.9]	n/c	216.4 [2.2]	215.7 [8.6]		213.8 ^d [−0.3]/203.3 ^e [−10.9]	215.0 ^d [7.9]/204.5 ^e [−2.6]	n/c
[2, 8]	270.8 [−8.5]	n/c	n/c	281.8 [2.5]	n/c		278.8 ^d [−0.4]/266.8 ^e [−12.5]	n/c	n/c

Deviations from the corresponding atomization enthalpy approach results at the same level of theory as provided in Table 1 are given in brackets. Values are in kcal/mol

^a Based on the reaction $[2, n]$ polyprismene + $4n$ CH₄ → $3n$ C₂H₆ using $\Delta_f H_{(g)}^\circ$ of −17.8 kcal/mol [88, 89] and −20.1 kcal/mol [88, 89] for methane and ethane, respectively

^b Based on the reaction $[2, n]$ polyprismene + $n/2$ cyclohexane → $n/2$ adamantane using $\Delta_f H_{(g)}^\circ$ of −29.8 kcal/mol [90] and −32.1 kcal/mol [66, 91] for cyclohexane and adamantane, respectively

^c Based on the reaction $(4/n)$ $[2, n]$ polyprismene → $[2, 4]$ polyprismene

^d Using an experimental $[2, 4]$ polyprismene $\Delta_f H_{(g)}^\circ$ of 148.7 kcal/mol

^e Using an experimental $[2, 4]$ polyprismene $\Delta_f H_{(g)}^\circ$ of 142.7 kcal/mol

^f Not applicable as the experimental $\Delta_f H_{(g)}^\circ$ for $[2, 4]$ polyprismene is used in the homodesmic reaction scheme

^g Not calculated due to computational expense for $[2, n]$ polyprismene

[71], with the small difference between the two values due to our use of the more recent 1993 group additivity value updates [69]. Using the pre-1993 additivity values, we also obtain a Benson $\Delta_f H_{(g)}^\circ$ of 142.0 kcal/mol for this compound.

For the $[m, n]$ polyprismanes, the ideal bond lengths and angles in the C–(H)(C)₃ and C–(C)₄ Benson constants do not accurately model the strained terminal C(CH)₃H (for $m = 2$) and C(C)(CH)₂H (for $m > 2$) groups or intrastack C(C)₂(CH)₂ (for units adjacent to the terminal rings where $m = 3$), C(C)₃(CH) (for units adjacent to the terminal rings where $m > 3$), and C(C)₄ (for units not adjacent to the terminal rings where $m > 5$; i.e., the central ring in [5, 3]polyprismene) functionalities present in these compounds. In addition, the cycloalkane rings present in $[m, n]$ polyprismanes are planar with flat cage faces, whereas experimentally, the corresponding C₄ and higher cycloalkane rings upon which the Benson additivity values are developed (i.e., cyclobutane, cyclopentane, cyclohexane, etc.) are non-planar (e.g., chair conformation of cyclohexane). Consequently, there is a failure to incorporate the known $[m, n]$ polyprismene non-classical bond angles and lengths in both the Benson carbon hybridization and ring approximation group additivity corrections, resulting in ever increasing deviations from high-level theoretical calculations using atomization, isodesmic, and/or homodesmic reaction schemes. Based on this rationale, the good agreement between the Benson and experimental $\Delta_f H_{(g)}^\circ$ for cubane is likely due to error cancelation, otherwise we would expect this additivity estimation approach to yield reasonably good agreement with the high level calculations for other $[2, n]$ polyprismanes.

Gas-phase enthalpies ($\Delta_{\text{acid}} H_{(g)}^\circ$) and free energies ($\Delta_{\text{acid}} G_{(g)}^\circ$) of acid dissociation were also calculated for the $[m, n]$ polyprismanes ($m = 2$; $m \times n \leq 16$) at the CBS-Q//B3, G4MP2, and G4 levels of theory (Table 4). All methods yielded $\Delta_{\text{acid}} H_{(g)}^\circ$ and $\Delta_{\text{acid}} G_{(g)}^\circ$ estimates within the corresponding experimental errors for cubane, the only polyprismene with experimental gas-phase acidity data. Among the several previous theoretical reports on the $\Delta_{\text{acid}} H_{(g)}^\circ$ for cubane, only the ‘best theoretical estimate’ by Ritchie et al. (406.3 kcal/mol; [35]) and the MP2/6-31 + G(d)//MP2/6-31 + G(d) estimate by Hare et al. (404.7 kcal/mol; [1]) are within the experimental error and in agreement with our CBS-Q//B3, G4MP2, and G4 estimates. To the best of our knowledge, no previous $\Delta_{\text{acid}} H_{(g)}^\circ$ / $\Delta_{\text{acid}} G_{(g)}^\circ$ estimates have been reported for the $[2, n]$ ($n = 3, 5, 6, 7, 8$) members, and no prior $\Delta_{\text{acid}} G_{(g)}^\circ$ theoretical estimate has been put forward for cubane.

Excellent agreement was also obtained between the three computational methods, with differences of no more than 1–1.5 kcal/mol for each compound. For the [2, 3] and [2, 4]polyprismanes with both CBS-Q//B3 and G4

Table 3 Gas-phase standard state (298.15 K, 1 atm) enthalpies of formation ($\Delta_f H_{(g)}^\circ$) for the $[m, n]$ polyprismanes ($m \geq 2$; $m \times n \leq 16$) estimated using the Benson group additivity values (GAVs) from Cohen and Benson [69] and Sabbe et al. [70]

	Cohen and Benson [69] GAVs	Sabbe et al. [70]
[2, 3]	121.4 [−14.7 _{CBS-Q//B3} , −10.8 _{G4MP2} , −11.8 _{G4}]	127.4 [−8.7 _{CBS-Q//B3} , −4.8 _{G4MP2} , −5.8 _{G4}]
[2, 4]	141.6 [−7.3 _{CBS-Q//B3} , −2.6 _{G4MP2} , −3.6 _{G4}]	148.2 [−0.7 _{CBS-Q//B3} , 4.0 _{G4MP2} , 3.0]
[2, 5]	124.2 [−0.2 _{CBS-Q//B3} , 4.7 _{G4MP2}]	132.8 [8.4 _{CBS-Q//B3} , 13.3 _{G4MP2}]
[2, 6]	133.4 [−25.5 _{CBS-Q//B3} , −19.5 _{G4MP2}]	143.1 [−15.8 _{CBS-Q//B3} , −9.8 _{G4MP2}]
[2, 7]	167.6 [−46.6 _{CBS-Q//B3} , −39.5 _{G4MP2}]	180.2 [−34.0 _{CBS-Q//B3} , −26.9 _{G4MP2}]
[2, 8]	196.6 [−82.7 _{CBS-Q//B3}]	210.8 [−68.5 _{CBS-Q//B3}]
[3, 3]	229.8 [−80.9 _{CBS-Q//B3} , −73.8 _{G4MP2} , −76.3 _{G4}]	239.2 [−71.5 _{CBS-Q//B3} , −64.4 _{G4MP2} , −66.9 _{G4}]
[3, 4]	276.0 [−107.2 _{CBS-Q//B3} , −97.9 _{G4MP2}]	286.4 [−96.8 _{CBS-Q//B3} , −87.5 _{G4MP2}]
[3, 5]	265.8 [−120.1 _{CBS-Q//B3} , −109.6 _{G4MP2}]	279.3 [−106.6 _{CBS-Q//B3} , −96.1 _{G4MP2}]
[4, 3]	338.2 [−139.9 _{CBS-Q//B3} , −129.5 _{G4MP2}]	351.0 [−127.1 _{CBS-Q//B3} , −116.7 _{G4MP2}]
[4, 4]	410.4 [−187.7 _{CBS-Q//B3} , −173.5 _{G4MP2}]	424.6 [−173.5 _{CBS-Q//B3} , −159.3 _{G4MP2}]
[5, 3]	446.6 [−196.5 _{CBS-Q//B3} , −182.9 _{G4MP2}]	462.8 [−180.3 _{CBS-Q//B3} , −166.7 _{G4MP2}]

Deviations from the corresponding atomization enthalpy approach results at the CBS-Q//B3, G4MP2, and G4 levels of theory as provided in Table 1 are given in brackets

Table 4 Gas-phase standard state (298.15 K, 1 atm) enthalpies ($\Delta_{\text{acid}} H_{(g)}^\circ$) and free energies ($\Delta_{\text{acid}} G_{(g)}^\circ$) of acid dissociation for the $[m, n]$ polyprismanes ($m = 2$; $m \times n \leq 16$) at the CBS-Q//B3, G4MP2, and G4 levels of theory

$\Delta_{\text{acid}} H_{(g)}^\circ$		$\Delta_{\text{acid}} G_{(g)}^\circ$	
Current work	Literature	Current Work	Literature
[2, 3] 401.4 _{CBS-Q//B3} , 399.8 _{G4MP2} , 401.1 _{G4}	n/a ^a	393.5 _{CBS-Q//B3} , 392.0 _{G4MP2} , 393.2 _{G4}	n/a
[2, 4] 407.2 _{CBS-Q//B3} , 406.0 _{G4MP2} , 406.9 _{G4}	404 ± 3 _{expt} [1], 411.6 _{AM1} [35], 442.5 _{SCF/6-31G} [35], 440.5 _{SCF/6-31G*} [35], 431.1 _{SCF/6-31 + G} [35], 429.2 _{SCF/6-31 + G*} [35], 416.9/406.7 _{B3LYP/6-311 + G**} [2], 416.3 _{MP2/6-31 + G} [35], 406.3 ^{best} theoretical estimate [35], 404.7 _{MP2/6-31 + G(d)/MP2/6-31 + G(d)} [1]	399.3 _{CBS-Q//B3} , 398.1 _{G4MP2} , 399.0 _{G4}	397 ± 3 _{expt} [1]
[2, 5] 406.6 _{CBS-Q//B3} , 405.4 _{G4MP2}	n/a	398.7 _{CBS-Q//B3} , 397.5 _{G4MP2}	n/a
[2, 6] 407.5 _{CBS-Q//B3} , 406.6 _{G4MP2}	n/a	399.5 _{CBS-Q//B3} , 398.6 _{G4MP2}	n/a
[2, 7] 406.3 _{CBS-Q//B3} , 405.3 _{G4MP2}	n/a	398.3 _{CBS-Q//B3} , 397.3 _{G4MP2}	n/a
[2, 8] 404.7 _{CBS-Q//B3}	n/a	396.6 _{CBS-Q//B3}	n/a

Previous theoretical and experimental values reported in the literature are also provided. Subscripts indicate levels of theory or experimental values. Values are in kcal/mol

^a Theoretical or experimental literature values not available


calculations, the two methods yielded respective $\Delta_{\text{acid}} H_{(g)}^\circ$ and $\Delta_{\text{acid}} G_{(g)}^\circ$ within 0.3 kcal/mol of each other. Thus, assuming G4 level $\Delta_{\text{acid}} H_{(g)}^\circ$ and $\Delta_{\text{acid}} G_{(g)}^\circ$ estimates to be the most accurate among the three methods, it appears that for the $[2, n]$ ($n \geq 5$) compounds where only CBS-Q//B3 and G4MP2 calculations were completed, the CBS-Q//B3 results may be more accurate. Where three or more stacked rings were present (i.e., $m > 2$), geometry optimizations for the deprotonated anions converged on cage-opened non-prismatic geometries. This suggests that $[\geq 3, n]$ polyprismanes may be structurally unstable to gas-phase deprotonation/protonation cycles, potentially precluding reliable experimental measurement or calculation of their acidities.

The $[2, 3]$ polyprismane is more acidic (G4 $\Delta_{\text{acid}} G_{(g)}^\circ = 393.2$ kcal/mol) than its larger cycle $[2, n]$ polyprismane ($n = 4$ –8) counterparts (computational and experimental

$\Delta_{\text{acid}} G_{(g)}^\circ \sim 397$ to 400 kcal/mol). This decreasing theoretical acidity trend between $n = 3$ –4, and lack of a clear trend at $n \geq 4$, is consistent with the corresponding experimental trends in percent s-character of the C–H bonds (36% for $[2, 3]$, 31% for $[2, 4]$, and 30% for $[2, 5]$) as estimated from the $J(^{13}\text{C-H})$ NMR data [18]. Although the $[2, 6]$ and higher $[2, n]$ -polyprismanes are—at present—experimentally unknown, based on the gas-phase acidity calculations herein, we would expect the percent s-character of their C–H bonds to be approximately constant at $\sim 30\%$.

Strain energies of $[m, n]$ polyprismanes and other non-classical forms of carbon are of broad interest for correlating the structures of organic compounds with their stability and reactivity [72, 73]. Minyaev et al. were the first to systematically evaluate the strain in $[m, n]$ polyprismanes using theoretical methods and a series of homodesmotic reactions, and provide a detailed molecular orbital-based

discussion of the values obtained [15]. Generalizing the stoichiometries of the homodesmotic scheme used by these authors yields the following equation for predicting strain energy in any $[m, n]$ polyprismane:



In Table 5, we present the strain energies (both total strain and normalized per C–C bond) for the $[m, n]$ polyprismanes ($m \geq 2$; $n = 3\text{--}8$; $m \times n \leq 16$) calculated at the CBS-Q//B3, G4MP2, and G4 levels, with comparison to previous reports in the literature. For the $[2, n]$ polyprismanes ($n = 2\text{--}6$), where at least several prior estimates (including some high level QCISD(T), CBS, and Gaussian-n calculations for the $[2, 3]$ and $[2, 4]$ members) were available, our values are in excellent agreement and generally lie in the midpoint of the literature range. With all compounds, we observe little difference in predicted strain energies between the CBS-Q//B3, G4MP2, and G4 methods. The previous strain energy estimates for the stacked $[m, n]$ polyprismanes ($m > 2$) at the B3LYP/6-311G(d,p) [15] ($[3, 3]$, $[3, 4]$, $[3, 5]$, $[4, 3]$, $[4, 4]$) and B3LYP/6-31G* [7] ($[3, 4]$, $[5, 3]$) levels of theory range from about 14 to 20 kcal/mol less than our current CBS-Q//B3 and G4MP2 predictions. We see the same structure-strain trends as reported by Lewars [7], namely that total strain energies (E_{str}) inevitably increase with larger $[m, n]$ polyprismane size, and strain energies per one C–C bond ($E_{\text{str}}^{\text{CC}}$) increase with increasing stack height within a given ring size series. Within a given stack height, $E_{\text{str}}^{\text{CC}}$ reaches a minimum with a ring size of five, increasing with either decreasing or increasing ring size away from this minima.

We know generally that while compounds with large strain energies are susceptible to exothermic reactions such as combustion and strain lowering intramolecular rearrangements and disproportionations (i.e., thermodynamic instability), there is no straightforward correlation between strain energy and kinetic stability [7]. Thus, cubane is known to be “extraordinarily stable” in contrast to many substantially less strained compounds [17, 74]. If $E_{\text{str}}^{\text{CC}}$ does correlate with kinetic stability for the $[m, n]$ polyprismanes, we would expect the $[2, 5]$, $[2, 6]$, $[2, 7]$, and $[2, 8]$ members to be significantly more stable than cubane, and some stacked members (e.g., $[3, 5]$) to be about as stable as prismane ($[2, 3]$). With regard to thermodynamic stability, the predicted relative invariance of gas-phase acidities for $m > 4$ with the $[2, m]$ polyprismanes results in no significant correlation with either E_{str} or $E_{\text{str}}^{\text{CC}}$. Thus, equilibrium constants for gas-phase deprotonation of $[m, n]$ polyprismanes do not appear to correlate with molecular strain.

As discussed by Wiberg [73], a priori expectations that increased strain leads to increased reactivity are not always

realized. The location of the activated complex along the reaction coordinate and the strain energies of reactants and products must be taken into consideration. For example, where the structure of the activated complex resembles the highly strained reactants, only a small quantity of strain energy is released in the rate determining step, and hence there is little change in rate constant with increasing reactant strain. In addition, where the products or intermediates are also highly strained, there is little driving force for the reaction (either in kinetic or thermodynamic terms). Upon deprotonation, substantial changes in geometry to the $[m, n]$ polyprismanic carbon cages and carbon-hydrogen bond lengths and angles are expected (Fig. 2 and Table 6). As with the calculations on the undissociated molecular forms, little difference in gas-phase anion geometry is predicted (bond lengths within $\sim \pm 0.001\text{--}0.002$ Å, bond angles within $\sim \pm 0.1\text{--}0.2^\circ$) between the CBS-Q//B3, G4MP2, and G4 levels of theory. The anions have significantly longer carbon-hydrogen bonds compared to the molecular species (~ 1.08 to 1.09 Å \rightarrow 1.10 to 1.11 Å), and this lengthening is approximately uniform around the cage. The hydrogen atoms are also generally angled away asymmetrically from the reaction center, an effect whose magnitude decreases with increasing through-bond distance from the carbanion.

Concomitant with these changes, the internal angles of the carbon framework contract about the reaction center for the $[2, 3]$ through $[2, 6]$ members, increasing the s-character of the carbanion to better accommodate the negative charge. In contrast, all the $[2, 8]$ polyprismane internuclear angles at the carbanion increase upon deprotonation. For the $[2, 7]$ member, the internuclear angle between the carbanion center and the adjacent carbon atoms in the same ring ($\angle \text{C}_2\text{C}_1\text{C}_n$) decreases, but the internuclear angle between the carbon center, either adjacent carbon atom in the same ring, and the carbon directly above the reaction center in the stack ($\angle \text{C}_2\text{C}_1\text{C}_{1'} = \angle \text{C}_n\text{C}_1\text{C}_{1'}$) increases. The CBS-Q//B3 $\angle \text{C}_2\text{C}_1\text{C}_n$ gas-phase angular contractions ($\Delta \angle \text{C}_2\text{C}_1\text{C}_n$) about the reaction center due to deprotonation are as follows: $[2, 3]$, -2.3° ; $[2, 4]$, -3.0° ; $[2, 5]$, -3.3° ; $[2, 6]$, -3.3° ; $[2, 7]$, -1.4° ; and $[2, 8]$, $+2.4^\circ$. We find excellent agreement in the CBS-Q//B3 $\Delta \angle \text{C}_2\text{C}_1\text{C}_4$ of -3.0° for cubane and the SCF/6-31+G* estimate of -3.5° by Ritchie and Bachrach [35], who also noted that unlike other formal sp^3 centers, cubane is anomalous and behaves more like a formal sp^2 center. Using the INDO-LMO method, So et al. [75] reported $\Delta \angle \text{C}_2\text{C}_1\text{C}_n$ of -0.4° $[2, 3]$, -2.9° $[2, 4]$, and -3.0° $[2, 5]$ following gas-phase deprotonation for these three $[2, n]$ polyprismanes, which—with the exception of prismane—are also in excellent agreement with our calculations.

In the molecular forms, all $[2, n]$ polyprismane ($n = 3\text{--}8$) $\angle \text{C}_2\text{C}_1\text{C}_{1'}/\angle \text{C}_n\text{C}_1\text{C}_{1'}$ are 90.0° . Deprotonation yields the

Table 5 Total strain energies (E_{str}) and strain energies per one C–C bond ($E_{\text{str}}^{\text{CC}}$) for the $[m, n]$ polypyrismenes ($m \geq 2$; $m \times n \leq 16$) at the CBS-Q/B3, G4MP2, and G4 levels of theory

E_{str}	$E_{\text{str}}^{\text{CC}}$	
	Current work	Literature
[2, 3]	142.0 _{CBS-Q/B3} , 142.3 _{G4MP2} , 141.8 _{G4}	136 _{B3LYP/6-311G(2df,p)} [15], 148.2 [29] ^a , 145.3 _{HF/6-31G*} [92], 148.7 _{SCF/6-31G*} [78], 145.1 _{G3B3LYP} [81], 161.6 _{G3MP2} [85], 143.8 _{QCISD(T)/6-311+(2df,p)/MP2/6-311(d,p)} [4], 121.7 _{B3LYP/6-311+(3df,2p)/B3LYP/6-311G(d,p)} [4], 146.3 _{MP2/6-311G(d,p)/MP2/6-311G(d,p)} [4], 146.1 _{MP4/6-311G(d,p)/MP4/6-311G(d,p)} [4], 124.0 _{CBS-4} [4], 142.1 _{CBS-Q} [4], 143.4 _{G1} [4], 141.9 _{G2} [4]
[2, 4]	157.0 _{CBS-Q/B3} , 157.7 _{G4MP2} , 156.9 _{G4}	157 ^b [47], 151 _{B3LYP/6-311G(2df,p)} [15], 157.9 [29], 154.7 [73] ^c , 158.6 _{HF/6-31G*} [92], 169.1 _{B3LYP/6-31G*} [93], 165.9 _{MM} [94], 166.9 _{MM} [94], 164.9 _{SCF/6-31G*} [78], 165.4 _{MP2/6-311G(d,p)} [4], 162.8 \pm 2.3 _{G2(MP2), G2, G3/B3LYP, G3[avg]isodesmic} [86], 160.3 \pm 2.2 _{G2(MP2), G2, G3/B3LYP, G3[avg]} [86], 163.8 _{MP2(FULL)/6-31G(d)} [86], 161.3 ^c [87], 153.7 _{B3LYP/6-31G*} [7]
[2, 5]	134.6 _{CBS-Q/B3} , 136.7 _{G4MP2}	129 _{B3LYP/6-311G(2df,p)} [15], 142.9 [29], 140.1 _{SCF/6-31G*} [78]
[2, 6]	171.2 _{CBS-Q/B3} , 173.6 _{G4MP2}	163 _{B3LYP/6-311G(2df,p)} [15], 180.1 [29], 177.7 _{SCF/6-31G*} [78]
[2, 7]	228.7 _{CBS-Q/B3} , 231.3 _{G4MP2}	242.1 [29], 240.7 _{SCF/6-31G*} [78]
[2, 8]	295.9 _{CBS-Q/B3}	315.5 [29], 315.9 _{SCF/6-31G*} [78]
[3, 3]	313.4 _{CBS-Q/B3} , 313.6 _{G4MP2} , 313.6 _{G4}	298 _{B3LYP/6-311G(d,p)} [15]
[3, 4]	387.0 _{CBS-Q/B3} , 387.4 _{G4MP2}	373 _{B3LYP/6-311G(d,p)} [15], 378.8 _{B3LYP/6-31G*} [7]
[3, 5]	390.7 _{CBS-Q/B3} , 392.5 _{G4MP2}	378 _{B3LYP/6-311G(d,p)} [15]
[4, 3]	477.6 _{CBS-Q/B3} , 477.7 _{G4MP2}	457 _{B3LYP/6-311G(d,p)} [15]
[4, 4]	597.5 _{CBS-Q/B3} , 597.4 _{G4MP2}	579 _{B3LYP/6-311G(d,p)} [15]
[5, 3]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 4]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 5]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 6]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 7]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 8]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 9]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 10]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 11]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 12]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 13]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 14]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 15]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 16]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 17]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 18]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 19]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 20]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 21]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 22]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 23]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 24]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 25]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 26]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 27]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 28]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 29]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 30]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 31]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 32]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 33]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 34]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 35]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 36]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 37]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 38]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 39]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 40]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 41]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 42]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 43]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 44]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 45]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 46]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 47]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 48]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 49]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 50]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 51]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 52]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 53]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 54]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 55]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 56]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 57]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 58]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 59]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 60]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 61]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 62]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 63]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 64]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 65]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 66]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 67]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 68]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 69]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 70]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 71]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 72]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 73]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 74]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 75]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 76]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 77]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 78]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 79]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 80]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 81]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 82]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 83]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 84]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 85]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 86]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 87]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 88]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 89]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 90]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 91]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 92]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 93]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 94]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 95]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 96]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 97]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 98]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 99]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a
[5, 100]	639.3 _{CBS-Q/B3} , 639.5 _{G4MP2}	619.5 _{B3LYP/6-31G*} [7] ^a

Previous estimates reported in the literature are also provided. Subscripts indicate levels of theory. Values are in kcal/mol

^a Calculated in [28] using theoretical $\Delta_f H_{(g)}^0$ (estimated at an unclear level of theory, either HF/6-31G* or B3LYP/6-31G*, or some combination thereof; the experimental $\Delta_f H_{(g)}^0$ was used for cubane by these authors) using Benson increments to the $\Delta_f H_{(g)}^0$ corresponding to various structural fragments in aliphatic hydrocarbons


^b Calculated using Franklin's method of group equivalents and the experimental $\Delta_f H_{(g)}^0$

^c Calculated from experimental $\Delta_f H_{(g)}^0$

^d Molecular mechanics with the Schleyer force field and Allinger force field

^e Calculated using the Domalski and Hearing strain-free group contribution values and the experimental $\Delta_f H_{(g)}^0$

Fig. 2 Structures with gas-phase standard state (298.15 K, 1 atm) CBS-Q//B3 optimized bond lengths and angles for the deprotonated $[m, n]$ polyprismane ($m = 2$; $n = 3–8$; $m \times n \leq 16$) anions



following $\Delta\angle C_2C_1C_1'/\Delta\angle C_nC_1C_1'$, which consistently increase with increasing ring size: [2, 3], -4.8° ; [2, 4], -3.0° ; [2, 5], -1.6° ; [2, 6], -0.6° ; [2, 7], $+1.2^\circ$; and [2, 8], $+2.9^\circ$. Consequently, upon gas-phase deprotonation, the CBS-QB//3 calculations (as well as G4MP2 [$n = 3–7$] and G4 [$n = 3, 4$]) show rehybridization at the anionic center to increased s-character for $[2, n]$ polyprismanes for $n = 3–6$, reflected in internuclear angular contraction about the carbanion, consistent with expectations previously discussed in the literature [35, 75, 76]. Further work is needed to better understand the predicted internuclear angle expansions about the reaction center during gas-phase deprotonation of the [2, 7] and [2, 8]polyprismanes, as well as studies of the larger ring members at $n \geq 9$. Overall, we observe no clear decrease in strain upon deprotonation, despite significant geometrical changes, consistent with the predicted insensitivity of gas-phase acidity toward variability in total and normalized strain among the $[m, n]$ polyprismanes.

4 Conclusions

Gas-phase standard state enthalpies of formation and enthalpies and free energies of acid dissociation calculated at the CBS-Q//B3, G4MP2, and G4 levels of theory for $[m, n]$ polyprismanes ($m \geq 2$; $n = 3–8$; $m \times n \leq 16$) yield qualitatively consistent intermethod results with modest intermethod quantitative variability that increases with increasing molecular size. In particular, the lower cost CBS-Q//B3 method yields effectively equivalent gas-phase acidity estimates as the G4MP2 and G4 methods for these compounds, facilitating its use for larger members of the $[m, n]$ polyprismane class. Enthalpies of formation calculated using the atomization method are increasingly overestimated by the CBS-Q//B3 method with larger $[m, n]$ polyprismane sizes, necessitating the use of G4MP2 or G4 calculations for accurate estimates. However, beyond $m \times n = 15$, G4MP2 calculations were prohibitively expensive, and G4 calculations were only feasible up to

Table 6 Estimated gas-phase standard state (298.15 K, 1 atm) carbon–hydrogen bond lengths (r_{C-H}) and bond angles at the CBS-Q/B3 level of theory for the deprotonated anion forms of the $[m, n]$ polypyrmanes ($m = 2; m \times n \leq 16$)

Stack	C ₁ –H		C ₂ –H		C ₃ –H		C ₄ –H		C ₅ –H		C ₆ –H		C ₇ –H		C ₈ –H	
	r_{C_1-H}	Angle	r_{C_2-H}	Angle	r_{C_3-H}	Angle	r_{C_4-H}	Angle	r_{C_5-H}	Angle	r_{C_6-H}	Angle	r_{C_7-H}	Angle	r_{C_8-H}	Angle
[2, 3] 1 Anion	1.093 Å	129.1° C ₁ –C ₂ –H	1.093 Å	129.1° C ₁ –C ₃ –H	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
2 1.093 Å	127.8° C ₂ –C ₁ –H	1.093 Å	129.1° C ₁ –C ₂ –H	1.093 Å	129.1° C ₁ –C ₃ –H	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
[2, 4] 1 Anion	1.103 Å	125.5° C ₁ –C ₂ –H	1.103 Å	126.8° C ₂ –C ₃ –H	1.103 Å	125.5° C ₁ –C ₄ –H	1.103 Å	125.5° C ₁ –C ₄ –H	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
2 1.103 Å	123.7° C ₂ –C ₁ –H	1.103 Å	126.8° C ₁ –C ₂ –H	1.103 Å	125.1° C ₂ –C ₃ –H	1.103 Å	126.8° C ₁ –C ₄ –H	1.103 Å	126.8° C ₁ –C ₄ –H	n/a	n/a	n/a	n/a	n/a	n/a	n/a
[2, 5] 1 Anion	1.103 Å	121.1° C ₁ –C ₂ –H	1.098 Å	120.8° C ₂ –C ₃ –H	1.098 Å	120.8° C ₂ –C ₃ –H	1.098 Å	120.8° C ₂ –C ₃ –H	1.103 Å	121.1° C ₁ –C ₅ –H	n/a	n/a	n/a	n/a	n/a	n/a
2 1.104 Å	118.3° C ₂ –C ₁ –H	1.101 Å	121.1° C ₁ –C ₂ –H	1.095 Å	119.4° C ₂ –C ₃ –H	1.095 Å	119.4° C ₂ –C ₃ –H	1.095 Å	119.4° C ₂ –C ₃ –H	1.101 Å	121.1° C ₁ –C ₅ –H	n/a	n/a	n/a	n/a	n/a
[2, 6] 1 Anion	1.106 Å	117.6° C ₁ –C ₂ –H	1.097 Å	116.3° C ₂ –C ₃ –H	1.095 Å	115.7° C ₃ –C ₄ –H	1.095 Å	115.7° C ₃ –C ₄ –H	1.097 Å	116.3° C ₄ –C ₅ –H	1.106 Å	117.6° C ₁ –C ₆ –H	n/a	n/a	n/a	n/a
2 1.109 Å	114.3° C ₂ –C ₁ –H	1.102 Å	116.6° C ₁ –C ₂ –H	1.095 Å	115.5° C ₂ –C ₃ –H	1.096 Å	115.4° C ₃ –C ₄ –H	1.095 Å	115.5° C ₄ –C ₅ –H	1.102 Å	116.6° C ₁ –C ₆ –H	n/a	n/a	n/a	n/a	n/a
[2, 7] 1 Anion	1.109 Å	107.7° C ₃ –C ₂ –H	1.099 Å	112.8° C ₂ –C ₃ –H	1.096 Å	112.7° C ₃ –C ₄ –H	1.096 Å	112.7° C ₃ –C ₄ –H	1.096 Å	112.7° C ₄ –C ₅ –H	1.099 Å	112.8° C ₅ –C ₆ –H	1.109 Å	116.4° C ₁ –C ₇ –H	n/a	n/a
2 1.113 Å	111.1° C ₂ –C ₁ –H	1.102 Å	113.5° C ₁ –C ₂ –H	1.095 Å	112.7° C ₂ –C ₃ –H	1.096 Å	112.4° C ₃ –C ₄ –H	1.096 Å	112.4° C ₃ –C ₄ –H	1.096 Å	112.7° C ₅ –C ₆ –H	1.102 Å	113.5° C ₁ –C ₇ –H	n/a	n/a	n/a
[2, 8] 1 Anion	1.111 Å	105.3° C ₃ –C ₂ –H	1.098 Å	110.0° C ₂ –C ₃ –H	1.096 Å	110.4° C ₃ –C ₄ –H	1.096 Å	110.4° C ₃ –C ₄ –H	1.096 Å	109.4° C ₄ –C ₅ –H	1.096 Å	110.4° C ₅ –C ₆ –H	1.098 Å	110.0° C ₆ –C ₇ –H	1.111 Å	115.8° C ₁ –C ₈ –H
2 1.119 Å	108.2° C ₂ –C ₁ –H	1.102 Å	111.2° C ₁ –C ₂ –H	1.095 Å	110.2° C ₂ –C ₃ –H	1.096 Å	110.2° C ₂ –C ₃ –H	1.096 Å	110.2° C ₂ –C ₃ –H	1.096 Å	110.2° C ₃ –C ₄ –H	1.096 Å	110.2° C ₃ –C ₄ –H	1.095 Å	110.2° C ₆ –C ₇ –H	1.102 Å
[2, 9] 1 Anion	1.120 Å	108.2° C ₈ –C ₇ –H	1.092 Å	109.2° C ₇ –C ₈ –H	1.092 Å	109.2° C ₇ –C ₈ –H	1.092 Å	109.2° C ₇ –C ₈ –H	1.092 Å	109.2° C ₇ –C ₈ –H	1.092 Å	109.2° C ₇ –C ₈ –H	1.092 Å	109.2° C ₇ –C ₈ –H	1.092 Å	109.2° C ₇ –C ₈ –H

The numbering system for carbon atoms proceeds starting at 1 (carbon) and increasing sequentially clockwise around each ring with “*” denotation for adjacent carbon atoms in the second stack

$m \times n = 9$. Various isodesmic and homodesmic reaction schemes at these levels of theory do not offer any clear prediction advantage over the atomization method. Good agreement with previous DFT formation enthalpy estimates in the literature obtained using homodesmic approaches suggests the thermochemistry of larger $[m, n]$ polyprismanes may be studied using these methods and lower cost levels of theory. Group additivity methods yield, at present, inaccurate enthalpies of formation for $[m, n]$ polyprismanes due to the absence of appropriate strain correction factors.

Geometry optimizations for the deprotonated anions converged on cage-opened non-prismatic geometries where three or more stacked $[m, n]$ polyprismane rings were present. These compounds may be structurally unstable to gas-phase deprotonation/protonation cycles, potentially preventing reliable experimental measurement or calculation of their acidities. Good correlations were found between the gas-phase acidities of the [2,3], [2,4], and [2,5]polyprismanes and percent s-character of the C–H bonds. Calculations indicate no substantial change in the gas-phase acidity for the $[2, n]$ polyprismanes at $n \geq 6$, suggesting no corresponding change in C–H bond s-character over this range. Larger $[m, n]$ polyprismanes have higher total strain energies. Increasing stack height within a given ring size series results in larger strain energies per one C–C bond, which also reach a minimum with a ring size of five within a given stack height. Gas-phase deprotonation results in substantial geometry changes between the molecular and deprotonated anionic forms. Contraction of the internuclear angles about the reaction center occurs for the [2, 3]–[2, 6] members as a result of rehybridization to increase s-character in the carbanion and better accommodate the negative charge. Neither total energy nor per carbon–carbon bond strain energies correlate with gas-phase acidity, likely because deprotonation does not relieve significant molecular strain.

Acknowledgments This work was made possible by the facilities of the Western Canada Research Grid (WestGrid: <http://www.westgrid.ca>; project 100185), the Shared Hierarchical Academic Research Computing Network (SHARCNET: <http://www.sharcnet.ca>; project sn4612), and Compute/Calcul Canada. We thank three anonymous reviewers for their insights which improved the quality of the manuscript.

References

- Hare M, Emrick T, Eaton PE, Kass SR (1997) Cubyl anion formation and an experimental determination of the acidity and C–H bond dissociation of cubane. *J Am Chem Soc* 119:237–238
- Alkorta I, Elguero J (1997) Carbon acidity and ring strain: a hybrid HF-DFT approach (Becke3LYP/6–311++G**). *Tetrahedron* 53:9741–9748
- Streitwieser A, Xie L, Speers P, Williams PG (1998) Kinetic acidities of some hydrocarbons by tritium NMR. *Magn Reson Chem* 36:S209–S211
- Walker JE, Adamson PA, Davis SR (1999) Comparison of calculated hydrocarbon strain energies using ab initio and composite methods. *J Mol Struct Theochem* 487:145–150
- Earley CW (2000) Ab initio investigation of strain in group 14 polyhedrane clusters (M_nH_n ; $n = 4, 6, 8, 10, 12, 16, 20, 24$). *J Phys Chem A* 104:6622–6627
- Moran D, Manoharan M, Heine T, Schleyer PVR (2003) σ -Antiaromaticity in cyclobutane, cubane, and other molecules with saturated four-membered rings. *Org Lett* 5:23–26
- Lewars EG (2008) Modeling marvels. Springer, New York
- Minyaev RM, Minkin VI (2008) Nonclassical carbon: from theory to experiment. *Russ J Gen Chem* 78:732–749
- Jefford CW (1976) Syntheses and rearrangements of cage molecules related to cubane. *J Chem Ed* 53:477–482
- Smith LR (1978) Schemes and transformations in the $(CH)_8$ series. *J Chem Ed* 55:569–570
- Balaban AT, Banciu M (1984) Schemes and transformations in the $(CH)_{2k}$ series. *J Chem Ed* 61:766–770
- Schrivier GW, Gerson DJ (1990) Energies and isomerizations of $(CH)_{12}$ hydrocarbons. *J Am Chem Soc* 112:4723–4728
- Engelke R (1992) Calculated properties of the 22 carbon/nitrogen cubanoids. *J Org Chem* 57:4841–4846
- Engelke R (1993) Ab initio calculations of ten carbon/nitrogen cubanoids. *J Am Chem Soc* 115:2961–2967
- Minyaev RM, Minkin VI, Gribanova TN, Starikov AG (2003) Poly[n]prismanes: a family of stable cage structures with half-planar carbon centers. *J Org Chem* 68:8588–8594
- Schmidt MW, Gordon MS, Boatz JA (2000) Cubic fuels? *Int J Quant Chem* 76:434–446
- Eaton PE, Cole TW (1964) Cubane. *J Am Chem Soc* 86:3157–3158
- Eaton PE, Or YS, Branca SJ (1981) Pentaprismene. *J Am Chem Soc* 103:2134–2136
- Katz TJ, Acton N (1973) Synthesis of prismane. *J Am Chem Soc* 95:2738–2739
- Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Scalmani G, Barone V, Mennucci B, Petersson GA, Nakatsuji H, Caricato M, Li X, Hratchian HP, Izmaylov AF, Bloino J, Zheng G, Sonnenberg JL, Hada M, Ehara M, Toyota K, Fukuda R, Hasegawa J, Ishida M, Nakajima T, Honda Y, Kitao O, Nakai H, Vreven T, Montgomery JA, Peralta JE, Ogliaro F, Bearpark M, Heyd JJ, Brothers E, Kudin KN, Staroverov VN, Kobayashi R, Normand J, Raghavachari K, Rendell A, Burant JC, Iyengar SS, Tomasi J (2009) Gaussian 09. Gaussian, Inc., Wallingford
- Montgomery JA, Frisch MJ, Ochterski JW, Petersson GA (1999) A complete basis set model chemistry. VI. Use of density functional geometries and frequencies. *J Chem Phys* 110:2822–2827
- Montgomery JA, Frisch MJ, Ochterski JW, Petersson GA (2000) A complete basis set model chemistry. VII. Use of the minimum population localization method. *J Chem Phys* 112:6532–6542
- Curtiss LA, Redfern PC, Raghavachari K (2007) Gaussian-4 theory using reduced order perturbation theory. *J Chem Phys* 127:124105–124108
- Curtiss LA, Redfern PC, Raghavachari K (2007) Gaussian-4 theory. *J Chem Phys* 126:84108–84112
- Stewart J (2007) Optimization of parameters for semiempirical methods V: Modification of NDDO approximations and application to 70 elements. *J Mol Model* 13:1173–1213
- Miller MA, Schulman JM (1988) AM1, MNDO and MM2 studies of concatenated cyclobutanes: Prismanes, ladderanes, and asteranes. *J Mol Struct Theochem* 163:133–141

27. Ball DW (1996) On the ΔH_f values of tetrahedrane and cubane: Density functional theory calculations. *J Mol Struct Theochem* 364:183–188
28. Li Z, Rogers DW, McLafferty FJ, Mandziuk M, Podosenin AV (1999) Ab initio calculations of enthalpies of hydrogenation, isomerization, and formation of cyclic C_6 hydrocarbons. Benzene isomers. *J Phys Chem A* 103:426–430
29. Karpushenkava LS, Kabo GJ, Bazyleva AB (2009) Structure, frequencies of normal vibrations, thermodynamic properties, and strain energies of the cage hydrocarbons C_nH_n in the ideal-gas state. *J Mol Struct Theochem* 913:43–49
30. Fleischer EB (1964) X-ray structure determination of cubane. *J Am Chem Soc* 86:3889–3890
31. Jenkins SJ, King DA (2000) Pentaprismane and hypostrophene from first-principles, with plane waves. *Chem Phys Lett* 317:381–387
32. Politzer P, Seminario JM (1989) Computational determination of the structures and some properties of tetrahedrane, prismane, and some of their aza analogues. *J Phys Chem* 93:588–592
33. Seidl ET, Schaefer HF (1991) Face-fused dicubane ($C_{12}H_8$) and its dicubane isomer: achievable synthetic targets? *J Am Chem Soc* 113:1915–1917
34. Alkorta I, Elguero J, Rozas I, Balaban AT (1990) Theoretical studies of aza analogues of platonic hydrocarbons: Part 1. Cubane and its aza derivatives. *J Mol Struct Theochem* 206:67–75
35. Ritchie JP, Bachrach SM (1990) Comparison of the calculated acidity of cubane with that of other strained and unstrained hydrocarbons. *J Am Chem Soc* 112:6514–6517
36. Allinger NL, Eaton PE (1983) The geometries of pentaprismane and hexaprismane: Insights from molecular mechanics. *Tet Lett* 24:3697–3700
37. Dailey WP (1987) The structures and energies of pentaprismane and hexaprismane—An ab initio study. *Tet Lett* 28:5787–5790
38. Honegger E, Eaton PE, Ravi Shankar BK, Heilbronner E (1982) The electronic structure of pentaprismane ($C_{10}H_{10}$), as revealed by its photoelectron spectrum. *Helv Chim Acta* 65:1982–1989
39. Engelke R, Hay PJ, Kleier DA, Wadt WR (1984) Theoretical study of dimeric forms of ground-state benzene molecules. *J Am Chem Soc* 106:5439–5446
40. Karl RR, Wang YC, Bauer SH (1975) The gas phase structures of the isomers of benzene. III. Hexamethylprismane and hexamethylbenzene. *J Mol Struct* 25:17–34
41. Xu WG, Liu XF, Lu SX (2008) Theoretical investigation on the heats of formation and the interactions among the azido groups in polyazidoprismanes $C_6H_{6-n}(N_3)_n$ ($n = 1-6$). *J Mol Struct Theochem* 864:80–84
42. Wingert H, Maas G, Regitz M (1986) Syntheses with cyclobutadienes. 15. Prismane/dewar benzene isomerization—X-ray crystal structure of tert-butyl-3, 4, 5-tri-tert-butyl-2, 6-bis(trifluoromethyl)prismane-1-carboxylate. *Tetrahedron* 42:5341–5353
43. Grimme S (1992) Theoretical study of [4]paracyclophane and its Dewar benzene and prismane valence isomers. *J Am Chem Soc* 114:10542–10547
44. Herrera B, Valencia F, Romero AH, Kiwi M, Ramirez R, Toro-Labbe A (2006) Cubane oligomers: a density functional theory study. *J Mol Struct Theochem* 769:183–187
45. Saeys M, Reyniers MF, Marin GB, Van Speybroeck V, Waroquier M (2003) Ab initio calculations for hydrocarbons: enthalpy of formation, transition state geometry, and activation energy for radical reactions. *J Phys Chem A* 107:9147–9159
46. Van Speybroeck V, Gani R, Meier RJ (2010) The calculation of thermodynamic properties of molecules. *Chem Soc Rev* 39:1764–1779
47. Kybett BD, Carroll S, Natalis P, Bonnell DW, Margrave JL, Franklin JL (1966) Thermodynamic properties of cubane. *J Am Chem Soc* 88:626
48. Diky VV, Frenkel M, Karpushenkava LS (2003) Thermodynamics of sublimation of cubane: Natural anomaly or experimental error? *Thermochim Acta* 408:115–121
49. Bashir-Hashemi A, Chickos JS, Hanshaw W, Zhao H, Farivar BS, Liebman JF (2004) The enthalpy of sublimation of cubane. *Thermochim Acta* 424:91–97
50. Afeefy HY, Liebman JF, Stein SE (2010) Neutral thermochemical data. In: Linstrom PJ, Mallard WG (eds) NIST Chemistry WebBook, NIST standard reference database number 69. National Institute of Standards and Technology, Gaithersburg
51. Lias SG, Bartmess JE, Liebman JF, Holmes JL, Levin RD, Mallard WG (2010) Ion energetics data. In: Linstrom PJ, Mallard WG (eds) NIST Chemistry WebBook, NIST standard reference database number 69. National Institute of Standards and Technology, Gaithersburg
52. Rayne S, Forest K (2010) Gas phase isomerization enthalpies of organic compounds: A semiempirical, density functional theory, and ab initio post-Hartree–Fock theoretical study. *J Mol Struct Theochem* 948:102–107
53. Rayne S, Forest K (2010) Computational note on a G4MP2 study into the gas phase enthalpies of formation and isomerization for the $(CH)_{2n}$ ($n = 1-6$) isomers. *J Mol Struct Theochem* 948:111–112
54. Rayne S, Forest K (2010) Performance of Gaussian-3 and Gaussian-4 level theoretical methods in estimating gas phase enthalpies of formation for representative C_1 and C_2 chlorofluorocarbons and hydrochlorofluorocarbons. *J Mol Struct Theochem* 953:47–48
55. Wang H, Law CK (1997) Thermochemistry of benzvalene, dihydrobenzvalene, and cubane: a high-level computational study. *J Phys Chem B* 101:3400–3403
56. Cox JD, Wagman DD, Medvedev VA (1984) CODATA key values for thermodynamics. Hemisphere, New York
57. Roth WA, Banse H (1932) Die verbrennungs- und bildungswärme von kohlenoxyd und methan. *Arch Eisenhütten* 6:43–46
58. Cox JD, Pilcher G (1970) Thermochemistry of organic and organometallic compounds. Academic Press, New York
59. Chase MW (1998) NIST-JANAF thermochemical tables (4th ed). *J Phys Chem Ref Data* 9:1–1951
60. Pittam DA, Pilcher G (1972) Measurements of heats of combustion by flame calorimetry. Part 8. Methane, ethane, propane, n-butane and 2-methylpropane. *J Chem Soc Faraday Trans 1* 68:2224–2229
61. Prosen EJ, Rossini FD (1945) Heats of combustion and formation of the paraffin hydrocarbons at 25 C. *J Res NBS* 36:263–267
62. Prosen EJ, Johnson WH, Rossini FD (1946) Heats of formation and combustion of the normal alkylcyclopentanes and cyclohexanes and the increment per CH_2 group for several homologous series of hydrocarbons. *J Res NBS* 37:51–56
63. Moore GE, Renquist ML, Parks GS (1940) Thermal data on organic compounds. XX. Modern combustion data for two methylnonanes, methyl ethyl ketone, thiophene and six cycloparaffins. *J Am Chem Soc* 62:1505–1507
64. Butler RS, Carson AS, Laye PG, Steele WV (1971) The enthalpy of formation of adamantane. *J Chem Thermodyn* 3:277–280
65. Baroody EE, Carpenter GA (1972) Heats of formation of propellant compounds (Command task 331–003/067–1/UR2402–001). Naval Ordnance Station, Indian Head
66. Clark T, Knox TMO, McKervey MA, Mackle H, Rooney JJ (1979) Thermochemistry of bridged-ring substances. Enthalpies of formation of some diamondoid hydrocarbons and of perhydroquinacene. Comparisons with data from empirical force field calculations. *J Am Chem Soc* 101:2404–2410
67. Benson SW, Buss JH (1958) Additivity rules for the estimation of molecular properties. Thermodynamic properties. *J Chem Phys* 29:546–572

68. Benson SW (1976) Thermochemical kinetics: methods for the estimation of thermochemical data, 2nd edn. Wiley, New York
69. Cohen N, Benson SW (1993) Estimation of heats of formation of organic compounds by additivity methods. *Chem Rev* 93: 2419–2438
70. Sabbe MK, Saeys M, Reyniers MF, Marin GB, Van Speybroeck V, Waroquier M (2005) Group additive values for the gas phase standard enthalpy of formation of hydrocarbons and hydrocarbon radicals. *J Phys Chem A* 109:7466–7480
71. Hassenruck K, Martin HD, Walsh R (1989) Consequences of strain in $(\text{CH})_8$ hydrocarbons. *Chem Rev* 89:1125–1146
72. Greenberg A, Liebman JF (1978) Strained organic molecules: organic chemistry series, vol 38. Academic Press, New York
73. Wiberg K (1986) The concept of strain in organic chemistry. *Angew Chem Int Ed Engl* 25:312–322
74. Griffin GW, Marchand AP (1989) Synthesis and chemistry of cubanes. *Chem Rev* 89:997–1010
75. So SP, Wong MH, Luh TY (1985) Hybridization of the lone pair electrons in carbanions. *J Org Chem* 50:2632–2635
76. Luh TY, Stock LM (1974) Kinetic acidity of cubane. *J Am Chem Soc* 96:3712–3713
77. Rogers DW, McLafferty FJ, Fang W, Qi Y (1993) Molecular enthalpies 3: ground-state thermodynamic and computed enthalpies of the valence isomers of benzene. *Struct Chem* 4:161–166
78. Disch RL, Schulman JM (1988) Ab initio heats of formation of medium-sized hydrocarbons. 7. The $[n]$ prismanes. *J Am Chem Soc* 110:2102–2105
79. Cheung YS, Wong CK, Li WK (1998) Ab initio calculations of the heats of formation for $(\text{CH})_6$ isomers. *J Mol Struct Theochem* 454:17–24
80. Cheung TS, Law CK, Li WK (2001) Gaussian-3 heats of formation for $(\text{CH})_6$ isomers. *J Mol Struct Theochem* 572:243–247
81. Novak I (2003) Substituent effects on steric strain. *Chem Phys Lett* 380:258–262
82. Novak I, Pratt LM (2004) Computational study of lithioprismanes. *Chem Phys Lett* 400:558–562
83. Schulman JM, Disch RL (1984) Ab initio heats of formation of medium-sized hydrocarbons. The heat of formation of dodecahedrane. *J Am Chem Soc* 106:1202–1204
84. Reddy VP, Jemmis ED (1986) A D_{4d} structure for [8]-prismane. *Tet Lett* 27:3771–3774
85. Rogers DW (2000) G3(MP2) calculation of the enthalpies of formation, isomerization and hydrogenation of cubane and cyclooctatetraene. *J Mol Struct* 556:207–215
86. Roux MV, Davalos JZ, Jimenez P, Notario R, Castano O, Chirkos JS, Hanshaw W, Zhao H, Rath N, Liebman JF, Farivar BS, Bashir-Hashemi A (2005) Cubane, cuneane, and their carboxylates: a calorimetric, crystallographic, calculational, and conceptual coinvestigation. *J Org Chem* 70:5461–5470
87. Kirklin DR, Churney KL, Domalski ES (1989) Enthalpy of combustion of 1, 4-dimethylcubane carboxylate. *J Chem Thermodyn* 21:1105–1113
88. Manion JA (2002) Evaluated enthalpies of formation of the stable closed shell C_1 and C_2 chlorinated hydrocarbons. *J Phys Chem Ref Data* 31:123–172
89. Gurvich LV, Veyts IV, Alcock CB (1992) Thermodynamic properties of individual substances, vol 2, 4th edn. Hemisphere, New York
90. Spitzer R, Huffman HM (1947) The heats of combustion of cyclopentane, cyclohexane, cycloheptane and cyclooctane. *J Am Chem Soc* 69:211–213
91. Pedley JB, Naylor RD, Kirby SP (1986) Thermochemical data of organic compounds. Chapman and Hall, New York
92. Nagase S (1989) Much less strained cubane analogues with Si, Ge, Sn, and Pb skeletons. *Angew Chem Int Ed Engl* 28:329–330
93. Fan XW, Ju XH, Xia QY, Xiao HM (2008) Strain energies of cubane derivatives with different substituent groups. *J Haz Mater* 151:255–260
94. Engler EM, Andose JD, Schleyer PVR (1973) Critical evaluation of molecular mechanics. *J Am Chem Soc* 95:8005–8025