

In the treatment of
relapsed ovarian cancer...

Hycamtin
A first choice
for a
second chance

Hycamtin is indicated for your patients with metastatic carcinoma of the ovary after failure of initial or subsequent chemotherapy.

HYCAMTIN
(topotecan HCl) FOR INJECTION

Hycamtin is contraindicated in patients with a history of hypersensitivity to topotecan or to any of its ingredients. *Hycamtin* should not be used in patients who are pregnant or breast-feeding or in those with severe bone marrow depression. The most common serious adverse event is myelosuppression.

Please see adjacent page for brief summary of prescribing information, including warnings, precautions and adverse events.

gsk GlaxoSmithKline

Possibilities

10.23


96
1

GYNECOLOGIC ONCOLOGY

VOLUME 96, NUMBER 1, JANUARY 2005

PAGES 1-268

ELSEVIER



Volume 96, Number 1, January 2005

ISSN 0090-8258

Gynecologic Oncology

EDITOR-IN-CHIEF
David M. Gershenson

EDITORS
Larry J. Copeland
William J. Hoskins
Karl C. Podratz

The Official Publication of



Also available on
SCIENCE @ DIRECT®
www.sciencedirect.com

•CYAN •MAGENTA •YELLOW •BLACK •PMS 3405CVC •PMS 534CVC •PMS BLACKCVC