

Overlapped Metabolic and Therapeutic Links between Alzheimer and Diabetes

Waqar Ahmad

Received: 23 August 2012 / Accepted: 12 September 2012 / Published online: 26 September 2012
© Springer Science+Business Media New York 2012

Abstract Alzheimer's disease (AD) and diabetes are among the most common diseases associated with ageing. The pathology of AD is strongly associated with accumulated misfolding proteins that results in neuronal dysfunction within the brain. Diabetes, on the contrary, is characterised by altered insulin signaling that results in reduced glucose uptake, metabolic suppression of energy consuming cells and conversion of glucose to fat in the liver. Despite distinguishing features, these diseases share common elements and may in fact be viewed as fundamentally similar disorders that differ in magnitude of specific traits, primarily affected tissues and time of onset. In this review, we outline the fundamental basis of each of the two diseases and highlight similarities in their pathophysiology. Further ahead we will discuss these features in relation to the development of drugs to treat these two diseases, particularly AD, for which the development of therapeutic chemicals has proven to be particularly difficult. We conclude with comments on efforts to develop a simple organism, *Caenorhabditis elegans*, as a genetic model to be used to study the systems biology of diabetes and AD.

Keywords Alzheimer's disease · Diabetes · Pathophysiology · Impaired metabolism · Therapies · *C. elegans*

Introduction

Alzheimer's disease (AD) and diabetes are both age-associated diseases. AD is the most common form of dementia in the elderly and is clinically categorized by a progressive loss of cognitive and memory functions. Post-mortem examination of brains of AD sufferers reveals accumulation of extracellular plaques of amyloid beta ($A\beta$) peptides and intracellular neurofibrillary tangles (NFTs) consisting of hyperphosphorylated tau protein. In addition, there is a decrease in synapse density and a corresponding decrease in brain volume due to neurodegeneration [1–3]. On the other hand, diabetes mellitus is associated with decreased body insulin production than required (type 1) and impaired insulin signaling (type 2/T2DM). Another hallmark of diabetes is the formation of Human islet amyloid polypeptide (hIAPP, amylin) that leads to pancreatic β -cells dysfunction. The resulting metabolic disturbance leads to chronic hyperglycemia, which is the immediate cause of many of the symptoms of diabetes [4–6].

Even though the pathologies of these two diseases appear superficially unrelated, diabetes sufferers have an increased risk of development of age-associated neurodegenerative diseases, including AD. In 2005, Suzanne de la Monte [7] introduced the term "type 3 diabetes" in reference to AD. Her team observed insulin impairment associated with AD patient during post-mortem. They observed an inverse correlation between insulin receptor abundance and the Braak score of AD brains, with 80 % reduced IRS levels in the most extreme cases. They observed reduced mRNA levels of IGF-1, polypeptides and increased tau protein levels regulated by these receptors [7, 8]. Further studies on model organisms confirmed the impaired insulin signaling and oxidative insult of rat/mice brains leading to reduced brain size and neurodegeneration when treated with streptozotocin (STZ), a chemical which causes diminution of insulin, and induces an insulin-

W. Ahmad (✉)
Paul R. Ebert Laboratory, School of Biological Sciences,
University of Queensland,
Brisbane 4072, Australia
e-mail: waqar.ahmad@uqconnect.edu.au

W. Ahmad
e-mail: waqarchemist@hotmail.com

resistant brain state [9, 10]. Moreover, the success of diabetic drugs to reduce AD complexities also opens new doors for researchers to understand their mechanisms in depth. In the following discussion, we review the common pathways linking AD and diabetes, and recent advancements in these fields.

Common Features in Alzheimer's Disease and Diabetes

As described earlier, formation of A β and NFTs in AD, while hIAPP and insulin resistance in diabetes are characteristic hallmarks for these diseases. Both AD and diabetes are degenerative diseases involving neuronal loss and β -cells destruction, respectively [11–14]. The main systematic link between these diseases is impaired insulin signaling leading to neurodegeneration and cognitive damages [15]. Ott and colleagues in 1999 [16] first observed the association of diabetes with dementia and further studies found 2-fold higher risk of AD in diabetic patients. Diabetes was also found to accelerate the onset of AD rather than only a risk factor in prospective and cross-sectional study [17]. Moreover, in diabetic patients with ApoE4 allele (that is genetically associated with AD), chances of AD were 2-fold higher than in non-diabetic patients with ApoE4 allele. ApoE4 is associated with APP trafficking, inhibition of A β clearance and lipid (mainly cholesterol) transportation. Insulin regulates cholesterol biosynthesis and high cholesterol levels were found not only to be a risk factor for diabetes but also for AD [18, 19]. Recent studies also elaborate the role of impaired insulin signaling in tau hyperphosphorylation, revealing another association between AD and diabetes [20].

AD and diabetes not only shares common pathways but also several enzymes (glutamic acid decarboxylase, dopa-decarboxylase), growth factor receptors (p75 receptors, neuronal growth factor receptors and thyrotrophin-releasing hormone) and second messenger abnormalities such as dysregulated protein phosphorylation and glycogen synthase kinase 3 (GSK3) overactivity [15, 21–23]. Furthermore, mitochondrial dysfunctions resulting in increased oxidative stress and induced glyceraldehyde-derived advanced glycation end products (AGEs) are also associated with AD and diabetes [10, 24]. This review contains a brief outline about these common features and recent advancements in these fields.

Epidemiological/Environmental Risk Factors

Aging is associated with an increased risk of AD and diabetes. While aging, the decrease in mitochondrial function and oxidative capacity is observed and such is the case in AD and diabetic patients [25, 26]. In the USA, approximately

one in eight persons aged 65 and older (13 %), and more than 45 % of people >85 years have AD. Meanwhile 42 % of the population over 65 are afflicted with diabetes [27–29]. The aging process is frequently associated with environmental factors especially the diet of the person. Dietary restriction or caloric restriction (CR) is linked to increased lifespan and numerous model systems showing longer life spans in CR studies. Moreover, CR has also shown to delay several age associated diseases such as neurodegeneration, diabetes, cancer and heart disease [30–33]. The reverse of CR is obesity, which is mainly associated with excess body fat (adiposity) and excessive energy intake.

Obesity is the most highlighted risk factor linked with diabetes and AD. Obesity usually refers to higher amounts of fat in the body than normal, although the normal threshold of adiposity is not clear and is associated with body height. Whereas, increased risk of insulin resistance, diabetes, dyslipidemia, hypertension, heart diseases cancer, and respiratory disease are associated with adiposity [34, 35]. Body mass index and waist circumference are two methods to measure adiposity. Several studies have reported higher risk of diabetes and AD with obesity and individuals with midlife body mass index >30 have greater chances of AD [36, 37]. Increased cognitive impairment has also been reported in patients as well as in experimental models with obesity and/or dyslipidemia [38]. Furthermore, elevated waist circumference than cut off value (cut-off value is 102 cm for men and 88 cm for women) is also associated with diabetes as well AD [39].

Several epidemiological studies performed in different countries have shown the positive correlation of diabetes with AD, where diabetic patients exhibit an increased risk of AD development when compared to non-diabetic individuals. Recent population based studies showed 2–5 times higher incidence of AD in patients suffering from diabetes and a lot of literature is available on AD and diabetic association [40–43]. Furthermore, recent studies reported AD as a form of diabetes and termed as "type 3 diabetes" due to overlapping of molecular and biochemical features that are discussed in detail later on in this review. [44, 45].

Misfolding of Processed Amyloid Peptides to Form Plaques

One of the characteristic features of AD is amyloidogenesis (changing of soluble protein into insoluble fibrillar protein aggregates to form A β plaques) in various brain regions that also pathologically represent AD [46]. Extensive study of A β formation and its after affects has led to the formulation of A β cascade hypothesis, which places A β as the primary cause of AD pathogenesis [47]. This process begins with the proteolytic processing of the precursor protein into peptides

predominantly of 40 or 42 amino acids ($A\beta_{1-40}$ and $A\beta_{1-42}$) [48]. These $A\beta$ oligomers can directly inhibit the hippocampal long-term potentiating (LTP) component of memory by impairment of synaptic plasticity [49]. Ultimately the peptides aggregate to form extracellular protein plaques (also called senile plaques) that are cytotoxic resulting in neurological dysfunction [48]. $A\beta$ may undergo conditional conformational changes (from native random coil structure to alpha-helical or beta-pleated strands) under slight alterations in pH, peptide concentration, metal ion concentration and environmental composition. These changes may result in diverse $A\beta$ aggregation; however as to which confirmation is neurotoxicity directly linked to is still unknown. The neurotoxicity of $A\beta$ is discussed later in detail with relation to other mechanisms [50, 51].

A related protein, hIAPP (human amylin), is produced in β -islet cells of the pancreas [52]. It is a 37 residues protein synergistically acting with insulin to control glycemia [53] and regulates food intake by slowing gastric emptying, signaling the brain to decrease meal size thus reducing blood glucose levels [54]. Additionally, it is also involved in the regulation of calcium homeostasis, vasodilation and renal filtration [22]. Damage of β -cells is early onset dysfunction in diabetes with deposition of misfolded hIAPP as amyloid fibrils. As hIAPP is co-secreted with insulin, insulin resistance leads to overproduction of hIAPP that accumulates as amyloid fibrils on β -cells [53, 55]. Reduced β -cells masses with the presence of β -sheet amyloid fibres of hIAPP are the characteristic of T2DM. In isolated islets these amyloids were associated with increased apoptosis with decreased area and viability of β -cells [56].

Thus, $A\beta$ and hIAPP both can aggregate to form extracellular plaques, but this occurs in two distinct organs, brain and pancreatic β -cells, respectively [52]. The distinct tissue plaque formation is associated with the unique pathology of each disease, but whether this is a causal relationship is still unknown [50]. Despite the dissimilarity in amino acid homology, these amyloids share structural similarities. More than 90 %, and are believed to share common mode of toxicity [57]. A recent study by Fu et al. [58] found that $A\beta$ and hIAPP both were involved in triggering multiple cellular pathways such as MAPK, Akt, cFos and signal transduction mediator protein kinase A by increasing cytosolic cAMP and Ca^{+2} levels in HEK293 cell line through amylin receptor-3 (AMY3). This change was blocked using AMY3 antagonist AC253. Meanwhile, in a study by Gazit group small aromatic molecules naphthoquinones termed as NQT_{rp}^{23} and $Cl-NQT_{rp}^{31}$ were observed as potential inhibitors of $A\beta$, in vitro as well as in fly and murine model systems of AD. Furthermore, they found that these molecules were also capable to disaggregate amyloid formed by hIAPP with unknown mechanisms [59]. These results not only strengthen the AD–diabetes association but also point

out new therapeutic target for AD and diabetes. Moreover, human Ca^{+2} binding protein nucleobindin 1 (NUCB1) was found be useful in inhibiting hIAPP fibril formation as well as disaggregation of pre-existing hIAPP fibrils through unknown mechanism [60]. We purpose that NUCB1 should be checked as potential inhibitor for $A\beta$ plaques formation in AD patients.

Impaired Insulin and Glucose Metabolism in AD and Diabetes

Insulin is a hormone that has an important role in glucose homeostasis in a dynamic relationship with both feeding and fasting as well as in growth and development of body tissues [61, 62]. Insulin is mainly synthesized by pancreatic β -cells and secreted to peripheral circulatory system. Although insulin is transported to brain through the blood–brain barrier (BBB), evidence indicates that the brain can synthesize insulin locally. In the central nervous system, insulin also normalizes and maintains cognitive functions by regulating key processes such as neuronal survival and longevity, learning, and memory. Defective insulin signaling leads to energy deficient neurons resulting in various metabolic insults and impaired synaptic plasticity which is discussed later [63, 64].

Insulin Resistance in Peripheral System and Central Nervous System

Insulin resistance may be defined as a diminished ability of cells or tissues to respond to physiological levels of insulin. Insulin resistance can be associated with defects in insulin receptor function, insulin signal transduction, glucose metabolism, transport, glycogen synthesis, hyperinsulinemia, hyperglycemia, inflammation and lipids metabolism [65]. Increased lipid content induces insulin secretion referred to as hyperinsulinemia for maintaining normoglycemia. Any defects in insulin secretion (low insulin levels, type 1 diabetes) and signaling (insulin resistance at cellular levels, type 2 diabetes) may cause hyperglycemia or high blood sugar levels. In this situation, diabetes is a disease originated from defects in the body's ability to control insulin and glucose homeostasis due to hyperglycemia, insulin resistance and β -cell failure [66, 67]. Insulin resistance also has a pivotal role in the progression of neurodegenerative diseases especially in AD. Many studies have found neuroprotective effects of insulin actions in the brain [9, 68, 69]. Insulin deficiency, resistance and hyperinsulinemia are involved in cognitive impairments observed in patients with diabetes [70]. AD is associated with reduced insulin and insulin mRNA as well as decrease level of insulin receptors [8]. In vitro studies have shown the impaired insulin signaling,

hyperphosphorylated tau proteins, and neural loss in T2DM affected AD animal models [70–73].

Increased cerebrospinal fluid (CSF) insulin levels are correlated with high peripheral insulin in AD patients [13]. Insulin also has effects on A β metabolism. Hyperinsulinemia can increase extracellular A β concentration by stimulating its trafficking from A β generating sites (trans-Golgi network and endoplasmic reticulum) that results in reduced intracellular concentrations of A β . Insulin-degrading enzyme (IDE) plays an important role in A β degradation. Insulin can competitively inhibit IDE that results in reduced A β degradation [74, 75]. Induced activity of IDE in transgenic mice brain results in decreased levels of A β thereby preventing cognitive decline [76]. Furthermore, persistent peripheral hyperinsulinemia leads to decreased transportation of insulin to the brain and causes an insulin-resistance brain state resulting in reduced insulin levels in CSF [7]. Brain insulin resistance consequently decreases the IDE levels in AD patients resulting in increased A β neurotoxicity [77]. Interestingly, A β also has the ability to reduce insulin signaling and receptor autophosphorylation suggesting that A β competitively inhibits insulin binding. Moreover, A β is also reported to reduce insulin receptors substrates signaling in cultured cells [74, 78]. These findings suggest an important role of insulin dysfunction in neurodegenerative diseases.

Insulin Signaling Mechanism

Insulin implements its actions through insulin signaling pathway by interacting with insulin receptors. A brief outline is presented here. Preproinsulin an inactive single chain precursor with a signal sequence is the primary form of insulin secreted by pancreatic β -cells. After removal of signal sequence by proteolysis, preproinsulin changed to proinsulin that is further converted to insulin by special proteases when blood glucose or amino acid concentration increases. Active insulin is consisted of two chains, bound together with disulfide bonds [79]. Insulin performs its duty by binding to its specific receptors known as insulin receptors (IR). Insulin binds to the α -subunit of the receptor and activates the tyrosine phosphorylation of β -subunit of the receptor [80, 81]. This process activates two important signaling pathways: Akt (also known as protein kinase B [PKB]) and MAPK (mitogen activated protein kinase) [82, 83]. Akt signaling regulates cell growth, proliferation, survival, and protein synthesis; while MAPK signaling activates cell differentiation, proliferation and death (Fig. 1).

Insulin Signaling in Diabetes and AD

Activation of Akt results in phosphorylation of 3'-OH group of inositol thus generating PIP2 and PIP3 (phosphoinositide

phosphates) that further activates serine/threonine kinases, 3'-phosphoinositide-dependent kinase-1 (PDK-1), PKB/Akt, and MAPK regulatory pathways [84, 85]. As Akt and MAPK are major pathways involved during insulin signaling through tyrosine phosphorylation on insulin receptors, defects in insulin receptors, insulin receptor substrates (IRS) and these pathways lead to complications that are hall mark of diabetes, AD and their related abnormalities [86, 87].

Defects in Insulin Receptors

A decrease in tyrosine kinase activity of insulin receptors was observed in T2DM suggesting a post-receptor defect with reduced association of the insulin receptor with p85 subunit of Akt leading to impairment of insulin stimulating glucose disposal and glycogen synthesis. This also leads to defects in glucose transportation and lower level of GLUT4 mRNA in muscles. Glucose may undergo glycolytic pathway or glycogen synthesis after phosphorylation through hexokinases (HK-I to -IV) [67, 88, 89]. Increased insulin induces glycogen synthesis, while decreased glycogen synthesis in muscle cells due to insulin resistance is hallmark in T2DM especially in skeletal muscles [67, 90]. Rare mutations on IRS especially IRS-2 may lead to insulin resistance in T2DM patients as chances of diabetes were more in IRS-2 mouse mutants compared to IRS-1 [91]. Insulin receptors found in the brain's hippocampus and cerebral cortex play important role in memory and learning [8, 92]. Mutated mice lacking IRS-2 show improved premature mortality and A β deposition lead to the IR role in AD pathology [93, 94]. A reduced insulin signaling was observed in brains of AD patients. A β also has the ability to induce insulin resistance by down regulating the IRS expression [74].

Akt Signaling

The binding of Akt with insulin receptors is affected by serine phosphorylation of IRS that decreases the tyrosine phosphorylation of IRS due to several mechanisms such as hyperglycemia, inflammation, hyperinsulinemia, hyperlipidemia and mitochondrial dysfunction. It may lead to disassociation of Akt from IRS or IRS from insulin receptors [95, 96]. Circulating free fatty acids (FFA) and tumour necrosis factor (TNF) may induce serine phosphorylation in muscle cells leading to IRS dysfunctioning [97, 98]. Although the mechanism of insulin signal impairment through FFA is still unclear, the intracellular lipid metabolites such as diacylglycerol (DAG) and acetyl CoA lead to insulin resistance through IRS serine/threonine phosphorylation via protein kinase C (PKC) and may result in endothelial dysfunction in T2DM [99, 100]. DAG that induced the PKC is also observed in muscles during lipid infusions and fat feeding. FFA also induces PKC phosphorylation independent of




Fig. 1 Linking insulin-signaling mechanism with diabetes and AD. Impaired insulin signaling due to insulin resistance or defects in insulin receptors and Akt phosphorylation lead to activation of all pathways that remain block during normal signaling. Deactivation of Akt triggers

several transcription factors that resulted in abnormal cell functions, apoptosis, neuronal cell death via NFT production, tissue atrophy and inflammation. These are major hallmarks in patients with AD and diabetes

PDK1 and translocates PKC to the nucleus and inhibits IR gene transcription [101, 102]. A recent study by Liu et al. [103, 104] has observed the decreased levels of insulin signaling components, increased hyperphosphorylation of tau and reduced *O*-GlcNAcylation (OGlcNAc) levels in the brain of patient suffering from T2DM. Decreased brain glucose levels are resulted in reduced *O*-GlcNAc levels that regulate phosphorylation of tau inversely. Interestingly, *O*-GlcNAc modification also regulates the AKT phosphorylation [103, 104]. It is interesting to know that at first insulin induced IDE expression in hippocampal neurons, while IDE decrease is associated with reduction of Akt p85 subunit. This observation lead to the hypothesis that IDE reduction by insufficient insulin action might increase the chances of AD, as brain IDE activity reduced in hyperinsulinemia conditions [105, 106].

Akt signaling may be blocked by PTEN (tensin homologue deleted on chromosome ten) and tumour suppressor protein phosphatase resulted in dephosphorylation of PIP2 and PIP3 [107, 108]. Imbalances among the Akt subunits p85 and p110, and their heterodimer p85–p110 are also responsible for insulin resistance. Akt p85 unit competes

with p85–p110 (as this heterodimer unit is responsible for Akt activity) and the p85 induction is caused by human placental growth hormone. This type of resistance is observed in pregnant women with T2DM [109–111]. Akt triggers downstream effects leading to glycogen synthesis, hepatic glucose production and on/off switch of many transcriptional factors. Studies in mice have shown that deletion of Akt leads to insulin resistance ending up with diabetes phenotype as humans. Upon activation, Akt blocks apoptosis related genes such as BAD, caspase 9, GSK3, FoxO family transcription factors, CREB and NF- κ B. It also blocks the MDM2 activity by blocking p53 [106, 112]. Furthermore, Akt phosphorylation allows the translocation of GLUT4 (glucose transporter) to the plasma membrane by AS160 phosphorylation [113]. The mutated Akt was unable to phosphorylate downstream targets and inhibition of gluconeogenic through phosphoenolpyruvate carboxykinase (PEPCK) [114, 115]. Another factor regulating gluconeogenesis and its related genes PEPCK and G-6-pase is FoxO protein 1 and 3 in patients with hyperglycemia. A reduction in Akt activity leads to decreased phosphorylation on FoxO allowing its entrance to the nucleus and activation of its

transcription factors in T2DM. Phosphorylation of FoxO regulates gene transcription and inhibits apoptosis to promote neural growth and survival through phosphorylation of BAD (Bcl-2 associated death promoter protein) [116–118].

Akt also initiates protein synthesis by p70s6k phosphorylation that may shutdown IRS activity by direct phosphorylation of IRS through mTOR (overactive mammalian target of rapamycin) signaling triggering by Rheb (GTPase RAS homolog enriched in brain). Akt also regulates the binding of TSC1 protein from TSC2 (tuberous sclerosis protein 1 and 2). Upon phosphorylation, TSC2 is released and activates Rheb and mTOR pathway [119, 120]. mTOR further activates initiation factor 4E-BPs (4E-binding proteins) phosphorylation and inhibits its binding with eIF (eukaryotic initiation factor) and promotes eIF4E cap-dependent translation that control cell growth and survival. This event also activates adipocyte differentiation transcription factors, PPAR γ , CCAAT enhancer binding protein α (E/EBP), etc., that leads to adipogenesis, obesity and insulin resistance if not properly controlled [121–124]. Activation of p70S6 protein regulates ribosomal function by phosphorylating ribosomal proteins and initiates amino acid addition to newly synthesized peptides. In T2DM, effected kidney tissues, induced mTOR levels were observed with matrix expansion and renal hypertrophy [125, 126].

Under normal conditions Akt signaling phosphorylate the glycogen synthase kinase 3B (GSK3B) and inactivates glycogen synthase. Insulin resistance leads to dephosphorylation and activation of GSK3B. Increased expression of GSK3 was observed in patients with neurodegeneration and was associated with Tau hyperphosphorylation [127, 128]. A study by Schubert et al. reported high tau phosphorylation with significantly reduced phosphorylation of Akt and GSK3 in NIRKO mouse brains [129, 130]. Hyper tau phosphorylation via GSK3 impairment is associated with NFT formation in brain, which is an important hallmark of AD [131].

In brief, defects in Akt signaling not only lead to defects in glycogen synthesis, gluconeogenesis, glucogenolysis, and amino acid synthesis but also promote AD through hyperphosphorylation of tau and reduction of *O*-GlcNAcylation.

MAPK Signaling in Diabetes and AD

MAPK signaling activates intracellular enzymes that have ability to respond to stimuli from extracellular environment. MAPK pathway runs parallel to Akt and stimulates extracellular signal-regulated kinases 1/2 (ERK1/2) that induce activation of several transcription factors such as Elk-1 and c-Myc that are important in cell growth regulation [132, 133]. The Ras-ERK/MAPK activated cascade is thought to be involved in memory formation by synaptic plasticity [134]. MAPK signaling also activates MK2

(MAPK-activated protein kinase 2) that is further involved in mediating inflammatory response to cellular stress. It is shown by cell culture studies showed that MAPK activation resulted in expression of inflammation-associated iNOS, TNF α and IL-1 β genes [135]. MAPK activation also up-regulates cytokine production by direct phosphorylation of transcription factors such as CREB (cAMP response element binding). CREB has direct effect on structural changes associated with memory formation [136].

MAPK pathway is involved in excessive tau phosphorylation, neuroinflammation and synaptic plasticity in AD patients. Induced MAPK expression is reported in hippocampal and cortical regions of brain in AD patients when compared with aged-matched healthy individuals. Furthermore, greater immunoreactivity of MAPK is observed in AD post-mortem brains [137, 138]. Moreover, in transgenic mice, tau hyperphosphorylation was found to be directly associated with phosphorylated MAPK. Furthermore, A β fibrils in microglia also able to activate MAPK signaling that resulted in inflammatory gene expression and elevated proinflammatory cytokines [139]. Briefly, impaired MAPK signaling is not only associated with neuroinflammation but also leads to the promotion of A β and NFTs generation.

Abnormal Glucose Metabolism

Abnormal glucose metabolism is a characteristic of T2DM and is mechanistically linked with AD [52]. Metabolic abnormalities due to impaired glucose consumption and energy metabolism in AD almost resemble with T2DM. In addition, impaired glucose metabolism and insulin resistance also lead to memory and synaptic dysfunction in patients suffering from diabetes (Fig. 2) [140, 141]. Furthermore, increased ratio of plasma insulin levels to decreased CSF was reported in AD patients [13]. These findings lead to the conclusion that normal glucose metabolism is associated with standard cognitive response and any defects in glucose metabolism lead to cognitive system decline. Brain neurons are unable to produce and store glucose, and require continuous glucose transportation through the BBB by glucose transporters (GLUTs). Glucose transporters, isoforms 1, 3, 4 and 8 are abundant in the brain. GLUT 1 expressed on BBB endothelial cells and cortical membranes, GLUT 3 expressed on neurons while GLUT 4 and GLUT 8 expressed in intracellular compartments of neurons [10, 142, 143]. Reduced levels of GLUT1 observed in AD patients as well as in transgenic mice lead to reduced supply of glucose from peripheral transport system [8]. Reduced GLUT 1 and 3 levels and decreased glucose utilisation are likely to be associated with tau hyperphosphorylation, increased density of NFTs, and reduction in *O*-GlcNAcylation [144–146].

Functional neuroimaging technique positron emission tomography (PET) has been found best to monitor neuronal




Fig. 2 Hyperglycemia-prompted activation of molecular pathways associated with AD, and diabetes. Defects in insulin signaling pathway and/or in production of insulin lead to hyperglycemia. Hyperglycemia promote production of AGES, oxidative stress, and induce levels of various growth factors and cytokines that accumulatively endorse the AD and diabetic complications

activity and specific biological processes at tissue levels in vivo. PET with 2- ^{18}F fluoro-2-deoxy-D-glucose (FDG) is used to estimate qualitative and quantitative changes in cerebral metabolic rate of glucose (CMRglc) [147]. Several studies have reported the cerebral metabolism decline before the attenuation of cognitive functions. In AD, reduced CMRglc has been reported in hippocampus, parietal-temporal and prefrontal regions of brain when compared with age-matched health individuals [148, 149]. Decreased CMRglc metabolism may lead to mitochondrial dysfunction, hyperglycemia and inflammation within the brain. A 55–65 % decrease in cerebral glucose utilization was observed not only in AD patients and in transgenic models but also in individuals carrying the apolipoprotein 4 allele. ApoE4 allele can lead to reduced glucose metabolism and energy production within brain [21, 150, 151]. Reduced activity of mitochondrial proteins mainly associated with TCA cycle such as α -ketoglutarate dehydrogenase, pyruvate dehydrogenase, and isocitrate dehydrogenase was observed in AD patient's brain tissues. This reduced activity was resulted in the release of cytochrome *c* from mitochondria, and caspase-3 activation leading to neuronal cell death. Moreover, these changes were positively associated with the degree of clinical disability in AD [152, 153]. Furthermore 50 % decrease in ATP production from glucose was also reported in early AD. All these findings suggest that

ATP-dependent energy generation is vital for normal cell performance [154].

Although glucose is the least reactive reducing sugar, it may lead to non-enzymatic Amadori products through Schiff base formation by reacting with free amino groups of proteins, lipids and nucleic acids. These Amadori products accumulate on proteins and initiates process of advanced glycation [155, 156]. Hyperglycemia resulting from insulin resistance promotes the production of AGEs that are involved in AD pathogenesis. Although this process occurs in normal aging, it becomes highly accelerated in diabetes. AGEs are accepted as active contributors in AD progression and induce A β and tau glycation that lead to A β aggregation and NFTs formation in the brain [157, 158].

AGEs implement their functions and induce several biological processes through binding with RAGEs (receptors for AGEs). RAGEs not only interact with AGEs but also bind and interact with A β . Activation of RAGEs through binding with these ligands may initiate ROS production and inflammation [159, 160]. Levels of RAGEs were increased in RAGE-bearing cells from AD patients when incubated with A β that resulted in increased oxidative stress through NADPH oxidase-like mechanism, and activated NF- κ B pathway [161]. Furthermore, A β transportation across the BBB was associated with RAGE that was blocked by using anti RAGE IgG or sRAGE (soluble RAGE). Moreover, RAGE associated transport of A β and induction of tumour necrosis factor alpha (TNF- α), interleukin-6 (IL-6) and heme oxygenase type 1 was also observed in mice neuronal cells. Mice administrated with sRAGE showed decreased A β production and accumulation in brain parenchymal regions [162].

It is suggested that vascular remodelling in diabetes and AD linked cerebrovascular amyloid angiopathy (CAA) are both associated with RAGE expression [163]. A β -RAGE interaction also has effects on cerebral blood flow in murine models. Infusion of A β into peripheral circulation resulted in reduced cerebral blood flow that was prevented by sRAGE, RAGE IgG or RAGE specific inhibitor (FPS-ZM1) administration [162, 164]. Furthermore, induced levels of vasoconstrictor mediator endothelin-1 were induced to block A β -associated suppressed cerebral blood flow. A recent study by Xiong et al. [165] reported that RAGE expression in human umbilical vein endothelial cell line (ECV-304) had prevented the endothelial cell membrane repair by activating β -catenin levels that resulted in reduced F-actin stress fibres and attenuated plasma membrane resealing [165].

Impaired glucose metabolism also plays a very important role in NFTs formation by promoting hyperphosphorylation of tau protein. Tau protein undergoes many posttranslational modifications (PTMs) particularly phosphorylation and glycosylation. Tau phosphorylation has been thought to inhibit

the binding of tau protein from microtubules, while hyperphosphorylation leads to self-aggregation of tau filaments and NFTs formation. *O*-glycosylation (addition of one molecule of *N*-acetylglucosamine to serine or threonine residue) is thought to regulate tau phosphorylation reciprocally by competing phosphate groups on single or proximal Ser/Thr residues [166, 167]. Although exact mechanism by which *O*-glycosylation regulates tau phosphorylation is unknown, *O*-glycosylation is thought to be associated with glucose metabolism. Furthermore, STZ-induced diabetic rat showed reduced *O*-glycosylation levels that were restored after insulin treatment [168]. Decreased GLUTs levels and *O*-glycosylation have also been reported in the AD brain when compared to normal [20, 169]. A recent study by Yuzwa et al. [170] has observed that *O*-glycosylation at tau protein not only delays neurodegeneration but also hinders tau oligomerization. These observations suggest *O*-glycosylation as a potential therapeutic target to slow down AD progression and this mechanism also provides evidence of shared pathology between diabetes and AD.

Inflammation

Insulin resistance in diabetes and AD also leads to inflammation. Vascular inflammation mediated by RAGE has been proposed as a possible mechanism for vascular dysfunctions in AD and diabetes. Furthermore, the levels of inflammatory mediators IL-6 and C-reactive proteins were raised in diabetic, and AD patients [171]. It is reported that diabetes accelerates memory dysfunction via promoting A β aggregation and cerebrovascular inflammation by up regulating RAGE in diabetic mice AD models [2]. Ramasamy et al. [172] and Srikanth et al. [173] extensively reviewed the role of AGEs and RAGE in development of AD. Briefly, RAGE can be activated by many ligands as described earlier. AGEs and A β synergistically induce the expression of the proinflammatory cytokines IL-6, TNF- α and M-CSF (macrophage colony-stimulating factor). A β -mediated persistent activation of microglial cells due to proinflammatory state is resulted in neuronal cell death and disease progression in AD. Interestingly, inflammation induced levels of both A β and AGEs. Stimulation of TNF- α accelerates BACE1 expression and resulting in enhanced amyloidic processing of APP in astrocytes and formation of A β in diabetic models of AD [172, 173].

Summary of the Section

Impaired insulin signaling and glucose metabolism is not only risk for diabetes but also for AD. Peripheral insulin dysfunctions induce impairments to insulin functions in brain that result in A β aggregation and NFTs formation. Furthermore, abnormal glucose metabolism not only promotes oxidative

stress, A β aggregation and NFTs formation but is also responsible for vascular angiopathy and inflammation in AD as well as in diabetes. Additionally, AGEs and RAGE both promote AD and diabetic complications and provide a common link between these two diseases. In the words of Kroner et al. [174], “co-existence of brain insulin deficiency and resistance suggest that AD might be form of diabetes i.e., type 3 diabetes.”

Linking Lipids with AD and Diabetes

AD and diabetes are not only characterized by insulin metabolism impairments but also by dyslipidemia. Lipid-mediated signaling regulates many physiological processes like trafficking and proteolytic activities of membrane bound protein [175, 176]. Fatty acids (FAs) are mostly synthesized in all body cells, stored and circulated as triglycerides. FAs can be degraded by lipases such as phospholipase (PL) and lipoxygenase (LOX), and resynthesized in the form of phospholipids [177]. Increased levels of FFA, triglycerides, cholesterol and low density lipoproteins (LDL), and reduced levels of high density lipoproteins (HDL) are well-described risk factors not only for diabetes but also for neurodegenerative diseases like AD [178–181]. Induced FFAs levels in brain were also linked with mitochondrial dysfunction, and IDE inhibition that resulted in increased A β production [182, 183].

Phospholipase isoform A2 (PLA2) hydrolyses the phospholipids and converts them to arachidonic acid (AA) which further can be converted to various signaling molecules required for cellular metabolism [177]. Several studies have found that induced levels of LOX and PLA2 are associated with higher intensity of A β , and cognitive decline in transgenic mice models [184, 185]. Furthermore AA also activates NADPH, a major source of ROS production in neurons that consequently activates cytoplasmic PLA2 (cPLA2) [186, 187]. cPLA2 regulates AA release at synapses. In neurons, AA is involved in synaptic functions and it's over production can trigger depolarization of neuronal cells via calcium dependent apoptosis [188]. FAs metabolism through LOX also produces ROS and is very sensitive to glutathione (GSH) a major antioxidant in mitochondria. GSH can be depleted under high concentrations of LOX [189]. A reduction in A β production has been reported in LOX deleted transgenic mice [184]. Moreover, LOX metabolites such as 5-hydroperoxy eicosatetraenoic acid and leukotriene have been reported to positively induce A β production in cultured cell lines [190]. AA can also be converted to inflammation mediators, eicosanoids; such as prostaglandins by the action of cyclooxygenase (COX) [191]. Elevated levels of PLA2, LOX and COX are reported in patients with AD and diabetes [192]. These data suggest

that diabetes can mediate AD through lipid metabolism as increased FFAs levels are reported in diabetes and derivatives of AA may be responsible for A β production in AD [131].

Hypercholesterolemia also contributes to the pathology of AD and diabetes. Hypercholesterolemia is associated with increased risk of type 2 diabetes, and more than 70 % patients diagnosed with type 2 diabetes have hypercholesterolemia. High levels of cholesterol and other fatty acids lead to atherosclerosis and vascular lesions [193, 194]. In the brain, unesterified cholesterol is present in cellular membranes and myelin sheets of astrocytes and neurons. As the majority of lipoproteins cannot cross BBB, cholesterol in the brain is derived from *de novo* synthesis [195]. In the brain excess cholesterol is converted to cholesteryl esters by enzyme acyl CoA: cholesterol acyltransferase 1 (ACAT1) or oxysterol 24S-hydroxycholesterol (24OHC). 24OHC can be easily transported to peripheral circulation through BBB. Moreover, 27S-hydroxycholesterol (27OHC) which is produced outside the brain has the ability to cross the BBB and is reported to increase in AD patients [196–198]. Brain lipoproteins that transport lipids resemble to HDL particles present in CSF and apolipoprotein E is the most studied lipid carrier in AD pathology discussed later [199]. Induced levels of ACAT1 associated with increased A β generation and vice versa in mouse models suggests direct association of cholesterol esters with A β production and AD promotion [200, 201]. Furthermore, cholesterol also modulates the BACE1 and secretase activities. Reduction in cholesterol levels lead to decreased activities of BACE1 and secretase that resulted in reduced A β production. Moreover, association of BACE1 with lipid rafts is found to be positively linked with amyloid processing of APP, and production of A β is decreased after cholesterol depletion [202, 203].

ApoE is well-established risk for the late-onset AD. ApoE that is significant for lipid transport also plays an important role in A β clearance as well as stimulation of IDE which is also involved in A β degradation [204]. Normal individuals have 77 % ApoE3 allele and 15 % ApoE4. ApoE4 allele is significantly higher in AD patients, and normal individuals carrying ApoE4 allele have 3- to 4-fold more chances to develop AD when compared to non carriers [193]. Furthermore, diabetic patients who are ApoE4 allele carrier have 2-fold more chances to develop AD than non-diabetic individuals [205]. A β clearance in the brain is ApoE isoform specific (ApoE4<ApoE3<ApoE2). High levels of ApoE4 were associated with induced A β generation [206]. Moreover, transgenic over expression of ApoE4 in neurons increases the tau phosphorylation suggesting that ApoE4 is also involved in development of NFTs by unknown mechanism [207]. Diabetes and ApoE4 synergistically increase the risk for AD. Various studies linked type 2 diabetes to brain pathology particularly in individuals with

ApoE4 allele (reviewed by Luchsinger et al. [208]). Low lipid clearance characteristic of ApoE4 may also make it an independent risk factor for type 2 diabetes. A study by Chaudhary et al. [209] has found that ApoE4 allele has influence on lipid plasma levels and is associated with diabetes with or without coronary artery disease.


Intracellular cholesterol trafficking is mainly thought to be modulate by ATP-binding cassette (ABC) transporter family proteins such as ABCA [210]. ABCA1 primary function is the transportation of cholesterol efflux onto lipid-poor ApoE. Mutations in human ABCA1 allele result in Tangier disease (absence of HDL in plasma with increased cardiovascular risk) and cause significantly low levels of ABCA1 and ApoE in brain [211]. ABCA1 deficiency in mice lead to 80 % decrease in brain ApoE levels. Thus poor lipidation of ApoE due to decreased ABCA1 levels is resulted in increased A β production and vice versa [212]. A recent study by Wang et al. has reported high cholesterol levels in STZ-induced rat brain with decreased expression of ApoE mRNA. However, they have observed decreased cholesterol levels and high ABCA1 expressions in peripheral tissues [213]. These results suggest that in case of diabetes, the body tends to decrease cholesterol levels by expressing ABCA1. Moreover diabetes also results in high cholesterol levels in the brain.

The above data suggests that impaired lipid metabolism is as important as insulin dysfunctions in AD and diabetes. Although much attention is given to ApoE but still lipid metabolism association between AD and diabetes has not been fully addressed till date. More research is required to explore the role of fatty acids in the formation of A β plaques and NFTs in AD, as this mechanism is still unclear. Figure 3 represents a way how impaired fatty acid metabolism may be responsible for diabetes and AD promotion. Secondly, very limited studies are available on any correlation between human amylin and ApoE in diabetes as well as in AD. Although one study found that amylin fibrillation could be prevented by lowering the ApoE4 levels in T2DM [214], no further evidence came from this side.

Mitochondrial Dysfunctions

Aerobic organisms mostly produce energy in the form of ATP utilizing oxygen and nutrients. Mitochondria produce over 90 % of cellular ATP through oxidative phosphorylation, which is also the major source of ROS production. Furthermore mitochondria also regulate other physiological processes and cellular functions like cell survival and death, intracellular calcium homeostasis, cell cycle regulation and synaptic plasticity [215, 216]. AD and diabetes are associated with mitochondrial dysfunctions. Especially as neurons require high energy to perform their functions, limited glycolytic activity

Fig. 3 Lipid metabolism in AD and diabetes. Production/accumulation of excessive free fatty acid and cholesterol lead towards several complication like oxidative stress, inflammation, atherosclerosis, vascular lesions, and neuronal loss that are major hallmarks of these diseases



of neuronal cells makes them highly dependent on mitochondrial energy production and any changes associated with mitochondrial impairment lead to neuronal dysfunction and neurodegeneration [217, 218]. Mitochondrial abnormalities like over oxidative stress and impaired calcium homeostasis are reported in AD and diabetes.

Oxidative Stress

According to Perez-Matute et al. [218], "living with the risk of oxidative stress is a price that aerobic organisms must pay for more efficient bioenergetics." Reactive oxygen species (ROS) and reactive nitrogen species (RNS) are important oxidants in living organisms. These ROS produced can be converted to many forms and have the ability to permeate through inner and outer mitochondrial membranes (OMM) to enter in cytoplasm. However, most ROS produced in mitochondria can be converted into water or oxygen in the presence of metal-dismutases either in mitochondria or in cytoplasm as described later [219, 220]. The oxygen-derived pro-oxidants (ROS) can be classified into two groups; radicals and nonradicals. The radicals are superoxide ($O_2^{\cdot-}$), alkoxyl (RO^{\cdot}), peroxy (ROO^{\cdot}), hydroxyl (OH^{\cdot}), hydroperoxyl (HO_2^{\cdot}) and nitric oxide (NO^{\cdot}). The reactive, oxygen-containing nonradicals include hydrogen peroxide (H_2O_2), organic peroxides ($ROOH$), aldehydes ($HCOR$), hydrochlorous acid ($HOCL$), and peroxynitrite ($ONOOH/ONOO^{\cdot}$) [221–224].

Several hypotheses have been proposed to explain the role of oxidative stress in AD and diabetes pathophysiology, while the combination of high-energy demand, relatively low antioxidant levels and a high concentration of iron

makes the brain particularly vulnerable to oxidative stress [221, 225]. In AD pathology, decline in synaptic activities and defects in low energy metabolism with comparatively increased amount of ROS are observed [226, 227]. Moreover, reduced levels of antioxidants enzymes like Cu/Zn SOD, glutathione (GSH) and catalase are found in frontal and temporal cortex. Furthermore presence of $A\beta$ and NFTs lead to mitochondrial dysfunctions, generation of oxidative stress and neuronal cell death [78, 228–230]. Meanwhile, diabetic patients exhibit a number of oxidative stress markers including F_2 -isoprostane and nitrotyrosine in urine, plasma and tissue, leading to the hypothesis that oxidative stress is also a hallmark of diabetes [231, 232]. Although mitochondria are capable of generating ROS/RNS by themselves however, in AD and diabetes, other sources/mechanisms like misfolded proteins ($A\beta$ and hIAPP), accumulated NFTs (tau plaques), hyperglycemia and AGEs also promote ROS generation [56, 233, 234]. We will briefly discuss the generation of ROS by these sources.

Generation of Oxidative Stress by Mitochondrion Itself

Many authors have reviewed the production of ROS and RNS in mitochondria in detail [235–239]. Here we briefly describe the ROS/RNS generation in mitochondria. In total, nine different types of enzymes in mitochondria have the capacity to produce ROS by converting molecular oxygen to either superoxide anion or hydrogen peroxide. Cytochrome *b5* reductase and monoamine oxidases (MAO) are located on OMM, whereas dihydroorotate dehydrogenase (DHOH, α -glycerophosphate dehydrogenase (α -GDH), succinate dehydrogenase (SDH), aconitase, α -ketoglutarate dehydrogenase

complex (KGDHC), complex I and complex III are present on inner mitochondrial membrane (IMM). MAO, DHOH, α -GDH and KGDHC produce H_2O_2 via direct or indirect biochemical reactions, while cytochrome b5, complex I and complex III produce superoxides. Aconitase localized in the mitochondrial matrix generates hydroxyl radical upon oxidation of its iron–sulphur cluster by superoxide. PDHC and KGDHC have also been reported to generate superoxide and hydrogen peroxide when isolated and purified from bovine heart, and mouse brain mitochondria.

Generation of Oxidative Stress by A β , Tau and hIAPP

A β insertion in mitochondria has been reported to disrupt the electron transport chain, reducing energy production and increased ROS production [240]. Reduced activity of complex IV in AD patients is well documented and A β is believed to inhibit the functions of complex IV. A β has been shown to produce hydrogen peroxide (H_2O_2) and release thiobarbituric acid reactive substances (TBARS) mainly associated with hydroxyl radicals (OH) via metal ion reduction [241, 242]. Furthermore, A β can also act directly on mitochondria. A β plaques interact with A β -binding alcohol dehydrogenase (ABAD) resulting in increased mitochondrial membrane permeability and reduced activities of respiratory proteins [243–245]. The oxidation of proteins at lysine, arginine, proline and histidine residues via peroxynitrite generates protein carbonyls and nitrile, both of which cause increase in NFTs, non-tangle bearing neurons and glia of AD patients [243].

Even though, several studies report the association of A β and oxidative stress, few studies have looked at the role of hIAPP. However, two studies have reported the elevation of oxidative stress markers using immortalised beta cell lines when treated with hIAPP exogenously [246, 247]. Furthermore, β -cells of patients with diabetes have shown higher oxidative stress with increased levels of hIAPP. A study by Zraika et al. [56] reported that amyloid deposition on mice islets is associated with increased ROS levels and beta cell apoptosis. Although treatment with antioxidants prevents ROS generation, it does not reduce amyloid formation. In contrast, amyloid inhibition reduces ROS generation as well as beta cell apoptosis. This information leads to the conclusion that hIAPP also induce oxidative stress and apoptosis although the exact mechanism is still unknown.

Hyperglycemia Induced ROS Generation

Hyperglycemia due to insulin impairment in diabetic patients is associated with many biochemical pathways including glucose mediated ROS production, protein kinase activation, formation of AGEs and cytokine secretion [233]. In glucose autooxidation, glucose forms enediol radicals and

is converted to reactive ketoaldehydes and superoxide; consequently hydroxyl radicals are produced in presence of transition metals via H_2O_2 if not degraded by catalase or glutathione peroxidase [248, 249]. Hyperglycemia drives the inner mitochondrial membrane potential upward through generation of excessive electron donors in the Krebs cycle [250]. Hyperpolarization of the mitochondrial inner membrane leads to an increased ATP/ADP ratio, which inhibits electron transport to complex III leaving coenzyme Q in a reduced state. Coenzyme Q can pass electrons to molecular oxygen to generate the partially reduced derivative, superoxide [155, 251]. This superoxide overproduction causes a 66 % decrease in GAPDH (glyceraldehyde-3-phosphate dehydrogenase) activity, PARP (poly-ADP ribose polymerase) activation, and NAD^+ depletion [252]. Overproduction of superoxidase radicals is countered by superoxide dismutases (SODs) and by uncoupling proteins (UCPs). In hyperglycemia, over expression of UCPs blocks glucose induced cell death by preventing mitochondrial hyperpolarization and ROS formation [253, 254]. Hyperglycemia was found to be responsible for cognitive decline in diabetic patients. Impaired antioxidant system with increased oxidative/nitrosative stress in hyperglycemic conditions has been observed in brain of STZ diabetic rats. Authors have also reported about reduced activity of ETC. complexes III, IV and V, and ATP synthesis [255]. Moreover in another study, 12-week older STZ-induced diabetic rat also showed lower ATP contents and calcium accumulation ability. A recent study by Cardoso et al. and Raza et al. also has reported increased ROSRNS levels, reduced antioxidant activity and decreased activities of mitochondrial enzymes such as complex III, complex IV in cortical and hippocampal mitochondria as well as pancreas of STZ induced diabetic rats [256, 257].

AGEs Induced ROS Generation

AGEs found in senile plaques also produce free radicals by chemical oxidation and degradation, by binding with their receptors (RAGE) or interacting with microglia that surround senile plaques and results in respiratory blast and production of ROS and RNS. AGEs particularly produce superoxides and hydrogen peroxides, and their production is associated with related proteins and sugars oxidative insult. Furthermore, AGEs also produce ROS by the metal-catalysed Fenton reaction that result in site-specific attack on proteins and lipid peroxidation [173, 258–260].

Consequences of Oxidative Stress in AD and Diabetes

Impaired Insulin Signaling

Some studies reported that high concentrations of H_2O_2 are able to activate insulin signaling and induce insulin-

associated metabolic actions leading to increased glucose uptake, stimulation of GLUT4 translocation and lipid synthesis [261, 262]. Furthermore, under oxidative stress, stress-associated signaling cascades like MAP kinase become activated and induce phosphorylation of insulin receptors (IRS) resulting in protein degradation and the release of IRS from membrane pools [263, 264]. All these processes accumulatively result in impaired insulin signaling.

Lipids and Protein Oxidation

Lipids and proteins are important constituents of nutrition and growing body of literature points out changes in their respective metabolism in AD and diabetes. Several lipid and protein oxidative products and their derivatives are consequences of oxidative stress in AD and diabetes, and might be able to predict the onset and progression of these diseases. Cell lipids especially cholesterol and polyunsaturated fatty acids (PUFAs) are most susceptible to oxidative stress and their oxidized derivatives are more reactive than parent compounds [265, 266]. Oxidation of cells, plasma and tissue proteins is also very prominent in AD and diabetes. Oxidative damage to proteins promote unfolding and conformational changes that lead to loss of proteins functions and aggregation of cross-linked protein structures [267]. Protein carbonylation, nitration, glutathionylation, lipid–protein interactions and AGEs formation are well described oxidative end products, and their accumulation results in the disruption of cellular functions and pathways leading to apoptosis and necrosis [243].

Oxysterols are important oxidation products of cholesterol in brain and are associated with proinflammatory, proapoptotic and profibrogenic effects [268, 269]. Furthermore, oxysterols are also involved in up-regulating APP and BACE1, and induction of A β peptides when studied in SH-SY5Y human neuroblastoma cells [270].

PUFAs are important constituents of membrane phospholipids and play an important role in enzyme activities and membrane fluidity. Two forms of PUFAs; PUFA ω -3 and ω -6 are essential for brain homeostasis and decreased ω -3/ ω -6 ratio are early markers of AD [271, 272]. α -Linolenic acid (ALA), eicosapentaenoic acid (EPA) and docosahexaenoic (DHA) are well known types of ω -3 PUFAs while linoleic acid (LA) and arachidonic acid (AA) are important forms of ω -6 PUFAs [265]. Non-enzymatic peroxidation of PUFAs resulted in unstable peroxy radical intermediates, which further converts to endoperoxides. Further reduction of chain after the insertion of second oxygen yielded in different isoforms of isoprostanes (IsoPs) like F2-IsoPs, F3-IsoPs and F4-IsoPs, and aldehydes such as malondialdehyde (MDA), acrolein and hydroxyalkenals especially 4-hydroxynonenal (HNE) in AD as well as in diabetes [273, 274].

Induced levels of F2-IsoPs and F4-IsoPs have been reported in hippocampus, frontal, temporal, parietal, and occipital lobes of AD patients when compared with normal individuals. As IsoPs increased to the same extent in mild cognitive impairment as well as in late one AD, these are not considered as reliable indicators of disease progression [275, 276].

Although MDA are popular oxidative stress markers, these are not considered as specific AD markers due to their presence in mild cognitive impairment as well as in healthy elderly persons. However, MDA was reported to induce oxidative stress by inhibiting ETC. complexes I, II and V, and other respiratory enzymes such as pyruvate dehydrogenase, α -ketoglutarate dehydrogenase and superoxidodismutase [277, 278].

Other oxidative aldehyde products acrolein and HNE have been detected in the brains of AD patients and are thought to be molecular mediators in AD pathogenesis [279]. Acrolein adducts are usually found in NFTs and induce tau oligomerization that results in aggregation of paired helical filaments. Furthermore, tau oligomerization is promoted during tau phosphorylation; and acrolein is thought as NFTs promoter [280, 281].

Most attention is given to HNE as it is highly reactive and makes bonds with amino and thiol groups of proteins [282]. These protein-bound HNE are considered as important contributors in formation of NFTs [283]. Moreover, HNE are also involved in production of short A β peptides which are also a source of oxidative stress in AD [284]. Furthermore, HNE also disrupts the binding of histones to DNA and increase chances of DNA oxidation in AD brain [285].

The oxidation of proteins at lysine (Lys), arginine (Arg), proline (Pro), threonine (Thr) and histidine (His) residues via peroxynitrite generates protein carbonyls and nitrile, both of which have shown increase in NFTs, non-tangle bearing neurons and glia of AD patients [243].

Amino acids like His, Lys, Arg, Pro and Thr are major targets for carbonylation and resulted in the formation of AGEs and products of lipid peroxidation. Protein carbonyls are generally stable compounds and used as markers to determine the degree and damage due to oxidative modifications in vitro and in vivo [286, 287].

Nitrosive stress is also reported in patients with AD. Amino acids Cys, Met, Tyr and Phe are major targets for protein nitration thus causing impaired redox cell signaling, inflammatory response and protein phosphorylation [288, 289].

Protein glutathionylation resulted in disulfide bonding of thiol with protein Cys residue is important in redox signaling. Although it occurs in normal conditions, excessive glutathionylation due to oxidative stress may cause impairment in cell sense and stress responses [290, 291].

Novel advanced oxidation protein products (AOPPs) that induce proinflammatory cytokines are also being used

as reliable markers of protein oxidation in AD as well as for diabetes. These are structurally similar to AGEs and generate during oxidation with chlorinated oxidants [292–294].

Oxidation of LDL is also important mechanism in AD and elevated LDL is associated with brain A β levels. Furthermore, paraoxonase 1 (PON1) that contributes as antioxidant for LDL has also been reported significantly low in AD patients [295, 296].

DNA Oxidation

DNA is also susceptible to oxidative stress and their oxidative damage plays an important role in aging as well as AD. Mitochondrial DNA (mtDNA) and nuclear DNA oxidation has been reported in parietal cortex of AD patients. ROS attack on DNA can be result in strand break, sister chromosome exchange, DNA–protein cross-linking, translocation and formation of more than 20 oxidized base adducts [297–299]. DNA mutations due to base modifications can lead to impaired protein synthesis and functions. Several studies have reported the induction of oxidized base adducts such as 8-hydroxyguanine (8-OHG), 8-hydroxyadenine (8-OHA), 5-hydroxyuracil (5-OHU) and 5-hydroxycytosine (5-OHC). Furthermore, failure of 8-OHG repairs has also been reported in patients with AD [298, 300, 301].

RNA Oxidation

The bases in RNA are more prone to oxidation than those in DNA as they are more exposed in the single-stranded RNA molecule and are not as well protected as DNA, which uses histones as, dedicated packaging proteins. The noncoding RNAs are also involved in synapses, neuronal specification and differentiation, and regulation of dendritic spine development. So their damage due to oxidative stress contributes to development of neurodegenerative diseases, especially AD [302–304]. Nunomura et al. [303] extensively has reviewed the RNA oxidation in neurodegenerative diseases and has discussed the biological significance and cellular mechanisms that protect against RNA oxidation.

Antioxidant Therapy for AD and Diabetes

An effective antioxidant treatment can minimize the cellular damage and reduce the burden of oxidative insult. Antioxidant mechanism may involve enzymatic or non-enzymatic approach. Several antioxidant such as vitamins, glutathione, catalase, SOD, α -lipoic acid, coenzyme Q₁₀, carotenoids, flavonides, minerals (zinc, manganese, copper and selenium) and cofactors (folic acid) have been tested to reduce the after-effects of oxidative stress in AD and diabetes. Several controversial studies are available on the role of antioxidants to inhibit oxidative stress and reduced AD and diabetes

incidence. Unfortunately, till date all large-scale clinical trials have failed to demonstrate any influential benefits for AD and diabetic patients [225, 305–307].


Impaired Calcium Homeostasis

Normal intracellular calcium homeostasis is important for neuronal functions and survival. Impaired calcium homeostasis is also a consequence of mitochondrial dysfunction and intracellular calcium levels in neurons of AD patients containing NFTs were reported higher than normal. Furthermore, neurons containing NFTs susceptible to degeneration also exhibit increased levels of calcium-dependent protease protein kinase II [308]. Calcium homeostasis also regulates transglutaminase activity. Transglutaminases catalyse the formation of covalent bond between proteins that exhibit high resistance to proteolytic degradation. Higher levels of transglutaminase enzyme in prefrontal cortex of AD patients lead to cross linking of tau proteins that resulted in NFTs [309]. Calcium homeostasis abnormalities also lead to induce production of A β peptides and increased calcium levels in AD patients which are thought to be involved in enhancement of proteolytic cleavage of APP to produce A β [310]. A β formation in AD patients leads to mPTP opening also induces mitochondrial calcium uptake and can, in turn, facilitate the A β formation and aggregation [78]. Abnormal calcium homeostasis is commonly observed in diabetic patients and high calcium levels in diabetic pancreatic β -cells were found to disrupt insulin secretion [311]. Furthermore, several studies have proposed the impaired calcium homeostasis in diabetic sensory neurons and have observed a steady increase in intracellular calcium levels. Moreover, depolarization of inner mitochondrial membrane was observed in sensory neurons of STZ-diabetic rats under increased calcium flux [312, 313]. These studies confirmed that impaired calcium homeostasis is also an important reason for mitochondrial damage in diabetes and AD.

Summary of the Section

AD and diabetes both result in increased oxidative stress and impaired antioxidant systems. Although many studies suggest oxidative stress as a main cause for AD and diabetes, from the above observations it seems that oxidative stress is a consequence of these diseases and not a cause. Oxidative stress can lead to damage in several metabolic processes and once it starts can promote AD and diabetes which is summarized in Fig. 4. Although several anti-oxidative stress therapies have been proposed, are zero so far have been useful in the treatment of AD and diabetic patients. This may be due to the involvement and overlapping of several mechanisms at a time such as abnormal

Fig. 4 Schematic description of cellular events linked with mitochondrial dysfunctions especially oxidative stress in diabetes and AD. Oxidative stress not only promotes inflammation and apoptosis but also foster the synaptic damage via membrane damage and enhancing production of NFTs, and amyloid beta. Furthermore, oxidative stress also modulates protein and lipid oxidation that resulted in microvascular complications and endothelial dysfunction in AD and diabetes



glucose, proteins and lipids metabolism, defects in body immune system and inflammation, etc.

Common Therapy for Diabetes and AD

Our above discussion on shared pathology between diabetes and AD leads to the fact that drugs used for diabetes treatment should be helpful for AD patients as rendered by other authors also. Noninsulin hypoglycemic agents such as thiazolidinediones that stimulates peroxisome proliferator-activated receptor- γ (PPAR γ) and control lipid and glucose metabolism by regulating insulin-sensitive genes and reduction in insulin resistance; are being used to treat AD, and an improvement in glucose utilization and neuronal mitochondrial biogenesis is observed in AD patients [174, 314]. A recent study by Sato et al. [315] has reported reduced fasting plasma insulin levels, increased insulin sensitivity and improved cognition in mild AD patients with type 2 diabetes when treated with PPAR γ agonist pioglitazone. These results have showed that PPAR γ agonist might be useful for pre-initial and mild AD patients as well as for AD patients with diabetes [315]. Meanwhile, PPAR γ are also involved in regulation of APP-cleaving enzyme BACE1. BACE1 overexpression leads to the formation of amyloidic proteolysis of APP and generation of A β plaques [316]. On the other hand, PPAR γ agonist ginsenoside Rg1 extracted from ginseng was found to inhibit the transcription and translation of BACE1, overwhelm the BACE1 activity and eventually reduce A β plaques generation. Degeneration of oligodendroglia (oligodendrocytes) associated with neuronal support and insulation is important hallmark of AD.

Recent research found that these oligodendrocytes require intact insulin/IGF signaling to maintain their functions and survival. PPAR γ agonist induces the MAG-1 expression corresponding to oligodendrocytes and enhances DNA repairing [317]. Current studies have showed that PPAR γ agonists are also useful in reducing oxidative stress, inflammation and apoptosis in AD patients [318, 319].

Acetylcholine (ACh), a critical neurotransmitter in cognitive function is emerging possible link between AD and diabetes. Impaired insulin mechanism reduced the ACh levels and acetylcholine transferase (ChAT), an enzyme responsible for ACh production. PPAR γ agonist induced expression of ChAT that brings improvement in learning, and memory in patients with AD [8, 320].

Acetylcholinesterase inhibitors are widely used in treatment of AD. A recent has study observed that tacrine (a drug form of acetylcholinesterase inhibitors) might reduce the diabetes induced cognitive deficits in mice model by dysfunction of central cholinergic system [321].

A latest study in 2012 has showed that antineoplastic agents such as bexarotene previously used for lung and breast cancer have been found to reduce A β plaques more than 50 % within 72 h treatment [322]. ApoE gene expression that normally promotes the A β plaques generation is regulated by ligand activated nuclear receptors PPAR γ and LXRs (liver x receptors) and their binding with RXRs (retinoid X receptors) [323, 324]. Bexarotene is RXR agonist and reduced A β plaques in an ApoE manner by increasing its expression. We hypothesize the use of Bexarotene will be useful for diabetic patients as RXR agonists also functions as insulin sensitizers and can reduce hyperinsulinemia and hyperglycemia [325].

Metformin is a famous anti diabetic drug and have some contradictory results on AD pathogenesis. Chen et al. [326] observed the induced A β formation through AMPK pathway in N2a695 cell line. This induction was inhibited through AMPK pathway pharmacological inhibitor Compound C [326]. On the other hand metformin was shown to be induce activity of PP2A and reduction in tau phosphorylation in human tau transgenic mice and the authors suggest a beneficial role of metformin for AD therapy [327]. A recent study by Gupta et al. [328] showed that metformin restored acetylcholine esterase activity and inhibited GSK3 β and ERK activity, and reduced A β production in patients with AD by sensitizing neuronal insulin resistance.

ABAD as described earlier has an important role in mitochondrial membrane permeability, converting estradiol to estrone. Optimal levels of estradiol are important for neuronal survival. A current study has found that inhibition of A β using AG18051; restores the A β induced deregulation of estradiol and reduced ROS. Furthermore, when AG18051 was tested for hIAPP toxicity, partial protection was observed in type 2 diabetic pancreas. This study also pointed out a shared pathological relationship between AD and diabetes [244].

In summary, the above information lead to the assumption that AD and diabetes shared common pathology; and treatment for one disease may be useful for the other due to the overlapping of shared metabolic pathways.

Future Directions


The fact that some drugs developed to treat diabetes also have a positive effect on Alzheimer's patients supports the view that there is a common metabolic basis for both disorders. The suggestion that Alzheimer's should be referred

to “type-3 diabetes” was first proposed in 2005 but the name was not widely accepted. The resurgence of metabolic research in the form of metabolomics has coincided with renewed interest in a metabolic basis of AD and there is a question mark that whether it represents a neural form of diabetes or not.

A system biology approach to study the animal metabolism will benefit our understanding of diseases of old age, but the mouse, which is the traditional model for diabetes and Alzheimer's, is not the most amenable to systems biology research. In contrast, *Caenorhabditis elegans* may provide an excellent model for system biology of metabolism, but is not typically used as a model of these diseases. However, the use of *C. elegans* in age related diseases like diabetes and AD has gained greater importance in recent years due to shorter lifespan and ease to handle [329]. The formation of misfolded proteins, AGEs, mitochondrial disorders, generation of oxidative stress, insulin resistance and abnormal glucose metabolism are main hallmarks of diabetes and AD. *C. elegans* showed the presence of AGE formation, high oxidative stress and reduced lifespan when exposed under high glucose concentrations. In addition, high caloric diet also induces the A β formation in *C. elegans* [330]. Moreover insulin signaling pathway present in *C. elegans* is well defined and many groups studied the genes and subpathways involved in insulin signaling in *C. elegans* and found that reducing insulin signaling decreases A β toxicity [73]. In recent years, several studies used *C. elegans* as model of drug research/screening for diabetes and AD. They studied the effect of diabetic drugs on AD *C. elegans* models and vice versa. Here, we briefly describe the results of these studies.

Traditional Chinese medicine *Ginkgo biloba* is found to reduce insulin resistance by lowering glucose levels by significant increase and improved insulin concentration in

Fig. 5 Schematic illustration of the common links between diabetes and AD. These common links lead toward the hypothesis that AD may be an another type of diabetes “type-3 diabetes”



patients with diabetes. It was also capable to reduce lipid and protein oxidation in diabetic patients [331]. *G. biloba* extract increases stress resistance and extends life span of *C. elegans*. Study using *G. biloba* leaf extracts showed reduction in A β induced paralysis in transgenic *C. elegans* [332]. Furthermore, increase in lifespan was observed when metformin was used in type 2 diabetic *C. elegans* model system [333]. A recent study using *C. elegans* as model by Saharia et al. [334] found that reserpine (FDA-approved drug for AD whose mode of action and pathways activation was unknown) acts through acetylcholine mechanism. This information lead to the conclusion that *C. elegans* not only used to study the known drugs effects but may also help to elucidate the mechanisms and actions of newly developed drugs and formation of new drugs on basis of these results.

The above discussion reveals that diabetes especially T2DM and AD follow the same pathological mechanisms resulting in misfolded proteins, insulin impairment, abnormal glucose metabolism, abnormal fatty acid metabolism, mitochondrial dysfunction, and high oxidative stress. These shared metabolic profiles, and diabetes as extreme risk factor for AD lead to the assumption that AD may reflect type-3 diabetes (Fig. 5). Recent studies also have observed the mechanistic links of the other pathways such as ApoE 4 allele, decreased acetylcholine synthesis, PPAR γ activation, and activation of inflammatory genes also shares their roles in AD and diabetes pathology. Conclusively, both AD and diabetes are resulted from metabolic abnormalities, and impaired insulin mechanism is the main reason for these defects. Future studies need deep exploration of energy metabolism associated with these mechanisms as recent studies observed direct link between lower energy metabolisms, induced lifespan, and reduced aging. Although model system *C. elegans* has some drawbacks and may not perfectly stand for the human diseases pathophysiology, it may be used as powerful model for screening drugs, and studying diabetes and AD pathophysiology due to conserved insulin signaling and other metabolic pathways.

Acknowledgements I highly acknowledge Dr. Paul R Ebert (Associate Professor, School of Biological Sciences, University of Queensland, Australia), Mr. Andrew Johnston (School of Biological Sciences, The University of Queensland, Australia) and Sultan Asad (Centre of Excellence in Molecular Biology, Lahore, Pakistan) for innovating suggestions and proof reading.

Conflict of interests The author declares that he has no conflict of interest.

References

1. Citron M (2004) Strategies for disease modification in Alzheimer's disease. *Nat Rev Neurosci* 5(9):677–685. doi:10.1038/nrn1495
2. Takeda S, Sato N, Uchio-Yamada K, Sawada K, Kunieda T, Takeuchi D, Kurinami H, Shinohara M, Rakugi H, Morishita R (2010) Diabetes-accelerated memory dysfunction via cerebrovascular inflammation and A β deposition in an Alzheimer mouse model with diabetes. *Proc Natl Acad Sci U S A* 107(15):7036–7041. doi:10.1073/pnas.1000645107
3. Iqbal K, Grundke-Iqbal I (2006) Discoveries of tau, abnormally hyperphosphorylated tau and others of neurofibrillary degeneration: a personal historical perspective. *J Alzheimers Dis* 9(3 Suppl):219–242
4. Lin EH, Rutter CM, Katon W, Heckbert SR, Ciechanowski P, Oliver MM, Ludman EJ, Young BA, Williams LH, McCulloch DK, Von Korff M (2010) Depression and advanced complications of diabetes: a prospective cohort study. *Diabetes Care* 33(2):264–269. doi:10.2337/dc09-1068
5. Jakicic JM, Jaramillo SA, Balasubramanyam A, Bancroft B, Curtis JM, Mathews A, Pereira M, Regensteiner JG, Ribisl PM, Look ASG (2009) Effect of a lifestyle intervention on change in cardiorespiratory fitness in adults with type 2 diabetes: results from the Look AHEAD Study. *Int J Obes* 33(3):305–316. doi:10.1038/ijo.2008.280
6. Virkamaki A, Ueki K, Kahn CR (1999) Protein–protein interaction in insulin signaling and the molecular mechanisms of insulin resistance. *J Clin Invest* 103(7):931–943. doi:10.1172/JCI6609
7. Steen E, Terry BM, Rivera EJ, Cannon JL, Neely TR, Tavares R, Xu XJ, Wands JR, de la Monte SM (2005) Impaired insulin and insulin-like growth factor expression and signaling mechanisms in Alzheimer's disease—is this type 3 diabetes? *J Alzheimers Dis* 7(1):63–80
8. Rivera EJ, Goldin A, Fulmer N, Tavares R, Wands JR, de la Monte SM (2005) Insulin and insulin-like growth factor expression and function deteriorate with progression of Alzheimer's disease: link to brain reductions in acetylcholine. *J Alzheimers Dis* 8(3):247–268
9. Umegaki H (2012) Neurodegeneration in diabetes mellitus. *Adv Exp Med Biol* 724:258–265. doi:10.1007/978-1-4614-0653-219
10. Correia SC, Santos RX, Perry G, Zhu X, Moreira PI, Smith MA (2011) Insulin-resistant brain state: the culprit in sporadic Alzheimer's disease? *Ageing Res Rev* 10(2):264–273. doi:10.1016/j.arr.2011.01.001
11. Ristow M (2004) Neurodegenerative disorders associated with diabetes mellitus. *J Mol Med* 82(8):510–529. doi:10.1007/s00109-004-0552-1
12. Roche E, Reig JA, Campos A, Paredes B, Isaac JR, Lim S, Calne RY, Soria B (2005) Insulin-secreting cells derived from stem cells: clinical perspectives, hypes and hopes. *Transpl Immunol* 15(2):113–129. doi:10.1016/j.trim.2005.09.008
13. Watson GS, Craft S (2004) Modulation of memory by insulin and glucose: neuropsychological observations in Alzheimer's disease. *Eur J Pharmacol* 490(1–3):97–113. doi:10.1016/j.ejphar.2004.02.048
14. Sun MK, Alkon DL (2006) Links between Alzheimer's disease and diabetes. *Drugs today* 42(7):481–489. doi:10.1358/dot.2006.42.7.973588
15. Hoyer S (2004) Glucose metabolism and insulin receptor signal transduction in Alzheimer disease. *Eur J Pharmacol* 490(1–3):115–125. doi:10.1016/j.ejphar.2004.02.049
16. Ott A, Stolk RP, van Harskamp F, Pols HA, Hofman A, Breteler MM (1999) Diabetes mellitus and the risk of dementia: the Rotterdam Study. *Neurology* 53(9):1937–1942
17. Pasquier F, Boulogne A, Leys D, Fontaine P (2006) Diabetes mellitus and dementia. *Diabetes Metabol* 32(5 Pt 1):403–414
18. Sjogren M, Mielke M, Gustafson D, Zandi P, Skoog I (2006) Cholesterol and Alzheimer's disease—is there a relation? *Mech Ageing Dev* 127(2):138–147. doi:10.1016/j.mad.2005.09.020

19. Leoni V, Solomon A, Kivipelto M (2010) Links between ApoE, brain cholesterol metabolism, tau and amyloid beta-peptide in patients with cognitive impairment. *Biochem Soc Trans* 38 (4):1021–1025. doi:10.1042/BST0381021
20. Deng Y, Li B, Liu Y, Iqbal K, Grundke-Iqbal I, Gong CX (2009) Dysregulation of insulin signaling, glucose transporters, O-GlcNAcylation, and phosphorylation of tau and neurofilaments in the brain: Implication for Alzheimer's disease. *Am J Pathol* 175 (5):2089–2098. doi:10.2353/ajpath.2009.090157
21. Hoyer S, Nitsch R (1989) Cerebral excess release of neurotransmitter amino acids subsequent to reduced cerebral glucose metabolism in early-onset dementia of Alzheimer type. *J Neural Transm* 75(3):227–232
22. Li L (2007) Is Glucagon-like peptide-1, an agent treating diabetes, a new hope for Alzheimer's disease? *Neurosci Bull* 23(1):58–65
23. Perry T, Greig NH (2002) The glucagon-like peptides: a new genre in therapeutic targets for intervention in Alzheimer's disease. *J Alzheimers Dis* 4(6):487–496
24. Li L, Holscher C (2007) Common pathological processes in Alzheimer disease and type 2 diabetes: a review. *Brain Res Rev* 56(2):384–402. doi:10.1016/j.brainresrev.2007.09.001
25. Chouliaras L, Rutten BP, Kenis G, Peerbooms O, Visser PJ, Verhey F, van Os J, Steinbusch HW, van den Hove DL (2010) Epigenetic regulation in the pathophysiology of Alzheimer's disease. *Prog Neurobiol* 90(4):498–510. doi:10.1016/j.pneurobio.2010.01.002
26. Ling C, Groop L (2009) Epigenetics: a molecular link between environmental factors and type 2 diabetes. *Diabetes* 58(12):2718–2725. doi:10.2337/db09-1003
27. Hebert LE, Scherr PA, Bienias JL, Bennett DA, Evans DA (2003) Alzheimer disease in the US population: prevalence estimates using the 2000 census. *Arch Neurol* 60(8):1119–1122. doi:10.1001/archneur.60.8.1119
28. Lopez OL (2011) The growing burden of Alzheimer's disease. *Am J Manage Care* 17(Suppl 13):S339–S345
29. Selvin E, Coresh J, Brancati FL (2006) The burden and treatment of diabetes in elderly individuals in the U.S. *Diabetes Care* 29 (11):2415–2419. doi:10.2337/dc06-1058
30. Colman RJ, Anderson RM, Johnson SC, Kastman EK, Kosmatka KJ, Beasley TM, Allison DB, Cruzen C, Simmons HA, Kemnitz JW, Weindruch R (2009) Caloric restriction delays disease onset and mortality in rhesus monkeys. *Science* 325(5937):201–204. doi:10.1126/science.1173635
31. Ford D, Ions LJ, Alatawi F, Wakeling LA (2011) The potential role of epigenetic responses to diet in ageing. *Proc Nutr Soc* 70 (3):374–384. doi:10.1017/S0029665111000851
32. Cruzen C, Colman RJ (2009) Effects of caloric restriction on cardiovascular aging in non-human primates and humans. *Clin Geriatr Med* 25(4):733–743. doi:10.1016/j.cger.2009.07.001
33. Gonzalez O, Tobia C, Ebersole J, Novak MJ (2012) Caloric restriction and chronic inflammatory diseases. *Oral Dis* 18 (1):16–31. doi:10.1111/j.1601-0825.2011.01830.x
34. Mandal S, Hart N (2012) Respiratory complications of obesity. *Clin Med* 12(1):75–78
35. Reaven GM (2011) Insulin resistance: the link between obesity and cardiovascular disease. *Med Clin North Am* 95(5):875–892. doi:10.1016/j.mcna.2011.06.002
36. Anstey KJ, Cherbuin N, Budge M, Young J (2011) Body mass index in midlife and late-life as a risk factor for dementia: a meta-analysis of prospective studies. *Obes Rev* 12(5):e426–e437. doi:10.1111/j.1467-789X.2010.00825.x
37. Slomko H, Heo HJ, Einstein FH (2012) Minireview: epigenetics of obesity and diabetes in humans. *Endocrinology* 153(3):1025–1030. doi:10.1210/en.2011-1759
38. Wolf PA, Beiser A, Elias MF, Au R, Vasan RS, Seshadri S (2007) Relation of obesity to cognitive function: importance of central obesity and synergistic influence of concomitant hypertension. The Framingham Heart Study. *Curr Alzheimer Res* 4(2):111–116
39. Luchsinger JA (2008) Adiposity, hyperinsulinemia, diabetes and Alzheimer's disease: an epidemiological perspective. *Eur J Pharmacol* 585(1):119–129. doi:10.1016/j.ejphar.2008.02.048
40. Cheng D, Noble J, Tang MX, Schupf N, Mayeux R, Luchsinger JA (2011) Type 2 diabetes and late-onset Alzheimer's disease. *Dement Geriatr Cogn Disord* 31(6):424–430. doi:10.1159/000324134
41. Iglseder B (2011) Diabetes mellitus and cognitive decline. *Wien Med Wochenschr* 161(21–22):524–530. doi:10.1007/s10354-011-0003-x
42. Llibre Jde J, Valhuerdi A, Calvo M, Garcia RM, Guerra M, Laucerique T, Lopez AM, Llibre JC, Noriega L, Sanchez IY, Porto R, Arencibia F, Marcheco B, Moreno C (2011) Dementia and other chronic diseases in older adults in Havana and Matanzas: the 10/66 study in Cuba. *MEDICC rev* 13(4):30–37
43. Tripathi M, Vibha D, Gupta P, Bhatia R, Srivastava MV, Vivekanandhan S, Bhushan Singh M, Prasad K, Dergalust S, Mendez MF (2012) Risk factors of dementia in North India: a case–control study. *Aging Ment Health* 16(2):228–235. doi:10.1080/13607863.2011.583632
44. Accardi G, Caruso C, Colonna-Romano G, Camarda C, Monastero R, Candore G (2012) Can Alzheimer disease be a form of type 3 diabetes? *Rejuvenation Res* 15(2):217–221. doi:10.1089/rej.2011.1289
45. de la Monte SM, Wands JR (2008) Alzheimer's disease is type 3 diabetes-evidence reviewed. *J Diabetes Sci Tech* 2(6):1101–1113
46. Soto C (2003) Unfolding the role of protein misfolding in neurodegenerative diseases. *Nat Rev Neurosci* 4(1):49–60. doi:10.1038/nrn1007
47. Reitz C (2012) Alzheimer's disease and the amyloid cascade hypothesis: a critical review. *Int J Alzheimers Dis* 2012:369808. doi:10.1155/2012/369808
48. Milojevic J, Melacini G (2011) Stoichiometry and affinity of the human serum albumin-Alzheimer's Aβ peptide interactions. *Biophys J* 100(1):183–192. doi:10.1016/j.bpj.2010.11.037
49. Shankar GM, Li S, Mehta TH, Garcia-Munoz A, Shepardson NE, Smith I, Brett FM, Farrell MA, Rowan MJ, Lemere CA, Regan CM, Walsh DM, Sabatini BL, Selkoe DJ (2008) Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat Med* 14(8):837–842. doi:10.1038/nm1782
50. DeToma AS, Salamekh S, Ramamoorthy A, Lim MH (2012) Misfolded proteins in Alzheimer's disease and type II diabetes. *Chem Soc Rev* 41(2):608–621. doi:10.1039/c1cs15112f
51. Haass C, Selkoe DJ (2007) Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* 8(2):101–112. doi:10.1038/nrm2101
52. Correia SC, Santos RX, Carvalho C, Cardoso S, Candeias E, Santos MS, Oliveira CR, Moreira PI (2012) Insulin signaling, glucose metabolism and mitochondria: major players in Alzheimer's disease and diabetes interrelation. *Brain Res* 1441:64–78. doi:10.1016/j.brainres.2011.12.063
53. Luca S, Yau WM, Leapman R, Tycko R (2007) Peptide conformation and supramolecular organization in amylin fibrils: constraints from solid-state NMR. *Biochemistry* 46(47):13505–13522. doi:10.1021/bi701427q
54. Woods SC, Lutz TA, Geary N, Langhans W (2006) Pancreatic signals controlling food intake; insulin, glucagon and amylin. *Philos Trans R Soc B Biol Sci* 361(1471):1219–1235. doi:10.1098/rstb.2006.1858

55. Haataja L, Gurlo T, Huang CJ, Butler PC (2008) Islet amyloid in type 2 diabetes, and the toxic oligomer hypothesis. *Endocr Rev* 29(3):303–316. doi:10.1210/er.2007-0037
56. Zraika S, Hull RL, Udayasankar J, Aston-Mourney K, Subramanian SL, Kisilevsky R, Szarek WA, Kahn SE (2009) Oxidative stress is induced by islet amyloid formation and time-dependently mediates amyloid-induced beta cell apoptosis. *Diabetologia* 52(4):626–635. doi:10.1007/s00125-008-1255-x
57. Janson J, Laedtke T, Parisi JE, O'Brien P, Petersen RC, Butler PC (2004) Increased risk of type 2 diabetes in Alzheimer disease. *Diabetes* 53(2):474–481
58. Fu W, Ruangkittisakul A, MacTavish D, Shi JY, Ballanyi K, Jhamandas JH (2012) Amyloid beta (A β) peptide directly activates amylin-3 receptor subtype by triggering multiple intracellular signaling pathways. *J Biol Chem* 287(22):18820–18830. doi:10.1074/jbc.M111.331181
59. Scherzer-Attali R, Shaltiel-Karyo R, Adalist YH, Segal D, Gazit E (2012) Generic inhibition of amyloidogenic proteins by two naphthoquinone–tryptophan hybrid molecules. *Proteins* 80(8):1962–1973. doi:10.1002/prot.24080
60. Gupta R, Kapoor N, Raleigh DP, Sakmar TP (2012) Nucleobindin 1 caps human islet amyloid polypeptide protofibrils to prevent amyloid fibril formation. *J Mol Biol* 421(2–3):378–389. doi:10.1016/j.jmb.2012.04.017
61. Sesti G (2006) Pathophysiology of insulin resistance. *Best Pract Res Clin Endocrinol Metab* 20(4):665–679. doi:10.1016/j.beem.2006.09.007
62. Yang XL, Chan JC (2012) Diabetes, insulin and cancer risk. *World J Diabetes* 3(4):60–64. doi:10.4239/wjd.v3.i4.60
63. Banks WA (2004) The source of cerebral insulin. *Eur J Pharmacol* 490(1–3):5–12. doi:10.1016/j.ejphar.2004.02.040
64. Bosco D, Fava A, Plastino M, Montalcini T, Pujia A (2011) Possible implications of insulin resistance and glucose metabolism in Alzheimer's disease pathogenesis. *J Cell Mol Med* 15(9):1807–1821. doi:10.1111/j.1582-4934.2011.01318.x
65. Wang Q, Jin T (2009) The role of insulin signaling in the development of beta-cell dysfunction and diabetes. *Islets* 1(2):95–101. doi:10.4161/isl.1.2.9263
66. Thorens B (2012) Sensing of glucose in the brain. *Handb Exp Pharmacol* 209:277–294. doi:10.1007/978-3-642-24716-3_12
67. Pansuria M, Xi H, Li L, Yang XF, Wang H (2012) Insulin resistance, metabolic stress, and atherosclerosis. *Front Biosci* 4:916–931
68. Collino M, Aragno M, Castiglia S, Tomasinelli C, Thiemermann C, Boccuzzi G, Fantozzi R (2009) Insulin reduces cerebral ischemia/reperfusion injury in the hippocampus of diabetic rats: a role for glycogen synthase kinase-3 β . *Diabetes* 58(1):235–242. doi:10.2337/db08-0691
69. Talbot K, Wang HY, Kazi H, Han LY, Bakshi KP, Stucky A, Fuino RL, Kawaguchi KR, Samoyedny AJ, Wilson RS, Arvanitakis Z, Schneider JA, Wolf BA, Bennett DA, Trojanowski JQ, Arnold SE (2012) Demonstrated brain insulin resistance in Alzheimer's disease patients is associated with IGF-1 resistance, IRS-1 dysregulation, and cognitive decline. *J Clin Invest* 122(4):1316–1338. doi:10.1172/JCI59903
70. Park SA (2011) A common pathogenic mechanism linking type-2 diabetes and Alzheimer's disease: evidence from animal models. *J Clin Neurol* 7(1):10–18. doi:10.3988/jcn.2011.7.1.10
71. Li ZG, Zhang W, Sima AA (2007) Alzheimer-like changes in rat models of spontaneous diabetes. *Diabetes* 56(7):1817–1824. doi:10.2337/db07-0171
72. Shingo AS, Kanabayashi T, Murase T, Kito S (2012) Cognitive decline in STZ-3 V rats is largely due to dysfunctional insulin signalling through the dentate gyrus. *Behav Brain Res* 229(2):378–383. doi:10.1016/j.bbr.2012.01.034
73. Zemva J, Schubert M (2011) Central insulin and insulin-like growth factor-1 signaling: implications for diabetes associated dementia. *Curr Diabetes Rev* 7(5):356–366
74. Zhao WQ, De Felice FG, Fernandez S, Chen H, Lambert MP, Quon MJ, Krafft GA, Klein WL (2008) Amyloid beta oligomers induce impairment of neuronal insulin receptors. *FASEB J* 22(1):246–260. doi:10.1096/fj.06-7703com
75. Gasparini L, Gouras GK, Wang R, Gross RS, Beal MF, Greengard P, Xu H (2001) Stimulation of beta-amyloid precursor protein trafficking by insulin reduces intraneuronal beta-amyloid and requires mitogen-activated protein kinase signaling. *J Neurosci* 21(8):2561–2570
76. Leissring MA, Farris W, Chang AY, Walsh DM, Wu X, Sun X, Frosch MP, Selkoe DJ (2003) Enhanced proteolysis of beta-amyloid in APP transgenic mice prevents plaque formation, secondary pathology, and premature death. *Neuron* 40(6):1087–1093
77. Leal MC, Fernandez Gamba A, Morelli L, Castano EM (2009) Cerebral proteolysis of amyloid- β peptide: relevance of insulin-degrading enzyme in Alzheimer's disease. *Medicina* 69(4):466–472
78. Parihar MS, Brewer GJ (2010) Amyloid- β as a modulator of synaptic plasticity. *J Alzheimers Dis* 22(3):741–763. doi:10.3233/JAD-2010-101020
79. Brezar V, Carel JC, Boitard C, Mallone R (2011) Beyond the hormone: insulin as an autoimmune target in type 1 diabetes. *Endocr Rev* 32(5):623–669. doi:10.1210/er.2011-0010
80. Al-Qatati A, Winter PW, Wolf-Ringwall AL, Chatterjee PB, Van Orden AK, Crans DC, Roess DA, Barisas BG (2012) Insulin receptors and downstream substrates associate with membrane microdomains after treatment with insulin or chromium(III) picolinate. *Cell Biochem Biophys* 62(3):441–450. doi:10.1007/s12013-011-9326-x
81. Denley A, Carroll JM, Brierley GV, Cosgrove L, Wallace J, Forbes B, Roberts CT Jr (2007) Differential activation of insulin receptor substrates 1 and 2 by insulin-like growth factor-activated insulin receptors. *Mol Cell Biol* 27(10):3569–3577. doi:10.1128/MCB.01447-06
82. Hulmi JJ, Silvennoinen M, Lehti M, Kivela R, Kainulainen H (2012) Altered REDD1, myostatin, and Akt/mTOR/FoxO/MAPK signaling in streptozotocin-induced diabetic muscle atrophy. *Am J Physiol Endocrinol Metab* 302(3):E307–E315. doi:10.1152/ajpendo.00398.2011
83. Machado-Neto JA, Favaro P, Lazarini M, Costa FF, Olalla Saad ST, Traina F (2011) Knockdown of insulin receptor substrate 1 reduces proliferation and downregulates Akt/mTOR and MAPK pathways in K562 cells. *Biochim Biophys Acta* 1813(8):1404–1411. doi:10.1016/j.bbamcr.2011.04.002
84. Duarte AI, Moreira PI, Oliveira CR (2012) Insulin in central nervous system: more than just a peripheral hormone. *J Aging Res* 2012:384017. doi:10.1155/2012/384017
85. Schultze SM, Jensen J, Hemmings BA, Tschopp O, Niessen M (2011) Promiscuous affairs of PKB/AKT isoforms in metabolism. *Arch Physiol Biochem* 117(2):70–77. doi:10.3109/13813455.2010.539236
86. Bayascas JR (2008) Dissecting the role of the 3-phosphoinositide-dependent protein kinase-1 (PDK1) signalling pathways. *Cell Cycle* 19:2978–2982
87. Maeda H, Rajesh KG, Maeda H, Suzuki R, Sasaguri S (2004) Epidermal growth factor and insulin inhibit cell death in pancreatic beta cells by activation of PI3-kinase/AKT signaling pathway under oxidative stress. *Transplant Proc* 36(4):1163–1165. doi:10.1016/j.transproceed.2004.04.018
88. Ohtsubo K, Chen MZ, Olefsky JM, Marth JD (2011) Pathway to diabetes through attenuation of pancreatic beta cell glycosylation

- and glucose transport. *Nat Med* 17(9):1067–1075. doi:10.1038/nm.2414
89. Bornfeldt KE, Tabas I (2011) Insulin resistance, hyperglycemia, and atherosclerosis. *Cell Metab* 14(5):575–585. doi:10.1016/j.cmet.2011.07.015
 90. Henriksen EJ (2010) Dysregulation of glycogen synthase kinase-3 in skeletal muscle and the etiology of insulin resistance and type 2 diabetes. *Curr Diabetes Rev* 6(5):285–293
 91. Saini V (2010) Molecular mechanisms of insulin resistance in type 2 diabetes mellitus. *World J Diabetes* 1(3):68–75. doi:10.4239/wjd.v1.i3.68
 92. Szeman B, Nagy G, Varga T, Veres-Szekely A, Sasvari M, Fitala D, Szollosi A, Katonai R, Kotyuk E, Somogyi A (2012) Changes in cognitive function in patients with diabetes mellitus. *Orv Hetil* 153(9):323–329. doi:10.1556/OH.2012.29319
 93. Freude S, Hettich MM, Schumann C, Stohr O, Koch L, Kohler C, Udelhoven M, Leiser U, Muller M, Kubota N, Kadowaki T, Krone W, Schroder H, Bruning JC, Schubert M (2009) Neuronal IGF-1 resistance reduces Abeta accumulation and protects against premature death in a model of Alzheimer's disease. *FASEB J* 23(10):3315–3324. doi:10.1096/fj.09-132043
 94. Killick R, Scales G, Leroy K, Causevic M, Hooper C, Irvine EE, Choudhury AI, Drinkwater L, Kerr F, Al-Qassab H, Stephenson J, Yilmaz Z, Giese KP, Brion JP, Withers DJ, Lovestone S (2009) Deletion of Irs2 reduces amyloid deposition and rescues behavioural deficits in APP transgenic mice. *Biochem Biophys Res Commun* 386(1):257–262. doi:10.1016/j.bbrc.2009.06.032
 95. Dong X, Park S, Lin X, Copps K, Yi X, White MF (2006) Irs1 and Irs2 signaling is essential for hepatic glucose homeostasis and systemic growth. *J Clin Invest* 116(1):101–114. doi:10.1172/JCI25735
 96. Shah OJ, Wang Z, Hunter T (2004) Inappropriate activation of the TSC/Rheb/mTOR/S6K cassette induces IRS1/2 depletion, insulin resistance, and cell survival deficiencies. *Curr Biol* 14(18):1650–1656. doi:10.1016/j.cub.2004.08.026
 97. Draznin B (2006) Molecular mechanisms of insulin resistance: serine phosphorylation of insulin receptor substrate-1 and increased expression of p85alpha: the two sides of a coin. *Diabetes* 55(8):2392–2397. doi:10.2337/db06-0391
 98. White MF (2002) IRS proteins and the common path to diabetes. *Am J Physiol Endocrinol Metab* 283(3):E413–E422. doi:10.1152/ajpendo.00514.2001
 99. Samuel VT, Shulman GI (2012) Mechanisms for insulin resistance: common threads and missing links. *Cell* 148(5):852–871. doi:10.1016/j.cell.2012.02.017
 100. Zhang H, Dellsperger KC, Zhang C (2012) The link between metabolic abnormalities and endothelial dysfunction in type 2 diabetes: an update. *Basic Res Cardiol* 107(1):237. doi:10.1007/s00395-011-0237-1
 101. Dey D, Basu D, Roy SS, Bandyopadhyay A, Bhattacharya S (2006) Involvement of novel PKC isoforms in FFA induced defects in insulin signaling. *Mol Cell Endocrinol* 246(1–2):60–64. doi:10.1016/j.mce.2005.12.014
 102. Timmers S, de Vogel-van den Bosch J, Hesselink MK, van Beurden D, Schaart G, Ferraz MJ, Losen M, Martinez-Martinez P, De Baets MH, Aerts JM, Schrauwen P (2011) Paradoxical increase in TAG and DAG content parallel the insulin sensitizing effect of unilateral DGAT1 overexpression in rat skeletal muscle. *PLoS One* 6(1):e14503. doi:10.1371/journal.pone.0014503
 103. Liu Y, Liu F, Grundke-Iqbal I, Iqbal K, Gong CX (2009) Brain glucose transporters, O-GlcNAcylation and phosphorylation of tau in diabetes and Alzheimer's disease. *J Neurochem* 111(1):242–249. doi:10.1111/j.1471-4159.2009.06320.x
 104. Liu Y, Liu F, Grundke-Iqbal I, Iqbal K, Gong CX (2011) Deficient brain insulin signalling pathway in Alzheimer's disease and diabetes. *J Pathol* 225(1):54–62. doi:10.1002/path.2912
 105. Jolivald CG, Lee CA, Beiswenger KK, Smith JL, Orlov M, Torrance MA, Masliah E (2008) Defective insulin signaling pathway and increased glycogen synthase kinase-3 activity in the brain of diabetic mice: parallels with Alzheimer's disease and correction by insulin. *J Neurosci Res* 86(15):3265–3274. doi:10.1002/jnr.21787
 106. Waite K, Eickholt BJ (2010) The neurodevelopmental implications of PI3K signaling. *Curr Top Microbiol Immunol* 346:245–265. doi:10.1007/82_2010_82
 107. Peyrou M, Bourgoin L, Foti M (2010) PTEN in liver diseases and cancer. *World J Gastroenterol* 16(37):4627–4633
 108. Rosivatz E, Matthews JG, McDonald NQ, Mulet X, Ho KK, Lossi N, Schmid AC, Mirabelli M, Pomeranz KM, Erneux C, Lam EW, Vilar R, Woscholski R (2006) A small molecule inhibitor for phosphatase and tensin homologue deleted on chromosome 10 (PTEN). *ACS Chem Biol* 1(12):780–790. doi:10.1021/cb600352f
 109. Barbour LA, Shao J, Qiao L, Leitner W, Anderson M, Friedman JE, Draznin B (2004) Human placental growth hormone increases expression of the p85 regulatory unit of phosphatidylinositol 3-kinase and triggers severe insulin resistance in skeletal muscle. *Endocrinology* 145(3):1144–1150. doi:10.1210/en.2003-1297
 110. Kirwan JP, Varastehpour A, Jing M, Presley L, Shao J, Friedman JE, Catalano PM (2004) Reversal of insulin resistance postpartum is linked to enhanced skeletal muscle insulin signaling. *J Clin Endocrinol Metab* 89(9):4678–4684. doi:10.1210/jc.2004-0749
 111. Ueki K, Fruman DA, Brachmann SM, Tseng YH, Cantley LC, Kahn CR (2002) Molecular balance between the regulatory and catalytic subunits of phosphoinositide 3-kinase regulates cell signaling and survival. *Mol Cell Biol* 22(3):965–977
 112. Liao Y, Hung MC (2010) Physiological regulation of Akt activity and stability. *Am J Transl Res* 2(1):19–42
 113. Tan SX, Ng Y, Meoli CC, Kumar A, Khoo PS, Fazakerley DJ, Junutula JR, Vali S, James DE, Stockli J (2012) Amplification and demultiplexing in insulin-regulated Akt protein kinase pathway in adipocytes. *J Biol Chem* 287(9):6128–6138. doi:10.1074/jbc.M111.318238
 114. George S, Rochford JJ, Wolfrum C, Gray SL, Schinner S, Wilson JC, Soos MA, Murgatroyd PR, Williams RM, Acerini CL, Dunger DB, Barford D, Umpleby AM, Wareham NJ, Davies HA, Schafer AJ, Stoffel M, O'Rahilly S, Barroso I (2004) A family with severe insulin resistance and diabetes due to a mutation in AKT2. *Science* 304(5675):1325–1328. doi:10.1126/science.1096706
 115. Xu N, Lao Y, Zhang Y, Gillespie DA (2012) Akt: a double-edged sword in cell proliferation and genome stability. *J Oncol* 2012:951724. doi:10.1155/2012/951724
 116. Sakamaki J, Daitoku H, Ueno K, Hagiwara A, Yamagata K, Fukamizu A (2011) Arginine methylation of BCL-2 antagonist of cell death (BAD) counteracts its phosphorylation and inactivation by Akt. *Proc Natl Acad Sci U S A* 108(15):6085–6090. doi:10.1073/pnas.1015328108
 117. Wolfrum C, Besser D, Luca E, Stoffel M (2003) Insulin regulates the activity of forkhead transcription factor Hnf-3beta/Foxa-2 by Akt-mediated phosphorylation and nuclear/cytosolic localization. *Proc Natl Acad Sci U S A* 100(20):11624–11629. doi:10.1073/pnas.1931483100
 118. Zhao X, Gan L, Pan H, Kan D, Majeski M, Adam SA, Unterman TG (2004) Multiple elements regulate nuclear/cytoplasmic shuttling of FOXO1: characterization of phosphorylation- and 14-3-3-dependent and -independent mechanisms. *Biochem J* 378(Pt 3):839–849. doi:10.1042/BJ20031450
 119. Manning BD, Cantley LC (2007) AKT/PKB signaling: navigating downstream. *Cell* 129(7):1261–1274. doi:10.1016/j.cell.2007.06.009

120. Sarbassov DD, Guertin DA, Ali SM, Sabatini DM (2005) Phosphorylation and regulation of Akt/PKB by the rictor-mTOR complex. *Science* 307(5712):1098–1101. doi:[10.1126/science.1106148](https://doi.org/10.1126/science.1106148)
121. Cheng Z, Tseng Y, White MF (2010) Insulin signaling meets mitochondria in metabolism. *Trends Endocrinol Metab* 21(10):589–598. doi:[10.1016/j.tem.2010.06.005](https://doi.org/10.1016/j.tem.2010.06.005)
122. Fraenkel M, Ketzinel-Gilad M, Ariav Y, Pappo O, Karaca M, Castel J, Berthault MF, Magnan C, Cerasi E, Kaiser N, Leibowitz G (2008) mTOR inhibition by rapamycin prevents beta-cell adaptation to hyperglycemia and exacerbates the metabolic state in type 2 diabetes. *Diabetes* 57(4):945–957. doi:[10.2337/db07-0922](https://doi.org/10.2337/db07-0922)
123. 2Le Bacquer O, Petroulakis E, Paglialunga S, Poulin F, Richard D, Cianflone K, Sonenberg N (2007) Elevated sensitivity to diet-induced obesity and insulin resistance in mice lacking 4E-BP1 and 4E-BP2. *J Clin Invest* 117(2):387–396. doi:[10.1172/JCI29528](https://doi.org/10.1172/JCI29528)
124. Ma XM, Blenis J (2009) Molecular mechanisms of mTOR-mediated translational control. *Nat Rev Mol Cell Biol* 10(5):307–318. doi:[10.1038/nrm2672](https://doi.org/10.1038/nrm2672)
125. Mariappan MM (2012) Signaling mechanisms in the regulation of renal matrix metabolism in diabetes. *Exp Diabetes Res* 2012:749812. doi:[10.1155/2012/749812](https://doi.org/10.1155/2012/749812)
126. Proud CG (2007) Signalling to translation: how signal transduction pathways control the protein synthetic machinery. *Biochem J* 403(2):217–234. doi:[10.1042/BJ20070024](https://doi.org/10.1042/BJ20070024)
127. Kaidanovich-Beilin O, Woodgett JR (2011) GSK-3: functional Insights from Cell Biology and Animal Models. *Front Mol Neurosci* 4:40. doi:[10.3389/fnmol.2011.00040](https://doi.org/10.3389/fnmol.2011.00040)
128. Medina M, Garrido JJ, Wandosell FG (2011) Modulation of GSK-3 as a therapeutic strategy on tau pathologies. *Front Mol Neurosci* 4:24. doi:[10.3389/fnmol.2011.00024](https://doi.org/10.3389/fnmol.2011.00024)
129. Poljsak B (2011) Strategies for reducing or preventing the generation of oxidative stress. *Oxidative Med Cell Longev* 2011:194586. doi:[10.1155/2011/194586](https://doi.org/10.1155/2011/194586)
130. Schubert M, Brazil DP, Burks DJ, Kushner JA, Ye J, Flint CL, Farhang-Fallah J, Dikkes P, Warot XM, Rio C, Corfas G, White MF (2003) Insulin receptor substrate-2 deficiency impairs brain growth and promotes tau phosphorylation. *J Neurosci* 23(18):7084–7092
131. Maher PA, Schubert DR (2009) Metabolic links between diabetes and Alzheimer's disease. *Expert Rev Neurother* 9(5):617–630. doi:[10.1586/ern.09.18](https://doi.org/10.1586/ern.09.18)
132. Besnard A, Galan-Rodriguez B, Vanhoutte P, Caboche J (2011) Elk-1 a transcription factor with multiple facets in the brain. *Front Neurosci* 5:35. doi:[10.3389/fnins.2011.00035](https://doi.org/10.3389/fnins.2011.00035)
133. Jeanneteau F, Deinhardt K (2011) Fine-tuning MAPK signaling in the brain: the role of MKP-1. *Commun Integr Biol* 4(3):281–283. doi:[10.4161/cib.4.3.14766](https://doi.org/10.4161/cib.4.3.14766)
134. Sharma SK, Carew TJ (2004) The roles of MAPK cascades in synaptic plasticity and memory in Aplysia: facilitatory effects and inhibitory constraints. *Learn Mem* 11(4):373–378. doi:[10.1101/lm.81104](https://doi.org/10.1101/lm.81104)
135. Kim SH, Smith CJ, Van Eldik LJ (2004) Importance of MAPK pathways for microglial pro-inflammatory cytokine IL-1 beta production. *Neurobiol Aging* 25(4):431–439. doi:[10.1016/S0197-4580\(03\)00126-X](https://doi.org/10.1016/S0197-4580(03)00126-X)
136. Hu JF, Niu F, Ning N, Duan WZ, Chu SF, Xue W, Yuan YH, Chen NH, Zhang JT (2012) Activation of ERK1/2-CREB pathway during potentiating synaptic transmission of (-)-clausenamide in rat dentate gyrus. *J Asian Nat Prod Res* 14(3):256–262. doi:[10.1080/10286020.2011.650885](https://doi.org/10.1080/10286020.2011.650885)
137. Chico LK, Van Eldik LJ, Watterson DM (2009) Targeting protein kinases in central nervous system disorders. *Nat Rev Drug Discov* 8(11):892–909. doi:[10.1038/nrd2999](https://doi.org/10.1038/nrd2999)
138. Munoz L, Ammit AJ (2010) Targeting p38 MAPK pathway for the treatment of Alzheimer's disease. *Neuropharmacology* 58(3):561–568. doi:[10.1016/j.neuropharm.2009.11.010](https://doi.org/10.1016/j.neuropharm.2009.11.010)
139. Kelleher I, Garwood C, Hanger DP, Anderton BH, Noble W (2007) Kinase activities increase during the development of tauopathy in htau mice. *J Neurochem* 103(6):2256–2267. doi:[10.1111/j.1471-4159.2007.04930.x](https://doi.org/10.1111/j.1471-4159.2007.04930.x)
140. Baker LD, Cross DJ, Minoshima S, Belongia D, Watson GS, Craft S (2011) Insulin resistance and Alzheimer-like reductions in regional cerebral glucose metabolism for cognitively normal adults with prediabetes or early type 2 diabetes. *Arch Neurol* 68(1):51–57. doi:[10.1001/archneurol.2010.225](https://doi.org/10.1001/archneurol.2010.225)
141. Zhao WQ, Townsend M (2009) Insulin resistance and amyloidogenesis as common molecular foundation for type 2 diabetes and Alzheimer's disease. *Biochim Biophys Acta* 1792(5):482–496. doi:[10.1016/j.bbadis.2008.10.014](https://doi.org/10.1016/j.bbadis.2008.10.014)
142. Scheepers A, Joost HG, Schurmann A (2004) The glucose transporter families SGLT and GLUT: molecular basis of normal and aberrant function. *J Parenter Enter Nutr* 28(5):364–371
143. Schulingkamp RJ, Pagano TC, Hung D, Raffa RB (2000) Insulin receptors and insulin action in the brain: review and clinical implications. *Neurosci Biobehav Rev* 24(8):855–872
144. Gomez Ravetti M, Rosso OA, Berretta R, Moscato P (2010) Uncovering molecular biomarkers that correlate cognitive decline with the changes of hippocampus' gene expression profiles in Alzheimer's disease. *PLoS One* 5(4):e10153. doi:[10.1371/journal.pone.0010153](https://doi.org/10.1371/journal.pone.0010153)
145. Hooijmans CR, Graven C, Dederen PJ, Tanila H, van Groen T, Kiliaan AJ (2007) Amyloid beta deposition is related to decreased glucose transporter-1 levels and hippocampal atrophy in brains of aged APP/PS1 mice. *Brain Res* 1181:93–103. doi:[10.1016/j.brainres.2007.08.063](https://doi.org/10.1016/j.brainres.2007.08.063)
146. Liu Y, Liu F, Iqbal K, Grundke-Iqbal I, Gong CX (2008) Decreased glucose transporters correlate to abnormal hyperphosphorylation of tau in Alzheimer disease. *FEBS Lett* 582(2):359–364. doi:[10.1016/j.febslet.2007.12.035](https://doi.org/10.1016/j.febslet.2007.12.035)
147. Cavado E, Frisoni GB (2011) The dynamic marker hypothesis of Alzheimer's disease and its implications for clinical imaging. *Q J Nucl Med Mol Imaging* 55(3):237–249
148. Garrido GE, Furuie SS, Buchpiguel CA, Bottino CM, Almeida OP, Cid CG, Camargo CH, Castro CC, Glabus MF, Busatto GF (2002) Relation between medial temporal atrophy and functional brain activity during memory processing in Alzheimer's disease: a combined MRI and SPECT study. *J Neurol Neurosurg Psychiatry* 73(5):508–516
149. Small GW, Ercoli LM, Silverman DH, Huang SC, Komo S, Bookheimer SY, Lavretsky H, Miller K, Siddarth P, Rasgon NL, Mazziotta JC, Saxena S, Wu HM, Mega MS, Cummings JL, Saunders AM, Pericak-Vance MA, Roses AD, Barrio JR, Phelps ME (2000) Cerebral metabolic and cognitive decline in persons at genetic risk for Alzheimer's disease. *Proc Natl Acad Sci U S A* 97(11):6037–6042. doi:[10.1073/pnas.090106797](https://doi.org/10.1073/pnas.090106797)
150. Reiman EM, Chen K, Alexander GE, Caselli RJ, Bandy D, Osborne D, Saunders AM, Hardy J (2004) Functional brain abnormalities in young adults at genetic risk for late-onset Alzheimer's dementia. *Proc Natl Acad Sci U S A* 101(1):284–289. doi:[10.1073/pnas.2635903100](https://doi.org/10.1073/pnas.2635903100)
151. Valla J, Gonzalez-Lima F, Reiman EM (2008) FDG autoradiography reveals developmental and pathological effects of mutant amyloid in PDAPP transgenic mice. *Int J Dev Neurosci* 26(3–4):253–258. doi:[10.1016/j.jdevneu.2008.02.003](https://doi.org/10.1016/j.jdevneu.2008.02.003)
152. Bubber P, Haroutunian V, Fisch G, Blass JP, Gibson GE (2005) Mitochondrial abnormalities in Alzheimer brain: mechanistic implications. *Ann Neurol* 57(5):695–703. doi:[10.1002/ana.20474](https://doi.org/10.1002/ana.20474)
153. Huang HM, Ou HC, Xu H, Chen HL, Fowler C, Gibson GE (2003) Inhibition of alpha-ketoglutarate dehydrogenase complex

- promotes cytochrome *c* release from mitochondria, caspase-3 activation, and necrotic cell death. *J Neurosci Res* 74(2):309–317. doi:10.1002/jnr.10756
154. Moreira PI, Santos MS, Oliveira CR (2007) Alzheimer's disease: a lesson from mitochondrial dysfunction. *Antioxid Redox Signal* 9(10):1621–1630. doi:10.1089/ars.2007.1703
 155. Brownlee M (2001) Biochemistry and molecular cell biology of diabetic complications. *Nature* 414(6865):813–820. doi:10.1038/414813a
 156. Yamagishi S (2011) Role of advanced glycation end products (AGEs) in osteoporosis in diabetes. *Curr Drug Targets* 12(14):2096–2102
 157. Goh SY, Cooper ME (2008) Clinical review: the role of advanced glycation end products in progression and complications of diabetes. *J Clin Endocrinol Metab* 93(4):1143–1152. doi:10.1210/jc.2007-1817
 158. Takuma K, Fang F, Zhang W, Yan S, Fukuzaki E, Du H, Sosunov A, McKhann G, Funatsu Y, Nakamichi N, Nagai T, Mizoguchi H, Ibi D, Hori O, Ogawa S, Stern DM, Yamada K, Yan SS (2009) RAGE-mediated signaling contributes to intraneuronal transport of amyloid-beta and neuronal dysfunction. *Proc Natl Acad Sci U S A* 106(47):20021–20026. doi:10.1073/pnas.0905686106
 159. Biessels GJ, van der Heide LP, Kamal A, Bley RL, Gispen WH (2002) Ageing and diabetes: implications for brain function. *Eur J Pharmacol* 441(1–2):1–14
 160. Yan SD, Bierhaus A, Nawroth PP, Stern DM (2009) RAGE and Alzheimer's disease: a progression factor for amyloid-beta-induced cellular perturbation? *J Alzheimers Dis* 16(4):833–843. doi:10.3233/JAD-2009-1030
 161. Yan SD, Chen X, Fu J, Chen M, Zhu H, Roher A, Slattey T, Zhao L, Nagashima M, Morser J, Migheli A, Nawroth P, Stern D, Schmidt AM (1996) RAGE and amyloid-beta peptide neurotoxicity in Alzheimer's disease. *Nature* 382(6593):685–691. doi:10.1038/382685a0
 162. Deane R, Du Yan S, Subramanyam RK, LaRue B, Jovanovic S, Hogg E, Welch D, Manness L, Lin C, Yu J, Zhu H, Ghiso J, Frangione B, Stern A, Schmidt AM, Armstrong DL, Arnold B, Liliensiek B, Nawroth P, Hofman F, Kindy M, Stern D, Zlokovic B (2003) RAGE mediates amyloid-beta peptide transport across the blood–brain barrier and accumulation in brain. *Nat Med* 9(7):907–913. doi:10.1038/nm890
 163. Sato N, Takeda S, Uchio-Yamada K, Ueda H, Fujisawa T, Rakugi H, Morishita R (2011) Role of insulin signaling in the interaction between Alzheimer disease and diabetes mellitus: a missing link to therapeutic potential. *Cur Aging Sci* 4(2):118–127
 164. Deane R, Singh I, Sagare AP, Bell RD, Ross NT, LaRue B, Love R, Perry S, Paquette N, Deane RJ, Thiagarajan M, Zarcone T, Fritz G, Friedman AE, Miller BL, Zlokovic BV (2012) A multimodal RAGE-specific inhibitor reduces amyloid beta-mediated brain disorder in a mouse model of Alzheimer disease. *J Clin Invest* 122(4):1377–1392. doi:10.1172/JCI58642
 165. Xiong F, Leonov S, Howard AC, Xiong S, Zhang B, Mei L, McNeil P, Simon S, Xiong WC (2011) Receptor for advanced glycation end products (RAGE) prevents endothelial cell membrane resealing and regulates F-actin remodeling in a beta-catenin-dependent manner. *J Biol Chem* 286(40):35061–35070. doi:10.1074/jbc.M111.261073
 166. Liu F, Iqbal K, Grundke-Iqbal I, Hart GW, Gong CX (2004) O-GlcNAcylation regulates phosphorylation of tau: a mechanism involved in Alzheimer's disease. *Proc Natl Acad Sci U S A* 101(29):10804–10809. doi:10.1073/pnas.0400348101
 167. Robertson LA, Moya KL, Breen KC (2004) The potential role of tau protein O-glycosylation in Alzheimer's disease. *J Alzheimers Dis* 6(5):489–495
 168. Majumdar G, Wright J, Markowitz P, Martinez-Hernandez A, Raghov R, Solomon SS (2004) Insulin stimulates and diabetes inhibits O-linked N-acetylglucosamine transferase and O-glycosylation of Sp1. *Diabetes* 53(12):3184–3192
 169. Liu F, Shi J, Tanimukai H, Gu J, Gu J, Grundke-Iqbal I, Iqbal K, Gong CX (2009) Reduced O-GlcNAcylation links lower brain glucose metabolism and tau pathology in Alzheimer's disease. *Brain* 132(Pt 7):1820–1832. doi:10.1093/brain/awp099
 170. Yuzwa SA, Shan X, Macauley MS, Clark T, Skorobogatko Y, Vosseller K, Vocadlo DJ (2012) Increasing O-GlcNAc slows neurodegeneration and stabilizes tau against aggregation. *Nat Chem Biol* 8(4):393–399. doi:10.1038/nchembio.797
 171. Akter K, Lanza EA, Martin SA, Myronyuk N, Rua M, Raffa RB (2011) Diabetes mellitus and Alzheimer's disease: shared pathology and treatment? *Br J Clin Pharmacol* 71(3):365–376. doi:10.1111/j.1365-2125.2010.03830.x
 172. Ramasamy R, Yan SF, Schmidt AM (2011) Receptor for AGE (RAGE): signaling mechanisms in the pathogenesis of diabetes and its complications. *Ann N Y Acad Sci* 1243:88–102. doi:10.1111/j.1749-6632.2011.06320.x
 173. Srikanth V, Maczurek A, Phan T, Steele M, Westcott B, Juskiw D, Munch G (2011) Advanced glycation endproducts and their receptor RAGE in Alzheimer's disease. *Neurobiol Aging* 32(5):763–777. doi:10.1016/j.neurobiolaging.2009.04.016
 174. Kroner Z (2009) The relationship between Alzheimer's disease and diabetes: type 3 diabetes? *Altern Med Rev* 14(4):373–379
 175. Di Paolo G, Kim TW (2011) Linking lipids to Alzheimer's disease: cholesterol and beyond. *Nat Rev Neurosci* 12(5):284–296. doi:10.1038/nrn3012
 176. Taskinen MR (2002) Diabetic dyslipidemia. *Atheroscler Suppl* 3(1):47–51
 177. Roland I, De Leval X, Evrard B, Pirotte B, Dogne JM, Delattre L (2004) Modulation of the arachidonic cascade with omega3 fatty acids or analogues: potential therapeutic benefits. *Mini Rev Med Chem* 4(6):659–668
 178. Eckel RH, Grundy SM, Zimmet PZ (2005) The metabolic syndrome. *Lancet* 365(9468):1415–1428. doi:10.1016/S0140-6736(05)66378-7
 179. Hayashi H (2011) Lipid metabolism and glial lipoproteins in the central nervous system. *Biol Pharm Bull* 34(4):453–461
 180. Jenkins AJ, Best JD, Klein RL, Lyons TJ (2004) Lipoproteins, glycoxidation and diabetic angiopathy. *Diabetes Metabol Res Rev* 20(5):349–368. doi:10.1002/dmrr.491
 181. Lutjohann D (2006) Cholesterol metabolism in the brain: importance of 24S-hydroxylation. *Acta Neurol Scand Suppl* 185:33–42. doi:10.1111/j.1600-0404.2006.00683.x
 182. Hamel FG, Upward JL, Bennett RG (2003) In vitro inhibition of insulin-degrading enzyme by long-chain fatty acids and their coenzyme A thioesters. *Endocrinology* 144(6):2404–2408
 183. Hillered L, Chan PH (1988) Role of arachidonic acid and other free fatty acids in mitochondrial dysfunction in brain ischemia. *J Neurosci Res* 20(4):451–456. doi:10.1002/jnr.490200407
 184. Firuzi O, Zhuo J, Chinnici CM, Wisniewski T, Pratico D (2008) 5-Lipoxygenase gene disruption reduces amyloid-beta pathology in a mouse model of Alzheimer's disease. *FASEB J* 22(4):1169–1178. doi:10.1096/fj.07-9131.com
 185. Sanchez-Mejia RO, Newman JW, Toh S, Yu GQ, Zhou Y, Halabisky B, Cisse M, Scearce-Levie K, Cheng IH, Gan L, Palop JJ, Bonventre JV, Mucke L (2008) Phospholipase A2 reduction ameliorates cognitive deficits in a mouse model of Alzheimer's disease. *Nat Neurosci* 11(11):1311–1318. doi:10.1038/nn.2213
 186. Cherny VV, Henderson LM, Xu W, Thomas LL, DeCoursey TE (2001) Activation of NADPH oxidase-related proton and electron currents in human eosinophils by arachidonic acid. *J Physiol* 535(Pt 3):783–794
 187. Zhu D, Lai Y, Shelat PB, Hu C, Sun GY, Lee JC (2006) Phospholipases A2 mediate amyloid-beta peptide-induced

- mitochondrial dysfunction. *J Neurosci Off J Soc Neurosci* 26 (43):11111–11119. doi:[10.1523/JNEUROSCI.3505-06.2006](https://doi.org/10.1523/JNEUROSCI.3505-06.2006)
188. Penzo D, Petronilli V, Angelin A, Cusan C, Colonna R, Scorrano L, Pagano F, Prato M, Di Lisa F, Bernardi P (2004) Arachidonic acid released by phospholipase A(2) activation triggers Ca(2+)-dependent apoptosis through the mitochondrial pathway. *J Biol Chem* 279(24):25219–25225. doi:[10.1074/jbc.M310381200](https://doi.org/10.1074/jbc.M310381200)
 189. Tan S, Sagara Y, Liu Y, Maher P, Schubert D (1998) The regulation of reactive oxygen species production during programmed cell death. *J Cell Biol* 141(6):1423–1432
 190. Puccio S, Chu J, Pratico D (2011) Involvement of 5-lipoxygenase in the corticosteroid-dependent amyloid beta formation: in vitro and in vivo evidence. *PLoS One* 6(1):e15163. doi:[10.1371/journal.pone.0015163](https://doi.org/10.1371/journal.pone.0015163)
 191. Claria J (2003) Cyclooxygenase-2 biology. *Curr Pharm Des* 9 (27):2177–2190
 192. Sun GY, Horrocks LA, Farooqui AA (2007) The roles of NADPH oxidase and phospholipases A2 in oxidative and inflammatory responses in neurodegenerative diseases. *J Neurochem* 103(1):1–16. doi:[10.1111/j.1471-1459.2007.04670.x](https://doi.org/10.1111/j.1471-1459.2007.04670.x)
 193. Sims-Robinson C, Kim B, Rosko A, Feldman EL (2010) How does diabetes accelerate Alzheimer disease pathology? *Nat Rev Neurol* 6(10):551–559. doi:[10.1038/nrneurol.2010.130](https://doi.org/10.1038/nrneurol.2010.130)
 194. Yaffe K (2007) Metabolic syndrome and cognitive decline. *Curr Alzheimer Res* 4(2):123–126
 195. Dietschy JM, Turley SD (2001) Cholesterol metabolism in the brain. *Curr Opin Lipidol* 12(2):105–112
 196. Bjorkhem I (2006) Crossing the barrier: oxysterols as cholesterol transporters and metabolic modulators in the brain. *J Intern Med* 260(6):493–508. doi:[10.1111/j.1365-2796.2006.01725.x](https://doi.org/10.1111/j.1365-2796.2006.01725.x)
 197. Chang TY, Chang CC, Ohgami N, Yamauchi Y (2006) Cholesterol sensing, trafficking, and esterification. *Annu Rev Cell Dev Biol* 22:129–157. doi:[10.1146/annurev.cellbio.22.010305.104656](https://doi.org/10.1146/annurev.cellbio.22.010305.104656)
 198. Heverin M, Meaney S, Lutjohann D, Diczfalussy U, Wahren J, Bjorkhem I (2005) Crossing the barrier: net flux of 27-hydroxycholesterol into the human brain. *J Lipid Res* 46 (5):1047–1052. doi:[10.1194/jlr.M500024-JLR200](https://doi.org/10.1194/jlr.M500024-JLR200)
 199. Ladu MJ, Reardon C, Van Eldik L, Fagan AM, Bu G, Holtzman D, Getz GS (2000) Lipoproteins in the central nervous system. *Ann N Y Acad Sci* 903:167–175
 200. Bhattacharyya R, Kovacs DM (2010) ACAT inhibition and amyloid beta reduction. *Biochim Biophys Acta* 1801(8):960–965. doi:[10.1016/j.bbalip.2010.04.003](https://doi.org/10.1016/j.bbalip.2010.04.003)
 201. Hutter-Paier B, Huttunen HJ, Puglielli L, Eckman CB, Kim DY, Hofmeister A, Moir RD, Domnitz SB, Frosch MP, Windisch M, Kovacs DM (2004) The ACAT inhibitor CP-113,818 markedly reduces amyloid pathology in a mouse model of Alzheimer's disease. *Neuron* 44(2):227–238. doi:[10.1016/j.neuron.2004.08.043](https://doi.org/10.1016/j.neuron.2004.08.043)
 202. Simons M, Keller P, De Strooper B, Beyreuther K, Dotti CG, Simons K (1998) Cholesterol depletion inhibits the generation of beta-amyloid in hippocampal neurons. *Proc Natl Acad Sci U S A* 95(11):6460–6464
 203. Vetrivel KS, Thinakaran G (2010) Membrane rafts in Alzheimer's disease beta-amyloid production. *Biochim Biophys Acta* 1801 (8):860–867. doi:[10.1016/j.bbalip.2010.03.007](https://doi.org/10.1016/j.bbalip.2010.03.007)
 204. Jiang Q, Lee CY, Mandrekar S, Wilkinson B, Cramer P, Zelcer N, Mann K, Lamb B, Willson TM, Collins JL, Richardson JC, Smith JD, Comery TA, Riddell D, Holtzman DM, Tontonoz P, Landreth GE (2008) ApoE promotes the proteolytic degradation of Abeta. *Neuron* 58(5):681–693. doi:[10.1016/j.neuron.2008.04.010](https://doi.org/10.1016/j.neuron.2008.04.010)
 205. Peila R, Rodriguez BL, Launer LJ, Honolulu-Asia Aging S (2002) Type 2 diabetes, APOE gene, and the risk for dementia and related pathologies: the Honolulu-Asia Aging Study. *Diabetes* 51(4):1256–1262
 206. Belinson H, Lev D, Masliah E, Michaelson DM (2008) Activation of the amyloid cascade in apolipoprotein E4 transgenic mice induces lysosomal activation and neurodegeneration resulting in marked cognitive deficits. *J Neurosci Off J Soc Neurosci* 28(18):4690–4701. doi:[10.1523/JNEUROSCI.5633-07.2008](https://doi.org/10.1523/JNEUROSCI.5633-07.2008)
 207. Brecht WJ, Harris FM, Chang S, Tesseur I, Yu GQ, Xu Q, Dee Fish J, Wyss-Coray T, Buttini M, Mucke L, Mahley RW, Huang Y (2004) Neuron-specific apolipoprotein e4 proteolysis is associated with increased tau phosphorylation in brains of transgenic mice. *J Neurosci Off J Soc Neurosci* 24(10):2527–2534. doi:[10.1523/JNEUROSCI.4315-03.2004](https://doi.org/10.1523/JNEUROSCI.4315-03.2004)
 208. Luchsinger JA (2012) Type 2 diabetes and cognitive impairment: linking mechanisms. *J Alzheimers Dis* 30:S185–S198. doi:[10.3233/JAD-2012-111433](https://doi.org/10.3233/JAD-2012-111433)
 209. Chaudhary R, Likidilid A, Peerapatdit T, Tresukosol D, Srisuma S, Ratanamaneechat S, Sriratanasathavorn C (2012) Apolipoprotein E gene polymorphism: effects on plasma lipids and risk of type 2 diabetes and coronary artery disease. *Cardiovasc Diabetol* 11:36. doi:[10.1186/1475-2840-11-36](https://doi.org/10.1186/1475-2840-11-36)
 210. Hayden MR, Clee SM, Brooks-Wilson A, Genest J Jr, Attie A, Kastelein JJ (2000) Cholesterol efflux regulatory protein, Tangier disease and familial high-density lipoprotein deficiency. *Curr Opin Lipidol* 11(2):117–122
 211. Koldamova R, Fitz NF, Lefterov I (2010) The role of ATP-binding cassette transporter A1 in Alzheimer's disease and neurodegeneration. *Biochim Biophys Acta* 1801(8):824–830. doi:[10.1016/j.bbalip.2010.02.010](https://doi.org/10.1016/j.bbalip.2010.02.010)
 212. Hirsch-Reinshagen V, Zhou S, Burgess BL, Bernier L, McIsaac SA, Chan JY, Tansley GH, Cohn JS, Hayden MR, Wellington CL (2004) Deficiency of ABCA1 impairs apolipoprotein E metabolism in brain. *J Biol Chem* 279(39):41197–41207. doi:[10.1074/jbc.M407962200](https://doi.org/10.1074/jbc.M407962200)
 213. Wang XT, Li J, Liu L, Hu N, Jin S, Liu C, Mei D, Liu XD (2012) Tissue cholesterol content alterations in streptozotocin-induced diabetic rats. *Acta Pharmacol Sin* 33(7):909–917. doi:[10.1038/aps.2012.50](https://doi.org/10.1038/aps.2012.50)
 214. Lei P, Wu WH, Li RW, Ma JW, Yu YP, Cui W, Zhao YF, Li YM (2008) Prevention and promotion effects of apolipoprotein E4 on amylin aggregation. *Biochem Biophys Res Commun* 368 (2):414–418. doi:[10.1016/j.bbrc.2008.01.103](https://doi.org/10.1016/j.bbrc.2008.01.103)
 215. Mattson MP, Gleichmann M, Cheng A (2008) Mitochondria in neuroplasticity and neurological disorders. *Neuron* 60(5):748–766. doi:[10.1016/j.neuron.2008.10.010](https://doi.org/10.1016/j.neuron.2008.10.010)
 216. Moreira PI, Duarte AI, Santos MS, Rego AC, Oliveira CR (2009) An integrative view of the role of oxidative stress, mitochondria and insulin in Alzheimer's disease. *J Alzheimers Dis* 16(4):741–761. doi:[10.3233/JAD-2009-0972](https://doi.org/10.3233/JAD-2009-0972)
 217. Chen H, Chan DC (2009) Mitochondrial dynamics—fusion, fission, movement, and mitophagy—in neurodegenerative diseases. *Hum Mol Genet* 18(R2):R169–R176. doi:[10.1093/hmg/ddp326](https://doi.org/10.1093/hmg/ddp326)
 218. Perez-Matute P, Zulet MA, Martinez JA (2009) Reactive species and diabetes: counteracting oxidative stress to improve health. *Curr Opin Pharmacol* 9(6):771–779. doi:[10.1016/j.coph.2009.08.005](https://doi.org/10.1016/j.coph.2009.08.005)
 219. Jomova K, Valko M (2011) Advances in metal-induced oxidative stress and human disease. *Toxicology* 283(2–3):65–87. doi:[10.1016/j.tox.2011.03.001](https://doi.org/10.1016/j.tox.2011.03.001)
 220. Malinouski M, Zhou Y, Belousov VV, Hatfield DL, Gladyshev VN (2011) Hydrogen peroxide probes directed to different cellular compartments. *PLoS One* 6(1):e14564. doi:[10.1371/journal.pone.0014564](https://doi.org/10.1371/journal.pone.0014564)
 221. Jiang D, Li X, Williams R, Patel S, Men L, Wang Y, Zhou F (2009) Ternary complexes of iron, amyloid-beta, and nitrotriacetic acid: binding affinities, redox properties, and relevance to iron-induced oxidative stress in Alzheimer's disease. *Biochemistry* 48(33):7939–7947. doi:[10.1021/bi900907a](https://doi.org/10.1021/bi900907a)

222. Lipinski B (2011) Hydroxyl radical and its scavengers in health and disease. *Oxidative Med Cell Longev* 2011:809696. doi:10.1155/2011/809696
223. Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J (2007) Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol* 39 (1):44–84. doi:10.1016/j.biocel.2006.07.001
224. Flora SJ (2007) Role of free radicals and antioxidants in health and disease. *Cell Mol Biol* 53(1):1–2
225. Bonda DJ, Wang X, Perry G, Nunomura A, Tabaton M, Zhu X, Smith MA (2010) Oxidative stress in Alzheimer disease: a possibility for prevention. *Neuropharmacology* 59(4–5):290–294. doi:10.1016/j.neuropharm.2010.04.005
226. Hampel H, Prvulovic D, Teipel S, Jessen F, Luckhaus C, Frolich L, Riepe MW, Dodel R, Leyhe T, Bertram L, Hoffmann W, Faltraco F, German Task Force on Alzheimer's D (2011) The future of Alzheimer's disease: the next 10 years. *Prog Neurobiol* 95(4):718–728. doi:10.1016/j.pneurobio.2011.11.008
227. Pratico D, Delanty N (2000) Oxidative injury in diseases of the central nervous system: focus on Alzheimer's disease. *Am J Med* 109(7):577–585
228. Obel LF, Muller MS, Walls AB, Sickmann HM, Bak LK, Waagepetersen HS, Schousboe A (2012) Brain glycogen-new perspectives on its metabolic function and regulation at the subcellular level. *Front Neuroener* 4:3. doi:10.3389/fnene.2012.00003
229. Readnower RD, Sauerbeck AD, Sullivan PG (2011) Mitochondria, amyloid beta, and Alzheimer's disease. *Int J Alzheimers Dis* 2011:104545. doi:10.4061/2011/104545
230. Sultana R, Butterfield DA (2011) Identification of the oxidative stress proteome in the brain. *Free Radic Biol Med* 50(4):487–494. doi:10.1016/j.freeradbiomed.2010.11.021
231. Ceriello A, Mercuri F, Quagliaro L, Assaloni R, Motz E, Tonutti L, Taboga C (2001) Detection of nitrotyrosine in the diabetic plasma: evidence of oxidative stress. *Diabetologia* 44(7):834–838. doi:10.1007/s001250100529
232. Douglas G, Bendall JK, Crabtree MJ, Tatham AL, Carter EE, Hale AB, Channon KM (2012) Endothelial-specific Nox2 overexpression increases vascular superoxide and macrophage recruitment in ApoE(–)/(–) mice. *Cardiovasc Res* 94(1):20–29. doi:10.1093/cvr/cvs026
233. Rolo AP, Palmeira CM (2006) Diabetes and mitochondrial function: role of hyperglycemia and oxidative stress. *Toxicol Appl Pharmacol* 212(2):167–178. doi:10.1016/j.taap.2006.01.003
234. Zhu X, Perry G, Smith MA, Wang X (2012) Abnormal mitochondrial dynamics in the pathogenesis of Alzheimer's disease. *J Alzheimers Dis*. doi:10.3233/JAD-2012-129005
235. Andreyev AY, Kushnareva YE, Starkov AA (2005) Mitochondrial metabolism of reactive oxygen species. *Biochem Biokhim* 70 (2):200–214
236. Brand MD (2010) The sites and topology of mitochondrial superoxide production. *Exp Gerontol* 45(7–8):466–472. doi:10.1016/j.exger.2010.01.003
237. Murphy MP (2009) How mitochondria produce reactive oxygen species. *Biochem J* 417(1):1–13. doi:10.1042/BJ20081386
238. Selivanov VA, Votyakova TV, Pivtoraiko VN, Zeak J, Sukhomlin T, Trucco M, Roca J, Cascante M (2011) Reactive oxygen species production by forward and reverse electron fluxes in the mitochondrial respiratory chain. *PLoS Comput Biol* 7(3):e1001115. doi:10.1371/journal.pcbi.1001115
239. Ten VS, Starkov A (2012) Hypoxic–ischemic injury in the developing brain: the role of reactive oxygen species originating in mitochondria. *Neurol Res Int* 2012:542976. doi:10.1155/2012/542976
240. Spuch C, Ortolano S, Navarro C (2012) New insights in the amyloid-beta interaction with mitochondria. *J Aging Res* 2012:324968. doi:10.1155/2012/324968
241. Drechsel DA, Estevez AG, Barbeito L, Beckman JS (2012) Nitric oxide-mediated oxidative damage and the progressive demise of motor neurons in ALS. *Neurotox Res*. doi:10.1007/s12640-012-9322-y
242. Smith MA, Zhu X, Tabaton M, Liu G, McKeel DW Jr, Cohen ML, Wang X, Siedlak SL, Dwyer BE, Hayashi T, Nakamura M, Nunomura A, Perry G (2010) Increased iron and free radical generation in preclinical Alzheimer disease and mild cognitive impairment. *J Alzheimers Dis* 19(1):363–372. doi:10.3233/JAD-2010-1239
243. Di Domenico F, Coccia R, Butterfield DA, Perluigi M (2011) Circulating biomarkers of protein oxidation for Alzheimer disease: expectations within limits. *Biochim Biophys Acta* 1814 (12):1785–1795. doi:10.1016/j.bbapap.2011.10.001
244. Lim YA, Grimm A, Giese M, Mensah-Nyagan AG, Villafranca JE, Ittner LM, Eckert A, Gotz J (2011) Inhibition of the mitochondrial enzyme ABAD restores the amyloid-beta-mediated deregulation of estradiol. *PLoS One* 6(12):e28887. doi:10.1371/journal.pone.0028887
245. Lustbader JW, Cirilli M, Lin C, Xu HW, Takuma K, Wang N, Caspersen C, Chen X, Pollak S, Chaney M, Trinchese F, Liu S, Gunn-Moore F, Lue LF, Walker DG, Kuppusamy P, Zewier ZL, Arancio O, Stern D, Yan SS, Wu H (2004) ABAD directly links Abeta to mitochondrial toxicity in Alzheimer's disease. *Science* 304(5669):448–452. doi:10.1126/science.1091230
246. Janciauskiene S, Ahren B (2000) Fibrillar islet amyloid polypeptide differentially affects oxidative mechanisms and lipoprotein uptake in correlation with cytotoxicity in two insulin-producing cell lines. *Biochem Biophys Res Commun* 267(2):619–625. doi:10.1006/bbrc.1999.1989
247. Konarkowska B, Aitken JF, Kistler J, Zhang S, Cooper GJ (2005) Thiol reducing compounds prevent human amylin-evoked cytotoxicity. *FEBS J* 272(19):4949–4959. doi:10.1111/j.1742-4658.2005.04903.x
248. Folli F, Guzzi V, Perego L, Coletta DK, Finzi G, Placidi C, La Rosa S, Capella C, Socci C, Lauro D, Tripathy D, Jenkinson C, Paroni R, Orsenigo E, Cighetti G, Gregorini L, Staudacher C, Secchi A, Bachi A, Brownlee M, Fiorina P (2010) Proteomics reveals novel oxidative and glycolytic mechanisms in type 1 diabetic patients' skin which are normalized by kidney–pancreas transplantation. *PLoS One* 5(3):e9923. doi:10.1371/journal.pone.0009923
249. Gersten RA, Gretebeck LM, Hildick-Smith G, Sandwick RK (2010) Maillard reaction of ribose 5-phosphate generates superoxide and glycation products for bovine heart cytochrome c reduction. *Carbohydr Res* 345(17):2499–2506. doi:10.1016/j.carres.2010.09.013
250. Kassab A, Piwowar A (2012) Cell oxidant stress delivery and cell dysfunction onset in type 2 diabetes. *Biochimie* 94(9):1837–1848. doi:10.1016/j.biochi.2012.01.020
251. Munusamy S, MacMillan-Crow LA (2009) Mitochondrial superoxide plays a crucial role in the development of mitochondrial dysfunction during high glucose exposure in rat renal proximal tubular cells. *Free Radic Biol Med* 46(8):1149–1157. doi:10.1016/j.freeradbiomed.2009.01.022
252. Pacher P, Szabo C (2005) Role of poly(ADP-ribose) polymerase-1 activation in the pathogenesis of diabetic complications: endothelial dysfunction, as a common underlying theme. *Antioxid Redox Signal* 7(11–12):1568–1580. doi:10.1089/ars.2005.7.1568
253. Perry SW, Norman JP, Barbieri J, Brown EB, Gelbard HA (2011) Mitochondrial membrane potential probes and the proton gradient: a practical usage guide. *Biotechniques* 50(2):98–115. doi:10.2144/000113610
254. Vincent AM, Olzmann JA, Brownlee M, Sivitz WI, Russell JW (2004) Uncoupling proteins prevent glucose-induced neuronal oxidative stress and programmed cell death. *Diabetes* 53(3):726–734

255. Mastrocola R, Restivo F, Vercellinatto I, Danni O, Brignardello E, Aragno M, Boccuzzi G (2005) Oxidative and nitrosative stress in brain mitochondria of diabetic rats. *J Endocrinol* 187(1):37–44. doi:10.1677/joe.1.06269
256. Cardoso S, Carvalho C, Santos R, Correia S, Santos MS, Seica R, Oliveira CR, Moreira PI (2011) Impact of STZ-induced hyperglycemia and insulin-induced hypoglycemia in plasma amino acids and cortical synaptosomal neurotransmitters. *Synapse* 65(6):457–466. doi:10.1002/syn.20863
257. Raza H, Prabu SK, John A, Avadhani NG (2011) Impaired mitochondrial respiratory functions and oxidative stress in streptozotocin-induced diabetic rats. *Int J Mol Sci* 12(5):3133–3147. doi:10.3390/ijms12053133
258. Giacco F, Brownlee M (2010) Oxidative stress and diabetic complications. *Circ Res* 107(9):1058–1070. doi:10.1161/CIRCRESAHA.110.223545
259. Rahmadi A, Steiner N, Munch G (2011) Advanced glycation endproducts as gerontotoxins and biomarkers for carbonyl-based degenerative processes in Alzheimer's disease. *Clin Chem Lab Med* 49(3):385–391. doi:10.1515/CCLM.2011.079
260. Thomas MC (2011) Advanced glycation end products. *Contrib Nephrol* 170:66–74. doi:10.1159/000324945
261. Higaki Y, Mikami T, Fujii N, Hirshman MF, Koyama K, Seino T, Tanaka K, Goodyear LJ (2008) Oxidative stress stimulates skeletal muscle glucose uptake through a phosphatidylinositol 3-kinase-dependent pathway. *Am J Physiol Endocrinol Metab* 294(5):E889–E897. doi:10.1152/ajpendo.00150.2007
262. Pitocco D, Zaccardi F, Di Stasio E, Romitelli F, Santini SA, Zuppi C, Ghirlanda G (2010) Oxidative stress, nitric oxide, and diabetes. *Rev Diabet Stud* 7(1):15–25. doi:10.1900/RDS.2010.7.15
263. Pessler D, Rudich A, Bashan N (2001) Oxidative stress impairs nuclear proteins binding to the insulin responsive element in the GLUT4 promoter. *Diabetologia* 44(12):2156–2164. doi:10.1007/s001250100024
264. Powell DJ, Hajdich E, Kular G, Hundal HS (2003) Ceramide disables 3-phosphoinositide binding to the pleckstrin homology domain of protein kinase B (PKB)/Akt by a PKCzeta-dependent mechanism. *Mol Cell Biol* 23(21):7794–7808
265. Corsinovi L, Biasi F, Poli G, Leonarduzzi G, Isaia G (2011) Dietary lipids and their oxidized products in Alzheimer's disease. *Mol Nutr Food Res* 55(Suppl 2):S161–S172. doi:10.1002/mnfr.201100208
266. Yang H, Jin X, Kei Lam CW, Yan SK (2011) Oxidative stress and diabetes mellitus. *Clin Chem Lab Med CCLM/FESCC* 49(11):1773–1782. doi:10.1515/CCLM.2011.250
267. Butterfield DA, Perluigi M, Reed T, Muharib T, Hughes CP, Robinson RA, Sultana R (2012) Redox proteomics in selected neurodegenerative disorders: from its infancy to future applications. *Antioxid Redox Signal*. doi:10.1089/ars.2011.4109
268. Otaegui-Arrazola A, Menendez-Carreno M, Ansorena D, Astiasaran I (2010) Oxysterols: a world to explore. *Food Chem Toxicol Int J Publ Br Ind Biol Res Assoc* 48(12):3289–3303. doi:10.1016/j.fct.2010.09.023
269. Sottero B, Gamba P, Gargiulo S, Leonarduzzi G, Poli G (2009) Cholesterol oxidation products and disease: an emerging topic of interest in medicinal chemistry. *Curr Med Chem* 16(6):685–705
270. Famer D, Meaney S, Mousavi M, Nordberg A, Bjorkhem I, Crisby M (2007) Regulation of alpha- and beta-secretase activity by oxysterols: cerebrosterol stimulates processing of APP via the alpha-secretase pathway. *Biochem Biophys Res Commun* 359(1):46–50. doi:10.1016/j.bbrc.2007.05.033
271. Chen CT, Green JT, Orr SK, Bazinet RP (2008) Regulation of brain polyunsaturated fatty acid uptake and turnover. *Prostaglandins Leukot Essent Fat Acids* 79(3–5):85–91. doi:10.1016/j.plefa.2008.09.003
272. Florent-Bechard S, Desbene C, Garcia P, Allouche A, Youssef I, Escanyé MC, Koziel V, Hanse M, Malaplate-Armand C, Stenger C, Kriem B, Yen-Potin FT, Olivier JL, Pillot T, Oster T (2009) The essential role of lipids in Alzheimer's disease. *Biochimie* 91(6):804–809. doi:10.1016/j.biochi.2009.03.004
273. Schneider C, Porter NA, Brash AR (2008) Routes to 4-hydroxynonenal: fundamental issues in the mechanisms of lipid peroxidation. *J Biol Chem* 283(23):15539–15543. doi:10.1074/jbc.R800001200
274. Spickett CM, Wiswedel I, Siems W, Zarkovic K, Zarkovic N (2010) Advances in methods for the determination of biologically relevant lipid peroxidation products. *Free Radic Res* 44(10):1172–1202. doi:10.3109/10715762.2010.498476
275. Markesbery WR, Kryscio RJ, Lovell MA, Morrow JD (2005) Lipid peroxidation is an early event in the brain in amnesic mild cognitive impairment. *Ann Neurol* 58(5):730–735. doi:10.1002/ana.20629
276. Mufson EJ, Leurgans S (2010) Inability of plasma and urine F2A-isoprostane levels to differentiate mild cognitive impairment from Alzheimer's disease. *Neurodegener Dis* 7(1–3):139–142. doi:10.1159/000289224
277. Greilberger J, Fuchs D, Leblhuber F, Greilberger M, Wintersteiger R, Tafeit E (2010) Carbonyl proteins as a clinical marker in Alzheimer's disease and its relation to tryptophan degradation and immune activation. *Clin Lab* 56(9–10):441–448
278. Sun L, Shen W, Liu Z, Guan S, Liu J, Ding S (2010) Endurance exercise causes mitochondrial and oxidative stress in rat liver: effects of a combination of mitochondrial targeting nutrients. *Life Sci* 86(1–2):39–44. doi:10.1016/j.lfs.2009.11.003
279. Bradley MA, Markesbery WR, Lovell MA (2010) Increased levels of 4-hydroxynonenal and acrolein in the brain in preclinical Alzheimer disease. *Free Radic Biol Med* 48(12):1570–1576. doi:10.1016/j.freeradbiomed.2010.02.016
280. Dang TN, Arseneault M, Murthy V, Ramassamy C (2010) Potential role of acrolein in neurodegeneration and in Alzheimer's disease. *Curr Mol Pharmacol* 3(2):66–78
281. Singh M, Dang TN, Arseneault M, Ramassamy C (2010) Role of by-products of lipid oxidation in Alzheimer's disease brain: a focus on acrolein. *J Alzheimers Dis* 21(3):741–756. doi:10.3233/JAD-2010-100405
282. Bacot S, Bernoud-Hubac N, Baddas N, Chantegrel B, Deshayes C, Doutheau A, Lagarde M, Guichardant M (2003) Covalent binding of hydroxy-alkenals 4-HDDE, 4-HHE, and 4-HNE to ethanolamine phospholipid subclasses. *J Lipid Res* 44(5):917–926. doi:10.1194/jlr.M200450-JLR200
283. Butterfield DA, Bader Lange ML, Sultana R (2010) Involvements of the lipid peroxidation product, HNE, in the pathogenesis and progression of Alzheimer's disease. *Biochim Biophys Acta* 1801(8):924–929. doi:10.1016/j.bbalip.2010.02.005
284. Siegel SJ, Bieschke J, Powers ET, Kelly JW (2007) The oxidative stress metabolite 4-hydroxynonenal promotes Alzheimer protofibril formation. *Biochemistry* 46(6):1503–1510. doi:10.1021/bi061853s
285. Trachootham D, Lu W, Ogasawara MA, Nilsa RD, Huang P (2008) Redox regulation of cell survival. *Antioxid Redox Signal* 10(8):1343–1374. doi:10.1089/ars.2007.1957
286. Stadtman ER, Berlett BS (1998) Reactive oxygen-mediated protein oxidation in aging and disease. *Drug Metab Rev* 30(2):225–243. doi:10.3109/03602539808996310
287. Widmer R, Ziaja I, Grune T (2006) Protein oxidation and degradation during aging: role in skin aging and neurodegeneration. *Free Radic Res* 40(12):1259–1268. doi:10.1080/10715760600911154
288. Schopfer FJ, Baker PR, Freeman BA (2003) NO-dependent protein nitration: a cell signaling event or an oxidative inflammatory

- response? *Trends Biochem Sci* 28(12):646–654. doi:10.1016/j.tibs.2003.10.006
289. Sultana R, Poon HF, Cai J, Pierce WM, Merchant M, Klein JB, Markesbery WR, Butterfield DA (2006) Identification of nitrated proteins in Alzheimer's disease brain using a redox proteomics approach. *Neurobiol Dis* 22(1):76–87. doi:10.1016/j.nbd.2005.10.004
 290. Dalle-Donne I, Rossi R, Giustarini D, Colombo R, Milzani A (2007) S-glutathionylation in protein redox regulation. *Free Radic Biol Med* 43(6):883–898. doi:10.1016/j.freeradbiomed.2007.06.014
 291. Giustarini D, Rossi R, Milzani A, Colombo R, Dalle-Donne I (2004) S-Glutathionylation: from redox regulation of protein functions to human diseases. *J Cell Mol Med* 8(2):201–212
 292. Komosinska-Vashev K, Olczyk P, Wysz-Szczotka K, Kuznik-Trocha K, Klimek K, Olczyk K (2011) Age- and gender-related alteration in plasma advanced oxidation protein products (AOPP) and glycosaminoglycan (GAG) concentrations in physiological ageing. *Clin Chem Lab Med*, doi:10.1515/CCLM.2011.789
 293. Kovacic P, Somanathan R (2011) Cell signaling and receptors in toxicity of advanced glycation end products (AGEs): alpha-dicarbonyls, radicals, oxidative stress and antioxidants. *J Recept Signal Transduct Res* 31(5):332–339. doi:10.3109/10799893.2011.607171
 294. Kalousova M, Skrha J, Zima T (2002) Advanced glycation end-products and advanced oxidation protein products in patients with diabetes mellitus. *Physiol Res Acad Sci Bohemoslovaca* 51(6):597–604
 295. Aldred S, Bennett S, Mecocci P (2010) Increased low-density lipoprotein oxidation, but not total plasma protein oxidation, in Alzheimer's disease. *Clin Biochem* 43(3):267–271. doi:10.1016/j.clinbiochem.2009.08.021
 296. Wehr H, Bednarska-Makaruk M, Graban A, Lipczynska-Lojkowska W, Rodo M, Bochynska A, Ryglewicz D (2009) Paraoxonase activity and dementia. *J Neurol Sci* 283(1–2):107–108. doi:10.1016/j.jns.2009.02.317
 297. Malinski T (2007) Nitric oxide and nitroxidative stress in Alzheimer's disease. *J Alzheimers Dis* 11(2):207–218
 298. Moreira PI, Nunomura A, Nakamura M, Takeda A, Shenk JC, Aliev G, Smith MA, Perry G (2008) Nucleic acid oxidation in Alzheimer disease. *Free Radic Biol Med* 44(8):1493–1505. doi:10.1016/j.freeradbiomed.2008.01.002
 299. Wang J, Markesbery WR, Lovell MA (2006) Increased oxidative damage in nuclear and mitochondrial DNA in mild cognitive impairment. *J Neurochem* 96(3):825–832. doi:10.1111/j.1471-4159.2005.03615.x
 300. Lovell MA, Markesbery WR (2008) Oxidatively modified RNA in mild cognitive impairment. *Neurobiol Dis* 29(2):169–175. doi:10.1016/j.nbd.2007.07.030
 301. Morocz M, Kalman J, Juhasz A, Sinko I, McGlynn AP, Downes CS, Janka Z, Rasko I (2002) Elevated levels of oxidative DNA damage in lymphocytes from patients with Alzheimer's disease. *Neurobiol Aging* 23(1):47–53
 302. Lovell MA, Markesbery WR (2007) Oxidative DNA damage in mild cognitive impairment and late-stage Alzheimer's disease. *Nucleic Acids Res* 35(22):7497–7504. doi:10.1093/nar/gkm821
 303. Nunomura A, Hofer T, Moreira PI, Castellani RJ, Smith MA, Perry G (2009) RNA oxidation in Alzheimer disease and related neurodegenerative disorders. *Acta Neuropathol* 118(1):151–166. doi:10.1007/s00401-009-0508-1
 304. Tal TL, Tanguay RL (2012) Non-coding RNAs—novel targets in neurotoxicity. *Neurotoxicology* 33(3):530–544. doi:10.1016/j.neuro.2012.02.013
 305. Golbidi S, Ebadi SA, Laher I (2011) Antioxidants in the treatment of diabetes. *Curr Diabetes Rev* 7(2):106–125
 306. Johansen JS, Harris AK, Rychly DJ, Ergul A (2005) Oxidative stress and the use of antioxidants in diabetes: linking basic science to clinical practice. *Cardiovasc Diabetol* 4(1):5. doi:10.1186/1475-2840-4-5
 307. Vina J, Lloret A, Giraldo E, Badia MC, Alonso MD (2011) Antioxidant pathways in Alzheimer's disease: possibilities of intervention. *Curr Pharm Des* 17(35):3861–3864
 308. Chakroborty S, Stutzmann GE (2011) Early calcium dysregulation in Alzheimer's disease: setting the stage for synaptic dysfunction. *Sci China Life Sci* 54(8):752–762. doi:10.1007/s11427-011-4205-7
 309. Jeitner TM, Pinto JT, Krasnikov BF, Horswill M, Cooper AJ (2009) Transglutaminases and neurodegeneration. *J Neurochem* 109(Suppl 1):160–166. doi:10.1111/j.1471-4159.2009.05843.x
 310. Kim S, Rhim H (2011) Effects of amyloid-beta peptides on voltage-gated L-type Ca(V)1.2 and Ca(V)1.3 Ca(2+) channels. *Mol Cells* 32(3):289–294. doi:10.1007/s10059-011-0075-x
 311. Wiederkehr A, Wollheim CB (2008) Impact of mitochondrial calcium on the coupling of metabolism to insulin secretion in the pancreatic beta-cell. *Cell Calcium* 44(1):64–76. doi:10.1016/j.ceca.2007.11.004
 312. Fernyhough P, Calcutt NA (2010) Abnormal calcium homeostasis in peripheral neuropathies. *Cell Calcium* 47(2):130–139. doi:10.1016/j.ceca.2009.11.008
 313. Verkhratsky A, Fernyhough P (2008) Mitochondrial malfunction and Ca²⁺ dyshomeostasis drive neuronal pathology in diabetes. *Cell Calcium* 44(1):112–122. doi:10.1016/j.ceca.2007.11.010
 314. Strum JC, Shehee R, Virley D, Richardson J, Mattie M, Selley P, Ghosh S, Nock C, Saunders A, Roses A (2007) Rosiglitazone induces mitochondrial biogenesis in mouse brain. *J Alzheimers Dis* 11(1):45–51
 315. Sato T, Hanyu H, Hirao K, Kanetaka H, Sakurai H, Iwamoto T (2011) Efficacy of PPAR-gamma agonist pioglitazone in mild Alzheimer disease. *Neurobiol Aging* 32(9):1626–1633. doi:10.1016/j.neurobiolaging.2009.10.009
 316. Chesik D, De Keyser J, Wilczak N (2008) Insulin-like growth factor system regulates oligodendroglial cell behavior: therapeutic potential in CNS. *J Mol Neurosci* 35(1):81–90. doi:10.1007/s12031-008-9041-2
 317. Chen LM, Lin ZY, Zhu YG, Lin N, Zhang J, Pan XD, Chen XC (2012) Ginsenoside Rg1 attenuates beta-amyloid generation via suppressing PPARgamma-regulated BACE1 activity in N2a-APP695 cells. *Eur J Pharmacol* 675(1–3):15–21. doi:10.1016/j.ejphar.2011.11.039
 318. Nicolakakis N, Hamel E (2011) Neurovascular function in Alzheimer's disease patients and experimental models. *J Cereb Blood Flow Metab Off J Int Soc Cereb Blood Flow Metab* 31(6):1354–1370. doi:10.1038/jcbfm.2011.43
 319. Sodhi RK, Singh N, Jaggi AS (2011) Neuroprotective mechanisms of peroxisome proliferator-activated receptor agonists in Alzheimer's disease. *Naunyn Schmiedeberg's Arch Pharmacol* 384(2):115–124. doi:10.1007/s00210-011-0654-6
 320. Hanyu H, Sato T, Kiuchi A, Sakurai H, Iwamoto T (2009) Pioglitazone improved cognition in a pilot study on patients with Alzheimer's disease and mild cognitive impairment with diabetes mellitus. *J Am Geriatr Soc* 57(1):177–179. doi:10.1111/j.1532-5415.2009.02067.x
 321. Zhao Q, Matsumoto K, Tsuneyama K, Tanaka K, Li F, Shibahara N, Miyata T, Yokozawa T (2011) Diabetes-induced central cholinergic neuronal loss and cognitive deficit are attenuated by tacrine and a Chinese herbal prescription, kangen-karyu: elucidation in type 2 diabetes db/db mice. *J Pharmacol Sci* 117(4):230–242
 322. Cramer PE, Cirrito JR, Wesson DW, Lee CY, Karlo JC, Zinn AE, Casali BT, Restivo JL, Goebel WD, James MJ, Brunden KR, Wilson DA, Landreth GE (2012) ApoE-directed therapeutics rapidly clear beta-amyloid and reverse deficits in AD mouse models. *Science* 335(6075):1503–1506. doi:10.1126/science.1217697

323. Chawla A, Boisvert WA, Lee CH, Laffitte BA, Barak Y, Joseph SB, Liao D, Nagy L, Edwards PA, Curtiss LK, Evans RM, Tontonoz P (2001) A PPAR gamma-LXR-ABCA1 pathway in macrophages is involved in cholesterol efflux and atherogenesis. *Mol Cell* 7(1):161–171
324. Lefebvre P, Benomar Y, Staels B (2010) Retinoid X receptors: common heterodimerization partners with distinct functions. *Trends Endocrinol Metab* 21(11):676–683. doi:[10.1016/j.tem.2010.06.009](https://doi.org/10.1016/j.tem.2010.06.009)
325. Feige JN, Gelman L, Michalik L, Desvergne B, Wahli W (2006) From molecular action to physiological outputs: peroxisome proliferator-activated receptors are nuclear receptors at the cross-roads of key cellular functions. *Prog Lipid Res* 45(2):120–159. doi:[10.1016/j.plipres.2005.12.002](https://doi.org/10.1016/j.plipres.2005.12.002)
326. Chen Y, Zhou K, Wang R, Liu Y, Kwak YD, Ma T, Thompson RC, Zhao Y, Smith L, Gasparini L, Luo Z, Xu H, Liao FF (2009) Antidiabetic drug metformin (GlucophageR) increases biogenesis of Alzheimer's amyloid peptides via up-regulating BACE1 transcription. *Proc Natl Acad Sci U S A* 106(10):3907–3912. doi:[10.1073/pnas.0807991106](https://doi.org/10.1073/pnas.0807991106)
327. Kickstein E, Krauss S, Thornhill P, Rutschow D, Zeller R, Sharkey J, Williamson R, Fuchs M, Kohler A, Glossmann H, Schneider R, Sutherland C, Schweiger S (2010) Biguanide metformin acts on tau phosphorylation via mTOR/protein phosphatase 2A (PP2A) signaling. *Proc Natl Acad Sci U S A* 107(50):21830–21835. doi:[10.1073/pnas.0912793107](https://doi.org/10.1073/pnas.0912793107)
328. Gupta A, Bisht B, Dey CS (2011) Peripheral insulin-sensitizer drug metformin ameliorates neuronal insulin resistance and Alzheimer's-like changes. *Neuropharmacology* 60(6):910–920. doi:[10.1016/j.neuropharm.2011.01.033](https://doi.org/10.1016/j.neuropharm.2011.01.033)
329. Markaki M, Tavernarakis N (2010) Modeling human diseases in *Caenorhabditis elegans*. *Biotechnol J* 5(12):1261–1276. doi:[10.1002/biot.201000183](https://doi.org/10.1002/biot.201000183)
330. Morcos M, Hutter H (2009) The model *Caenorhabditis elegans* in diabetes mellitus and Alzheimer's disease. *J Alzheimers Dis* 16(4):897–908. doi:[10.3233/JAD-2009-0977](https://doi.org/10.3233/JAD-2009-0977)
331. Wu Z, Smith JV, Paramasivam V, Butko P, Khan I, Cypser JR, Luo Y (2002) *Ginkgo biloba* extract EGb 761 increases stress resistance and extends life span of *Caenorhabditis elegans*. *Cell Mol Biol* 48(6):725–731
332. Dostal V, Link CD (2010) Assaying beta-amyloid toxicity using a transgenic *C. elegans* model. *Journal of visualized experiments : JoVE* (44). doi:[10.3791/2252](https://doi.org/10.3791/2252)
333. Onken B, Driscoll M (2010) Metformin induces a dietary restriction-like state and the oxidative stress response to extend *C. elegans* Healthspan via AMPK, LKB1, and SKN-1. *PLoS One* 5(1):e8758. doi:[10.1371/journal.pone.0008758](https://doi.org/10.1371/journal.pone.0008758)
334. Saharia K, Arya U, Kumar R, Sahu R, Das CK, Gupta K, Dwivedi H, Subramaniam JR (2012) Reserpine modulates neurotransmitter release to extend lifespan and alleviate age-dependent Abeta proteotoxicity in *Caenorhabditis elegans*. *Exp Gerontol* 47(2):188–197. doi:[10.1016/j.exger.2011.12.006](https://doi.org/10.1016/j.exger.2011.12.006)