

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/11264487>

# Oligonucleotides containing 7-vinyl-7-deazaguanine as a facile strategy for expanding the functional diversity of DNA

ARTICLE in BIOORGANIC & MEDICINAL CHEMISTRY LETTERS · SEPTEMBER 2002

Impact Factor: 2.42 · DOI: 10.1016/S0960-894X(02)00334-7 · Source: PubMed

CITATIONS

9

READS

15

## 4 AUTHORS, INCLUDING:



**Akimitsu Okamoto**

The University of Tokyo

**134** PUBLICATIONS **2,430** CITATIONS

SEE PROFILE



**Kazuki Tainaka**

RIKEN

**49** PUBLICATIONS **709** CITATIONS

SEE PROFILE



**Isao Saito**

Nihon University

**239** PUBLICATIONS **5,172** CITATIONS

SEE PROFILE



Pergamon

Bioorganic &amp; Medicinal Chemistry Letters 12 (2002) 1895–1896

BIOORGANIC &  
MEDICINAL  
CHEMISTRY  
LETTERS

# Oligonucleotides Containing 7-Vinyl-7-deazaguanine as a Facile Strategy for Expanding the Functional Diversity of DNA

Akimitsu Okamoto, Toshiji Taiji, Kazuki Tainaka and Isao Saito\*

*Department of Synthetic Chemistry and Biological Chemistry, Faculty of Engineering, Kyoto University, SORST, Japan Science and Technology Corporation, Kyoto 606-8501, Japan*

Received 17 April 2002; accepted 11 May 2002


**Abstract**—A modified nucleobase 7-vinyl-7-deazaguanine (<sup>V</sup>G) produced adducts with maleimides through Diels–Alder cycloaddition under very mild conditions. By this method, post-synthetic modification to oligonucleotides with diverse functionality (carboxylic acid, pyrene, benzophenone, succinimidyl ester, nitroxide and biotin) was accomplished. © 2002 Elsevier Science Ltd. All rights reserved.

Assembling of functionalized oligonucleotides (ODN) into complementary nucleic acids with target sequences via the formation of Watson–Crick base pairs has led to a rich variety of technologies exploiting new functionalities of ODN.<sup>1–5</sup> Thus, the development of a new synthetic method for ODN possessing diverse functionalities is of great interest. A number of methods for incorporating functionalities to ODN by means of post-synthetic modification have been described.<sup>6–9</sup> Commonly employed methods for the preparation of ODN bioconjugates include the introduction of the chemical modifier during solid-phase ODN synthesis or by post-synthetic modification using reactive handles already incorporated during ODN synthesis. These methods are often accompanied by undesirable operational complexity such as protection/deprotection and site-specific chemical activation.

The Diels–Alder reaction is a very attractive approach for bioconjugation due to the remarkable acceleration of the reaction in aqueous systems.<sup>10,11</sup> Only a few examples of nucleic acid modifications utilizing the Diels–Alder reaction have been reported so far.<sup>12–15</sup> However, most of these methods required long reaction times and/or a specific sequence that may catalyze the reaction. Furthermore, currently available Diels–Alder bioconjugation methods are restricted to only strand ends.


Herein, we report a facile method for the incorporation of functionalized groups into ODN via the Diels–Alder reaction using a novel nucleobase 7-vinyl-7-deazaguanine (<sup>V</sup>G, **1** in Fig. 1). We found that the reaction of <sup>V</sup>G with *N*-substituted maleimides proceeded exceedingly rapidly. By this technique, post-synthetic modification of ODNs was achieved under very mild aqueous conditions.

Prior to post-synthetic modification of <sup>V</sup>G-containing ODNs, we first investigated the reaction of protected <sup>V</sup>G nucleoside **2** with *N*-methylmaleimide. The preparation of **2** was accomplished by the method reported earlier.<sup>16</sup> The reaction mixture of **2** and *N*-methylmaleimide in methanol was incubated at room temperature (Scheme 1). Within 5 min, **2** was completely converted to maleimide-adduct **3** [calcd 459.1992 for (M+H)<sup>+</sup>, found 459.2004] via Diels–Alder cycloaddition and subsequent [1,3] H-shift. This experiment indicates that an exocyclic vinyl group at C7 and an endocyclic C7–C8 double bond of **2** serve as a very effective acceptor for the dienophile in the Diels–Alder reaction.


**1** (<sup>V</sup>G)

**Figure 1.** 7-Vinyl-7-deazaguanine (**1**, <sup>V</sup>G) used in this study.

\*Corresponding author. Tel.: +81-75-753-5656; fax: +81-75-753-5676; e-mail: saito@sbchem.kyoto-u.ac.jp



**Scheme 1.** Diels–Alder reaction of *N*-protected <sup>V</sup>G (**2**) with *N*-methylmaleimide.



**Figure 2.** Functionalized maleimide derivatives incorporated into <sup>V</sup>G-containing ODN 5'-d(T<sup>V</sup>GACGTCA)-3' through Diels–Alder cycloaddition.

We next examined the post-synthetic modification of a <sup>V</sup>G-containing ODN 5'-d(T<sup>V</sup>GACGTCA)-3' by the Diels–Alder reaction. The preparation of d(T<sup>V</sup>GACGTCA) was accomplished by conventional solid-phase DNA synthesis.<sup>17,18</sup> The ODN was incubated with *N*-methylmaleimide (**4**) in phosphate buffer (pH 7.0) at 0 °C. The progress of the reaction was monitored by MALDI-TOF mass spectrometry. The reaction proceeded up to 80% conversion in 10 min, and after 1 h the starting ODN had completely disappeared. The product was proved to be a 1:1 adduct of <sup>V</sup>G-containing ODN and *N*-methylmaleimide by MALDI-TOF mass [calcd 2544.76 for [(M–H)<sup>–</sup>], found 2544.24].

Similarly, incorporation of other functionalized maleimides, into <sup>V</sup>G-containing ODN was examined in phosphate buffer (pH 7.0) at 0 °C. The corresponding adducts with <sup>V</sup>G-containing ODN were characterized by MALDI-TOF mass. The maleimide derivatives insoluble in phosphate buffer were added to the reaction mixture in a methanol solution. The 1 h incubation of the <sup>V</sup>G-containing ODN with various functionalized maleimides effectively afforded the corresponding adducts. These include maleimides containing carboxylic acid **5**, an activated ester **6**, benzophenone **7** for

photoaffinity labelling, pyrene **8** for a fluorophore, TEMPO **9** as a nitroxide spin label, and biotin **10** (Fig. 2). In these reactions starting ODN completely disappeared and was converted to the corresponding adduct without any additive or protection at 0 °C.<sup>19,20</sup>

In conclusion, a facile method for the incorporation of functionalized groups into ODN was achieved by the Diels–Alder reaction. The present post-synthetic modification proceeded quantitatively under exceedingly mild conditions (pH 7.0, 0 °C). Furthermore, 7-vinyl-7-deazaguanine <sup>V</sup>G, possesses a high compatibility for the incorporation of diverse functionalities through Diels–Alder cycloaddition. Thus, the post-synthetic modification of oligonucleotides containing <sup>V</sup>G as a functionally diversifiable nucleotide is promising and applicable to site-selective DNA labelling and bioconjugation with inherently labile functionalized groups.

## References and Notes

- Goodchild, J. *Bioconjugate Chem.* **1990**, *1*, 165.
- Uhlmann, E.; Peyman, A. *Chem. Rev.* **1990**, *90*, 543.
- Beaucage, S. L.; Iyer, R. P. *Tetrahedron* **1993**, *49*, 1925.
- Beaucage, S. L.; Iyer, R. P. *Tetrahedron* **1993**, *49*, 6123.
- Verma, S.; Eckstein, F. *Ann. Rev. Biochem.* **1998**, *67*, 99.
- Khan, S. I.; Grinstaff, M. W. *J. Am. Chem. Soc.* **1999**, *121*, 4704.
- Hwang, J.-T.; Greenberg, M. M. *Org. Lett.* **1999**, *1*, 2021.
- Shinozuka, K.; Kohgo, S.; Ozaki, H.; Sawai, H. *Chem. Commun.* **2000**, 59.
- Tilquin, J.-M.; Dechamps, M.; Sonveaux, E. *Bioconjugate Chem.* **2001**, *12*, 451.
- Rideout, D. C.; Breslow, R. *J. Am. Chem. Soc.* **1980**, *102*, 7816.
- Otto, S.; Blandamer, F.; Engberts, J. B. F. N. *J. Am. Chem. Soc.* **1996**, *118*, 7702.
- Tarasow, T. M.; Tarasow, S. L.; Eaton, B. E. *Nature* **1997**, *389*, 54.
- Seelig, B.; Jäschke, A. *Tetrahedron Lett.* **1997**, *38*, 7729.
- Seelig, B.; Jäschke, A. *Chem. Biol.* **1999**, *6*, 167.
- Hill, K. W.; Taunton-Rigby, J.; Carter, J. D.; Kropp, E.; Valge, K.; Pieken, W.; McGee, D. P. C.; Husar, G. M.; Leuck, M.; Anziano, D. J.; Sebesta, D. P. *J. Org. Chem.* **2001**, *66*, 5352.
- Okamoto, A.; Taiji, T.; Tanaka, K.; Saito, I. *Tetrahedron Lett.* **2000**, *41*, 10035.
- MALDI-TOF MS for 5'-d(T<sup>V</sup>GACGTCA)-3', [(M–H)<sup>–</sup>]: calcd 2433.66, found 2433.37.
- <sup>V</sup>G forms a stable base pair with C in duplex DNA. For the stabilization of the duplex containing <sup>V</sup>G, see ref 16.
- MALDI-TOF MS for d(T<sup>V</sup>GACGTCA)–maleimide adducts, [(M–H)<sup>–</sup>]: **5**, calcd 2602.80, found 2602.54; **6**, calcd 2713.89, found 2713.23; **7**, calcd 2710.93, found 2711.88; **8**, calcd 2730.97, found 2730.47; **9**, calcd 2684.96, found 2685.30; **10**, calcd 2959.28, found 2959.63.
- Incubation of <sup>V</sup>G-containing ODN with succinimidyl ester **6** effectively affords the corresponding adduct, but the methyl ester produced by the solvolysis was also detected in low amounts in mass spectroscopy.