

Synthesis of pyrido[2,3-d] [1,2,4]triazolo[4,3-a]pyrimidin-5-ones as potential antimicrobial agents

**Thoraya A. Farghaly & Huwaida
M. E. Hassaneen**

Archives of Pharmacal Research

ISSN 0253-6269

Volume 36

Number 5


Arch. Pharm. Res. (2013) 36:564-572

DOI 10.1007/s12272-013-0045-2

ISSN 0253-6269 (PRINT)
ISSN 1976-3786 (ONLINE)

**Archives of
Pharmacal Research**

Volume 36 • Number 5 • May 2013



 Springer

 The Pharmaceutical
Society of Korea

 Springer

Your article is protected by copyright and all rights are held exclusively by The Pharmaceutical Society of Korea. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

RESEARCH ARTICLE

Synthesis of pyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]pyrimidin-5-ones as potential antimicrobial agents

Thoraya A. Farghaly · Huwaida M. E. Hassaneen

Received: 4 October 2012 / Accepted: 28 January 2013 / Published online: 28 February 2013
© The Pharmaceutical Society of Korea 2013

Abstract Synthesis of new derivatives of pyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]pyrimidin-5-one via reaction of 7-(4-bromophenyl)-1,2-dihydro-5-(4-fluorophenyl)-2-thioxopyrido[2,3-*d*] pyrimidin-4(3*H*)-one with hydrazonoyl chlorides or reaction of 2-hydrazino-pyrido[2,3-*d*]pyrimidin-4(3*H*)-one with different aldehydes followed by cyclization of the products. All the newly synthesized compounds were evaluated for their antimicrobial activities and also their minimum inhibitory concentration against most of test organisms was performed. Amongst the tested compounds displayed excellent activity against all the tested microorganisms except SR and PA.

Keywords Pyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]pyrimidin-5-ones · Hydrazonoyl chlorides · Antimicrobial activity

Introduction

Infections caused by multi-drug resistant bacteria are of major health concern worldwide. Particularly important are infections caused by the Gram-positive bacteria *Staphylococcus aureus* and species of the genus *Enterococcus*, due to increasing incidence of infections caused by these microorganisms in hospitals and communities, and their ability of developing antibiotic resistance to multiple antibiotics. Due to some serious side effects in newly introduced antibacterial agents such as semi-synthetic streptogramins quinupristin/dalfopristin, daptomycin, the development of a diversified series of antimicrobials still remains a necessity (Leclercq et al. 2009).

Pyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]pyrimidin-5-ones are pharmacological scaffold that represent a wide range of biological activities (El-Gazzar et al. 2008, 2009; Abu-Zied et al. 2008). As a part of our ongoing research on the synthesis of biologically active heterocyclic compounds (Hassaneen et al. 2011; Gomha and Hassaneen 2011; Farghaly and Abdalla 2009; Abdel Hafez et al. 2010; Farghaly et al. 2010, 2012; Riyadh et al. 2010) herein we report an efficient methodology for synthesis of novel pyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]pyrimidin-5-one derivatives and investigate their antimicrobial activities.

Materials and methods

Melting points were determined on an electrothermal Galenkamp apparatus and are uncorrected. The IR spectra were recorded in KBr using a Pye Unicam SP-1000 spectrophotometer. ¹H NMR and ¹³C NMR spectra were obtained on a Varian EM-300 MHz spectrometer (¹H NMR 300 or 400 MHz) using DMSO-*d*₆, solvent with TMS as internal standard. ¹³C NMR spectra were measured on a Varian EM-300 MHz spectrometer (100 MHz). Mass spectra were recorded on an AEI MS 30 mass spectrometer operating at 70e V. Elemental analyses were carried out by the Microanalytical Center of Cairo University, Giza, Egypt. Antimicrobial activities were carried out at the Regional Center for Mycology and Biotechnology at Al-Azhar University, Cairo, Egypt.

Synthesis of 7-(4-bromophenyl)-1,2-dihydro-5-(4-fluorophenyl)-2-thioxo-pyrido-[2,3-*d*]pyrimidin-4(3*H*)-one (**3**)

A solution of 6-amino-2,3-dihydro-2-thioxo-pyrimidin-4(1*H*)-one (**2**) (0.5 g, 3.5 mmol) and chalcone **1** (1.06 g, 3.5 mmol)

T. A. Farghaly · H. M. E. Hassaneen (✉)
Department of Chemistry, Faculty of Science, Cairo University,
Giza 12613, Egypt
e-mail: huwaidahassaneen@yahoo.com

in 5 ml dimethylformamide was refluxed for 15 h. After the reaction completed (examined by TLC), the solution is cooled down to 5 °C and the precipitate formed is filtered off and crystallized from DMF to give compound **3** as yellow crystals; (80 % yield), mp 303–305 °C; IR (KBr): 3,150, 3,086 (2NH), 1,701 (C=O); ¹H NMR (DMSO-*d*₆): δ = 7.24 (d, *J* = 8 Hz, 2H, Ar-H), 7.51 (d, *J* = 8 Hz, 2H, Ar-H), 7.70 (s, 1H, pyridine-H), 7.75 (d, *J* = 8 Hz, 2H, Ar-H), 8.20 (d, *J* = 8 Hz, 2H, Ar-H), 12.30 (s, 1H, NH, D₂O-exchangeable), 13.0 (s, 1H, NH, D₂O-exchangeable); ¹³C NMR (DMSO-*d*₆): δ = 110.20, 114.54, 116.07, 121.05, 122.29, 123.02, 124.0, 130.97, 134.28, 143.20, 145.21, 158.82, 160.81, 161.64, 163.74. MS: *m/z* 430 (M⁺+2, 1), 429 (M⁺+ 1, 2), 428 (M⁺, 2), 183 (3), 155 (1), 95 (2), 73 (100). Anal. calcd. For C₁₉H₁₁BrFN₃OS (428.28) C, 53.28; H, 2.59; N, 9.81; Found: C, 53.14; H, 2.37; N, 9.74 %.

General procedure for the preparation of **7a–d**

Method A

To a mixture of equimolar amounts of compound **3** and the appropriate hydrazonoyl chlorides **4** (1 mmol of each) in dioxane (15 ml) was added triethylamine (0.14 ml, 1 mmol). The reaction mixture was refluxed till all of the starting materials have disappeared and hydrogen sulfide gas ceased to evolve (5 h, monitored by TLC). The solvent was evaporated and the residue was treated with methanol. The solid that formed was filtered off and recrystallized from dioxane to give products **7a–d**.

Method B

To a mixture of equimolar amounts of **9** and the appropriate hydrazonoyl chloride **4a** (1 mmol) in dioxane (15 ml) was added triethylamine (0.14 ml, 1 mmol). The reaction mixture was refluxed till all methanethiol gas ceased to evolve (10 h, monitored by TLC). The solvent was evaporated and the residue was treated with methanol. The solid that formed was filtered and recrystallized from dioxane to give product identical in all respects (mp, mixed mp and IR) with **7a**.

8-(4-Bromophenyl)-6-(4-fluorophenyl)-1,3-diphenylpyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]-pyrimidin-5-one (**7a**)

Yellow crystals; (72 % yield), mp 343–345 °C; IR (KBr): 1,705 (C=O); ¹H NMR (DMSO-*d*₆): δ = 6.98–7.84 (m, 18H, Ar-H), 7.66 (s, 1H, pyridine-H); ¹³C NMR (DMSO-*d*₆): δ = 110.05, 118.55, 119.63, 120.25, 121.65, 123.32, 124.05, 126.53, 127.65, 129.37, 130.31, 131.78, 132.62, 133.85, 136.35, 138.16, 139.05, 147.08, 149.95,

151.28, 154.65, 158.23, 159.12, 162.32. MS: *m/z* 590 (M⁺+2, 9), 589 (M⁺+ 1, 25), 588 (M⁺, 25), 587 (27), 586 (21), 220 (14), 103 (12), 91 (100), 75 (11). Anal. calcd. for C₃₂H₁₉BrFN₅O (588.43) C, 65.32; H, 3.25; N, 11.90; Found: C, 65.30; H, 3.16; N, 11.85 %.

3-Acetyl-8-(4-bromophenyl)-6-(4-fluorophenyl)-1-phenyl-pyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]pyrimidin-5-one (**7b**)

Yellow crystals; (78 % yield), mp 292–294 °C; IR (KBr): 1,720, 1,674 (2C=O); ¹H NMR (DMSO-*d*₆): δ = 2.45 (s, 3H, CH₃CO), 7.12–7.86 (m, 13H, Ar-H), 7.68 (s, 1H, pyridine-H); ¹³C NMR (DMSO-*d*₆): δ = 25.63, 110.54, 119.67, 120.28, 122.10, 123.68, 127.23, 129.63, 130.12, 132.13, 134.36, 136.54, 138.38, 138.78, 148.86, 150.31, 150.85, 151.11, 158.35, 158.22, 163.15, 190.32. MS: *m/z* 556 (M⁺+2, 23), 555 (M⁺+ 1, 77), 554 (M⁺, 78), 553 (72), 485 (32), 158 (16), 91 (46), 77 (100). Anal. calcd. for C₂₈H₁₇BrFN₅O₂ (554.37) C, 60.66; H, 3.09; N, 12.63; Found: C, 60.51; H, 3.01; N, 12.53 %.

8-(4-Bromophenyl)-3-ethoxycarbonyl-6-(4-fluorophenyl)-1-phenyl-pyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]pyrimidin-5-one (**7c**)

Yellow crystals; (74 % yield), mp 284–282 °C; IR (KBr): 1,751, 1,712 (2C=O); ¹H NMR (DMSO-*d*₆): δ = 1.30 (t, *J* = 7.2 Hz, 3H, CH₃), 4.43 (q, *J* = 7.2 Hz, 2H, CH₂), 6.97–7.87 (m, 13H, Ar-H), 7.66 (s, 1H, pyridine-H); ¹³C NMR (DMSO-*d*₆): δ = 14.23, 58.66, 109.66, 119.88, 121.56, 124.09, 127.44, 129.56, 130.12, 131.54, 133.56, 136.55, 137.65, 138.54, 145.21, 148.23, 150.94, 151.72, 152.64, 153.08, 158.35, 160.62, 163.62. MS: *m/z* 586 (M⁺+2, 33), 585 (M⁺+ 1, 100), 584 (M⁺, 96), 583 (94), 482 (33), 360 (19), 157 (22), 91 (99), 77 (99). Anal. calcd. For C₂₉H₁₉BrFN₅O₃ (584.40) C, 59.60; H, 3.28; N, 11.98; Found: C, 59.42; H, 3.16; N, 11.88 %.

8-(4-Bromophenyl)-6-(4-fluorophenyl)-1-phenyl-3-Phenylcarbonyl-pyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]pyrimidin-5-one (**7d**)

Yellow crystals; (70 % yield), mp 279–281 °C; IR (KBr): 3,063 (NH), 1,701, 1,680 (2C=O); ¹H NMR (DMSO-*d*₆): δ = 7.18–7.60 and 7.67–8.34 (m, 18H, Ar-H), 7.65 (s, 1H, pyridine-H), 12.42 (s, 1H, NH); ¹³C NMR (DMSO-*d*₆): δ = 111.14, 119.45, 118.33, 119.12, 120.16, 123.30, 124.48, 126.49, 126.99, 129.73, 131.20, 131.58, 132.60, 134.95, 136.15, 137.26, 139.44, 147.18, 149.0, 150.65, 153.95, 158.02, 158.69, 159.98, 161.47. MS: *m/z* 632 (M⁺+ 1, 11), 631 (M⁺, 18), 554 (36), 485 (29), 158 (32), 118 (35), 93 (100), 77 (56). Anal. calcd. for C₃₃H₂₀BrFN₆O₂ (631.45) C,

62.77; H, 3.19; N, 13.31; Found: C, 62.58; H, 3.02; N, 13.22 %.

Synthesis of 7-(4-bromophenyl)-5-(4-fluorophenyl)-2-methylthio-pyrido[2,3-*d*]-pyrimidin-4(3*H*)-one (**9**)

To a stirred solution of compound **3** (2.14 g, 5 mmol) in dimethylformamide (20 ml) was added anhydrous potassium carbonate (0.70 g, 5 mmol) and methyl iodide (0.71 g, 5 mmol). The reaction mixture was stirred overnight at room temperature, then, the mixture was diluted with water. The solid formed was filtered off, washed with water, dried and crystallized from ethanol to give compound **9** as white solid; (76 % yield), mp 300–302 °C; IR (KBr): 3,074 (NH), 1,658 (C=O); ¹H NMR (DMSO-*d*₆): δ = 2.72 (s, 3H, SCH₃), 7.24 (d, *J* = 8 Hz, 2H, Ar-H), 7.48 (d, *J* = 8 Hz, 2H, Ar-H), 7.73 (d, *J* = 8 Hz, 2H, Ar-H), 7.80 (s, 1H, pyridine-H), 8.24 (d, *J* = 8 Hz, 2H, Ar-H), 11.54 (s, 1H, NH, D₂O-exchangeable); ¹³C NMR (DMSO-*d*₆): δ = 12.05, 111.12, 113.21, 116.17, 120.47, 123.12, 123.11, 125.05, 129.19, 133.35, 143.54, 144.01, 156.28, 158.48, 159.78, 164.35. MS: *m/z* 444 (M⁺+2, 12), 443 (M⁺+1, 22), 442 (M⁺, 62), 260 (23), 158 (13), 80 (12). Anal. calcd. for C₂₀H₁₃BrFN₃OS (442.31) C, 54.31; H, 2.96; N, 9.50; Found: C, 54.21; H, 2.85; N, 9.64 %.

Synthesis of 7-(4-bromophenyl)-5-(4-fluorophenyl)-2-hydrazino-pyrido[2,3-*d*]-pyrimidin-4(3*H*)-one (**11**)

To thione **3** or its methylthio analogue **9** (5 mmol) in ethanol (20 ml) was added hydrazine hydrate (10 ml, 80 %). The reaction mixture was refluxed till all H₂S (30 h) or MeSH (5 h) ceased to evolve, and then cooled. The solid that precipitated was filtered off and crystallized from dilute dioxane to give compound **11** as white solid; (80 % yield), mp 276 °C; IR (KBr): 3,200, 3,150, 3,100 (NH₂, 2NH) in IR 3310 cm⁻¹, 1,689 (C=O); ¹H NMR (DMSO-*d*₆): δ = 2.0 (br.s, 2H, HNS), 6.13 (br., 1H, NH), 7.24–8.19 (m, 9H, Ar-H) 10.56 (br., 1H, NH); ¹³C NMR (DMSO-*d*₆): δ = 111.0, 114.68, 119.88, 120.18, 123.91, 124.26, 128.35, 130.55, 134.94, 135.41, 145.20, 147.70, 158.71, 159.69, 164.35. MS: *m/z* 428 (M⁺+2, 5), 427 (M⁺+1, 2), 426 (M⁺, 16), 395 (12), 156 (34), 121 (52), 96 (18), 76 (100). Anal. calcd. for C₁₉H₁₃BrFN₅O (426.24) C, 53.54; H, 3.07; N, 16.43; Found: C, 53.35; H, 3.05; N, 16.28 %.

General procedure for the preparation of **12a–c**

A mixture of hydrazine **11** (1.07 g, 2.5 mmol) and the appropriate aldehydes (2.5 mmol) in acetic acid (20 ml) and few drops of conc. hydrochloric acid (≈1 ml) was heated under reflux for 3 h. The reaction mixture was then cooled and diluted with water. The so-formed solid product

was then collected by filtration, dried and recrystallized from dimethylformamide to afford the corresponding hydrazones **12a–c**

7-(4-Bromophenyl)-5-(4-fluorophenyl)-2-[(phenylmethylene)hydrazino]-3*H*-pyrido[2,3-*d*]pyrimidin-4-one (**12a**)

Pale Yellow solid; (76 % yield), mp 336–338 °C; IR (KBr): 3,336, 3,100 (2NH), 1,681 (C=O); ¹H NMR (DMSO-*d*₆): δ = 7.23–7.27 and 7.50–7.53 (m, 9H, Ar-H), 7.41 (s, 1H, =CH), 7.71 (d, *J* = 8 Hz, 2H, Ar-H), 7.97 (s, 1H, pyridine-H), 8.15 (d, *J* = 8 Hz, 2H, Ar-H), 12.25 (br.s, 2H, 2NH, D₂O-exchangeable); ¹³C NMR (DMSO-*d*₆): δ = 112.26, 118.21, 120.05, 121.57, 122.99, 127.53, 129.15, 131.78, 131.28, 133.12, 134.50, 137.12, 138.95, 147.54, 150.17, 154.36, 155.01, 158.29, 159.45, 161.45. MS: *m/z* 515 (M⁺+1, 54), 514 (M⁺, 42), 513 (50), 437 (54), 201 (50), 199 (69), 158 (27), 105 (58), 104 (62), 89 (50), 77 (46), 76 (100). Anal. calcd. for C₂₆H₁₇BrFN₅O (514.35) C, 60.71; H, 3.33; N, 13.62; Found: C, 60.49; H, 3.26; N, 13.47 %.

7-(4-Bromophenyl)-5-(4-fluorophenyl)-2-[(4-florophenylmethylene)hydrazino]-3*H*-pyrido[2,3-*d*]pyrimidin-4-one (**12b**)

Yellow solid; (80 % yield), mp 340–342 °C; IR (KBr): 3,200, 3,143 (2NH), 1,666 (C=O); ¹H NMR (DMSO-*d*₆): δ = 7.24–8.09 (m, 12H, Ar-H), 8.14 (s, 1H, pyridine-H), 8.33 (s, 1H, =CH), 12.25 (br.s, 2H, 2NH, D₂O-exchangeable); ¹³C NMR (DMSO-*d*₆): δ = 110.89, 113.53, 114.17, 120.65, 121.44, 123.14, 127.35, 131.54, 131.97, 133.58, 137.23, 137.96, 147.58, 150.19, 153.70, 158.06, 159.15, 159.94, 160.24, 161.58. MS: *m/z* 533 (M⁺+1, 2), 532 (M⁺, 3), 89 (25), 76 (11). Anal. calcd. for C₂₆H₁₆BrF₂N₅O (532.34) C, 58.66; H, 3.03; N, 13.16; Found: C, 58.44; H, 3.14; N, 13.08 %.

7-(4-Bromophenyl)-5-(4-fluorophenyl)-2-[(4-nitrophenylmethylene)hydrazino]-3*H*-pyrido[2,3-*d*]pyrimidin-4-one (**12c**)

Yellow solid; (80 % yield), mp 328–330 °C; IR (KBr): 3,375, 3,100 (2NH), 1,701 (C=O); ¹H NMR (DMSO-*d*₆): δ = 7.18–7.69 (m, 12H, Ar-H), 7.71 (s, 1H, pyridine-H), 8.21 (s, 1H, =CH), 11.80 (br.s, 2H, 2NH, D₂O-exchangeable); ¹³C NMR (DMSO-*d*₆): δ = 111.27, 120.48, 122.88, 123.0, 126.34, 127.50, 129.58, 131.02, 131.47, 135.48, 137.48, 138.04, 148.25, 148.98, 151.75, 155.45, 158.78, 159.61, 162.01, 164.42. MS: *m/z* 560 (M⁺+2, 1), 559 (M⁺+1, 1), 558 (M⁺, 2), 531 (100), 436 (21), 109 (64), 95 (58), 76 (15). Anal. calcd. for C₂₆H₁₆BrFN₆O₃ (559.35) C,

55.83; H, 2.88; N, 15.02; Found: C, 55.72; H, 2.81; N, 15.21 %.

General procedure for the preparation of **13a–c**

Method A

Bromine (0.052 g, 1 mmol) in acetic acid (5 ml) in presence of sodium acetate (1 mmol) was added drop wise to a stirred solution of the appropriate hydrazone **12a–c** (1 mmol of each) in acetic acid (10 ml). The reaction mixture was then poured onto ice cold water, and the solid that precipitated was filtered off, washed with sodium bicarbonate solution and then with water, dried and crystallized from the dimethylformamide to give the respective compounds **13a, c**.

Method B

A mixture of 2-hydrazinopyrido[2,3-*d*]pyrimidine **11** (1 mmol) and aroyl chloride (1.5 mmol) in dry pyridine (8 ml) was heated under reflux for 20 h. the reaction mixture was cooled and the solid formed was filtered, washed with ethanol, dried and ethanol and crystallized from the appropriate solvent to give the respective compounds **13a, c**.

8-(4-Bromophenyl)-6-(4-fluorophenyl)-3-phenyl-pyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]pyrimidin-5(1*H*)-one (**13a**)

Yellow solid; (68 % yield), mp 326–328 °C; IR (KBr): 3,313 (NH), 1,708 (C=O); ¹H NMR (DMSO-*d*₆): δ = 6.98–7.98 (m, 13H, Ar–H), 8.05 (s, 1H, pyridine-H), 12.08 (br. 1H, NH, D₂O-exchangeable), ¹³C NMR (DMSO-*d*₆): δ = 110.33, 119.45, 120.34, 122.26, 123.19, 126.90, 129.68, 130.06, 132.35, 132.52, 134.48, 137.32, 138.68, 148.36, 151.98, 154.28, 156.21, 158.27, 160.53, 162.94. MS: *m/z* 512 (M⁺, 12), 511 (29), 106 (21), 104 (44), 97 (47), 91 (27), 84 (47), 77 (52), 55 (100). Anal. calcd. for C₂₆H₁₅BrFN₅O (512.33) C, 60.95; H, 2.95; N, 13.67; Found: C, 60.76; H, 3.57; N, 13.52 %.

8-(4-Bromophenyl)-6-(4-fluorophenyl)-3-(4-fluorophenyl)-pyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]pyrimidin-5(1*H*)-one (**13b**)

Yellow solid; (65 % yield, mp 328–330 °C; IR (KBr): 3,282 (NH), 1,701 (C=O); ¹H NMR (DMSO-*d*₆): δ = 7.16–8.18 (m, 12H, Ar–H), 8.09 (s, 1H, pyridine-H), 12.11 (br. 1H, NH, D₂O-exchangeable), ¹³C NMR (DMSO-*d*₆): δ = 110.33, 114.14, 115.81, 120.32, 122.34, 123.22, 129.82, 130.11, 132.15, 134.38, 137.37, 138.86, 148.28,

151.97, 154.35, 158.57, 160.54, 162.95, 164.24, 167.06. MS: *m/z* 534 (M⁺+4, 18), 533 (M⁺+ 3, 78), 532, (M⁺+ 2, 81), 531 (M⁺+ 1, 86), 530 (M⁺, 100), 529 (59), 122 (25), 108 (91), 75 (25). Anal. calcd. for C₂₆H₁₄BrF₂N₅O (530.32) C, 58.88; H, 2.66; N, 13.21; Found: C, 58.74; H, 2.48; N, 13.11 %.

8-(4-Bromophenyl)-6-(4-fluorophenyl)-3-(4-nitrophenyl)-pyrido[2,3-*d*][1,2,4]-triazolo[4,3-*a*]pyrimidin-5(1*H*)-one (**13c**)

Yellow solid; (66 % yield), mp 338–340 °C; IR (KBr): 3,200 (NH), 1,697 (C=O); ¹H NMR (DMSO-*d*₆): δ = 7.19–8.05 (m, 12H, Ar–H), 8.04 (s, 1H, pyridine-H), 12.14 (br. 1H, NH, D₂O-exchangeable), ¹³C NMR (DMSO-*d*₆): δ = 110.34, 120.34, 122.27, 123.23, 124.08, 126.23, 129.76, 130.15, 131.43, 134.33, 137.41, 138.64, 148.19, 148.85, 151.88, 154.29, 158.28, 160.63, 162.99, 164.46. MS: *m/z* 559, (M⁺+ 2, 12), 558 (M⁺+ 1, 15), 557 (M⁺, 40), 122 (16), 95 (25), 91 (86), 89 (58), 77 (100). Anal. calcd. for C₂₆H₁₄BrFN₆O₃ (557.33) C, 56.03; H, 2.53; N, 15.08; Found: C, 55.94; H, 2.45; N, 15.01.

Biological assay

Agar diffusion well method to determine the antimicrobial activity


The microorganism inoculums were uniformly spread using sterile cotton swab on a sterile Petri dish containing Malt extract agar (for fungi) and nutrient agar (for bacteria). Each sample (100 µl) was added to each well (6 mm diameter holes cut in the agar gel, 20 mm apart from one another). The systems were incubated for 24–48 h at 37 °C (for bacteria) and at 28 °C (for fungi). After incubation, microorganism growth was observed. Inhibition of the bacterial and fungal growth were measured in mm. Tests were performed in triplicate (Smania et al. 1999).

Results and discussion

The starting 7-(4-bromophenyl)-1,2-dihydro-5-(4-fluorophenyl)-2-thioxo-pyrido-[2,3-*d*] pyrimidin-4(3*H*)-one **3** was prepared by refluxing 6-amino-2-thiouracil **2** and 3-(4-fluorophenyl)-1-(4-bromophenyl)-2-propen-1-one **1** in dimethylformamide according to the procedure reported previously (Quiroga et al. 1992) (Scheme 1). The structure of compound **3** was established on the bases of spectral data (Ms, IR, ¹H NMR and ¹³C NMR) and elemental analysis (see Experimental part).

Reaction of compound **3** with hydrazonoyl chlorides **4a–d** in dioxane in the presence of triethylamine was yielded one isolable product in each case. Both spectroscopic

Scheme 1 8-(4-Bromophenyl)-6-(4-fluorophenyl)-1,3-aryl-pyrido[2,3-*d*][1,2,4]triazolo-[4,3-*a*]pyrimidin-5-one **7**



R: a, Ph; b, COCH₃; c, COOEt; d, CONHPh

data and elemental analyses were consistent with structure **7** or **8** (Scheme 1). An immediate distinction between these two structures was reached by comparison of the ¹³C NMR spectra with those of similar annulated pyrimidinones. Literature report (Reiter et al. 1987) has shown that the chemical shift for the carbonyl carbon in 4-pyrimidinone derivatives is markedly affected by the nature of the adjacent nitrogen (N3) (pyrrole type in structure **7** and pyridine type as in structure **8**). For example, the ¹³C NMR spectra of **7b** and **7c** taken as typical examples of the series prepared, revealed the signals of the carbonyl carbon of the pyrimidinone ring residue at δ 163.15 and 163.62 ppm, respectively. Such chemical shift values are similar to those of annulated pyrimidines with N3 pyrrole type rather than those of N3 pyridine type. On the basis of this similarity, the

isolated products were assigned structure **7** and the isomeric structure **8** was excluded.

As depicted in Scheme 1, the reaction proceeded through *S*-alkylation (Geis et al. 1991), to give *S*-alkylated products **5** followed by Smiles rearrangement (Elliott et al. 1975), afforded intermediate **6** which consumed in situ via elimination of hydrogen sulfide gas to give the desired products **7a–d**. The assignment for the structure products and reaction mechanism can be manifested by alternate synthesis. Thus, refluxing of methylthio-derivative **9** (prepared by the reaction of **3** with methyl iodide in dimethylformamide in presence of anhydrous potassium carbonate) with hydrazonoyl chloride **4a** under the same reaction conditions afforded product **7a** through the intermediate **10** with concurrent elimination of methanethiol.

The product **7a** found to be identical in all respects (mp, mixed mp and IR) with the product produced from the reaction of **3** with **4a** (Scheme 2).

Reacting compound **3** or **9** with hydrazine hydrate in absolute ethanol afforded 2-hydrazino-pyrido[2,3-*d*]pyrimidin-4(3*H*)-one **11**. The structure of **11** was evidenced by its spectra and elemental analysis (see experimental section). Condensation of the hydrazine derivative **11** with appropriate aldehydes afforded the required 2-(*N*-arylidene-hydrazino)-7-(4-bromophenyl)-5-(4-fluorophenyl)-3*H*-pyrido[2,3-*d*]pyrimidin-4-one derivatives **12** (Scheme 3). All such hydrazones have not been reported hitherto. Their structures were confirmed by their elemental analyses and spectral (Ms, IR and ¹H NMR) data. For example, their ¹H NMR spectra revealed in each case, a characteristic singlet signal in region 7.41–8.33 ppm assignable to the –N=CH– proton. Their IR spectra showed the characteristic band for the N–H stretch of the hydrazone group in region 3,375–3,100 cm^{–1}. The conversion of **12** into **13** is reminiscent of the oxidative cycloaddition of aldehyde *N*-heteroaryl hydrazone with bromine, which have been reported to proceed via generation of respective nitilimines, with undergo in situ 1,5-


electrocyclization to give the respective fused heterocycles (Shawali 2010).

Thus, in our hand, treatment of hydrazone **12** with equiv. quantity of bromine in acetic acid in the presence of sodium acetate gave, in each case, a single product as evidenced by TLC analysis. Elemental analyses and mass spectra revealed that each such isolated product has two hydrogens less than the respective hydrazone. This finding was confirmed by ¹H NMR spectra, which indicated the absence of the –N=CH– proton. On the basis of this finding the isolated products were pyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]pyrimidin-5(1*H*)-one derivatives **13**. The assigned structure of compounds **13** was further confirmed by chemical transformation. Thus, reaction of **11** with aroyl chloride in dry pyridine led to formation of products which are identical in all respects (mp, mixed mp and IR) with the products produced from the reaction of **12** with bromine/acetic acid (Scheme 3).


Antimicrobial activity

In vitro antimicrobial screening of compounds prepared in this study was carried out using cultures of four fungal

Scheme 2 Alternative synthesis of compound **7**



Scheme 3 Synthesis of compound **13**



strains, including *Aspergillus fumigatus* (RCMB 002003) (AF), *Syncephalastrum racemosum* (RCMB 005003) (SR), *Geotrichum candidum* (RB 052006) (GC), and *Candida albicans* (RCMB 005002) (CA) as well as four bacteria species, namely, Gram positive Bacteria, *S. aureus* (RCMB 000106) (SA) and *Bacillus subtilis* (RCMB 000107) (BS), Gram negative Bacteria, *Pseudomonas aeruginosa* (RCMB 000102) (PA) and *Escherichia coli* (RCMB 000103) (EC). Amphotericin B as an antifungal agent, Ampicillin as an antibacterial agent for gram (+) bacteria and Gentamicin as an antibacterial agent for gram (–) bacteria were used as references to evaluate the potency of the tested compounds under the same conditions. Most of newly synthesized compounds showed excellent with respect to control drugs. The results of antimicrobial activities were shown in Table 1. Data in Table 1 revealed that most of compounds have superior significant antibacterial potency to antifungal

potency. Compounds **3**, **7a**, **9**, **11**, **12a** and **13c** exhibited the highest potency against all tested organisms with respect to reference drugs except SR and PA. The other derivatives showed moderate activity against the microorganisms used.

Minimum inhibitory concentration (MIC)

The minimum inhibitory concentration (MIC) of the synthesized compounds against highly inhibited organism is reported in Table 2.

Compound **7a** revealed high MIC 500 µg/ml against *Aspergillus fumigatus* (RCMB 002003), *Candida albicans* (RCMB 005002) and *Escherichia coli* (RCMB 000103) respectively. Compound **13c** exhibited low MIC 0.12 µg/ml against *Aspergillus fumigatus* (RCMB 002003) and also 0.24 µg/ml against *S. aureus* (RCMB 000106).

Table 1 Antimicrobial activity expressed as inhibition diameter zones in millimeter (mm) of chemical compounds against the pathological strains based on well diffusion as assay

Compound no.	Fungi				Gram positive bacteria		Gram negative bacteria	
	<i>A. fumigatus</i>	<i>S. racemosum</i>	<i>G. candidum</i>	<i>Candida albicans</i>	<i>Staphylococcus aureus</i>	<i>B. subtilis</i>	<i>Pseudomonas aeruginosa</i>	<i>E. coli</i>
3	18.7	N.A.	19.3	15.2	16.2	19.4	N.A.	16.2
7a	21.9	N.A.	24.3	19.3	20.4	22.8	N.A.	18.9
7b	9.3	N.A.	10.4	8.4	10.0	12.3	N.A.	7.4
7c	N.A.	N.A.	N.A.	N.A.	10.2	12.1	N.A.	6.4
7d	16.2	N.A.	17.6	16.0	17.3	18.7	N.A.	17.9
9	18.9	N.A.	19.5	15.9	16.8	19.7	N.A.	17.4
11	19.0	N.A.	19.8	16.3	17.1	19.9	N.A.	18.2
12a	19.9	N.A.	21.3	18.2	20.0	21.2	N.A.	17.2
12b	12.8	N.A.	14.2	11.7	15.8	18.6	N.A.	12.3
12c	17.2	N.A.	19.8	16.1	18.8	20.3	N.A.	18.0
13a	15.4	N.A.	18.3	13.5	17.8	19.3	N.A.	15.7
13b	11.3	N.A.	12.8	10.2	13.4	15.3	N.A.	9.1
13c	21.3	N.A.	26.4	24.2	22.0	21.4	N.A.	18.9
Amphotericin B	23.7	19.7	28.7	25.4	N.A.	N.A.	N.A.	N.A.
Ampicillin	N.A.	N.A.	N.A.	N.A.	23.8	32.4	N.A.	N.A.
Gentamicin	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	17.3	19.9

The experiment was carried out in triplicate and average zone of inhibition was calculated; (N.A. = no activity)

Table 2 Minimum inhibitory concentration ($\mu\text{g/ml}$) against the pathological strains

Compound no.	Fungi				Gram positive bacteria		Gram negative bacteria	
	<i>A. fumigatus</i>	<i>S. racemosum</i>	<i>G. candidum</i>	<i>Candida albicans</i>	<i>Staphylococcus aureus</i>	<i>B. subtilis</i>	<i>Pseudomonas aeruginosa</i>	<i>E. coli</i>
7a	15.63	N.A.	62.5	62.5	62.5	7.82	N.A.	15.63
11	500	N.A.	250	500	250	125	N.A.	500
12a	7.82	N.A.	3.91	15.63	3.91	3.91	N.A.	31.25
12c	250	N.A.	125	250	125	62.5	N.A.	500
13c	0.12	N.A.	0.04	5.63	0.24	0.02	N.A.	2.01
Amphotericin B	0.12	1.95	0.03	0.06	N.A.	N.A.	N.A.	N.A.
Ampicillin	N.A.	N.A.	N.A.	N.A.	0.24	0.015	N.A.	N.A.
Gentamicin	N.A.	N.A.	N.A.	N.A.	N.A.	N.A.	7.82	1.95

Structure activity relationship

From the results depicted in Tables 1 and 2 we found that the parent pyridopyrimidine derivative **3** showed promising activity against the tested organisms. This encouraged derivatization of compound **3** hopping to improve the antimicrobial effect. Here, the chemistry represented two classes, the first one constituted pyridopyrimidine nucleus linked to different Schiff bases **12a–c**, while the second class the Schiff base included in linear cyclic structure forming triazolopyridopyrimidines **7a–d**, **13a–c**. They are tested against the previously mentioned microorganisms comparatively with the parent compound **3** and standard

drug for each microbial class. Recorded data clarified that, the importance of hydrophobic moieties for the antimicrobial activity. Replacing the $-\text{SH}$ group in compound **3** with $-\text{SMe}$ or $-\text{NHNH}_2$ groups in compounds **9** and **11** respectively, produced positive impact on the activity. On the other hand, Schiff base bicyclic systems **12a–c** showed negative effect on the activity except compound **12a**. While in triazolopyridopyrimidine system, direct linking of hydrophobic phenyl moiety at C3 in **7a** explored reliable antimicrobial activity that was dramatically diminished upon introducing CONH moiety as spacer in **7d**. The activity was almost abolished in **7b, c** upon replacing the phenyl group with COCH_3 and COOEt in **7b, c**

respectively. Moreover, compounds **13a–c** showed the vital role of *p*-NO₂ group in **13c**. Compound **13c** displayed strong antimicrobial activity which was diminished in phenyl and *p*-F phenyl analogues **13a, b**. Compared to the reference drug, triazolo derivatives **7a** and **13c** were the most effective compounds. They explored similar antimicrobial activity proving that substitution at C3 dramatically affects the activity rather than at N1. Compound **13c** carries H atom at N1 which was replaced by phenyl group in **7a**. Simultaneously, substituent at C3 in **7a** was phenyl group that replaced by *p*-NO₂phenyl group in **13c**. All the tested compounds are fully inactive against SR and PA organism.

Conclusion

In conclusion, we reported herein a simple and convenient routes for the synthesis of new derivatives of pyrido[2,3-*d*][1,2,4]triazolo[4,3-*a*]pyrimidin-5-one via two different methods. All the newly synthesized compounds were evaluated for antimicrobial activity and the result showed that compounds **3, 7a, 9, 11, 12a** and **13c** have excellent activity against all the tested microorganisms except SR and PA. The most effective derivative was compound **13c** which includes pharmacologically active 4-NO₂, 4-F, and 4-Br groups attached to phenyl ring at position 3, 6 and 8.

References

- Abdel Hafez, N.A., T.A. Farghaly, M.A. Al-Omar, and M.M. Abdalla. 2010. Synthesis of bioactive polyheterocyclic ring systems as 5 α -reductase inhibitors. *European Journal of Medicinal Chemistry* 45: 4838–4844.
- Abu-Zied, KhM, A.B.A. El-Gazzar, and N.A. Hassan. 2008. Synthesis and reactions of some novel triazolo-, azolo-, tetrazolo-pyridopyrimidine and their nucleoside derivatives. *Journal of the Chinese Chemical Society* 55: 209–216.
- El-Gazzar, A.B.A., and H.N. Hafez. 2008. A Simple synthesis and antimicrobial activity of sulfur-containing poly-condensed heterocyclic derivatives from 1,3-benzothiazole. *Acta Chimica Slovenica* 55: 359–371.
- El-Gazzar, B.A., H.N. Hafez, and G.A.M. Nawwar. 2009a. New acyclic nucleosides analogues as potential analgesic, anti-inflammatory, anti-oxidant and anti-microbial derived from pyrimido[4,5-*b*]quinolines. *European Journal of Medicinal Chemistry* 44: 1427–1436.
- El-Gazzar, A.B.A., M.M. Youssef, A.M.S. Youssef, A.A. Abu-Hashem, and F.A. Badria. 2009b. Design and synthesis of azolopyrimidoquinolines, pyrimidoquinazolines as anti-oxidant, anti-inflammatory and analgesic activities. *European Journal of Medicinal Chemistry* 44: 609–624.
- Elliott, A.J., P.D. Callaghan, M.S. Gibson, and S.T. Nemeth. 1975. Rearrangement of thiohydrazonates. *Canadian Journal of Chemistry* 53: 1484–1490.
- Farghaly, T.A., and M.M. Abdalla. 2009. Synthesis, tautomerism, and antimicrobial, anti-HCV, anti-SSPE, antioxidant, and antitumor activities of arylazobenzosuberones. *Bioorganic and Medicinal Chemistry* 17: 8012–8019.
- Farghaly, T.A., N.A. Abdel Hafez, E.A. Ragab, H.M. Awad, and M.M. Abdalla. 2010. Synthesis, anti-HCV, antioxidant, and peroxyinitrite inhibitory activity of fused benzosuberone derivatives. *European Journal of Medicinal Chemistry* 45: 492–500.
- Farghaly, T.A., S.M. Gomha, E.M. Abbas, and M.M. Abdalla. 2012. Hydrazonoyl halides as precursors for new fused heterocycles of 5 α -reductase inhibitors. *Archiv der Pharmazie* 345: 117–122.
- Geis, A.A., A.M. Kamal-Eldeen, A.A. Abdelhafez, and A.M. Gaber. 1991. Synthesis of some thiazolo(3,2-*a*) pyrimidines. *Phosphorus, Sulfur and Silicon and related elements* 56: 87–93.
- Gomha, S.M., and H.M.E. Hassaneen. 2011. Synthesis and antimicrobial activity of some new pyrazoles, fused pyrazolo[3,4-*d*]pyrimidine and 1,2-dihydroimidazo-[2,1-*c*][1,2,4]triazin-6-one derivatives. *Molecules* 16: 6549–6560.
- Hassaneen, H.M., H.M.E. Hassaneen, Y.S. Mohammed, and R.M. Pagni. 2011. Synthesis, reactions and antibacterial activity of 3-acetyl[1,2,4]triazolo[3,4-*a*]isoquinoline derivatives using chitosan as heterogeneous catalyst under microwave irradiation. *Zeitschrift für Naturforschung* 66b: 299–310.
- Leclercq, R. 2009. Epidemiological and resistance issues in multi-drug-resistant Staphylococci and Enterococci. *Clinical Microbiology and Infection* 15: 224–231.
- Quiroga, J., B. Insuasty, B. Sánchez, M. Noguera, and H. Meier. 1992. Synthesis of Pyrido[2,3-*d*]pyrimidines in the reaction of 6-Amino-2,3-dihydro-2-thioxo-4(1*H*)-pyrimidinone with Chalcones. *Journal of Heterocyclic Chemistry* 29: 1045–1048.
- Reiter, J., L. Bongo, and P. Dyortsok. 1987. On triazoles XI. Structure elucidation of isomeric 1,2,4-triazolopyrimidinones. *Tetrahedron* 43: 2497–2504.
- Riyadh, S.M., T.A. Farghaly, M.A. Abdallah, M.M. Abdalla, and M.R. Abd El-Aziz. 2010. New pyrazoles incorporating pyrazolylpyrazole moiety: synthesis, anti-HCV and antitumor activity. *European Journal of Medicinal Chemistry* 45: 1042–1055.
- Shawali, A. S.(2010) Tandem in situ generation and 1,5-electrocyclization of *N*-hetaryl nitrilimines. A facile methodology for synthesis of annulated 1,2,4-triazoles and their acyclo C-nucleosides *Arkivoc*, (i), 33–97.
- Smania, J.A., F.D. Monache, E.F.A. Smania, and R.S. Cuneo. 1999. Antibacterial activity of steroidal compounds isolated from *Ganoderma applanatum* (Pers.) Pat. (Aphyllphoromycetidae) Fruit Body. *International Journal of medicinal mushrooms* 1: 325–330.