

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/244230848>

A Practical and Eco-Friendly Synthesis of Stereocontrolled Alkylaminomethylidene Derivatives of 2-Thiohydantoins by Dimethylamine Substitution.

ARTICLE *in* TETRAHEDRON LETTERS · NOVEMBER 2002

Impact Factor: 2.38 · DOI: 10.1016/S0040-4039(02)02156-1

CITATIONS

12

READS

22

3 AUTHORS:



Jean-rene Cherouvrier

Université de La Rochelle

12 PUBLICATIONS 174 CITATIONS

SEE PROFILE



François Carreaux

Université de Rennes 1

89 PUBLICATIONS 1,054 CITATIONS

SEE PROFILE



Jean Pierre Bazureau

Université de Rennes 1

108 PUBLICATIONS 1,666 CITATIONS

SEE PROFILE



A practical and eco-friendly synthesis of stereocontrolled alkylaminomethylidene derivatives of 2-thiohydantoins by dimethylamine substitution

Jean-René Chérouvrier, François Carreaux and Jean Pierre Bazureau*

Université de Rennes 1, Institut de Chimie, Synthèse et Electrosynthèse Organiques 3, UMR 6510, Bât. 10A, Campus de Beaulieu, Avenue du Général Leclerc, CS 74205, 35042 Rennes Cedex, France


Received 29 August 2002; accepted 24 September 2002

Abstract—3-Alkyl-5-dimethylaminomethylidene-2-thioxo-imidazolidin-4-ones **3(a–c)** available in two steps from methyl glycinate hydrochloride, represent a useful synthetic tool for efficient and mild solventless preparations of new alkylaminomethylidene derivatives of 2-thiohydantoins **8(a–e)**, **10(a–c)** and **12(a–d)** by stereocontrolled transamination reactions under microwave irradiations. The ^1H , ^{13}C NMR spectrum and the (5*Z*)-conformation of some representatives products are also discussed. © 2002 Elsevier Science Ltd. All rights reserved.

Derivatives of 2-thiohydantoins play an important role in organic synthesis, especially as starting materials for the preparation of synthetic intermediates with a wide range of applications as therapeutics¹ as well as fungicides and herbicides.² Among these compounds the S-glucosylated hydantoins³ exhibit properties against the herpes simplex virus⁴ (HSV), the human immunodeficiency virus⁵ (HIV). Recently, a series of S-alkylated 4-ylidene thiohydantoins have

been prepared in our laboratory as versatile reagents for the synthesis of marine alkaloid 2-amino imidazolones⁶ derivatives for investigations of protein kinase C inhibition activities,⁷ using solventless Knoevenagel reaction conditions under microwave irradiations.

Owing to economic and ecological reasons, organic synthetic chemists in pharmaceutical industry face an



Scheme 1. Reagents and conditions: (i) TEA 1 equiv., R¹NCS 1 equiv., Et₂O or AcOEt, reflux, 15 h; (ii) DMF–DEA 1.05 equiv., $\mu\omega$ (in the Synthewave[®] 402 reactor), for **3a**: R¹ = Me, 70°C, 15–30 min (74%), for **3b**: R¹ = Bu, 80°C, 45 min (77%), for **3c**: Ph, 70°C, 30 min (75%); (iii) K₂CO₃ 0.55 equiv., EtI 1.1 equiv., 65°C, MeCN, 14 h; (iv) DMF–DEA 1 equiv., 70°C, 1 h.

Keywords: transamination reactions; dimethylamine substitutions; microwaves; thiohydantoins; alkylaminomethylidene-2-thioxo-imidazolidin-4-ones.

* Corresponding author. Fax: +(33) 02 23 23 63 74; e-mail: jean-pierre.bazureau@univ-rennes1.fr

increasing obligation to optimize the quantities of volatile organic solvents (VOCs) and toxic waste in chemical processes.⁸ Thus, the development of solvent-free organic synthesis under microwaves has received much attention.⁹

In the course of identifying new chemical structures derived from 2-thiohydantoin for their biological activities,¹⁰ we were interested to develop a new route to alkylamino derivatives of 2-thiohydantoin towards new, simple and efficient procedures.

Scheme 1 shows the route for the preparation of 3-alkyl-5-dimethylaminomethylene-2-thioxo-imidazolidin-4-ones **3(a–c)**. In the first step, the 3-substituted-2-thioxo-imidazolidin-4-ones **2(a–c)** ($R^1 = \text{Me, Bu, Ph}$) were readily available in large scale (up to 20 g) with good yields ($\sim 96\%$) by addition of commercial isothiocyanates to methyl glycinate hydrochloride in basic medium.⁶ For the second step, we have investigated the reactivities of 2-thioxo-imidazolidin-4-ones derivatives **2(a–c)** with *N,N*-dimethylformamide diethylacetal¹¹ (DMF–DEA) using solvent-free conditions under microwave irradiations. The microwave instrument (Synthewave^{  } 402 reactor¹²) comprises a monomode (sometimes also called single-mode) microwave cavity that operates at a frequency of 2.45 GHz with continuous microwave irradiation power from 0 to 300 W.

The reaction vial is a cylindrical quartz reactor ($\varnothing = 4$ cm) which was introduced into the Synthewave^{  } 402 microwave reactor. Inside the microwave cavity the vial was exposed to microwave irradiations. The temperature was measured with an IR captor¹³ (infrared thermometry). The software algorithm regulates the microwave output power so that the preselected maximum temperature is maintained for the desired reaction/irradiation time. After the irradiation period, the reaction vial is cooled rapidly to ambient temperature by compressed air (gas jet cooling).


The 2-thioxo-imidazolidin-4-ones **2(a–c)** were converted with DMF–DEA (1.05 equiv.) into the corresponding 5-dimethylaminomethylene-2-thioxo-imidazolidin-4-ones **3(a–c)** in yields ranging from 74 to 77% after a reaction time of ~ 30 minutes at $70\text{--}80^\circ\text{C}$ under

microwave irradiations.¹⁴ From **2c** ($R^1 = \text{Ph}$) we have also observed that ethylation at the exocyclic sulfur¹⁵ took place to give 5-dimethylaminomethylene-3-phenyl-2-ethylsulfanyl-3*H*-imidazol-4(5*H*)-one **5c** in $\sim 10\%$ yield via the intermediate **4c** which could not be isolated (Scheme 1). The structure of **5c** was confirmed by S-alkylation¹⁶ (with ethyliodide) of **2c** ($R^1 = \text{Ph}$) in basic medium (K_2CO_3 0.55 equiv.) which provided 2-ethylsulfanyl-3-phenyl-3,5-dihydro imidazol-4-one **6c** in 75% yield, then **6c** was transformed with DMF–DEA (1 equiv.) at 70°C into **5c** in 95% yield after 1 hour. The expected compounds **3(a–c)**, **5c** and **6c** were purified by recrystallization.

The 5-dimethylaminomethylene-3-substituted-2-thioxo-imidazolidin-4-ones **3(a–c)** and compound **5c** can exist in (5*Z*) and/or (5*E*) isomeric forms with respect to the exocyclic C=C double bond. In all cases, the compounds **3(a–c)** exist as single isomers, as shown by the presence of only one set of signals in each of ^1H and ^{13}C NMR, but differentiation between (*Z*)- and (*E*)-form is not possible on the basis of chemical shifts. However, the two isomeric forms are easily differentiated on the basis of the magnitude of the long range heteronuclear $^{13}\text{C}\text{--}^1\text{H}$ coupling constants, $^3J_{\text{CH}}$ which have been used for determination of configuration in various systems.¹⁷ Generally, the magnitude of coupling constant for *cis*-configuration around the C=C double bond is smaller (2–6 Hz) than for *trans*-oriented nuclei (8–12 Hz). In the case of compound **3a**¹⁸ ($R^1 = \text{Me}$), the magnitude of coupling constant $^3J_{\text{CH}} = 3.4$ Hz showed that **3a** exist in the (*Z*) form.

With compounds **3(a–c)** in hand, we then studied their reactivity in transamination reactions¹⁹ with various primary aliphatic amines **7(a–c)** using solvent-free technique under microwave irradiations²⁰ (Scheme 2). Several experiments were performed with **3a** ($R^1 = \text{Me}$), at various powers and irradiations times, with an excess of amine **7** (2–10 equiv.) in order to find the most adequate reaction conditions under microwave. The optimized reaction conditions were summarized in Table 1.


The expected compounds **8(a–e)** were isolated from the crude reaction mixture (after elimination of the excess



Scheme 2. Reagents and conditions: (i) **7** 2–10 equiv., $\mu\omega$ (in the Synthewave^{  } 402 reactor), 15–30 min, $50\text{--}70^\circ\text{C}$; (ii) **9** 5 equiv., $\mu\omega$, 80°C , 30 min; (iii) **11** 1 equiv., MeCO_2H , $\mu\omega$, 95°C , 90 min.

Table 1. 5-Alkylaminomethylidene-3-methyl-2-thioxo-imidazolidin-4-ones **8(a–e)** and **10(a–c)** prepared by transamination reactions from **3a** ($R^1 = \text{Me}$), primary amines **7(a–e)** and cycloalkylamines **9(a–c)**

Amine 7 or 9	Ratio 3a /amine 7 or 9	Product 8 or 10	Reaction conditions ^a		Yield of 8 or 10 (%) ^b
			Reaction time (min)	Temperature (�C)	
7a	1:10	8a	15	50	88
7b	1:10	8b	60 + 30	70	68
7c	1:4	8c	30	70	67
7d	1:2	8d	30	70	56
7e	1:5	8e	30	60	58
9a	1:5	10a	30	80	63
9b	1:5	10b	30	80	53
9c	1:5	10c	30	80	72



^a The reactions were performed under microwave irradiation in the Synthewave[ ] 402 reactor.

^b Isolated yield of **8** or **10**.

of amine **7** and dimethylamine in vacuo) by crystallization and repeated washings with ether or pentane in yields ranging from 56 to 88% (Table 1). The structure of the new 5-alkylaminomethylidene-2-thioxo-imidazolidin-4-ones **8(a–e)** were substantiated by the ¹H, ¹³C and HRMS analyses.

A characteristic feature of the ¹H NMR spectra of 5-alkylaminomethylidene-3-methyl-2-thioxo-imidazolidin-4-ones **8(a–e)** is the downfield shift of the doublet for the exocyclic C=CH double bond (**8a**: $\delta_{\text{vinyl}} = 6.93$ ppm). The coupling constant $^3J = 13.2$ Hz between the aminoproton NH and exocyclic vinyl proton CH suggests the *trans* (antiperiplanar) orientation²¹ of these hydrogens. Furthermore, the shift of H-1 (NH) has been found at low field (**8a**: $\delta_{\text{H-1}} = 11.1$ ppm). The shift of the CH signal for the exocyclic double bond C=CH is confirmed in the ¹H resonance-coupled ¹³C NMR spectra by identification of a doublet centered at $\delta_{\text{CH}} = 130.80$ ppm for **8a**²² ($J = 173$ Hz).

In order to define the ability of the 5-dimethylaminomethylidene-2-thioxo-imidazolidin-4-ones **3** for transamination reaction, we have also evaluated the reactivity of **3a** ($R^1 = \text{Me}$) with secondary amines **9(a–c)** using the same reaction conditions (Scheme 2). For this study, the cyclic secondary amines employed were, i.e. pyrrolidine **9a**, piperidine **9b** and morpholine **9c**. The results obtained and the isolated yields of the new 5-cycloalkylaminomethylidene-3-methyl-2-thioxo-imidazolidin-4-ones **10(a–c)** are given in Table 1. The expected compounds **10(a–c)** were produced in yields ranging from 53 to 72% and required a reaction time of 30 minutes at 80 C under microwave irradiations.

During the course of our work, we have also found that the dimethylamino group in compounds **3(a,b)** can formally be substituted with N-nucleophiles derived from aminoester hydrochlorides²³ **11(a,b)**. The following N-nucleophiles used were methyl glycinate hydrochloride **11a** ($n = 0$), and methyl 4-aminobutanoate hydrochloride **11b** ($n = 2$). They were treated with an equimolar amount of **3** in glacial acetic acid at 95 C under microwave irradiations. After heating for 90 minutes, derivatives of methyl 4-[(1-alkyl-5-oxo-2-thioxo-imidazolidin-4-ylidenemethyl)amino]acetate **12(a,b)**²⁴ and methyl 4-[(1-alkyl-5-oxo-2-thioxo-imidazolidin-4-ylidenemethyl)amino]butanoate **12(c,d)** were isolated in moderate yield (14–47%) (Table 2).



In summary, the major significance of these results is the development of a straightforward access to 5-yliden-3-alkyl-2-thioxo-imidazolidin-4-ones²⁵ using the eco-friendly solventless methodology assisted by microwave²⁶ heating. From the 2-thiohydantoin **3(a–c)**, the transamination reactions gave good results with N-nucleophiles derived from non-sterically hindered primary amines **7(a–c)**, cyclic secondary amines **9(a–c)** and moderate yields with aminoesters **11(a,b)**. Work is now in progress to study the biological potentialities²⁷ of these new 5-alkylaminomethylidene-3-alkyl-2-thioxo-imidazolidin-4-ones.

Acknowledgements

One of us (J.R.C.) wishes to thank the ‘Conseil R  gional de Bretagne’ for a research fellowship (contrat N  99CBQ-4). We also thank Professor Jack Hamelin for fruitful discussions.

Table 2. Preparation of methyl 4-[(1-alkyl-5-oxo-2-thioxo-imidazolidin-4-ylidenemethyl)-amino]acetate **12(a,b)** and butanoate **12(c,d)** from **3(a,b)** and aminoesters **11(a,b)**

Aminoester 11	<i>n</i>	Product 12	R ¹	Reaction conditions ^a		Yield of 12 (%) ^b
				Time (h)	Temp. (�C)	
11a	0	12a	Me	1.5	95	16
11a	0	12b	Bu	1.5	95	47
11b	2	12c	Me	1.5	95	40
11b	2	12d	Bu	1.5	95	30



^a The reactions were performed under microwave irradiations in the Synthwave[ ] 402 reactor.

^b Isolated yield.

References

- (a) Mehat, N.; Risinger, C. A.; Soroko, F. E. *J. Med. Chem.* **1981**, *24*, 465; (b) Wessels, F. L.; Schwan, T. J.; Pong, S. F. *J. Pharm. Sci.* **1980**, *69*, 1102; (c) Cadwell, A. G.; Harris, C. J.; Stepney, R.; Wittaker, N. *J. Chem. Soc., Perkin. Trans. 1* **1980**, 495.
- Cremlyn, R. J.; Elias, R. S.; Geoghagan, M. J. A.; Braunholtz, J. L. Brit. 1964,166,967. *Chem. Abstr.* **1965**, *62*, 7768g.
- Khodair, A. I. *Carbohydr. Res.* **2001**, *331*, 445.
- El-Barbary, A. A.; Khodair, A. I.; Pedersen, E. B.; Nielsen, C. *J. Med. Chem.* **1994**, *37*, 73.
- Khodair, A. I.; El-Subbagh, H. I.; El-Eman, A. A. *Bull. Soc. Chim. Farm.* **1997**, *136*, 561.
- Ch  rouvrier, J. R.; Carreaux, F.; Bazureau, J. P. *Tetrahedron Lett.* **2002**, *43*, 3581.
- Borgne, A.; Meijer, L. *Med. Sci.* **1999**, *4*, 496.
- Green Chemistry: Challenging Perspectives*, P. Tundo, P.T. Anastas (eds), Oxford University Press, Oxford 1999.
- (a) Wathey, B.; Tierney, J.; Lidstr  m, P.; Westman, J. *Drug Discovery Today* **2002**, *7*, 373; (b) Lidstr  m, P.; Tierney, J.; Wathey, B.; Westman, J. *Tetrahedron* **2001**, *57*, 9225; (c) Varma, R. S. *Green Chem.* **1999**, *1*, 43; (d) Loupy, A.; Petit, A.; Hamelin, J.; Texier-Boullet, F.; Jacquault, P.; Math  , D. *Synthesis* **1998**, 1213; (e) Cad-dick, S. *Tetrahedron* **1995**, *51*, 10403.
- (a) Szymanska, E.; Kiec-Kononowicz, K.; Bialecka, A.; Kasproicz, A. *Farmaco* **2002**, *57*, 39; (b) Szymanska, E.; Kiec-Kononowicz, K. *Il Farm.* **2002**, *57*, 355.
- Abdulla, R. F.; Brinkmeyer, R. S. *Tetrahedron* **1979**, *35*, 1675.
- (a) Commarmot, R.; Didenot, R.; Gardais, J. F.; *Fr Demande*, 25 560 529, 1985, *Chem. Abstr.* **1986**, *105*, 17442; (b) For description of commercial microwave devices available with adequate mixing and control of reaction parameters, see sites: <http://www.cem.com> and <http://www.personalchemistry.com>
- Temperature measured by an IR captor: Prolabo, French Patent 622 410, 14669 Fr, 1991.
- (a) Dahmani, Z.; Rahmouni, M.; Brugidou, R.; Bazureau, J. P.; Hamelin, J. *Tetrahedron Lett.* **1998**, *39*, 8453; (b) M  ziane, M. A  t; Amer, M.; Rahmouni, M.; Bazureau, J. P.; Hamelin, J. *Synthesis* **1998**, 967.
- Jokse, R.; Reenik, S.; Svete, J.; Golobic, A.; Galic, L.; Stanovnik, B. *Tetrahedron* **2001**, *57*, 8395.
- Ch  rouvrier, J. R.; Boissel, J.; Carreaux, F.; Bazureau, J. P. *Green Chem.* **2001**, *3*, 165.
- (a) Fischer, P.; Schweizer, E.; Langner, J.; Schmidt, U. *Magn. Reson. Chem.* **1994**, *32*, 567; (b) Golic Grdadolnik, S.; Stanovnik, B. *Magn. Reson. Chem.* **1997**, *35*, 482; (c) Ando Koseki, M.; Toia, R. F.; Casida, J. E. *Magn. Reson. Chem.* **1999**, *31*, 90.
- Selected spectral data of 5-dimethylaminomethylidene-3-methyl-2-thioxo-imidazolidin-4-one (**3a**): ¹H NMR (300 MHz, DMSO-*d*₆, TMS)   3.09 (s, 3H), 3.11 (s, 3H), 6.79 (s, 1H, CH=), 11.10 (br s, 1H, NH). ¹³C NMR (75 MHz, DMSO-*d*₆, TMS)   26.90 (q, *J*=141 Hz), 42.40 (q, *J*=158 Hz), 102.00 (d, *J*=7 Hz, C-5), 132.60 (dt, *J*=170, 3.4 Hz, CH=), 163.50 (dd, *J*=3.7, 2.4 Hz, C-4, C=O), 169.90 (q, *J*=3.7 Hz, C-2, C=S). HRMS, *m/z*=185.0623 found (calculated for C₇H₁₁N₃OS requires 185.0684). mp=245–246 C.
- For reviews see: (a) Stanovnik, B.; Svete, J.; *Synlett* **2000**, *8*, 1077; (b) Stanovnik, B. *Molecules* **1999**, *1*, 123.
- (a) Meddad, N.; Rahmouni, M.; D  r  ur, A.; Bazureau, J. P.; Hamelin, J. *Synthesis* **2001**, 581; (b) Brugidou, R.; Bazureau, J. P.; Hamelin, J.; Dahmani, Z.; Rahmouni, M. *Heteroatom Chem.* **1999**, *10*, 446.
- Tagaki, K.; Aotsuka, T.; Morita, H.; Okamoto, Y. *J. Heterocycl. Chem.* **1986**, *23*, 1443.
- Selected spectral data of 3-methyl-5-propylaminomethylidene-2-thioxo-imidazolidin-4-one (**8a**): ¹H NMR (300 MHz, DMSO-*d*₆, TMS)   0.86 (t, 3H, *J*=7 Hz), 1.48 (q, 2H, *J*=7 Hz), 3.06 (s, 3H), 3.24 (d, 2H, *J*=4.9 Hz), 6.93 (d, 1H, *J*=13.2 Hz, CH=), 6.97 (br d, *J*=13.2 Hz, NH), 11.1 (br s, 1H, H-1). ¹³C NMR (75 MHz, DMSO-*d*₆, TMS)   10.70 (qt, *J*=125, 8.4 Hz), 23.60 (tm, *J*=127 Hz), 26.60 (q, *J*=141 Hz), 49.60 (tm, *J*=137 Hz), 103.30 (m, C-5), 130.80 (dm, *J*=173 Hz, CH), 162.10 (m, C-4, C=O), 168.60 (d, *J*=8.4 Hz, C-2, C=S). HRMS, *m/z*=199.0782 found (calculated for C₈H₁₃N₃OS requires 199.0779). mp>260 C.
- (a) Sorsak, G.; Grolic Grdadolnik, S.; Stanovnik, B. *ACH-Models in Chemistry* **1998**, *135*, 613; (b) Selic, L.; Grolic Grdadolnik, S.; Stanovnik, B. *Helv. Chim. Acta* **1997**, *80*, 2418.

24. Preparation of methyl 4-[(1-methyl-5-oxo-2-thioxo-imidazolidin-4-ylidenemethyl)-amino]acetate (**12a**): In the Synthewave[®] 402 microwave reactor ($\varnothing=4$ cm), an equimolar mixture of **3a** (1 g, 5.4 mmol.) and methyl glycinate hydrochloride **11a** (0.68 g, 5.54 mmol.) in glacial acetic acid (2 ml) was heated at 95°C under nitrogen with vigorous magnetic stirring during 1 hour under microwave irradiations. Then, the reaction mixture was allowed to cool down. After addition of MeOH (10 ml), the insoluble compound **12a** was filtered off, washed twice with Et₂O (10 ml) and dried in a dessicator over CaCl₂ which gave **12a** in 16% yield as white needles (mp>260°C). ¹H NMR (300 MHz, DMSO-*d*₆, TMS) δ 3.07 (s, 3H), 3.67 (s, 3H), 4.22 (d, 2H, $J=5.7$ Hz), 6.90 (d, 1H, $J=13.3$ Hz, CH=), 7.08–7.14 (m, 1H, NH), 11.33 (br s, 1H, NH). ¹³C NMR (75 MHz, DMSO-*d*₆, TMS) δ 26.60 (q, $J=141$ Hz), 48.4 (t, $J=141$ Hz), 51.90 (q, $J=147$ Hz), 104.40 (d, $J=5$ Hz, C-5), 130.10 (dm, $J=175$ Hz, CH), 162.50 (m, C-4), 169.70 (m, C=O), 170.50 (m, C-2, C=S). HRMS, $m/z=226.6524$ found (calculated for C₈H₁₁N₃O₃S requires 226.6521).
25. Part of this work was presented at ‘Le D  fi des Nouvelles Technologies en Chimie Mol  culaire’, Universit   de Rennes 1, Campus de Beaulieu, France, 15–18 Avril 2002, Poster Abstracts, P-4, see site <http://ntc2002.univ-rennes1.fr>
26. When the same reaction mixture was heated in an oil bath previously set at the same boiling point for the same reaction time, the yields were lower (**8a**: 85% in oil bath and 88% under microwave).
27. The new 5-alkylaminomethylidene-3-alkyl-2-thioxo-imidazolidin-4-ones **8**, **10** and **12** will be evaluated in a drug discovery program (protein kinase C inhibition activities) at the ‘Station Biologique de Roscoff, BP 74-29682 Roscoff Cedex, France’.