See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/51674273

# Eponym

# ARTICLE in EUROPEAN JOURNAL OF PEDIATRICS · SEPTEMBER 2011

Impact Factor: 1.89 · DOI: 10.1007/s00431-011-1575-6 · Source: PubMed

**CITATIONS** 

9

READS

34

# **7 AUTHORS**, INCLUDING:



# Atsuhito Takeda

Hokkaido University

24 PUBLICATIONS 95 CITATIONS

SEE PROFILE



# Hirokuni Yamazawa

Hokkaido University Hospital

17 PUBLICATIONS 38 CITATIONS

SEE PROFILE



# Akira Sudo

Sapporo City General Hospital

32 PUBLICATIONS 423 CITATIONS

SEE PROFILE



# Ichizo Nishino

National Center of Neurology and Psychiatry

**589** PUBLICATIONS **13,305** CITATIONS

SEE PROFILE

#### **REVIEW**

# **Eponym**

# Barth syndrome

Atsuhito Takeda • Akira Sudo • Masafumi Yamada • Hirokuni Yamazawa • Gaku Izumi • Ichizo Nishino • Tadashi Ariga

Received: 5 September 2011 / Accepted: 7 September 2011 / Published online: 23 September 2011 © Springer-Verlag 2011



Fig. 1 Dr. Peter G. Barth

**Abstract** Barth syndrome (OMIM #302060) (BTHS) is an X-linked disorder of lipid metabolism characterized by

A. Takeda ((() · M. Yamada · H. Yamazawa · T. Ariga Department of Pediatrics, Hokkaido University Graduate School of Medicine, North 15, West 7, Kita-ku, Sapporo 060-8638, Japan e-mail: a-takeda@med.hokudai.ac.jp

A. Sudo

Department of Pediatrics, Sapporo City General Hospital, 1-1, North 11, West 13, Chuo-ku, Sapporo 060-8604, Japan

G. Izumi Department of Pediatric Cardiology, Tokyo Women's Medical University, 8-1 Kawada-cho, Shinjukuku, Tokyo 162-8666, Japan

I. Nishino
National Institute of Neuroscience,
National Center of Neurology and Psychiatry,
4-1-1 Ogawa-Higashi, Kodaira,
Tokyo 187-8502, Japan

skeletal myopathy, neutropenia, growth delay, and cardiomyopathy. It is caused by mutations in the tafazzin gene (TAZ), which lead to decreased production of an enzyme required to produce cardiolipin, a component of the inner mitochondrial membrane necessary for proper functioning of the electron transport chain. The most common initial presentation of BTHS is significant heart failure due to cardiomyopathy, which is the main cause of death in infancy or childhood. On the other hand, some patients have limited clinical features of BTHS. These patients may be overlooked or misdiagnosed with unclassified congenital myopathy, especially when heart failure is not clinically significant. However, these patients could also develop significant heart failure or life-threatening arrhythmias during or even after childhood. Heart failure in BTHS is often responsive to standard medical therapy, indicating early diagnosis is critical. Diagnostic clues of BTHS in the subclinical stage of heart failure include family histories, findings of lipid storage myopathy in the skeletal muscle biopsy, and elevated plasma brain natriuretic peptide levels. The genetic analysis of TAZ is the only confirmatory method for the diagnosis of BTHS. Conclusion: physicians should be aware of the possibility of this disease and carry out genetic studies when it is considered.

**Keywords** Barth syndrome (BTHS) · Lipid storage myopathy · Brain natriuretic peptide (BNP) · Isolated non-compaction of the ventricular myocardium (INVM) · Cardiomyopathy

# Introduction and historical overview of the disorder

In 1983, Dr. Peter G. Barth (Fig. 1), a pediatric neurologist in the Netherlands, first described with his colleagues a large pedigree of X-linked recessive disorder characterized



by skeletal myopathy, neutropenia, growth delay, and cardiomyopathy [3]. The affected patients have high mortality during infancy from severe heart failure or serious bacterial infections owing to neutropenia. Abnormalities of ultrastructure and respiratory chain were observed in their postmortem mitochondria samples. In 1991, Kelley et al. reported seven boys with an X-linked syndrome of dilated cardiomyopathy, growth retardation, neutropenia, and persistently elevated urinary levels of 3-methylglutaconate [15]. Since then, this disorder has been referred to as Barth syndrome (BTHS). In 1996, Bione et al. reported the identification of mutations in Xq28-linked gene, termed G4.5 or TAZ, in five families with BTHS [6]. In 1997, Neuwald et al. reported that tafazzin, the protein products of TAZ, has a phospholipid acyltransferase function and seems to have an important role in remodeling cardiolipin, a component of the inner mitochondrial membrane necessary for proper functioning of the electron transport chain [4, 18]. So far, over 100 different disease-causing TAZ mutations have been reported in all exons.

#### Clinical presentation

BTHS is associated with a variety of clinical features, including cardiomyopathy, neutropenia, skeletal myopathy, and growth delay (Table 1). These features can change with age and vary significantly between patients.

Heart failure due to cardiomyopathy is the most common initial presentation and the main cause of death in infancy. Survivors beyond infancy could also develop significant heart failure or life-threatening arrhythmias during or even after childhood. Various types of cardiomyopathy have been reported including dilated cardiomyopathy, hypertrophic cardiomyopathy, endocardial fibroelastosis, and isolated non-compaction of the ventricular myocardium (INVM) [21]. Neutropenia may be present at birth and lead to neonatal sepsis, which is the second cause of death in BTHS. However, more common presentation is localized infections of the skin and mucosal membranes [1]. The histology of bone marrow in BTHS often shows a maturation arrest at the myelocyte stage [3].

Table 1 Clinical features in patients with Barth syndrome

Cardiomyopathy (dilated cardiomyopathy, endocardial fibroelastosis, and INVM)

Neutropenia (chronic, cyclic, or intermittent)

Skeletal myopathy (lipid storage myopathy)

Growth delay

3-Methylglutaconic aciduria (normal, <10 µmol/mmol creatinine)

INVM isolated non-compaction of the ventricular myocardium



Patients with BTHS may present with muscular hypotonia due to skeletal myopathy soon after birth and manifest delays in motor development within the first year [5]. However, muscular hypotonia usually improves and does not progress toward respiratory insufficiency or wheelchair dependence. Growth delay is also a common presentation in patients with BTHS, but nearly normal adult height may be achieved in the late teenage years.

#### Diagnostic clues and workup

Clinical diagnosis of BTHS can be made when there is a combination of clinical features as listed in Table 1. Most patients are diagnosed during infancy after the development of overt heart failure. In contrast, patients in the subclinical stage of heart failure are often overlooked or misdiagnosed even if other features of BTHS are noticed. There are several potential clues to the early diagnosis of BTHS, especially before the development of clinically significant heart failure. A positive family history of BTHS can be the most obvious clue. A family history of early death from cardiomyopathy in males also strongly suggests BTHS [9, 10, 24]. Skeletal muscle biopsy may be another clue to the diagnosis of BTHS; lipid storage myopathy is the characteristic pathological finding of BTHS [3], although the differential diagnoses of lipid storage myopathy alone are extensive [17]. Detection of 3-methylglutaconic aciduria may support the diagnosis of BTHS, although it is not always present.

Additionally, an elevated plasma brain natriuretic peptide level could be one of the most important clues to the early diagnosis of BTHS, especially when it is observed in patients with additional clinical features of BTHS. In such cases, morphological evaluation using echocardiography or cardiac magnetic resonance imaging could be the next step for the diagnosis of BTHS because INVM is the typical finding of BTHS [7, 13].

Thus far, the genetic analysis of *TAZ* is the only confirmatory method for the diagnosis of BTHS [8]. Recently, a tetralinoleoyl cardiolipin high-pressure liquid chromatography—tandem mass spectrometry blood test has become available [16]. Tetralinoleoyl cardiolipin, a metabolite of the acyltransferase that *TAZ* encodes, is significantly reduced in BTHS [12, 19, 20]. In the future, the application of this method to newborn screening will contribute to the early detection of BTHS.

## Genetic background

The causative mutations are in *TAZ* [6, 8], which is located at Xq28 and consists of 11 exons. This gene encodes the tafazzin protein, which has a phospholipid acyltransferase

function and seems to play an important role in remodeling cardiolipin, a component of the inner mitochondrial membrane [11]. So far, over 100 different disease-causing *TAZ* mutations including nonsense, frameshift, and missense mutations have been reported in all exons. There was no genotype—phenotype correlation in a study of 14 BTHS pedigrees [14].

#### **Treatment**

The treatment strategy depends on the clinical spectrum and severity of the disease. Heart failure in BTHS is mostly responsive to standard medical therapy, such as angiotensin-converting enzyme inhibitors, beta-blockers, and diuretics in a cross-sectional study of the large cohort (age range, 1.2–22.6 years) [21]. Successful heart transplantation was also reported in patients with severe heart failure, although the numbers are limited [2]. Placement of an internal cardiac defibrillator for life-threatening arrhythmias in BTHS was documented to prevent from sudden death [22]. Neutropenia can be usually managed with parenteral antibiotics and administration of granulocyte colony-stimulating factor [23]. Growth hormone therapy is discouraging for growth-retarded children with BTHS. No specific treatment is required for muscle hypotonia because it usually improves spontaneously.

## Prognosis and outcome

Infantile BTHS has a high mortality rate from heart failure and severe bacterial infections owing to neutropenia. However, these two manifestations in BTHS are often responsive to standard medical therapy, indicating early diagnosis of BTHS and management of heart failure, and bacterial infections are critical for a better prognosis [7, 21]. Although some patients may have limited clinical features, they could also develop significant heart failure or lifethreatening arrhythmias during or even after childhood [21]. Thus, physicians should be aware of the possibility of this disease and carry out genetic studies when it is considered.

**Conflict of interest** The authors declare that they have no conflict of interest to report.

#### References

- Ades LC, Gedeon AK, Wilson MJ et al (1993) Barth syndrome: clinical features and confirmation of gene localisation to distal Xq28. Am J Med Genet 45(3):327–334. doi:10.1002/ajmg.1320450309
- Adwani SS, Whitehead BF, Rees PG et al (1997) Heart transplantation for Barth syndrome. Pediatr Cardiol 18(2):143–145

- Barth PG, Scholte HR, Berden JA et al (1983) An X-linked mitochondrial disease affecting cardiac muscle, skeletal muscle and neutrophil leucocytes. J Neurol Sci 62(1–3):327–355
- Barth PG, Valianpour F, Bowen VM et al (2004) X-linked cardioskeletal myopathy and neutropenia (Barth syndrome): an update. Am J Med Genet A 126A(4):349–354. doi:10.1002/ajmg.a.20660
- Barth PG, Wanders RJ, Vreken P et al (1999) X-linked cardioskeletal myopathy and neutropenia (Barth syndrome) (MIM 302060). J Inherit Metab Dis 22(4):555–567
- Bione S, D'Adamo P, Maestrini E et al (1996) A novel X-linked gene, G4.5. is responsible for Barth syndrome. Nat Genet 12 (4):385–389. doi:10.1038/ng0496-385
- Bleyl SB, Mumford BR, Thompson V et al (1997) Neonatal, lethal noncompaction of the left ventricular myocardium is allelic with Barth syndrome. Am J Hum Genet 61(4):868–872. doi:10.1086/514879
- Cantlay AM, Shokrollahi K, Allen JT et al (1999) Genetic analysis of the G4.5 gene in families with suspected Barth syndrome. J Pediatr 135(3):311–315
- D'Adamo P, Fassone L, Gedeon A et al (1997) The X-linked gene G4.5 is responsible for different infantile dilated cardiomyopathies. Am J Hum Genet 61(4):862–867. doi:10.1086/514886
- Gedeon AK, Wilson MJ, Colley AC, Sillence DO, Mulley JC (1995) X linked fatal infantile cardiomyopathy maps to Xq28 and is possibly allelic to Barth syndrome. J Med Genet 32(5):383–388
- Hauff KD, Hatch GM (2006) Cardiolipin metabolism and Barth syndrome. Prog Lipid Res 45(2):91–101. doi:10.1016/j. plipres.2005.12.001
- 12. Houtkooper RH, Rodenburg RJ, Thiels C et al (2009) Cardiolipin and monolysocardiolipin analysis in fibroblasts, lymphocytes, and tissues using high-performance liquid chromatography—mass spectrometry as a diagnostic test for Barth syndrome. Anal Biochem 387(2):230–237. doi:10.1016/j.ab.2009.01.032
- Ichida F, Tsubata S, Bowles KR et al (2001) Novel gene mutations in patients with left ventricular noncompaction or Barth syndrome. Circulation 103(9):1256–1263
- Johnston J, Kelley RI, Feigenbaum A et al (1997) Mutation characterization and genotype-phenotype correlation in Barth syndrome. Am J Hum Genet 61(5):1053–1058. doi:10.1086/301604
- Kelley RI, Cheatham JP, Clark BJ et al (1991) X-linked dilated cardiomyopathy with neutropenia, growth retardation, and 3methylglutaconic aciduria. J Pediatr 119(5):738–747
- Kulik W, van Lenthe H, Stet FS et al (2008) Bloodspot assay using HPLC-tandem mass spectrometry for detection of Barth syndrome. Clin Chem 54(2):371–378. doi:10.1373/clinchem.2007.095711
- Liang WC, Nishino I (2011) Lipid storage myopathy. Curr Neurol Neurosci Rep 11(1):97–103. doi:10.1007/s11910-010-0154-y
- Neuwald AF (1997) Barth syndrome may be due to an acyltransferase deficiency. Curr Biol 7(8):R465–R466
- Schlame M, Kelley RI, Feigenbaum A et al (2003) Phospholipid abnormalities in children with Barth syndrome. J Am Coll Cardiol 42(11):1994–1999
- Schlame M, Towbin JA, Heerdt PM et al (2002) Deficiency of tetralinoleoyl-cardiolipin in Barth syndrome. Ann Neurol 51 (5):634–637. doi:10.1002/ana.10176
- 21. Spencer CT, Bryant RM, Day J et al (2006) Cardiac and clinical phenotype in Barth syndrome. Pediatrics 118(2):e337–e346. doi:10.1542/peds.2005-2667
- Spencer CT, Byrne BJ, Gewitz MH et al (2005) Ventricular arrhythmia in the X-linked cardiomyopathy Barth syndrome. Pediatr Cardiol 26(5):632–637. doi:10.1007/s00246-005-0873-z
- Stein SM, Dale DC (2003) Molecular basis and therapy of disorders associated with chronic neutropenia. Curr Allergy Asthma Rep 3(5):385–388
- Steward CG, Newbury-Ecob RA, Hastings R et al (2010) Barth syndrome: an X-linked cause of fetal cardiomyopathy and stillbirth. Prenat Diagn 30(10):970–976. doi:10.1002/pd.2599

